

Modern Quantum Chemistry, Szabo & Ostlund

HW

WSF

July 2, 2020

Contents

5	Pair and Coupled-pair Theories	2
5.1	The Independent Electron Pair Approximation	2
	Ex 5.1	2
	Ex 5.2	2
	Ex 5.3	3
5.1.1	Invariance under Unitary Transformations: An Example	3
	Ex 5.4	3
	Ex 5.5	3
	Ex 5.6	4
	Ex 5.7	4
	Ex 5.8	6
	Ex 5.9	6
	Ex 5.10	6
5.1.2	Some Illustrative Calculations	6
5.2	Coupled-pair Theories	6
5.2.1	The Coupled-cluster Approximation	6
5.2.2	The Cluster Expansion of the Wave Function	6
	Ex 5.11	6
5.2.3	Linear CCA and the Coupled-Electron Pair Approximation	7
	Ex 5.12	7
5.2.4	Some Illustrative Calculations	7
5.3	Many-electron Theories with Single Particle Hamiltonians	7
5.3.1	The Relaxation Energy via CI, IEPA, CEPA and CCA	7
	Ex 5.14	7
	Ex 5.15	7
5.3.2	The Resonance Energy of Polyenes in Hückel Theory	7
	Ex 5.16	7
	Ex 5.17	7
	Ex 5.18	7
	Ex 5.19	7
	Ex 5.20	7
	Ex 5.21	7

5 Pair and Coupled-pair Theories

5.1 The Independent Electron Pair Approximation

Ex 5.1

a.

$$\begin{aligned}
 {}^1E_{\text{corr}}(\text{FO}) &= \frac{|\langle 1\bar{1} | 2\bar{2} \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} \\
 &= \frac{|\langle 1\bar{1} | 2\bar{2} \rangle - \langle 1\bar{1} | \bar{2}2 \rangle|^2}{2\varepsilon_1 - 2\varepsilon_2} \\
 &= \frac{|[12|\bar{1}\bar{2}] - [\bar{1}2|1\bar{2}]|^2}{2\varepsilon_1 - 2\varepsilon_2} \\
 &= \frac{K_{12}^2}{2(\varepsilon_1 - \varepsilon_2)}
 \end{aligned} \tag{5.1.1}$$

b.

$$\begin{aligned}
 {}^1E_{\text{corr}} &= \Delta - \Delta \sqrt{1 + \frac{K_{12}^2}{\Delta^2}} \\
 &= \Delta - \Delta \left(1 + \frac{K_{12}^2}{2\Delta^2} \right) \\
 &= -\frac{K_{12}^2}{2\Delta} \\
 &\approx \frac{K_{12}^2}{2(\varepsilon_1 - \varepsilon_2)}
 \end{aligned} \tag{5.1.2}$$

Ex 5.2 From Eq. 5.9a and 5.9b in the textbook, we get

$$\sum_{t < u} c_{1_i \bar{1}_i}^{tu} \langle \Psi_0 | \mathcal{H} | \Psi_{1_i \bar{1}_i}^{tu} \rangle = e_{1_i \bar{1}_i} \tag{5.1.3}$$

$$\langle \Psi_{1_i \bar{1}_i}^{rs} | \mathcal{H} | \Psi_0 \rangle + \sum_{t < u} \langle \Psi_{1_i \bar{1}_i}^{rs} | \mathcal{H} - E_0 | \Psi_{1_i \bar{1}_i}^{tu} \rangle c_{1_i \bar{1}_i}^{tu} = e_{1_i \bar{1}_i} c_{1_i \bar{1}_i}^{rs} \tag{5.1.4}$$

\therefore

$$c_{1_i \bar{1}_i}^{2_i \bar{2}_i} \langle \Psi_0 | \mathcal{H} | \Psi_{1_i \bar{1}_i}^{2_i \bar{2}_i} \rangle = e_{1_i \bar{1}_i} \tag{5.1.5}$$

$$\langle \Psi_{1_i \bar{1}_i}^{2_i \bar{2}_i} | \mathcal{H} | \Psi_0 \rangle + \sum_{t < u} \langle \Psi_{1_i \bar{1}_i}^{2_i \bar{2}_i} | \mathcal{H} - E_0 | \Psi_{1_i \bar{1}_i}^{tu} \rangle c_{1_i \bar{1}_i}^{tu} = e_{1_i \bar{1}_i} c_{1_i \bar{1}_i}^{2_i \bar{2}_i} \tag{5.1.6}$$

(5.1.5) gives

$$K_{12} c_{1_i \bar{1}_i}^{2_i \bar{2}_i} = e_{1_i \bar{1}_i} \tag{5.1.7}$$

(5.1.6) gives

$$K_{12} + \sum_{j \neq k} \langle \Psi_{1_i \bar{1}_i}^{2_i \bar{2}_i} | \mathcal{H} - E_0 | \Psi_{1_i \bar{1}_i}^{2_j \bar{2}_k} \rangle c_{1_i \bar{1}_i}^{2_j \bar{2}_k} = e_{1_i \bar{1}_i} c_{1_i \bar{1}_i}^{2_i \bar{2}_i} \tag{5.1.8}$$

Since

$$\langle \Psi_{1_i \bar{1}_i}^{2_i \bar{2}_i} | \mathcal{H} - E_0 | \Psi_{1_i \bar{1}_i}^{2_j \bar{2}_k} \rangle c_{1_i \bar{1}_i}^{2_j \bar{2}_k} = \begin{cases} 2\Delta & j = k = i \\ 0 & j = k \neq i \\ 0 & i = j \neq k \end{cases} \tag{5.1.9}$$

we have

$$K_{12} + 2\Delta c_{1_i \bar{1}_i}^{2_i \bar{2}_i} = e_{1_i \bar{1}_i} c_{1_i \bar{1}_i}^{2_i \bar{2}_i} \tag{5.1.10}$$

Ex 5.3

$$\begin{aligned}
{}^2E_{\text{corr}}(\text{FO}) &= \sum_i \frac{|\langle 1_i \bar{1}_i \| 2_i \bar{2}_i \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} \\
&= 2 \times \frac{K_{12}^2}{2(\varepsilon_1 - \varepsilon_2)} \\
&= \frac{K_{12}^2}{(\varepsilon_1 - \varepsilon_2)}
\end{aligned} \tag{5.1.11}$$

5.1.1 Invariance under Unitary Transformations: An Example

Ex 5.4

$$\begin{aligned}
|a\bar{a}b\bar{b}\rangle &= 2^{-1/2}(|1_1\bar{a}b\bar{b}\rangle + |1_2\bar{a}b\bar{b}\rangle) \\
&= 2^{-1}(|1_1\bar{1}_1b\bar{b}\rangle + |1_1\bar{1}_2b\bar{b}\rangle + |1_2\bar{1}_1b\bar{b}\rangle + |1_2\bar{1}_2b\bar{b}\rangle) \\
&= 2^{-2}(|1_1\bar{1}_11_1\bar{1}_1\rangle - |1_1\bar{1}_11_1\bar{1}_2\rangle - |1_1\bar{1}_11_2\bar{1}_1\rangle + |1_1\bar{1}_11_2\bar{1}_2\rangle \\
&\quad + |1_1\bar{1}_21_1\bar{1}_1\rangle - |1_1\bar{1}_21_1\bar{1}_2\rangle - |1_1\bar{1}_21_2\bar{1}_1\rangle + |1_1\bar{1}_21_2\bar{1}_2\rangle \\
&\quad + |1_2\bar{1}_11_1\bar{1}_1\rangle - |1_2\bar{1}_11_1\bar{1}_2\rangle - |1_2\bar{1}_11_2\bar{1}_1\rangle + |1_2\bar{1}_11_2\bar{1}_2\rangle \\
&\quad + |1_2\bar{1}_21_1\bar{1}_1\rangle - |1_2\bar{1}_21_1\bar{1}_2\rangle - |1_2\bar{1}_21_2\bar{1}_1\rangle + |1_2\bar{1}_21_2\bar{1}_2\rangle) \\
&= 2^{-2}(2|1_1\bar{1}_11_1\bar{1}_1\rangle + 2|1_1\bar{1}_11_2\bar{1}_2\rangle - 2|1_1\bar{1}_21_1\bar{1}_2\rangle - 2|1_1\bar{1}_21_2\bar{1}_1\rangle) \\
&= 2^{-2}(2|1_1\bar{1}_11_2\bar{1}_2\rangle - 2|1_1\bar{1}_11_2\bar{1}_2\rangle) \\
&= |1_1\bar{1}_11_2\bar{1}_2\rangle
\end{aligned} \tag{5.1.12}$$

Ex 5.5

$$\begin{aligned}
\langle \Psi_0 | \mathcal{H} | \Psi_{a\bar{a}}^{**} \rangle &= 2^{-1/2}(\langle \Psi_0 | \mathcal{H} | \Psi_{a\bar{a}}^{r\bar{r}} \rangle + \langle \Psi_0 | \mathcal{H} | \Psi_{a\bar{a}}^{s\bar{s}} \rangle) \\
&= 2^{-1/2}\left(2 \times \frac{1}{2}K_{12}\right) \\
&= 2^{-1/2}K_{12}
\end{aligned} \tag{5.1.13}$$

$$\begin{aligned}
\langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{**} \rangle &= 2^{-1}(\langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{r\bar{r}} \rangle + \langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{s\bar{s}} \rangle \\
&\quad + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{r\bar{r}} \rangle + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{s\bar{s}} \rangle) \\
&= 2^{-1}\left[\left(2h_{11} + 2h_{22} + \frac{1}{2}J_{11} + \frac{1}{2}J_{22} + 2J_{12} - K_{12}\right) - (4h_{11} + 2J_{11})\right. \\
&\quad \left.+ \frac{1}{2}J_{22} + \frac{1}{2}J_{22}\right. \\
&\quad \left.+ \left(2h_{11} + 2h_{22} + \frac{1}{2}J_{11} + \frac{1}{2}J_{22} + 2J_{12} - K_{12}\right) - (4h_{11} + 2J_{11})\right] \\
&= 2^{-1}\left(-2h_{11} + 2h_{22} - \frac{3}{2}J_{11} + J_{22} + 2J_{12} - K_{12}\right) \times 2 \\
&= -2h_{11} + 2h_{22} - \frac{3}{2}J_{11} + J_{22} + 2J_{12} - K_{12}
\end{aligned} \tag{5.1.14}$$

Since

$$\varepsilon_2 - \varepsilon_1 = h_{22} - h_{11} + 2J_{12} - K_{12} - J_{11} \tag{5.1.15}$$

we have

$$\langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{**} \rangle = 2(\varepsilon_2 - \varepsilon_1) - 2J_{12} + K_{12} + \frac{1}{2}J_{11} + J_{22} \tag{5.1.16}$$

Ex 5.6 Since

$$|\Psi_{ab}^{**}\rangle = 2^{-1/2}(|\Psi_{ab}^{r\bar{s}}\rangle + |\Psi_{ab}^{s\bar{r}}\rangle) \quad (5.1.17)$$

$$\begin{aligned} \langle \Psi_0 | \mathcal{H} | \Psi_{ab}^{**} \rangle &= 2^{-1/2} (\langle \Psi_0 | \mathcal{H} | \Psi_{ab}^{r\bar{s}} \rangle + \langle \Psi_0 | \mathcal{H} | \Psi_{ab}^{s\bar{r}} \rangle) \\ &= 2^{-1/2} (\langle a\bar{b} || r\bar{s} \rangle + \langle a\bar{b} || s\bar{r} \rangle) \\ &= 2^{-1/2} ((ar|bs) + (as|br)) \\ &= 2^{-1/2} K_{12} \end{aligned} \quad (5.1.18)$$

$$\begin{aligned} \langle \Psi_{ab}^{**} | \mathcal{H} - E_0 | \Psi_{ab}^{**} \rangle &= 2^{-1} (\langle \Psi_{ab}^{r\bar{s}} | \mathcal{H} - E_0 | \Psi_{ab}^{r\bar{s}} \rangle + \langle \Psi_{ab}^{r\bar{s}} | \mathcal{H} - E_0 | \Psi_{ab}^{s\bar{r}} \rangle \\ &\quad + \langle \Psi_{ab}^{s\bar{r}} | \mathcal{H} - E_0 | \Psi_{ab}^{r\bar{s}} \rangle + \langle \Psi_{ab}^{s\bar{r}} | \mathcal{H} - E_0 | \Psi_{ab}^{s\bar{r}} \rangle) \\ &= 2^{-1} \left[\left(2h_{11} + 2h_{22} + \frac{1}{2}J_{11} + \frac{1}{2}J_{22} + 2J_{12} - K_{12} \right) - (4h_{11} + 2J_{11}) \right. \\ &\quad \left. + \frac{1}{2}J_{22} + \frac{1}{2}J_{22} \right. \\ &\quad \left. + \left(2h_{11} + 2h_{22} + \frac{1}{2}J_{11} + \frac{1}{2}J_{22} + 2J_{12} - K_{12} \right) - (4h_{11} + 2J_{11}) \right] \\ &= \dots \\ &= 2(\varepsilon_2 - \varepsilon_1) - 2J_{12} + K_{12} + \frac{1}{2}J_{11} + J_{22} \equiv 2\Delta' \end{aligned} \quad (5.1.19)$$

Thus the equations determining $e_{a\bar{b}}$ are identical to that of $e_{a\bar{a}}$. Similarly, $e_{\bar{a}b}$ shares the same equations with them.

$\therefore e_{a\bar{b}} = e_{\bar{a}b} = e_{a\bar{a}}$.

Ex 5.7

a. As shown in Ex 5.5, 5.6

$$\langle \Psi_0 | \mathcal{H} | \Psi_{a\bar{a}}^{**} \rangle = \langle \Psi_0 | \mathcal{H} | \Psi_{a\bar{b}}^{**} \rangle = \langle \Psi_0 | \mathcal{H} | \Psi_{\bar{a}b}^{**} \rangle = 2^{-1/2} K_{12} \quad (5.1.20)$$

$$\langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{a}}^{**} \rangle = \langle \Psi_{a\bar{b}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle = \langle \Psi_{\bar{a}b}^{**} | \mathcal{H} - E_0 | \Psi_{\bar{a}b}^{**} \rangle = 2\Delta' \quad (5.1.21)$$

Similarly, we get

$$\langle \Psi_0 | \mathcal{H} | \Psi_{b\bar{b}}^{**} \rangle = 2^{-1/2} K_{12} \quad (5.1.22)$$

$$\langle \Psi_{b\bar{b}}^{**} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{**} \rangle = 2\Delta' \quad (5.1.23)$$

For the rest,

$$\begin{aligned} \langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{**} \rangle &= 2^{-1} (\langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{r\bar{r}} \rangle + \langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{s\bar{s}} \rangle \\ &\quad + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{r\bar{r}} \rangle + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{b\bar{b}}^{s\bar{s}} \rangle) \\ &= 2^{-1} [\langle b\bar{b} || a\bar{a} \rangle + 0 + 0 + \langle b\bar{b} || a\bar{a} \rangle] \\ &= (ab|ab) \\ &= \frac{1}{2} J_{11} \end{aligned} \quad (5.1.24)$$

$$\begin{aligned}
\langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle &= 2^{-1} (\langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{r\bar{s}} \rangle + \langle \Psi_{a\bar{a}}^{r\bar{r}} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{s\bar{r}} \rangle \\
&\quad + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{r\bar{s}} \rangle + \langle \Psi_{a\bar{a}}^{s\bar{s}} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{s\bar{r}} \rangle) \\
&= 2^{-1} [\langle \bar{r}\bar{b} | \bar{a}\bar{s} \rangle - \langle \bar{r}\bar{b} | \bar{s}\bar{a} \rangle + \langle \bar{s}\bar{b} | \bar{r}\bar{a} \rangle - \langle \bar{s}\bar{b} | \bar{a}\bar{r} \rangle] \\
&= 2^{-1} [(ra|bs) - (rs|ba) - (rs|ba) - (sr|ba) + (sa|br) - (sr|ba)] \\
&= 2^{-1} [(ra|bs) + (sa|br) - 4(ab|sr)] \\
&= 2^{-1} \left[2 \times \frac{1}{2} K_{12} - 4 \times \frac{1}{2} J_{12} \right] \\
&= \frac{1}{2} K_{12} - J_{12}
\end{aligned} \tag{5.1.25}$$

Similarly, we get

$$\langle \Psi_{a\bar{b}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle = \frac{1}{2} J_{11} \tag{5.1.26}$$

$$\langle \Psi_{a\bar{a}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle = \langle \Psi_{b\bar{b}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle = \langle \Psi_{b\bar{b}}^{**} | \mathcal{H} - E_0 | \Psi_{a\bar{b}}^{**} \rangle = \frac{1}{2} K_{12} - J_{12} \tag{5.1.27}$$

thus the DCI equation is

$$\begin{pmatrix} 0 & 2^{-1/2} K_{12} & 2^{-1/2} K_{12} & 2^{-1/2} K_{12} & 2^{-1/2} K_{12} \\ 2^{-1/2} K_{12} & 2\Delta' & \frac{1}{2} J_{11} & \frac{1}{2} K_{12} - J_{12} & \frac{1}{2} K_{12} - J_{12} \\ 2^{-1/2} K_{12} & \frac{1}{2} J_{11} & 2\Delta' & \frac{1}{2} K_{12} - J_{12} & \frac{1}{2} K_{12} - J_{12} \\ 2^{-1/2} K_{12} & \frac{1}{2} K_{12} - J_{12} & \frac{1}{2} K_{12} - J_{12} & 2\Delta' & \frac{1}{2} J_{11} \\ 2^{-1/2} K_{12} & \frac{1}{2} K_{12} - J_{12} & \frac{1}{2} K_{12} - J_{12} & \frac{1}{2} J_{11} & 2\Delta' \end{pmatrix} \begin{pmatrix} 1 \\ c_1 \\ c_2 \\ c_3 \\ c_4 \end{pmatrix} = {}^2E_{\text{corr}}(\text{DCI}) \begin{pmatrix} 1 \\ c_1 \\ c_2 \\ c_3 \\ c_4 \end{pmatrix} \tag{5.1.28}$$

b. By solving the DCI equation above (see **5-7.nb**), we get

$${}^2E_{\text{corr}}(\text{DCI}) = \frac{2\Delta' + \frac{1}{2} J_{11} + 2(\frac{1}{2} K_{12} - J_{12}) - \sqrt{16(2^{-1/2} K_{12})^2 + [2\Delta' + \frac{1}{2} J_{11} + 2(\frac{1}{2} K_{12} - J_{12})]^2}}{2} \tag{5.1.29}$$

and

$$c_1 = c_2 = c_3 = c_4 = \frac{2\Delta' + \frac{1}{2} J_{11} + 2(\frac{1}{2} K_{12} - J_{12}) + \sqrt{16(2^{-1/2} K_{12})^2 + [2\Delta' + \frac{1}{2} J_{11} + 2(\frac{1}{2} K_{12} - J_{12})]^2}}{8 \times 2^{-1/2} K_{12}} \tag{5.1.30}$$

Since

$$2\Delta' = 2(\varepsilon_2 - \varepsilon_1) - 2J_{12} + K_{12} + \frac{1}{2} J_{11} + J_{22} \tag{5.1.31}$$

$$2\Delta = 2(\varepsilon_2 - \varepsilon_1) + J_{11} + J_{22} - 4J_{12} + 2K_{12} \tag{5.1.32}$$

we have

$$2\Delta = 2\Delta' + \frac{1}{2} J_{11} - 2J_{12} + K_{12} \tag{5.1.33}$$

\therefore

$$\begin{aligned}
{}^2E_{\text{corr}}(\text{DCI}) &= \frac{2\Delta - \sqrt{8K_{12}^2 + (2\Delta)^2}}{2} \\
&= \Delta - \sqrt{2K_{12}^2 + \Delta^2}
\end{aligned} \tag{5.1.34}$$

$$\begin{aligned}
c_1 = c_2 = c_3 = c_4 &= \frac{2\Delta + \sqrt{8K_{12}^2 + (2\Delta)^2}}{4\sqrt{2}K_{12}} \\
&= \frac{\Delta + \sqrt{2K_{12}^2 + \Delta^2}}{2\sqrt{2}K_{12}}
\end{aligned} \tag{5.1.35}$$

Ex 5.8

$$\begin{aligned}
E_{\text{corr}}(\text{FO}) &= \sum_{A < B} \sum_{R < S} \frac{|\langle AB \| RS \rangle|^2}{\varepsilon_A + \varepsilon_B - \varepsilon_R - \varepsilon_S} \\
&= \frac{|\langle a\bar{a} \| r\bar{r} \rangle|^2 + |\langle a\bar{a} \| r\bar{s} \rangle|^2 + |\langle a\bar{a} \| s\bar{r} \rangle|^2 + |\langle a\bar{a} \| s\bar{s} \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} + \frac{|\langle a\bar{b} \| r\bar{r} \rangle|^2 + |\langle a\bar{b} \| r\bar{s} \rangle|^2 + |\langle a\bar{b} \| s\bar{r} \rangle|^2 + |\langle a\bar{b} \| s\bar{s} \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} \\
&\quad + \frac{|\langle b\bar{a} \| r\bar{r} \rangle|^2 + |\langle b\bar{a} \| r\bar{s} \rangle|^2 + |\langle b\bar{a} \| s\bar{r} \rangle|^2 + |\langle b\bar{a} \| s\bar{s} \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} + \frac{|\langle b\bar{b} \| r\bar{r} \rangle|^2 + |\langle b\bar{b} \| r\bar{s} \rangle|^2 + |\langle b\bar{b} \| s\bar{r} \rangle|^2 + |\langle b\bar{b} \| s\bar{s} \rangle|^2}{\varepsilon_1 + \varepsilon_1 - \varepsilon_2 - \varepsilon_2} \\
&= \frac{|(ar|ar)|^2 + |(ar|as)|^2 + |(as|ar)|^2 + |(as|as)|^2}{2(\varepsilon_1 - \varepsilon_2)} + \frac{|(ar|br)|^2 + |(ar|bs)|^2 + |(as|br)|^2 + |(as|bs)|^2}{2(\varepsilon_1 - \varepsilon_2)} \\
&\quad + \frac{|(br|ar)|^2 + |(br|as)|^2 + |(bs|ar)|^2 + |(bs|as)|^2}{2(\varepsilon_1 - \varepsilon_2)} + \frac{|(br|br)|^2 + |(br|bs)|^2 + |(bs|br)|^2 + |(bs|bs)|^2}{2(\varepsilon_1 - \varepsilon_2)} \\
&= \frac{|\frac{1}{2}K_{12}|^2 + 0 + 0 + |\frac{1}{2}K_{12}|^2}{2(\varepsilon_1 - \varepsilon_2)} + \frac{0 + |\frac{1}{2}K_{12}|^2 + |\frac{1}{2}K_{12}|^2 + 0}{2(\varepsilon_1 - \varepsilon_2)} \\
&\quad + \frac{0 + 0 + |\frac{1}{2}K_{12}|^2 + |\frac{1}{2}K_{12}|^2}{2(\varepsilon_1 - \varepsilon_2)} + \frac{|\frac{1}{2}K_{12}|^2 + 0 + 0 + |\frac{1}{2}K_{12}|^2}{2(\varepsilon_1 - \varepsilon_2)} \\
&= \frac{2K_{12}^2}{2(\varepsilon_1 - \varepsilon_2)} \tag{5.1.36}
\end{aligned}$$

Ex 5.9

a.

$$\begin{aligned}
{}^2E_{\text{corr}}(\text{EN(L)}) &= - \sum_{a < b} \sum_{r < s} \frac{|\langle \Psi_0 | \mathcal{H} | \Psi_{ab}^{rs} \rangle|^2}{\langle \Psi_{ab}^{rs} | \mathcal{H} - E_0 | \Psi_{ab}^{rs} \rangle} \\
&= - \tag{5.1.37}
\end{aligned}$$

Ex 5.10

5.1.2 Some Illustrative Calculations

5.2 Coupled-pair Theories

5.2.1 The Coupled-cluster Approximation

5.2.2 The Cluster Expansion of the Wave Function

Ex 5.11 Eq. 5.49 gives

$$\begin{aligned}
|\Phi_0\rangle &= |1_1 \bar{1}_1 1_2 \bar{1}_2\rangle + c_{1_1 \bar{1}_1}^{2_1 \bar{2}_1} |2_1 \bar{2}_1 1_2 \bar{1}_2\rangle + c_{1_2 \bar{1}_2}^{2_2 \bar{2}_2} |1_1 \bar{1}_1 2_2 \bar{2}_2\rangle + c_{1_1 \bar{1}_1 1_2 \bar{1}_2}^{2_1 \bar{2}_1 2_2 \bar{2}_2} |2_1 \bar{2}_1 2_2 \bar{2}_2\rangle \\
&= \left[1 + c_{1_1 \bar{1}_1}^{2_1 \bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2 \bar{1}_2}^{2_2 \bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2} + c_{1_1 \bar{1}_1 1_2 \bar{1}_2}^{2_1 \bar{2}_1 2_2 \bar{2}_2} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2} a_{\bar{1}_1} a_{1_1} \right] |1_1 \bar{1}_1 1_2 \bar{1}_2\rangle \tag{5.2.1}
\end{aligned}$$

while

$$\begin{aligned}
&\exp\left(c_{1_1 \bar{1}_1}^{2_1 \bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2 \bar{1}_2}^{2_2 \bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right) |1_1 \bar{1}_1 1_2 \bar{1}_2\rangle \\
&= \left[1 + \left(c_{1_1 \bar{1}_1}^{2_1 \bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2 \bar{1}_2}^{2_2 \bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right) + \left(c_{1_1 \bar{1}_1}^{2_1 \bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2 \bar{1}_2}^{2_2 \bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right)^2 + \dots \right] |1_1 \bar{1}_1 1_2 \bar{1}_2\rangle \tag{5.2.2}
\end{aligned}$$

since we cannot annihilate or create any orbital twice, the terms over 3rd power must be zero, thus

$$\begin{aligned}
& \exp\left(c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right) |1_1\bar{1}_1 1_2\bar{1}_2\rangle \\
&= \left[1 + \left(c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right) + \left(c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right)^2\right] |1_1\bar{1}_1 1_2\bar{1}_2\rangle \\
&= \left[1 + \left(c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right) + \left(c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1}\right)^2 + \left(c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2}\right)^2\right. \\
&\quad \left.+ c_{1_1\bar{1}_1}^{2_1\bar{2}_1} c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_1} a_{1_1} a_{\bar{1}_2} a_{1_2}\right] |1_1\bar{1}_1 1_2\bar{1}_2\rangle \\
&= \left[1 + c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2} + c_{1_1\bar{1}_1}^{2_1\bar{2}_1} c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_1} a_{1_1} a_{\bar{1}_2} a_{1_2}\right] |1_1\bar{1}_1 1_2\bar{1}_2\rangle \\
&= \left[1 + c_{1_1\bar{1}_1}^{2_1\bar{2}_1} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{\bar{1}_1} a_{1_1} + c_{1_2\bar{1}_2}^{2_2\bar{2}_2} a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_2} a_{1_2} + c_{1_1\bar{1}_1 1_2\bar{1}_2}^{2_1\bar{2}_1 2_2\bar{2}_2} a_{2_1}^\dagger a_{\bar{2}_1}^\dagger a_{2_2}^\dagger a_{\bar{2}_2}^\dagger a_{\bar{1}_1} a_{1_1} a_{\bar{1}_2} a_{1_2}\right] |1_1\bar{1}_1 1_2\bar{1}_2\rangle \quad (5.2.3)
\end{aligned}$$

5.2.3 Linear CCA and the Coupled-Electron Pair Approximation

Ex 5.12

5.2.4 Some Illustrative Calculations

5.3 Many-electron Theories with Single Particle Hamiltonians

5.3.1 The Relaxation Energy via CI, IEPA, CEPA and CCA

Ex 5.14

Ex 5.15

5.3.2 The Resonance Energy of Polyenes in Hückel Theory

Ex 5.16

Ex 5.17

Ex 5.18

Ex 5.19

Ex 5.20

Ex 5.21