

Modern Quantum Chemistry, Szabo & Ostlund

HW

WSF

May 15, 2020

Contents

| | | |
|----------|---|----------|
| A | Integral Evaluation with 1s Primitive Gaussians | 2 |
| B | 2-Electron Self-consistent-field Program | 2 |
| C | Analytic Derivative Methods and Geometry Optimization | 2 |
| C.1 | Introduction | 2 |
| C.2 | General Considerations | 2 |
| C.3 | Analytic Derivatives | 2 |
| C.4 | Optimization Techniques | 2 |
| C.5 | Some Optimization Algorithms | 2 |
| Ex C.1 | | 2 |
| Ex C.2 | | 2 |
| Ex C.3 | | 2 |
| Ex C.4 | | 2 |
| C.6 | Transition States | 3 |
| C.7 | Constrained Variation | 3 |
| D | Molecular Integrals for H₂ as a Function of Bond Length | 3 |

A Integral Evaluation with 1s Primitive Gaussians

B 2-Electron Self-consistent-field Program

C Analytic Derivative Methods and Geometry Optimization

C.1 Introduction

C.2 General Considerations

C.3 Analytic Derivatives

C.4 Optimization Techniques

C.5 Some Optimization Algorithms

Ex C.1

(a)

$$\begin{aligned}\mathbf{H} &= \begin{pmatrix} \frac{\partial^2 E}{\partial x^2} & \frac{\partial^2 E}{\partial x \partial y} \\ \frac{\partial^2 E}{\partial y \partial x} & \frac{\partial^2 E}{\partial y^2} \end{pmatrix} \\ &= \begin{pmatrix} K & K'' \\ K'' & K' \end{pmatrix}\end{aligned}\tag{C.5.1}$$

$$\begin{aligned}\mathbf{f}(\mathbf{X}) &= \begin{pmatrix} \frac{\partial E}{\partial x} \\ \frac{\partial E}{\partial y} \end{pmatrix} \\ &= \begin{pmatrix} K(x-a) + K''y \\ K'(y-b) + K''x \end{pmatrix} \\ &= \begin{pmatrix} K & K'' \\ K'' & K' \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} Ka \\ K'b \end{pmatrix} \\ &= \begin{pmatrix} K & K'' \\ K'' & K' \end{pmatrix} \mathbf{X} - \begin{pmatrix} Ka \\ K'b \end{pmatrix}\end{aligned}\tag{C.5.2}$$

$$\begin{aligned}\mathbf{q} &= -\mathbf{H}^{-1}\mathbf{f} \\ &= -\mathbf{X} + \begin{pmatrix} K & K'' \\ K'' & K' \end{pmatrix}^{-1} \begin{pmatrix} Ka \\ K'b \end{pmatrix} \\ &= -\mathbf{X} + \frac{1}{KK' - K''^2} \begin{pmatrix} K' & -K'' \\ -K'' & K \end{pmatrix} \begin{pmatrix} Ka \\ K'b \end{pmatrix} \\ &= -\mathbf{X} + \frac{1}{KK' - K''^2} \begin{pmatrix} KK'a - K'K''b \\ -KK''a + KK'b \end{pmatrix} \\ &= -\mathbf{X} + \frac{1}{KK' - K''^2} \begin{pmatrix} KK' & K'K'' \\ -KK'' & KK' \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}\end{aligned}\tag{C.5.3}$$

Ex C.2

Ex C.3

Ex C.4

C.6 Transition States

C.7 Constrained Variation

D Molecular Integrals for H_2 as a Function of Bond Length