$\begin{array}{c} \mathbf{Modern~Quantum~Chemistry,~Szabo~\&~Ostlund} \\ \mathbf{HW} \end{array}$

王石嵘

$March\ 1,\ 2020$

Contents

5	Pair	r and Coupled-pair Theories	2
	5.1	The Independent Electron Pair Approximation	2
		Ex 5.1	2

5 Pair and Coupled-pair Theories

5.1 The Independent Electron Pair Approximation

Ex 5.1

a.

$${}^{1}E_{\text{corr}}(\text{FO}) = \frac{|\langle 1\bar{1} \parallel 2\bar{2} \rangle|^{2}}{\varepsilon_{1} + \varepsilon_{1} - \varepsilon_{2} - \varepsilon_{2}}$$

$$= \frac{|\langle 1\bar{1} \mid 2\bar{2} \rangle - \langle 1\bar{1} \mid \bar{2}2 \rangle|^{2}}{2\varepsilon_{1} - 2\varepsilon_{2}}$$

$$= \frac{|[12|\bar{1}\bar{2}] - [1\bar{2}|\bar{1}2]|^{2}}{2\varepsilon_{1} - 2\varepsilon_{2}}$$

$$= \frac{K_{12}^{2}}{2(\varepsilon_{1} - \varepsilon_{2})}$$
(5.1.1)

b.

$${}^{1}E_{\text{corr}} = \Delta - \Delta \sqrt{1 + \frac{K_{12}^{2}}{\Delta^{2}}}$$

$$= \Delta - \Delta \left(1 + \frac{K_{12}^{2}}{2\Delta^{2}}\right)$$

$$= -\frac{K_{12}^{2}}{2\Delta}$$

$$\approx \frac{K_{12}^{2}}{2(\varepsilon_{1} - \varepsilon_{2})}$$
(5.1.2)