

**Problem 1.8.25**

Let  $\mathcal{B}$  be a Banach space in which the parallelogram law holds. We can define an inner product on  $\mathcal{B}$  by

$$4(f, g) = \|f + g\|^2 - \|f - g\|^2$$

Note that  $(\cdot, \cdot)$  induces the norm because

$$\begin{aligned} (f, f) &= \frac{1}{4} (\|f + f\|^2 - \|f - f\|^2) \\ &= \frac{1}{4} (4\|f\|^2) \end{aligned}$$

So we have that  $\|f\| = \sqrt{(f, f)}$ . We need to show that  $(\cdot, \cdot)$  is indeed an inner product because

$$\begin{aligned} 4(f, g) &= \|f + g\|^2 - \|f - g\|^2 \\ &= (\|f + g\| - \|f - g\|)(\|f + g\| + \|f - g\|) \\ &= (\|f + g\| + \|g - f\|)(\|f + g\| + \|f - g\|) \end{aligned}$$

Because the norm is non-negative, we have  $(\cdot, \cdot)$  is the product of positive terms and hence, positive. Symmetry is also clear because

$$\begin{aligned} 4(f, g) &= \|f + g\|^2 - \|f - g\|^2 \\ &= \|g + f\|^2 - (-1)^2 \|g - f\|^2 \\ &= 4(g, f) \end{aligned}$$

We are only left to verify linearity. In the real case we compute

$$(f, g + h) = \|f + g + h\|^2 - \|f - (g + h)\|^2$$

We can decompose the right side into the following quantities

$$\begin{aligned} A &= \frac{1}{2} \|f + g + h\|^2 + \frac{1}{2} \|f + g - h\|^2 - \|h\|^2 \\ B &= \frac{1}{2} \|f - g + h\|^2 + \frac{1}{2} \|f - g - h\|^2 - \|h\|^2 \\ C &= \frac{1}{2} \|f + g + h\|^2 + \frac{1}{2} \|f - g - h\|^2 - \|g\|^2 \\ D &= \frac{1}{2} \|f + g - h\|^2 + \frac{1}{2} \|f - g - h\|^2 - \|g\|^2 \end{aligned}$$

Then we have

$$\begin{aligned} \|f + g + h\|^2 - \|f - (g + h)\|^2 &= A - B + C - D \\ &= (\|f + g\|^2 - \|f - g\|^2) + (\|f + h\|^2 - \|f - h\|^2) \\ &= 4(f, g) + 4(f, h) \end{aligned}$$

So the operation is linear.

To get scalar multiplication we note that linearity immediately implies that

$$(nf, g) = (f + \cdots + f, g) = (f, g) + \cdots + (f, g) = n(f, g)$$

whenever  $n \in \mathbb{Z}$ . To get rational scalars we note that

$$(f, g) = (n(\frac{1}{n}f), g) = n(\frac{1}{n}f, g)$$

So for any  $q \in \mathbb{Q}$  we must have  $q(f, g) = (qf, g)$ . To extend this to real scalars we define a function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  given by

$$t \mapsto t(f, g) - (tf, g)$$

This function is clearly continuous because it is the difference of continuous functions. Moreover, it is 0 on  $\mathbb{Q}$ , which is dense in  $\mathbb{R}$  and therefore it must be zero on the whole line. Consequently, we must have that  $(tf, g) = t(f, g)$  for every  $t \in \mathbb{R}$ .

For complex Banach spaces we define the inner product by the formula

$$4(f, g) = (\|f + g\|^2 - \|f - g\|^2) + i(\|f + ig\|^2 - \|f - ig\|^2)$$

The computation to show that this indeed an inner product in analogous to the one above, but considers the real and imaginary parts of the vectors separately.

To see that  $L^p(\mathbb{R})$  is only a Hilbert space for  $p = 2$  consider the functions  $f(x) = \chi_{[0,1]}(x)$  and  $g(x) = \chi_{[1/2,1]}(x)$ . In order for the parallelogram law to hold in  $L^p(\mathbb{R})$  we must have that  $2(2)^{2/p} = 1(1)^{2/p}$ , which holds only when  $p = 2$ .

### Problem 1.8.30

It is clear that  $\mathcal{B}/\mathcal{S}$  is a vector space. Let  $[f]$  denote the equivalence class of  $f$  under the relation  $\sim$ . We define addition pointwise

$$[f] + [g] = [f + g]$$

and scalar multiplication via  $\alpha[f] = [\alpha f]$ . We then put the following norm on  $\mathcal{B}/\mathcal{S}$ ,

$$\|[f]\|_{\mathcal{B}/\mathcal{S}} = \inf_{f' \sim f} \|f'\|_{\mathcal{B}}$$

To see that this is indeed a norm we note that  $f' \sim f$  implies that  $f = f' + s$  for some non-zero  $s \in \mathcal{S}$ . We check scalar multiplication by noting that if  $\alpha \neq 0$  then

$$\begin{aligned} \|\alpha[f]\|_{\mathcal{B}/\mathcal{S}} &= \|[\alpha f]\|_{\mathcal{B}/\mathcal{S}} \\ &= \inf_{s \in \mathcal{S}} \|\alpha f + s\|_{\mathcal{B}} \\ &= \inf_{s \in \mathcal{S}} \|\alpha f + \alpha s\|_{\mathcal{B}} \\ &= |\alpha| \inf_{s \in \mathcal{S}} \|f + s\|_{\mathcal{B}} \\ &= |\alpha| \|[f]\|_{\mathcal{B}/\mathcal{S}} \end{aligned}$$

Equality still holds when  $\alpha = 0$  because  $[0] \in \mathcal{S}$  and  $\inf_{s \in \mathcal{S}} \|s\| = 0$ . Now we check the triangle inequality by computing

$$\begin{aligned} \|[f] + [g]\|_{\mathcal{B}/\mathcal{S}} &= \|[f + g]\|_{\mathcal{B}/\mathcal{S}} \\ &= \inf_{s \in \mathcal{S}} \|f + g + s\|_{\mathcal{B}} \\ &= \inf_{s, s' \in \mathcal{S}} \|f + s + g + s'\|_{\mathcal{B}} \\ &= \inf_{s \in \mathcal{S}} \|f + s\|_{\mathcal{B}} + \|g + s\|_{\mathcal{B}} \\ &= \|[f]\|_{\mathcal{B}/\mathcal{S}} + \|[g]\|_{\mathcal{B}/\mathcal{S}} \end{aligned}$$

Now we need only verify that  $\|[f]\| = 0$  implies that  $[f] = 0$ . Let  $f \in \mathcal{B}$  be such that  $\|[f]\| = 0$ . Then we must have that  $\inf_{s \in \mathcal{S}} \|f + s\|_{\mathcal{B}} = 0$ . So for each  $n > 0$  we can find an  $s_n \in \mathcal{S}$  such that  $\|f + s_n\|_{\mathcal{B}} < 1/n$ . Consequently, we must have that  $-s_n \rightarrow f$  as  $n \rightarrow \infty$ . Because  $\mathcal{S}$  is closed we must have that  $f \in \mathcal{S}$  and therefore  $[f] = 0$  in  $\mathcal{B}/\mathcal{S}$ .

Now we need to show that  $\mathcal{B}/\mathcal{S}$  is complete. Let  $\sum_n F_n$  be an absolutely convergent series in  $\mathcal{B}/\mathcal{S}$ . By definition of  $\|\cdot\|_{\mathcal{B}/\mathcal{S}}$  we can find  $f_n$  such that

$$\|f_n\|_{\mathcal{B}} \leq \|F_n\|_{\mathcal{B}/\mathcal{S}} + 2^{-n}$$

It is then clear that  $\sum_n |f_n|$  is bounded and hence converges and therefore  $\sum_n f_n$  converges because  $\mathcal{B}$  is a Banach space.

Let  $f = \sum_n f_n$  and let  $S_N(f)$  denote the  $N^{\text{th}}$  partial sum. We then see that

$$\begin{aligned} \|[S_N(f)] - [f]\|_{\mathcal{B}/\mathcal{S}} &= \|S_N([f_n - f])\|_{\mathcal{B}/\mathcal{S}} \\ &= \inf_{s \in \mathcal{S}} \|S_N(f) - f + s\|_{\mathcal{B}} \\ &\leq \|S_N(f) - f + S_N(s)\|_{\mathcal{B}} \\ &= \|S_N(f + m) - f\|_{\mathcal{B}} \end{aligned}$$

The last quantity clearly goes to 0 as  $N \rightarrow \infty$ . Thus, we have that  $S_N([f]) \rightarrow [f]$  as  $N \rightarrow \infty$  and so  $\mathcal{B}/\mathcal{S}$  is complete.

**Problem 1.8.32**

If  $\mathcal{B}^*$  is separable then we can find a countable dense subset  $\{\varphi_1, \varphi_2, \dots\}$ . Now choose a set of unit vectors  $f \in \mathcal{B}$  such that  $|\varphi_n(x_n)| \geq \frac{1}{2}\|\varphi_n\|$ . Let  $\mathcal{C}$  be the set of linear combinations of the  $x_n$ . We need to show that  $\mathcal{C}$  is dense in  $\mathcal{B}$ . Suppose not, then we have that  $\bar{\mathcal{C}}$  is a proper closed subspace of  $\mathcal{B}$ . So we can find some non-zero bounded linear functional  $L \in \mathcal{B}^*$  such that  $L(\bar{\mathcal{C}}) = 0$ . Since  $L \in \mathcal{B}^*$  and  $\{\varphi_n\}$  is dense in  $\mathcal{B}^*$  we must have some sequence  $\varphi_{n_k} \rightarrow L$  as  $n_k \rightarrow \infty$ . Hence  $\|L - \varphi_{n_k}\| \rightarrow 0$  as  $n_k \rightarrow \infty$ . But,

$$\begin{aligned} \|L - \varphi_{n_k}\| &\geq |(L - \varphi_{n_k})(x_{n_k})| \\ &= |\varphi_{n_k}(x_{n_k})| \\ &\geq \frac{1}{2}\|\varphi_{n_k}\| \end{aligned}$$

So we must have that  $\|\varphi_{n_k}\| \rightarrow 0$  as  $n_k \rightarrow \infty$ . However,  $\varphi_{n_k} \rightarrow L$  and so  $\|\varphi_{n_k}\| \rightarrow \|L\|$ , which means that  $\|L\| = 0$ . This is a contradiction, and therefore we have that  $\mathcal{C}$  is dense in  $\mathcal{B}$ .

**Problem 1.8.33**

Following the hint, we take  $u = \text{Re}(\ell_0)$ . We then apply Theorem 1.5.2 to extend  $u$  to a linear functional  $U : V \rightarrow \mathbb{R}$  such that  $U(f_0) = u(f_0)$  for every  $f_0 \in V_0$  and  $U(f) \leq p(f)$  for every  $f \in V$ . We then define

$$\ell(f) = U(f) - iU(if)$$

Then it is clear that  $\ell(f_0) = \ell_0(f_0)$  whenever  $f_0 \in V_0$ . Fix a  $g \in V$  and find a complex  $z$  such that  $|z| = 1$  and  $z\ell(g) = |\ell(g)|$ . Hence,

$$|\ell(g)| = z\ell(g) = \ell(zg)$$

which means that  $\ell(zg) \in \mathbb{R}$  and therefore  $\ell(zg) = U(zg)$ . As a result we see

$$|\ell(g)| = U(zg) \leq p(zg) = |z|p(g) = p(g)$$

And therefore  $\ell$  is the functional with the desired properties.

**Problem 1.8.34**

Consider the functional  $\ell : V \rightarrow \mathbb{R}$  defined by

$$\ell(f) = \frac{\|f - p(f)\|}{\|f_0 - p(f_0)\|}$$

where  $p(f)$  is the canonical projection of  $f$  onto  $\mathcal{S}$ . If  $f_0 \notin \mathcal{S}$  then the mapping is well-defined because  $f_0 \neq p(f_0)$  so the denominator is non-zero. Moreover, it is clear that  $\ell$  is continuous because the norm is continuous and the denominator is a non-zero scalar. If

$f \in \mathcal{S}$  then we have that  $f = p(f)$  and so  $\ell(f) = 0$ . To check the other conditions we compute

$$\ell(f_0) = \frac{\|f_0 - p(f_0)\|}{\|f_0 - p(f_0)\|} = 1$$

So  $\ell(f_0) = 1$ .

Now we need to verify that  $\|\ell\| = 1/d$  where  $d$  is the distance from  $f_0$  to  $\mathcal{S}$ . We begin by computing

$$\begin{aligned} \|\ell\| &= \sup_{f \neq 0} \frac{|\ell(f)|}{\|f\|} \\ &= \sup_{f \neq 0} \frac{\|f - p(f)\|}{\|f\|} \cdot \frac{1}{d} \\ &= \sup_{f \neq 0} \left( 1 - \frac{\|p(f)\|}{\|f\|} \right) \cdot \frac{1}{d} \\ &\leq \frac{1}{d} \end{aligned}$$

But we can find a case where equality holds by choosing  $f \in \ker p$ . Then we have

$$\begin{aligned} \frac{|\ell(f)|}{\|f\|} &= \sup_{f \neq 0} \frac{\|f - p(f)\|}{\|f\|} \cdot \frac{1}{d} \\ &= \sup_{f \neq 0} \frac{\|f - 0\|}{\|f\|} \cdot \frac{1}{d} \\ &= \frac{1}{d} \end{aligned}$$

Which means that  $\|\ell\| = 1/d$  as desired.