

Stat 235

Lab 4

WOOSAREE, Arun

Lab EL12

TA: Jessa Marley

November 20, 2018

1

1 a)

Keeping other parameters constant, changing the confidence level yields the following:

| Confidence Level | Margin of Error |
|------------------|-----------------|
| 0.90 | 0.300308 |
| 0.95 | 0.357839 |
| 0.99 | 0.470280 |

Table 1: Confidence levels and the corresponding Margins of Error

As seen in Table 1 above, the Margin of Error increases as the Confidence Level is increased. This makes sense because the margin of error depends on the z value, which also increases as the Confidence Level ($1 - \alpha$) increases.

1 b)

| Confidence Level | Observed Fraction of Intervals That Failed to Cover the Hypothesized Population Mean |
|------------------|--|
| 0.90 | 0.11 |
| 0.95 | 0.06 |
| 0.99 | 0.02 |

Table 2: Observed fraction of intervals that failed to cover the hypothesized population mean at various confidence levels.

Theoretically, the confidence level and the fraction of intervals that failed to cover the hypothesized mean should be add up to 1. Here, we see that the values are reasonable, adding up to 1.01 in all cases. Of course, a small difference is expected since we are using experimental data.

2

$$H_0 : \mu = 64 \quad vs. \quad H_A : \mu \neq 64$$

2 a)

| Level of Significance | Number of Samples That Led to the Rejection of H_0 | Observed Fraction of Samples |
|-----------------------|--|---------------------------------|
| 0.10 | 89 | 0.89 |
| 0.05 | 94 | 0.94 |
| 0.01 | 98 | 0.98 |

Table 3: Samples that led to the rejection of H_0 at different levels of significance.

As the level of significance increases, the number of samples that led to the rejection of H_0 also increases. This makes sense, because the the probability of making a Type I error is α , (or the level of significance) which happens when the null hypothesis is incorrectly rejected. With a higher tolerance for error, we would expect the amount of incorrect rejections to increase, which would increase the total fraction of samples rejected.

2 b)**90% confidence interval null hypothesis**

The 90% confidence interval is calculated as follows:

$$64 \pm 0.300308 = (63.699692, 64.300308)$$

Thus, the null hypothesis is:

$$H_0 : \mu \in (63.699692, 64.300308)$$

95% confidence interval null hypothesis

The 95% confidence interval is calculated as follows:

$$64 \pm 0.357839 = (63.642161, 64.357839)$$

Thus, the null hypothesis is:

$$H_0 : \mu \in (63.642161, 64.357839)$$

99% confidence interval null hypothesis

The 95% confidence interval is calculated as follows:

$$64 \pm 0.470280 = (63.52972, 64.47028)$$

Thus, the null hypothesis is:

$$H_0 : \mu \in (63.52972, 64.47028)$$

When is the null hypothesis rejected?

The null hypothesis H_0 is rejected when the sample mean does not fall within the confidence interval with significance α .

3**3 a)**

| Alloy 1 | |
|-------------------------|-------------|
| Mean | 65.09 |
| Standard Error | 0.360980466 |
| Median | 64.6 |
| Mode | 63.8 |
| Standard Deviation | 1.977171438 |
| Sample Variance | 3.909206897 |
| Kurtosis | 0.042639157 |
| Skewness | 0.718164135 |
| Range | 8.2 |
| Minimum | 61.7 |
| Maximum | 69.9 |
| Sum | 1952.7 |
| Count | 30 |
| Confidence Level(95.0%) | 0.738287948 |

Table 4: Summary statistics for Alloy 1

The confidence interval is calculated as follows:

$$65.09 \pm 0.738287948 \approx (64.352, 65.828)$$

| Alloy 2 | |
|-------------------------|-------------|
| Mean | 65.27333333 |
| Standard Error | 0.167601973 |
| Median | 65 |
| Mode | 64.9 |
| Standard Deviation | 0.917993815 |
| Sample Variance | 0.842712644 |
| Kurtosis | 9.565960304 |
| Skewness | 2.914366915 |
| Range | 4.5 |
| Minimum | 64.5 |
| Maximum | 69 |
| Sum | 1958.2 |
| Count | 30 |
| Confidence Level(95.0%) | 0.342784524 |

Table 5: Summary statistics for Alloy 2

The confidence interval is calculated as follows:

$$65.27333333 \pm 0.342784524 \approx (64.931, 65.616)$$

Alloy 2 appears to be stronger, since it has a higher mean, median and mode compared to Alloy 1.

3 b)

For both of the alloys, there isn't any strong evidence that the mean strength is below the required threshold value of 64. For both of the 95% confidence intervals, the lower bound is above 64, so the chance for a mean strength below 64 is low.

4

4 a)

Alloy 1

$$H_0 : \mu \leq 64 \quad vs. \quad H_A : \mu > 64$$

$$t_0 = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} = \frac{65.09 - 64}{0.917993815/\sqrt{30}} \approx t_{29}$$

$$p\text{-value} : pt(3.019553976, df = 29, lower.tail = FALSE) = 0.0026185$$

Because the p-value obtained is extremely small, we reject H_0 . i.e. the data strongly suggests that Alloy 1 exceeds the threshold value of 64.

Alloy 2

$$H_0 : \mu \leq 64 \quad vs. \quad H_A : \mu > 64$$

$$t_0 = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} = \frac{65.27333333 - 64}{0.917993815/\sqrt{30}} = 7.597364802 \sim t_{29}$$

$$p\text{-value} : pt(7.597364802, df = 29, lower.tail = FALSE) = 1.121033 \times 10^{-8}$$

Again, this p-value is extremely close to zero, so we reject H_0 . i.e. the data strongly suggests that Alloy 2 exceeds the threshold value of 64.

4 b)

The following assumptions must be true for a t-test:

1. The samples are independent and random
2. The samples come from a normal population, OR from a population with sample size 30 or greater (The Central Limit Theorem guarantees samples are normally distributed when the population size is ≥ 30)

The assumptions outlined above hold for the test above, as we were told that the rods were randomly selected, and the sample size of the population is 30.

5**5 a)**

We'll do the two-tailed t-test as follows:

Since, the population variances are unknown, we'll assume unequal variances.

$$H_0 : \mu_1 = \mu_2 \quad vs. \quad H_A : \mu_1 \neq \mu_2$$

Using the "t-Test: Two-Sample Assuming Unequal Variances" tool, we obtain the following:

| | Alloy 1 | Alloy 2 |
|------------------------------|--------------|----------|
| Mean | 65.09 | 65.27333 |
| Variance | 3.909206897 | 0.842713 |
| Observations | 30 | 30 |
| Hypothesized Mean Difference | 0 | |
| df | 41 | |
| t Stat | -0.460646232 | |
| $P(T \leq t)$ one-tail | 0.32374327 | |
| t Critical one-tail | 1.682878002 | |
| $P(T \leq t)$ two-tail | 0.647486541 | |
| t Critical two-tail | 2.01954097 | |

Table 6: Excel output of the t-test. A value of $\alpha = 0.05$ was used.

The test statistic $t_0 = -0.460646232$ follows a t-distribution, with corresponding p-value: 0.647486541 obtained from the table above.

$$t_{\alpha/2, \min\{n-1, m-1\}} = t_{0.05/2, 29} = pt(0.025, df = 29, lower.tail = TRUE) = 0.5098869$$

H_0 should be rejected if t_0 is greater than the value above, which isn't the case. Similarly, with the "judgement approach" we find that the p-value is above 0.1, which is weak to no evidence against H_0 . Therefore, we fail to reject H_0 . i.e. there is not sufficient evidence to support that there is a difference in the mean strengths of Alloy 1 and Alloy 2 rods.

5 b)

The assumptions to make the tests in part (a) valid are as follows:

1. The samples are independent and random
2. The samples come from a normal population, OR from a population with sample size 30 or greater (The Central Limit Theorem guarantees samples are normally distributed when the population size is ≥ 30)
3. the populations have unequal variances

The first two assumptions outlined hold for the test above, as we were told that the rods were randomly selected, and the sample size of the population is 30. **The population variances are unknown so we don't know if the 3rd assumption is valid?!**

6

6 a)

Do the data provide evidence that the treatment increased the mean strength of the ALLOY 2 rods? Answer the question by carrying out an appropriate

test in Excel. In particular, state the null and alternative hypotheses in terms of the population parameters, obtain the value of the test statistic, specify the distribution of the test statistic under the null hypothesis, and obtain the p-value of the test. What do you conclude?

We'll do the one-tailed t-test for paired data as follows:

$$H_0 : \mu_2 - \mu_{2+treatment} \geq 0 \quad vs. \quad H_A : \mu_2 - \mu_{2+treatment} < 0$$

Using the “t-Test: Paired Two Sample for Means” tool, we obtain the following:

| | Variable 1 | Variable 2 |
|------------------------------|------------|------------|
| Mean | 65.27333 | 66.8233333 |
| Variance | 0.842713 | 0.3521954 |
| Observations | 30 | 30 |
| Pearson Correlation | 0.86516 | |
| Hypothesized Mean Difference | 0 | |
| df | 29 | |
| t Stat | -16.9038 | |
| $P(T \leq t)$ one-tail | 7.41E-17 | |
| t Critical one-tail | 1.699127 | |
| $P(T \leq t)$ two-tail | 1.48E-16 | |
| t Critical two-tail | 2.04523 | |

Table 7: Excel output of the t-test. A value of $\alpha = 0.05$ was used.

The test statistic $t_0 = -16.9038$ follows a t-distribution, with corresponding p-value: 7.41×10^{-17} obtained from the table above.

We immediately notice that the p-value is so incredibly small that we reject H_0 . i.e. the data strongly suggests that the treatment increased the mean strength of the Alloy 2 rods.

6 b)

| | |
|--------------------------|----------|
| Mean | 1.55 |
| Standard Error | 0.091695 |
| Median | 1.6 |
| Mode | 2 |
| Standard Deviation | 0.502236 |
| Sample Variance | 0.252241 |
| Kurtosis | 0.942104 |
| Skewness | -0.9204 |
| Range | 2.2 |
| Minimum | 0.1 |
| Maximum | 2.3 |
| Sum | 46.5 |
| Count | 30 |
| Confidence Level (95.0%) | 0.187538 |

Table 8: My caption

The two-sided confidence interval is found as follows:

$$1.55 \pm 0.187538 \approx (1.362, 1.738)$$

This interval is consistent with the test outcome in part (a), because it indicates with high confidence that the population mean of Alloy 2 + treatment is between 1.362 and 1.738 ksi stronger than the population mean of Alloy 2 (sans treatment). This agrees with the test outcome in part (a), which indicated with high confidence that the population mean of Alloy 2 + treatment is higher than the population mean of Alloy 2 (without treatment).

6 c)

The following assumptions are necessary to make the test in part (a) and confidence interval in part (b) valid:

1. The samples are random
2. The samples come from a normal population, OR from a population with sample size 30 or greater (The Central Limit Theorem guarantees samples are normally distributed when the population size is ≥ 30)
3. The sample data is paired.

The first two assumptions hold, as we were told that the rods were randomly selected, and the sample size of the population is 30. We know the data is paired, since both samples have to do with Alloy 2, just that in one sample, Alloy 2 is treated, so the third assumption also holds.

6 d)




Figure 1: Plot of the strength of Alloy 2 vs. the effect of the treatment.

From the scatterplot above, we can clearly see that with higher strength, the effect is less. This suggests that the effect of the treatment is not independent of the initial strength of the rods. The relationship may be linear.