Overview of Distribution in Continuous Variables



$$P(event) = \frac{count \ of \ outcomes \ in \ Event}{count \ of \ outcomes \ in \ Sample \ Space}$$



$$P(event) = \frac{count \ of \ outcomes \ in \ Event}{count \ of \ outcomes \ in \ Sample \ Space}$$

For continuous variable?



$$P(event) = \frac{count \ of \ outcomes \ in \ Event}{count \ of \ outcomes \ in \ Sample \ Space}$$

- For continuous variable?
- Sample Space has infinite values



$$P(event) = \frac{count \ of \ outcomes \ in \ Event}{count \ of \ outcomes \ in \ Sample \ Space}$$

- For continuous variable?
- Sample Space has infinite values
- So Probability at a particular value is 0



$$P(event) = \frac{count \ of \ outcomes \ in \ Event}{count \ of \ outcomes \ in \ Sample \ Space}$$

- For continuous variable?
- Sample Space has infinite values
- So Probability at a particular value is 0
- Probability Density Function



Probability Density Function (PDF): Describes the probability distribution of continuous variables.




Probability Density Function (PDF): Describes the probability distribution of continuous variables.

$$P(a \le x \le b) = \int_a^b f(x) dx$$



Probability Density Function (PDF): Describes the probability distribution of continuous variables.

$$P(a \le x \le b) = \int_{a}^{b} f(x)dx$$



The graph of the density function is continuous over its defined domain.




- The graph of the density function is continuous over its defined domain.
- The entire area under the curve of the density function is equal to 1.



- The graph of the density function is continuous over its defined domain.
- The entire area under the curve of the density function is equal to 1.


$$P(a \le x \le b) = \int_{a}^{b} f(x)dx \ge 0$$






- The graph of the density function is continuous over its defined domain.
- The entire area under the curve of the density function is equal to 1.
- Probability is the area under the density function bounded by a and b.

$$P(a \le x \le b) = \int_a^b f(x)dx \ge 0$$







The uniform distribution is a continuous probability distribution and is concerned with events that are equally likely to occur.



The uniform distribution is a continuous probability distribution and is concerned with events that are equally likely to occur.

The Probability Density Function for Uniform Distribution is :-


$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } b \ge x \ge a \\ 0 & \text{otherwise} \end{cases}$$



The uniform distribution is a continuous probability distribution and is concerned with events that are equally likely to occur.

The Probability Density Function for Uniform Distribution is :-

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } b \ge x \ge a \\ 0 & \text{otherwise} \end{cases}$$






The uniform distribution is concerned with events that are equally likely to occur.

Ex. Suppose a flight is about to land within 30 minutes. Find the probability of getting flight land between 25 to 30 minutes?



The uniform distribution is concerned with events that are equally likely to occur.

Ex. Suppose a flight is about to land within 30 minutes. Find the probability of getting flight land between 25 to 30 minutes?





The exponential distribution is generally concerned with the amount of time until some specific event occurs.



The exponential distribution is generally concerned with the amount of time until some specific event occurs.

The Probability Density Function for Exponential Distribution is:-


$$f(x) = \begin{cases} (1/\mu) \cdot e^{-(1/\mu) x} & x \ge 0 \\ 0 & \text{otherwise} \end{cases}$$
 here μ is historic average waiting time



The exponential distribution is generally concerned with the amount of time until some specific event occurs.

The Probability Density Function for Exponential Distribution is:-

$$f(x) = \begin{cases} (1/\mu) \cdot e^{-(1/\mu)x} & x \ge 0 \\ 0 & \text{otherwise} \end{cases}$$
 here μ is historic average waiting time






The exponential distribution is generally concerned with the amount of time until some specific event occurs.

Ex. On average, your spouse makes you wait for 10 minutes before leaving for the party. What is the probability that he/she will make you wait for more than 7 minutes?



The exponential distribution is generally concerned with the amount of time until some specific event occurs.

Ex. On average, your spouse makes you wait for 10 minutes before leaving for the party. What is the probability that he/she will make you wait for more than 7 minutes?






Poisson Distribution

The exponential distribution is generally concerned with the amount of time until some specific event occurs.

Ex. On average, your spouse makes you wait for 10 minutes before leaving for the party. What is the probability that he/she will make you wait for more than 7 minutes?








The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables.



The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables. Ex. Birth Weight, IQ Graph, Stock Price









The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables.

Probability Density Function for Normal Distribution is :-

$$f(\mathbf{x}) = \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

 μ here is Mean Value . σ here is Variance.




The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables.

Probability Density Function for Normal Distribution is :-

$$f(\mathbf{x}) = \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

 μ here is Mean Value .

 σ here is Variance.






The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables.

Ex. With the mean value 29 and standard deviation 6, BMI is normally distributed, Find out the probability of a person having BMI less than 35?



The normal distribution is essentially a frequency distribution curve which is often formed naturally by continuous variables.

Ex. With the mean value 29 and standard deviation 6, BMI is normally distributed, Find out the probability of a person having BMI less than





Thank You!

