PATTERN_BNU_RR: B(ν,a) and T-phase fibres vs MF2

2026-10-24 (America/Chicago)

RESULTS: PATTERN_BNU_RR (Map-PBR1, Map-PBR2)

PREREG sha256 51050fe71935c8113088b4eea76cf5684177ace94b69a11f38798104ac7f33bb. Hashed after explore and before test (logs/test.log).

fibre	ρ_res (MF2)	PBR1: 10·B(ν,a) mod 8	PBR2: 140·T-phase mod 8
0	1/32	1/8	8/35
1	1/4	1/8	11/70
2	3/32	1/8	0
3	0	1/8	9/70
4	7/32	1/8	19/70
5	1/4	1/8	3/35
6	5/32	1/8	0
7	0	1/8	9/70

- PBR1: TV = 3/8. This is uniform on all 8 fibres (not uniform-on-6, so not VACUOUS by the PREREG definition), but it carries no information. **FAIL (Category A).**
- PBR2: TV = 71/140. Mass 9/35 sits on {3,7}, and fibres 2 and 6 are empty. **FAIL (Category A).**
- Both maps escape PS1a (fibres 1 and 5 get mass), but neither matches.

Bridge: **NO.** ## NOTES: PATTERN_BNU_RR ## Lemma PB1 (Category A) B(ν,a) = (7/2)ν2, since Aa = (0,7/2). With ν2 = α2 + n2 and 35α2 ∈ Z, 10·B(ν,a) = 35ν2 = 35α2 + 35n2. Since 35 ≡ 3 is a unit mod 8, each label's fibre is uniform on Z/8. So **every** non-negative label pool gives exactly (1/8,…,1/8). The B(ν,a) fibre therefore carries no label information mod 8, and it cannot produce zeros at 3 and 7. The signed w-content (explore, R=200) is also near-uniform (0.119–0.129).

Lemma PB2 (Category A)

The T-phase 140(Q(α)+B(α,w_c)) is one value per α, with β-independent fibre. So the 140-label pool is a fixed finite multiset: (16,11,0,9,19,6,0,9)/70. It is non-zero on {3,7} and zero on {2,6}, so no label pool under the T-fibre can match MF2's support pattern unless some labels get weight 0. Restricting to a subset of labels would be a selection made after looking, so it is not done.

Observation (post hoc, not a claim)

PBR2 empties the even fibres {2,6}. Earlier maps emptied {1,5,6} (PQ1/PS1). MF2 empties {3,7}. No written reduction maps one zero pattern to another.

Status of written Resonant fibre coordinates (all tested)

Coordinate	Map	Result
unary n^2	BATCH8	FAIL
shifted u^2	PF1	FAIL
140Q	PQ1, PS1	FAIL
10B(ν,a)	PB1	uniform, FAIL
T-phase	PB2	FAIL

Next pattern step

The remaining written structure is the **S-law** (l.274): the label mixing u_{α,β}(−1/w) = e(B(α,β))/√35 Σ_p u_{p−β,α}. - A pattern-forced test is the Markov-type step on labels given by $|S|^2$ (the transition matrix from S, the way MF2 is a Markov law on the residual side), with its stationary law pushed to a written fibre coordinate (T-phase or 140Q). This makes the comparison object a Markov stationary law on both sides. - Alternatively: Arvin's written Π / grading, if it is ever supplied.