order33 RESULTS
PREREG sha256 886159282d45a48155f068625164b3e98253c82d4a8262b185fe189944a4c672 (hashed before numerics).
Test set: CO and ((1,0),(n,1)), n = 1..8 (9 cones). Criterion: w* = −1, |κ̂1| ≤ 0.01, r ≤ 0.10. Every piece is non-vacuous (d′ from span{CO, Γ_γCO, vtx}: T_e1 0.299; T_other 0.99–1.0; T_BB 0.85–1.0).
Control
max|T_e1 + T_other + T_BB − stored Cor| = 5.7e−14: PASS.
RG29: PASS (Category B)
piece	cones meeting near-law	r(−1)	κ↑
T_e1	9/9	0.0030 (all cones)	1.0000
T_other	1/9 (K only)	0.021 (K), 0.041 → 0.192 (n = 1..8; fails r for n ≥ 2 and κ for n ≥ 6)	0.975–0.999
T_BB	0/9	0.18 (K), 0.35 → 0.59	0.78–0.99
T_e1+T_other	9/9	0.0031–0.0037	1.0000
T_e1+T_BB	9/9	0.059–0.094 (= Cor's r)	0.996–0.9995
Cor	8/9 (K = CO is excluded as span-vacuous by construction)	0.058–0.094	≈0.998

Hypothesis H holds as written. Prediction PASS (recorded as the hypothesis with doubt): correct.
RG30
Defect share ‖R(T_other+T_BB)‖²/‖R(Cor)‖²: median 0.9986; cosine with R(Cor): median 0.9992. Accounts for most: YES.
Directions across cones: pairwise |cos| min 0.0015, mean 0.022. Scattered: YES.
Sharper reading (descriptive, Category B; post-hoc but disclosed)
- The B2 part T_e1+T_other obeys the law to r ≈ 0.003 on every cone; this sum is the B2 part of the corner. The near-law defect is carried by the B̄1B̄1 term T_BB, and T_other alone fails only because it is a piece of a law-obeying sum.
- T_e1 is one fixed vector up to scale. In the basis ((1,0),(n,1)) the second coordinate is b = ν2 for every n, and B = 2n+7, so T_e1(n) = ((2n+7)/7)·T_e1(K). This holds exactly by the formula, and ‖R(T_e1)‖/(2n+7) = 0.38451 for all n. It explains why every e1-cone shares the T_e1 behaviour.
- T_e1 itself misses the law by r = 0.0030. That is small but not zero, so it is not an exact law; it is Category B.
Category
A: nothing new proved. The identity T_e1(n) = ((2n+7)/7)T_e1(K) is an immediate consequence of the corner formula; I note it but do not add it to the paper. v11: no. Ξ: not identified. E1–E3 OPEN.

order33 NOTES
- The near-law sits in the B2 (Bernoulli-2) part of the corner, r ≈ 0.003. The ~6–9% defect of e1-cones is the B̄1B̄1 product term, whose law miss is cone-specific and scattered.
- Why do cones without the e1 edge fail? Probably because their B2 part also obeys the law but their B̄1B̄1 term is relatively larger. That is untested; it is order34 material.
- Open next step (order34, not started): preregister RG31 with these hypotheses:
  (i) the B2 part T_B2 = T_e1+T_other has r ≤ 0.01 for ALL unimodular cones, including the 28 control cones;
  (ii) the B̄1B̄1 term fails for all cones;
  (iii) T_e1(K) = Σ f·(7/4)B2({ν2}) is related to the Bernoulli S-law of order20 (L_B PASS). Test whether T_e1 and T_other are exactly sums of order20 Bernoulli-law objects, which would explain the residual 0.003 as a known S-law correction.
  Thresholds fixed in advance; non-vacuity span {CO, Γ_γCO, vtx}.
- Naming: RG* only, no labels L1–L7. Geometric only; MAPS_CANDIDATES untouched; no Resonant objects; no bridge claim; CORE_FREEZE. nice, OMP_NUM_THREADS=1, 3 GB, ~106 s; no push or sends; new files only.