order35 RESULTS: RG32 (quantitative L_B leakage)
PREREG sha256 04ab3e7dddfee76d7ce7e4f089325a146b7e0b8a8bd515a9a61a4ae5bbfb8007 (hashed before numerics).
Grid: P7 = {1, 1/2, 2, 1/3, 3, 2/3, 3/2} × 105 companions, as fixed in PREREG. The full grid is infeasible for the exact L_B right-hand side.
Control D0: PASS
Exact L_B (ii) reconstruction S_in + S_out = C2b(−1/x): max error 3.3e−15. Terms: 42 875 inside (reduce to a companion) and 52 675 outside.
RG32: FAIL (Category B)
quantity	value
r_meas(P7) (direct, w = −1, κ free)	0.05853 (κ≙ 0.9977 − 0.0229i)
r_leak (outside mass ⊥ law direction, no fit)	0.3527
relative difference	5.03 (threshold PASS ≤ 1e−6, FAIL ≥ 1e−2)
H1: S_in ∈ span{X}	FAIL (perpendicular share 0.926)
outside share ‖S_out‖/‖S‖	0.926
k = 0 share of the inside part	0.325

Prediction FAIL (blind): correct.
Reading
- Classifying L_B terms by whether their characteristic reduces to a companion is not a classification by span. 93% of S sits in "outside" characteristics, yet S is still within ~6% of the fixed-M law direction on P7. The outside Gauss means at x must be largely linear combinations of companion means through relations beyond the reduction rules (Gauss-mean linear relations).
- So the outside-characteristic mass does not measure the law miss, and the near-law size is NOT an exact consequence of L_B leakage in this sense.
- On P7, r(C2b) = 0.0585, far above the full-grid 0.00305. The near-law quality depends strongly on which points are in the grid (the small-denominator points are worse). As disclosed in PREREG, 0.00305 itself could not be tested.
- (0,1)-edge cones (r ≈ 0.022): out of scope as preregistered. L_B is proved only for the ρ2 coordinate; nothing was tested and no new objects were introduced.
Category
A: none new. v11: no. Ξ: not identified. E1–E3 OPEN.

order35 NOTES
- Lesson: "outside the companions by characteristic" ≠ "outside the span of the 35-component law vector". The correct leakage measure is a span projection. It needs the companion-span of Gauss means at each x (rank 36 ambient; order20 RG7 joint ranks).
- Open next step (order36, not started): preregister RG33. At each grid point, project the exact L_B outside terms onto the span of the 105 companion Gauss means (G(x) = A0(x)/I, rank 36). Define the leakage as the residual of that projection, and predict r(C2b) from it with no fit (r_pred = ‖P⊥_span S‖-type quantity, together with the κ-direction bookkeeping), on P7 first. Also state in advance how grid dependence (P7 0.0585 vs full 0.00305) is to be handled, e.g. per-point r.
- CORE_FREEZE; new files only. Geometric only; MAPS_CANDIDATES untouched; no Resonant objects; no bridge claim. Naming: RG* only, no labels L1–L7. nice, OMP_NUM_THREADS=1, 3 GB, ~66 s; no push or sends.