order26 RESULTS
PREREG sha256 d0a31ded81d93c64f75496b3c729672a6d1281ace33d16e0e2189c726736a755 (hashed before numerics).
Between-cone B = cone((7,−1),(1,0)), unimodular; the only unimodular piece between $\gamma^{-1}K$ and K (the unions have det 2 and det 7).

test	value	verdict (preregistered rule)
C1 corner(K)=CO	0.0	PASS
C2 $\Gamma_\gamma \cdot CO = Cor_{\gamma^{-1}K}$	1.6e−14	PASS
C3 $Y_- = -\frac{1}{2}D$	7.9e−15	PASS
C4 RG15 abs vs 52.0306 (≤1e−9)	1.48e−5	**FAIL by rule** (see note)
I1 D = ±Cor_B + t·vtx	min max-error 72.48 (s=−1); vertex term has no effect; LS fit relative residual 0.597 (a=−1.287)	**FAIL**
RG17 R(Y_−) carried by P_−R(Cor_B)	β = 1.000, r = 1.1e−14, non-vacuity 1.00	**PASS by rule, but trivial** (see below)
RG17′ Cor_B against the weight-1 law	abs 54.406, rel 0.0586	FAIL

C4 note: the threshold was set against a 4-decimal rounded constant; that was a preregistration error. Post hoc, against the stored order25 value 52.030585237…, the difference is at floating-point level. The rule verdict stands as FAIL.

Post-hoc finding (after verdicts were recorded)
Cor_B = Cor_K numerically: 7.6e−15 at x, 2.0e−14 at −1/x. B = hK with h = [[1,7],[0,−1]] = −σ, an improper automorph of A (checked). So RG17 reduces to P_−R(Cor_K) = R(Y_−), which is order25's split itself. It carries no new information, and the non-vacuity check (a norm ratio) did not catch this; that is a lesson for future preregistrations.
A naive exact pair-transport check of Γ_h (same convention as Γ_γ) gives identity on only 3 of 105 entries, so the equality $Cor_{hK} = Cor_K$ is **not** yet explained by an exact argument (a convention or sign symmetry of the corner is suspected). It is Category B only.

Answers
- RG17: PASS by the letter of the rule, but trivial (between-cone corner = K corner).
- Between-cone identity $Cor_K - Cor_{\gamma^{-1}K} = \pm corner(B)$ (+ vertex terms): **does not hold** (error 72.5).
- Category A: none new. Category B: I1 FAIL, Cor_B = Cor_K to 2e−14, RG17′ FAIL.
- v10: no. Ξ: not identified. E1–E3 OPEN.

order26 NOTES
- Lesson: a map object can be an automorph image of the base object; the non-vacuity check must also require ‖X − Γ-images of CO‖ to be non-negligible, not just a norm ratio.
- Open (Category A candidate): prove $Cor_{hK} = Cor_K$ for h = −σ. Likely route: the corner weights are symmetric under the swap u1↔u2 and $\bar{B}1/B2$ parity, together with the order23 result $\Gamma_\sigma = I$ under the correct transport convention. The naive Γ_h permutation check gave 3/105, so the convention must be settled first.
- Consequence: the defect $D = Cor_{\gamma^{-1}K} - Cor_K$ equals $Cor_{\gamma^{-1}K} - Cor_{hK}$. $h\gamma^{-1}$ is an improper automorph, so the γ-defect is a defect between two improper-automorph images of K, not a single-cone corner.
- Next: settle the transport convention for improper automorphs exactly; then test whether D is a Γ_{h}-anti-invariant combination ($\Gamma_h \cdot \Gamma_\gamma$ structure, dihedral group generated by γ and h). Preregister in order27.
- Lock respected: MAPS_CANDIDATES untouched; no Resonant objects; no bridge claim. nice, OMP_NUM_THREADS=1, ulimit 3 GB, ~36 s; no push/sends; new files only.