order32 RESULTS

PREREG sha256 d1a77766ed79b5347646c9e4b90e9e0784efd504b37db07e8cc551f2ac1900ad (hashed before numerics; not rehashed).

Criterion (as in order31): w* = −1, |κ̂−1| ≤ 0.01, r(−1) ≤ 0.10. All cones non-vacuous (d′ from span{CO, Γ_γCO, vtx}: 0.32–0.999).

RG27: PASS (Category B)

set	n	criterion as predicted	details
E (edge e1)	10	10/10 satisfy	r 0.068–0.094, κ0.996 –0.9995, |Im ɬ̂ ≤ 0.005
C (no e1-orbit edge)	28	28/28 violate	r 0.247–0.567, κ0.81 –0.97
blind subset E (new)	6	6/6	((1,0),(n,1)) n = 5..8: r 0.0763, 0.0746, 0.0676, 0.0702
blind subset C (new)	2	2/2	((1,5),(0,1)) r 0.296; ((1,6),(0,1)) r 0.258

Disclosure (post-hoc): ((1,0),(8,−1)) = h((1,0),(1,1)) and ((1,0),(9,−1)) = h((1,0),(2,1)), since h = [[1,7],[0,−1]] fixes e1. Their values are identical to their preimages, so only 4 of the 6 "new" E cones are independent blind tests (all passed).

RG27G: PASS (consistency with proved transport)

γ⁻¹-images ((13,−2),(72,−11)) and ((13,−2),(85,−13)) satisfy the criterion; Δr ≤ 1.1e−16, Δκ̂≤ 3.3e−16 against their E preimages.

RG28: FAIL (scatter)

Over the 10 E cones the unit residual directions R(Cor_C)/‖R‖ have pairwise |cos| min 0.0021, mean 0.064, and top-singular-value share 0.20. Each e1-cone has a small relative miss (~7–9%), but the misses point in essentially orthogonal directions. There is no common defect direction. Prediction "intermediate" (blind): wrong (stronger scatter than expected).

Reading

The near-law is a property of having the minimal vector e1 (or its ⟨γ,h⟩-orbit) as an edge. This is confirmed on new cones (Category B). The defect itself is cone-specific, not one shared vector. Category A: none new. v11: no. Ξ: not identified. E1–E3 OPEN.

order32 NOTES

- RG27 confirms the e1-edge rule out of sample (4 independent new E cones, 2 new C cones); the h- and γ-images behave as transport predicts.
- RG28: defect directions scatter, so a single correction vector cannot explain the near-law across cones.
- Open next step (order33, not started): decompose Cor_C for e1-edge cones into its three pieces: (a) the e1-ray B2 term (B/(4Q(e1)))·B2(b), with Q(e1) = 1; (b) the other-ray B2 term; (c) the B̄1B̄1 term. Preregister which piece carries the near-law. Hypothesis: (a) is shared by all e1-cones up to the change of complement and obeys the law closely, while (b)+(c) produce the scattered defect. Thresholds fixed in advance; non-vacuity span {CO, Γ_γCO, vtx}.
- Run note: the background job ran once to completion (~3 min); the steering said not to poll, so I waited on the pid once.
- Naming: RG* only, no labels L1–L7. Geometric only; MAPS_CANDIDATES untouched; no Resonant objects; no bridge claim; CORE_FREEZE. nice, OMP_NUM_THREADS=1, 3 GB; no push or sends; new files only.