order43 RESULTS: exact k = 0 law and exact miss split
PREREG sha256 66dd47276a7ec06b01a7ad2f3ff7e14f8bf4710c053d83c2a95e3e78d223bdeb (hashed before numerics). 12 points × 105 companions; κând M fixed.
LD0: D0(−1/x) = (x√35)^{−1}·M_b·D0(x), with M_b = diag(B2(α2))·M·diag(B2(α2))^{−1}: PASS
Max error 2.4e−16 (threshold 1e−12); min|B2(α2)| = 1/150 > 0. Category A: an immediate corollary of the G S-law (Theorem STlaw), as noted in PREREG.
RG40: exact split R = (a) + (b): PASS (algebraic consolidation, as disclosed)
Max ‖R − R_pred‖/‖R‖ = 4.8e−15.

x	r(C2b)	‖R‖	‖(a)‖ k = 0 law mismatch	‖(b)‖ k ≠ 0 remainder	‖R‖/(‖a‖+‖b‖)
1	0.1408	0.256	0.205	0.154	0.714
2	0.0848	0.308	1.226	1.223	0.126
3	0.0639	0.348	1.874	1.874	0.093
4	0.0523	0.380	2.503	2.501	0.076
5	0.0446	0.405	3.131	3.121	0.065
6	0.0393	0.428	3.756	3.745	0.057
7	0.0353	0.450	4.384	4.369	0.051
8	0.0323	0.470	5.011	5.002	0.047
1/2	0.0847	0.154	0.613	0.611	0.126
1/3	0.0638	0.116	0.625	0.625	0.093
2/3	0.0370	0.135	0.342	0.360	0.191
3/2	0.0370	0.202	0.514	0.540	0.191

Reading
- The miss of the C2b near-law is the exact difference between an explicit k = 0 law mismatch (a) and the k ≠ 0 L_B remainder (b). They are nearly equal and opposite, with cancellation to 5–13% for x ≥ 2.
- Post-hoc observation (not a test): the cancellation ratio is the same at x and 1/x (2 vs ½: 0.12593; 3 vs ⅓: 0.09298; ⅔ vs 3/2: 0.19144). The norms scale by x between the pair.
Category, v11
A: LD0 (minor corollary of Theorem STlaw). B: the split numbers. v11: **no**. By the PREREG criterion LD0 alone does not warrant it; a non-trivial statement about the remainder is needed. order44: not run. The preregistered rule says no law for the remainder can be written without fitting. Ξ: not identified. E1–E3 OPEN.

order43 NOTES
- The near-law is now an exact algebraic cancellation, R = (a) + (b), with (a) fully explicit (M_b vs M̂, weight 1/x vs x).
- The unknown is the k ≠ 0 remainder law: why (b) ≈ −(a) to within 5–13%.
- Open next step (order44+, not started):
  (i) preregister the x ↔ 1/x invariance of the cancellation ratio ‖R‖/(‖a‖+‖b‖) as a blind test on new pairs (e.g. 5/7 ↔ 7/5, 4 ↔ ¼, 9/2 ↔ 2/9), with a fixed tolerance;
  (ii) seek an exact relation for the k ≠ 0 remainder through L_B (ii) and the G S-law at shifted characteristics β+λ_k. That is the Category A route that could warrant v11.
- Naming: RG* only, no labels L1–L7; "36" only as dim V. MAPS_CANDIDATES untouched; no Resonant objects; no new Bernoulli objects; no bridge claim; CORE_FREEZE. nice, OMP_NUM_THREADS=1, ~2 s; no push or sends; new files only.