Beautiful Days at the Movies ★

Submissions



RATE THIS CHALLENGE

Problem

Lily likes to play games with integers. She has created a new game where she determines the difference between a number and its reverse. For instance, given the number 12, its reverse is 21. Their difference is 9. The number 120 reversed is 21, and their difference is 99.

Editorial

She decides to apply her game to decision making. She will look at a numbered range of days and will only go to a movie on a beautiful day.

Leaderboard

Given a range of numbered days, $[i \dots j]$ and a number k, determine the number of days in the range that are beautiful. Beautiful numbers are defined as numbers where |i-reverse(i)| is evenly divisible by k. If a day's value is a beautiful number, it is a beautiful day. Return the number of beautiful days in the range.

Function Description

Complete the beautiful Days function in the editor below.

beautiful Days has the following parameter(s):

- int i: the starting day number
- int j: the ending day number
- int k: the divisor

Returns

• int: the number of beautiful days in the range

Input Format

A single line of three space-separated integers describing the respective values of i, j, and k.

Constraints

- $1 \le i \le j \le 2 \times 10^6$
- $1 \le k \le 2 \times 10^9$

Sample Input

20 23 6

Sample Output

2

Explanation

Lily may go to the movies on days 20, 21, 22, and 23. We perform the following calculations to determine which days are beautiful:

- Day 20 is beautiful because the following evaluates to a whole number: $\frac{|20-02|}{6}=\frac{18}{6}=3$
- Day 21 is not beautiful because the following doesn't evaluate to a whole number: $\frac{|21-12|}{6}=\frac{9}{6}=1.5$
- Day 22 is beautiful because the following evaluates to a whole number: $\frac{|\mathbf{22}-\mathbf{22}|}{6} = \frac{0}{6} = 0$
- Day 23 is not beautiful because the following doesn't evaluate to a whole number: $\frac{|23-32|}{6} = \frac{9}{6} = 1.5$

Only two days, 20 and 22, in this interval are beautiful. Thus, we print 2 as our answer.



```
18
        int reverse(int num) {
   19
   20
            int reverse_int = 0 ;
   21
   22
            while(num > 0) {
                 reverse_int = reverse_int*10 + num % 10 ;
   23
   24
                 num /= 10 ;
   25
   26
   27
             return reverse_int ;
   28
   29
       }
   30
   31
        int beautifulDays(int i, int j, int k) {
             int beautifulDays = 0;
   32
             for(int days = i ; days <= j ; days++) {</pre>
   33
                 int reverse_day = reverse(days) ;
   34
   35
                if(abs(reverse_day - days) % k == 0) {
   36
                     beautifulDays++;
   37
   38
            }
   39
   40
             return beautifulDays;
        }
   41
   42
        int main()
                                                                                                      Line: 105 Col: 1
\hat{\bot} Upload Code as File
```

Test against custom input

Run Code

Submit Code

Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy