Lecture 4: Probability and Statistics

Math for Machine Learning

October 26, 2026

Contents

Introduction **2**

1 Probability Spaces **3**
1.1 Conditional probability and Bayes' theorem 4
1.2 Independence 4

2 Random Variables and Their Distributions **5**
2.1 Transformations of random variables 6

3 Expectation and Variance **6**
3.1 Inequalities 8

4 Joint Distributions and Independence **8**
4.1 Covariance and correlation 9
4.2 Conditional expectation 10

5 Random Vectors and Covariance Matrices **11**

6 The Gaussian Distribution **12**

7 Limit Theorems and Monte Carlo Methods **14**
7.1 Monte Carlo sampling 16

8 Markov Chains **17**
8.1 Detailed balance and Markov chain Monte Carlo 20

9 Parameter Estimation **21**
9.1 The method of moments 22
9.2 Maximum likelihood estimation 22
9.3 Consistency and efficiency 23
9.4 Loss functions as negative log-likelihoods 24
9.5 Confidence intervals 25

10 Hypothesis Testing **25**
10.1 Multiple testing 26
10.2 The bootstrap and permutation tests 27

11 Evaluating Models and Experiments **27**
11.1 Cross-validation 27
11.2 A/B testing and experiment design 28

12 Bayesian Inference **29**
12.1 Coin flips and the Beta distribution 29
12.2 Maximum a posteriori estimation and regularization 30
12.3 Bayesian linear regression 31
12.4 Gaussian processes 31

13 Information Theory **33**

14 Stochastic Differential Equations **35**
14.1 Langevin dynamics and diffusion models 36

15 Key Takeaways **37**

16 Exercises **39**

17 Programming Exercises **42**

Introduction

Machine learning is about learning from data, and data is noisy, incomplete and random: measurements contain errors, labels can be wrong, and the training set is only a sample of all the data we might see. Probability theory is the mathematical language for uncertainty, and statistics is the art of drawing conclusions from random data. Both are everywhere in machine learning:

- classifiers such as logistic regression and softmax regression (Lecture 3) output probabilities;
- the loss functions of Lecture 3 are negative log-likelihoods: least squares corresponds to Gaussian noise, and cross-entropy to categorical labels;
- regularization corresponds to prior beliefs about the parameters;
- stochastic gradient descent uses random mini-batches, and works because averages of many random terms concentrate around their mean;
- generative models, such as normalizing flows (Lecture 3), learn probability distributions.

In this lecture, we introduce probability spaces, conditional probability and Bayes' theorem, random variables, expectation and variance, and random vectors with their covariance matrices, where the linear algebra of Lectures 1 and 2 enters. We then study the Gaussian distribution in detail, the law of large numbers and the central limit theorem, Monte Carlo sampling, and Markov chains. The second half of the lecture is about learning from data: estimation (the method of moments, maximum likelihood, and the Cramér–Rao bound), hypothesis testing (p-values, multiple testing, the bootstrap and permutation tests), the evaluation of models and experiments (cross-validation and A/B testing), Bayesian inference (including Gaussian processes), and information theory (entropy, cross-entropy and the Kullback–Leibler divergence). Finally, we add Gaussian noise to the ordinary differential equations of Lecture 3: Brownian motion, stochastic differential equations and the Ornstein–Uhlenbeck process, and Langevin dynamics, which samples from a distribution by noisy gradient descent and is behind diffusion models.

Notation. As in Lecture 3, vectors are bold. Random variables are denoted by capital letters such as X and Y, their values by small letters such as x and y, and random vectors by bold capital letters such as $\mathbf{X}$. We write $P(A)$ for the probability of an event A, $\mathbb{E}[X]$ for the expectation of X, and log for the natural logarithm.

1 Probability Spaces

A random experiment has a set of possible outcomes, and we want to assign probabilities to sets of outcomes.

Definition 1.1 (Probability space)**.** A probability space consists of a sample space Ω, the set of all possible outcomes, and a probability P, which assigns a number $P(A)$ to every event $A \subseteq \Omega$, such that:

1. $P(A) \geq 0$ for every event A;
2. $P(\Omega) = 1$;
3. if $A_1, A_2, \ldots$ are pairwise disjoint events, then $P(A_1 \cup A_2 \cup \cdots) = P(A_1) + P(A_2) + \cdots$.

Remark 1.2 (Measure theory). If Ω is uncountable, such as $\Omega = \mathbb{R}$, it is impossible to assign a probability to every subset of Ω in a consistent way. A rigorous treatment specifies a collection of events (a σ-algebra) and uses measure theory. All sets that we meet in practice, such as intervals, balls and their unions and complements, are events, and we ignore these technicalities in this lecture.

Proposition 1.3. *For all events A, B and $A_1, A_2, \ldots$:*

1. *$P(A^c) = 1 - P(A)$, where $A^c = \Omega \setminus A$, and $P(\emptyset) = 0$;*
2. *if $A \subseteq B$, then $P(A) \leq P(B)$;*
3. *$P(A \cup B) = P(A) + P(B) - P(A \cap B)$;*
4. *(union bound) $P(A_1 \cup A_2 \cup \cdots) \leq P(A_1) + P(A_2) + \cdots$.*

Proof. (1) A and A^c are disjoint with union Ω, so $P(A)+P(A^c) = 1$; for $A = \Omega$, this gives $P(\emptyset) = 0$. (2) B is the disjoint union of A and $B \setminus A$, so $P(B) = P(A) + P(B \setminus A) \geq P(A)$. (3) $A \cup B$ is the disjoint union of A and $B \setminus A$, and B is the disjoint union of $B \setminus A$ and $A \cap B$; so $P(A \cup B) = P(A) + P(B \setminus A) = P(A) + P(B) - P(A \cap B)$. (4) Let $B_1 = A_1$ and $B_k = A_k \setminus (A_1 \cup \cdots \cup A_{k-1})$. The B_k are disjoint, $B_k \subseteq A_k$, and they have the same union as the A_k; now use the third axiom and (2). □

Example 1.4 (Equally likely outcomes)**.** If Ω is finite and all outcomes are equally likely, then $P(A) = \frac{|A|}{|\Omega|}$. For two fair dice, $\Omega = \{1, \ldots, 6\}^2$ has 36 outcomes. The sum is 7 for the 6 outcomes $(1,6), (2,5), \ldots, (6,1)$, so $P(\text{sum} = 7) = \frac{6}{36} = \frac{1}{6}$. The event "at least one six" is the complement of "no six", which has $5 \cdot 5 = 25$ outcomes, so its probability is $1 - \frac{25}{36} = \frac{11}{36}$.

1.1 Conditional probability and Bayes' theorem

If we learn that an event B has happened, the probabilities of other events change: only the outcomes in B are still possible.

Definition 1.5 (Conditional probability)**.** If $P(B) > 0$, the conditional probability of A given B is

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}.$$

For fixed B, the function $A \mapsto P(A \mid B)$ is again a probability. Rearranging the definition gives the product rule $P(A \cap B) = P(A \mid B)\, P(B)$.

Theorem 1.6 (Total probability and Bayes' theorem)**.** *Let $B_1, \ldots, B_n$ be disjoint events with union Ω and $P(B_i) > 0$. Then for every event A,*

$$P(A) = \sum_{i=1}^{n} P(A \mid B_i)\, P(B_i),$$

and if $P(A) > 0$, then

$$P(B_j \mid A) = \frac{P(A \mid B_j)\, P(B_j)}{P(A)} = \frac{P(A \mid B_j)\, P(B_j)}{\sum_{i=1}^{n} P(A \mid B_i)\, P(B_i)}.$$

Proof. A is the disjoint union of the events $A \cap B_i$, so $P(A) = \sum_i P(A \cap B_i) = \sum_i P(A \mid B_i) P(B_i)$ by the product rule. For the second formula, use the product rule twice: $P(B_j \mid A) P(A) = P(A \cap B_j) = P(A \mid B_j) P(B_j)$. □

Bayes' theorem reverses conditional probabilities: from the probability of the evidence A given each hypothesis B_j, it computes the probability of each hypothesis given the evidence. The probabilities $P(B_j)$ are called prior probabilities, and the $P(B_j \mid A)$ posterior probabilities.

Example 1.7 (A medical test)**.** A disease affects 1% of a population. A test detects it in 99% of the sick people, but also gives a positive result for 5% of the healthy people. What is the probability that a person with a positive test is sick? With D = "sick" and $+$ = "positive test",

$$P(D \mid +) = \frac{P(+ \mid D) P(D)}{P(+ \mid D) P(D) + P(+ \mid D^c) P(D^c)} = \frac{0.99 \cdot 0.01}{0.99 \cdot 0.01 + 0.05 \cdot 0.99} = \frac{1}{6}.$$

Although the test is quite accurate, only one in six people with a positive test is sick, because the disease is rare: among 10 000 people, about 99 sick people and 495 healthy people test positive. Ignoring the prior probability is a common mistake, called the base rate fallacy. In machine learning, the same effect makes classifiers for rare classes (such as fraud detection) produce many false alarms.

1.2 Independence

Definition 1.8 (Independence)**.** Two events A and B are independent if $P(A \cap B) = P(A) P(B)$. Events $A_1, \ldots, A_n$ are (mutually) independent if $P(A_{i_1} \cap \cdots \cap A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$ for every choice of indices $i_1 < \cdots < i_k$.

If $P(B) > 0$, then A and B are independent if and only if $P(A \mid B) = P(A)$: learning that B happened does not change the probability of A. For example, for two fair dice, the events A = "the first die is even" and B = "the sum is 7" are independent: $A \cap B$ consists of the 3 outcomes $(2,5), (4,3), (6,1)$, so $P(A \cap B) = \frac{3}{36} = \frac{1}{2} \cdot \frac{1}{6} = P(A) P(B)$. Independence is not the same as disjointness: disjoint events with positive probabilities are never independent, since $P(A \cap B) = 0$.

2 Random Variables and Their Distributions

Definition 2.1 (Random variable)**.** A random variable is a function $X : \Omega \to \mathbb{R}$ on a probability space. Its distribution describes the probabilities $P(X \in I) = P(\{\omega : X(\omega) \in I\})$ for intervals I. Its cumulative distribution function (CDF) is $F(x) = P(X \le x)$.

The CDF is nondecreasing, tends to 0 as $x \to -\infty$ and to 1 as $x \to \infty$, and $P(a < X \le b) = F(b) - F(a)$. In practice, we rarely look at the sample space: a random variable is described by its distribution. Most distributions in machine learning are of one of two kinds.

Definition 2.2 (Discrete and continuous random variables)**.** 1. X is discrete if it takes only finitely or countably many values $x_1, x_2, \ldots$. Its probability mass function (pmf) is $p(x) = P(X = x)$, and $\sum_i p(x_i) = 1$.

2. X is continuous if there is a function $p \ge 0$, its probability density function (pdf), with

$$P(a \le X \le b) = \int_a^b p(x)\,dx \qquad \text{for all } a \le b, \qquad \text{and} \qquad \int_{-\infty}^{\infty} p(x)\,dx = 1.$$

For a continuous random variable, $P(X = x) = 0$ for every single value x, and the density is the derivative of the CDF, $p = F'$ (where p is continuous). A density is not a probability: it can be larger than 1. For a small interval, $P(x \le X \le x + h) \approx p(x)h$.

Example 2.3 (Discrete distributions)**.** 1. Bernoulli(p): $X \in \{0, 1\}$ with $P(X = 1) = p$; for example, a coin flip, or a binary label.

2. Binomial(n, p): the number of successes in n independent Bernoulli(p) trials,

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \qquad k = 0, 1, \ldots, n,$$

since each of the $\binom{n}{k}$ sequences with k successes has probability $p^k(1-p)^{n-k}$. For example, for $n = 10$ and $p = 0.3$, $P(X = 3) = 120 \cdot 0.3^3 \cdot 0.7^7 \approx 0.267$ (Figure 1).

3. Geometric(p): the number of independent Bernoulli(p) trials until the first success, $P(X = k) = (1-p)^{k-1}p$ for $k = 1, 2, \ldots$.

4. Poisson(λ): $P(X = k) = e^{-\lambda}\frac{\lambda^k}{k!}$ for $k = 0, 1, 2, \ldots$; it describes counts of rare events, such as the number of clicks on an ad in an hour. It is the limit of Binomial($n, \frac{\lambda}{n}$) as $n \to \infty$.

5. Categorical($\mathbf{p}$): $X \in \{1, \ldots, K\}$ with $P(X = k) = p_k$, where $p_k \ge 0$ and $\sum_k p_k = 1$; for example, the class predicted by a softmax classifier.

Example 2.4 (Continuous distributions)**.** 1. Uniform(a, b): $p(x) = \frac{1}{b-a}$ for $a \le x \le b$, and 0 otherwise.

2. Exponential(λ): $p(x) = \lambda e^{-\lambda x}$ for $x \ge 0$; then $P(X > t) = e^{-\lambda t}$. It describes waiting times, and is memoryless: $P(X > s + t \mid X > s) = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} = P(X > t)$.

3. Normal (Gaussian) $\mathcal{N}(\mu, \sigma^2)$, with mean μ and variance $\sigma^2 > 0$:

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\Big(-\frac{(x-\mu)^2}{2\sigma^2}\Big).$$

It integrates to 1 by the Gaussian integral of Lecture 3. The standard normal distribution $\mathcal{N}(0, 1)$ has the density $\varphi(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}$ and the CDF $\Phi(z) = \int_{-\infty}^{z} \varphi(t)\,dt$, which has no formula in terms of elementary functions.




Figure 1: Left: the pmf of the Binomial$(10, 0.3)$ distribution. Right: the densities of the standard normal distribution, the Exponential(1) distribution and the Uniform$(-1, 1)$ distribution.

2.1 Transformations of random variables

If X is a random variable and $g : \mathbb{R} \to \mathbb{R}$ is a function, then $Y = g(X)$ is again a random variable. For continuous random variables and invertible g, the density changes by the derivative of g, just as areas change by the Jacobian determinant in Lecture 3.

Proposition 2.5 (Change of variables)**.** *Let X have the density p_X, and let g be differentiable and strictly increasing (or strictly decreasing). Then $Y = g(X)$ has the density*

$$p_Y(y) = p_X(x)\,\frac{1}{|g'(x)|}, \qquad \text{where } x = g^{-1}(y).$$

Proof. For increasing g, $F_Y(y) = P(g(X) \le y) = P(X \le g^{-1}(y)) = F_X(g^{-1}(y))$. Differentiating with the chain rule gives $p_Y(y) = p_X(g^{-1}(y))\,(g^{-1})'(y) = \frac{p_X(x)}{g'(x)}$. For decreasing g, $F_Y(y) = 1 - F_X(g^{-1}(y))$, which gives the same formula with $|g'(x)|$. □

Example 2.6.

1. (Standardization.) If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Z = \frac{X-\mu}{\sigma}$ has the density $\sigma\, p_X(\mu + \sigma z) = \varphi(z)$, so $Z \sim \mathcal{N}(0, 1)$. Conversely, $X = \mu + \sigma Z$. Hence $P(|X - \mu| \le k\sigma) = P(|Z| \le k)$, which is 0.683, 0.954 and 0.997 for $k = 1, 2, 3$, and 0.95 for $k = 1.96$.
2. (Log-normal distribution.) If $X \sim \mathcal{N}(0, 1)$, then $Y = e^X$ has the density $\frac{\varphi(\log y)}{y}$ for $y > 0$.

Theorem 2.7 (Inverse transform sampling)**.** *Let F be a continuous, strictly increasing CDF, and let $U \sim$ Uniform$(0, 1)$. Then $X = F^{-1}(U)$ has the CDF F.*

Proof. $P(X \le x) = P(F^{-1}(U) \le x) = P(U \le F(x)) = F(x)$, since $P(U \le u) = u$ for $u \in [0, 1]$. □

So a computer that can generate uniform random numbers can generate random numbers with any such distribution. For example, the Exponential(λ) distribution has $F(x) = 1 - e^{-\lambda x}$, so $X = -\frac{1}{\lambda}\log(1 - U)$ is exponentially distributed. Normalizing flows (Lecture 3) generalize this idea to many dimensions: they transform simple random vectors by learned invertible maps.

3 Expectation and Variance

Definition 3.1 (Expectation)**.** The expectation (expected value, or mean) of a random variable X is

$$\mathbb{E}[X] = \sum_i x_i\, p(x_i) \qquad \text{or} \qquad \mathbb{E}[X] = \int_{-\infty}^{\infty} x\, p(x)\, dx,$$

for discrete and continuous X, respectively, provided that the sum or integral converges absolutely.

The expectation is the center of mass of the distribution: if we put the mass $p(x_i)$ at each point x_i, the balance point is $\mathbb{E}[X]$. To compute the expectation of a function of X, we do not need the distribution of $g(X)$; one can show that

$$\mathbb{E}[g(X)] = \sum_i g(x_i)\, p(x_i) \qquad \text{or} \qquad \mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)\, p(x)\, dx.$$

The expectation is linear: $\mathbb{E}[aX + bY + c] = a\mathbb{E}[X] + b\mathbb{E}[Y] + c$ for all random variables X, Y with finite expectations, and all numbers a, b, c (we prove this in Section 4 for discrete random variables).

Definition 3.2 (Variance)**.** The variance of X with mean $\mu = \mathbb{E}[X]$ is $\mathrm{Var}(X) = \mathbb{E}[(X-\mu)^2]$, and its standard deviation is $\sigma = \sqrt{\mathrm{Var}(X)}$.

The variance measures how far X typically is from its mean. Expanding the square and using linearity,

$$\mathrm{Var}(X) = \mathbb{E}[X^2 - 2\mu X + \mu^2] = \mathbb{E}[X^2] - 2\mu^2 + \mu^2 = \mathbb{E}[X^2] - \mathbb{E}[X]^2,$$

and $\mathrm{Var}(aX + b) = \mathbb{E}[(aX - a\mu)^2] = a^2 \mathrm{Var}(X)$: shifting does not change the spread, and scaling by a scales the standard deviation by $|a|$.

Proposition 3.3. *The distributions of Section 2 have the following means and variances:*

distribution	$\mathbb{E}[X]$	$\mathrm{Var}(X)$
Bernoulli(p)	p	$p(1-p)$
Binomial(n,p)	np	$np(1-p)$
Geometric(p)	$\frac{1}{p}$	$\frac{1-p}{p^2}$
Poisson(λ)	λ	λ
Uniform(a,b)	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
Exponential(λ)	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
$\mathcal{N}(\mu,\sigma^2)$	μ	σ^2

Proof of some cases. Bernoulli: $\mathbb{E}[X] = 0\cdot(1-p) + 1\cdot p = p$, and $X^2 = X$, so $\mathrm{Var}(X) = p - p^2$. Uniform: $\mathbb{E}[X] = \frac{1}{b-a}\int_a^b x\, dx = \frac{b^2-a^2}{2(b-a)} = \frac{a+b}{2}$, and similarly $\mathbb{E}[X^2] = \frac{a^2+ab+b^2}{3}$, so $\mathrm{Var}(X) = \frac{(b-a)^2}{12}$. Exponential: integrating by parts, $\mathbb{E}[X] = \int_0^\infty x\lambda e^{-\lambda x}\, dx = \big[-xe^{-\lambda x}\big]_0^\infty + \int_0^\infty e^{-\lambda x}\, dx = \frac{1}{\lambda}$. Poisson: $\mathbb{E}[X] = \sum_{k\ge1} k e^{-\lambda}\frac{\lambda^k}{k!} = \lambda \sum_{k\ge1} e^{-\lambda}\frac{\lambda^{k-1}}{(k-1)!} = \lambda$. Normal: by Example 2.6, $X = \mu + \sigma Z$ with $Z \sim \mathcal{N}(0,1)$, so it suffices to show $\mathbb{E}[Z] = 0$ and $\mathrm{Var}(Z) = 1$. The first holds since $z\varphi(z)$ is an odd function, and for the second, integration by parts with $\varphi'(z) = -z\varphi(z)$ gives

$$\mathbb{E}[Z^2] = \int_{-\infty}^{\infty} z\cdot z\varphi(z)\, dz = \big[-z\varphi(z)\big]_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \varphi(z)\, dz = 1.$$

The binomial case follows from the Bernoulli case and Proposition 4.6 below. □

Example 3.4. For a fair die, $\mathbb{E}[X] = \frac{1+2+\cdots+6}{6} = 3.5$, which is not a possible value, and $\mathbb{E}[X^2] = \frac{91}{6}$, so $\mathrm{Var}(X) = \frac{91}{6} - \frac{49}{4} = \frac{35}{12} \approx 2.92$.

3.1 Inequalities

The variance controls how likely large deviations from the mean are.

Proposition 3.5 (Markov and Chebyshev)**.** *1. (Markov's inequality.) If $X \geq 0$ and $a > 0$, then $P(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$.*

2. (Chebyshev's inequality.) If X has mean μ and variance σ^2, then $P(|X - \mu| \geq k) \leq \frac{\sigma^2}{k^2}$ for every $k > 0$.

Proof. (1) Since $X \geq 0$, we have $X \geq a\mathbf{1}_{X \geq a}$, where $\mathbf{1}_{X \geq a}$ is 1 if $X \geq a$ and 0 otherwise. Taking expectations, $\mathbb{E}[X] \geq aP(X \geq a)$. (2) Apply (1) to $(X - \mu)^2$ and $a = k^2$: $P(|X - \mu| \geq k) = P((X - \mu)^2 \geq k^2) \leq \frac{\sigma^2}{k^2}$. □

For example, every random variable is at least 2 standard deviations from its mean with probability at most $\frac{1}{4}$. This bound holds for all distributions, so it is weak for specific ones: for a normal distribution, the probability is only 0.046.

Theorem 3.6 (Jensen's inequality)**.** *If f is convex, then $f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)]$. If f is concave, then $f(\mathbb{E}[X]) \geq \mathbb{E}[f(X)]$.*

Proof. We prove this for differentiable f. Let $\mu = \mathbb{E}[X]$. By the first-order characterization of convexity in Lecture 3, $f(x) \geq f(\mu) + f'(\mu)(x - \mu)$ for all x. So $f(X) \geq f(\mu) + f'(\mu)(X - \mu)$, and taking expectations gives $\mathbb{E}[f(X)] \geq f(\mu) + f'(\mu) \cdot 0$. For concave f, apply this to $-f$. □

For example, $f(x) = x^2$ gives $\mathbb{E}[X]^2 \leq \mathbb{E}[X^2]$, i.e., $\operatorname{Var}(X) \geq 0$, and the concave function log gives $\mathbb{E}[\log X] \leq \log \mathbb{E}[X]$ for $X > 0$. We will use the latter to show that the Kullback–Leibler divergence is nonnegative (Section 13).

4 Joint Distributions and Independence

To describe how several random variables depend on each other, we need their joint distribution.

Definition 4.1 (Joint, marginal and conditional distributions)**.** The joint pmf of two discrete random variables X and Y is $p(x, y) = P(X = x, Y = y)$. Two continuous random variables have a joint density $p(x, y) \geq 0$ if $P((X, Y) \in A) = \iint_A p(x, y)\, dx\, dy$ for all (reasonable) sets $A \subseteq \mathbb{R}^2$. The marginal distributions of X and Y are

$$p_X(x) = \sum_y p(x, y) \qquad \text{or} \qquad p_X(x) = \int_{-\infty}^{\infty} p(x, y)\, dy,$$

and similarly for p_Y. The conditional distribution of Y given $X = x$ (for $p_X(x) > 0$) is

$$p(y \mid x) = \frac{p(x, y)}{p_X(x)}.$$

The formulas $p_X(x) = \sum_y p(x, y)$ (the sum rule) and $p(x, y) = p(y \mid x)\, p_X(x)$ (the product rule) are the two basic rules of probabilistic machine learning, and Bayes' theorem $p(x \mid y) = \frac{p(y|x)p_X(x)}{p_Y(y)}$ follows from them. The same definitions apply to more than two random variables.

Definition 4.2 (Independent random variables)**.** X and Y are independent if $p(x,y) = p_X(x)\,p_Y(y)$ for all x and y; equivalently, $p(y \mid x) = p_Y(y)$ whenever $p_X(x) > 0$. Random variables $X_1, \dots, X_n$ are independent if their joint pmf or density is the product of the marginal ones. They are independent and identically distributed (i.i.d.) if moreover they all have the same distribution.

The standard assumption in machine learning is that the training examples are i.i.d. samples from an unknown distribution: each example is drawn from the same distribution, independently of the others.

Example 4.3. Let $X, Y \in \{0,1\}$ have the joint pmf

	$Y=0$	$Y=1$	p_X
$X=0$	0.3	0.2	0.5
$X=1$	0.1	0.4	0.5
p_Y	0.4	0.6	

The marginals are the row and column sums. The conditional probabilities are $P(Y=1 \mid X=1) = \frac{0.4}{0.5} = 0.8$ and $P(Y=1 \mid X=0) = \frac{0.2}{0.5} = 0.4$. Since $p(1,1) = 0.4 \neq 0.5 \cdot 0.6 = p_X(1)p_Y(1)$, X and Y are not independent: learning X changes the probabilities of Y.

For a function $g(X,Y)$, the expectation is $\mathbb{E}[g(X,Y)] = \sum_{x,y} g(x,y)\,p(x,y)$, or the corresponding integral. In particular, linearity of expectation follows: for discrete X and Y,

$$\mathbb{E}[X+Y] = \sum_{x,y}(x+y)\,p(x,y) = \sum_x x \sum_y p(x,y) + \sum_y y \sum_x p(x,y) = \mathbb{E}[X] + \mathbb{E}[Y].$$

Proposition 4.4. *If X and Y are independent, then $\mathbb{E}[XY] = \mathbb{E}[X]\,\mathbb{E}[Y]$, and more generally $\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\,\mathbb{E}[h(Y)]$.*

Proof. For discrete random variables,

$$\mathbb{E}[g(X)h(Y)] = \sum_{x,y} g(x)h(y)p_X(x)p_Y(y) = \Big(\sum_x g(x)p_X(x)\Big)\Big(\sum_y h(y)p_Y(y)\Big);$$

for densities, the same computation works with integrals. □

4.1 Covariance and correlation

Definition 4.5 (Covariance and correlation)**.** The covariance of X and Y, with means μ_X and μ_Y, is

$$\operatorname{Cov}(X,Y) = \mathbb{E}[(X-\mu_X)(Y-\mu_Y)] = \mathbb{E}[XY] - \mathbb{E}[X]\,\mathbb{E}[Y],$$

and if both have positive variances, their correlation is $\operatorname{Corr}(X,Y) = \frac{\operatorname{Cov}(X,Y)}{\sigma_X \sigma_Y}$. If $\operatorname{Cov}(X,Y) = 0$, they are uncorrelated.

The covariance is positive if X and Y tend to be larger than their means at the same time, and negative if one tends to be large when the other is small. Note that $\operatorname{Cov}(X,X) = \operatorname{Var}(X)$, and the covariance is bilinear and symmetric. By Proposition 4.4, independent random variables are uncorrelated. The converse is false: if X is uniform on $\{-1,0,1\}$ and $Y = X^2$, then $\operatorname{Cov}(X,Y) = \mathbb{E}[X^3] - \mathbb{E}[X]\mathbb{E}[X^2] = 0$, but Y is a function of X. Correlation only measures linear dependence.

Proposition 4.6. $\operatorname{Var}(X+Y) = \operatorname{Var}(X) + \operatorname{Var}(Y) + 2\operatorname{Cov}(X,Y)$. *In particular, if* $X_1, \dots, X_n$ *are pairwise uncorrelated (for example, independent), then* $\operatorname{Var}(X_1 + \cdots + X_n) = \operatorname{Var}(X_1) + \cdots + \operatorname{Var}(X_n)$.

Proof. With $\tilde{X} = X - \mu_X$ and $\tilde{Y} = Y - \mu_Y$: $\operatorname{Var}(X+Y) = \mathbb{E}[(\tilde{X}+\tilde{Y})^2] = \mathbb{E}[\tilde{X}^2] + \mathbb{E}[\tilde{Y}^2] + 2\mathbb{E}[\tilde{X}\tilde{Y}]$. The second statement follows by induction. □

For example, a Binomial(n,p) random variable is the sum of n independent Bernoulli(p) random variables, so its mean is np and its variance is $np(1-p)$, as claimed in Proposition 3.3.

The covariance behaves like an inner product. For random variables U and V with finite variances, $\langle U, V\rangle = \mathbb{E}[UV]$ is bilinear, symmetric, and $\langle U, U\rangle = \mathbb{E}[U^2] \geq 0$, with equality only if $U = 0$ with probability 1. So, if we do not distinguish random variables that are equal with probability 1, it is an inner product as in Lecture 2. For centered random variables, the inner product is the covariance, the norm is the standard deviation, and the correlation is the cosine of the angle between them.

Theorem 4.7. $|\operatorname{Cov}(X,Y)| \leq \sigma_X \sigma_Y$, *i.e.,* $-1 \leq \operatorname{Corr}(X,Y) \leq 1$. *Moreover,* $|\operatorname{Corr}(X,Y)| = 1$ *if and only if* $Y = aX + b$ *with probability* 1, *for some* $a \neq 0$ *and* b; *the sign of the correlation is the sign of* a.

Proof. This is the Cauchy–Schwarz inequality for the centered random variables $\tilde{X}$ and $\tilde{Y}$. Directly: for every $t \in \mathbb{R}$,

$$0 \leq \mathbb{E}[(\tilde{Y} - t\tilde{X})^2] = \sigma_Y^2 - 2t\operatorname{Cov}(X,Y) + t^2\sigma_X^2.$$

A quadratic polynomial in t that is never negative has a discriminant ≤ 0, so $4\operatorname{Cov}(X,Y)^2 - 4\sigma_X^2\sigma_Y^2 \leq 0$. Equality holds if and only if the polynomial has a zero $t = a$, i.e., $\mathbb{E}[(\tilde{Y} - a\tilde{X})^2] = 0$, i.e., $\tilde{Y} = a\tilde{X}$ with probability 1; then $Y = aX + (\mu_Y - a\mu_X)$, and $\operatorname{Cov}(X,Y) = a\sigma_X^2$. □

Example 4.8. In Example 4.3, $\mathbb{E}[XY] = P(X=1, Y=1) = 0.4$, so $\operatorname{Cov}(X,Y) = 0.4 - 0.5 \cdot 0.6 = 0.1$. With $\operatorname{Var}(X) = 0.25$ and $\operatorname{Var}(Y) = 0.24$, the correlation is $\frac{0.1}{\sqrt{0.06}} \approx 0.41$.

4.2 Conditional expectation

Definition 4.9 (Conditional expectation)**.** The conditional expectation of Y given $X = x$ is the mean of the conditional distribution,

$$m(x) = \mathbb{E}[Y \mid X = x] = \sum_y y\, p(y \mid x) \qquad \text{or} \qquad \int_{-\infty}^{\infty} y\, p(y \mid x)\, dy.$$

The random variable $\mathbb{E}[Y \mid X] = m(X)$ is the conditional expectation of Y given X.

Theorem 4.10. *1. (Tower property.)* $\mathbb{E}\big[\mathbb{E}[Y \mid X]\big] = \mathbb{E}[Y]$.

2. (Best prediction.) For every function g, $\mathbb{E}[(Y - g(X))^2] \geq \mathbb{E}[(Y - m(X))^2]$, *with* $m(x) = \mathbb{E}[Y \mid X = x]$. *More precisely,*

$$\mathbb{E}[(Y - g(X))^2] = \mathbb{E}[(Y - m(X))^2] + \mathbb{E}[(m(X) - g(X))^2].$$

Proof. We give the proof for discrete random variables. (1) $\mathbb{E}[m(X)] = \sum_x p_X(x) \sum_y y\, p(y \mid x) = \sum_{x,y} y\, p(x,y) = \mathbb{E}[Y]$. (2) For every function h, the error $Y - m(X)$ is orthogonal to $h(X)$:

$$\mathbb{E}[(Y - m(X))h(X)] = \sum_x p_X(x)h(x) \sum_y (y - m(x))p(y \mid x) = \sum_x p_X(x)h(x)\big(m(x) - m(x)\big) = 0.$$

Now write $Y - g(X) = (Y - m(X)) + (m(X) - g(X))$, expand the square, and use this with $h = m - g$; the mixed term vanishes, as in the theorem of Pythagoras. □

So $\mathbb{E}[Y \mid X]$ is the best prediction of Y from X in the sense of the mean squared error: it is the orthogonal projection of Y onto the space of functions of X, with respect to the inner product $\mathbb{E}[UV]$ (compare with orthogonal projections in Lecture 2). A regression model trained with the squared loss on enough data therefore estimates the conditional mean $\mathbb{E}[Y \mid X = \mathbf{x}]$. In Example 4.3, $\mathbb{E}[Y \mid X = 0] = 0.4$ and $\mathbb{E}[Y \mid X = 1] = 0.8$, and indeed $0.5 \cdot 0.4 + 0.5 \cdot 0.8 = 0.6 = \mathbb{E}[Y]$.

5 Random Vectors and Covariance Matrices

A random vector $\mathbf{X} = (X_1, \dots, X_d)^\top$ is a vector of random variables, such as the features of a randomly chosen data point. Its expectation is taken entrywise.

Definition 5.1 (Mean and covariance matrix)**.** The mean of a random vector $\mathbf{X}$ is $\boldsymbol{\mu} = \mathbb{E}[\mathbf{X}] = (\mathbb{E}[X_1], \dots, \mathbb{E}[X_d])^\top$, and its covariance matrix is the $d \times d$ matrix

$$\Sigma = \mathrm{Cov}(\mathbf{X}) = \mathbb{E}\big[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top\big], \qquad \text{with entries } \Sigma_{ij} = \mathrm{Cov}(X_i, X_j).$$

The diagonal entries of Σ are the variances $\mathrm{Var}(X_i)$. Since $(\mathbf{X}-\boldsymbol{\mu})(\mathbf{X}-\boldsymbol{\mu})^\top$ is an outer product, linear algebra enters immediately.

Proposition 5.2. *Let $\mathbf{X}$ have mean $\boldsymbol{\mu}$ and covariance matrix Σ.*

1. $\Sigma = \mathbb{E}[\mathbf{X}\mathbf{X}^\top] - \boldsymbol{\mu}\boldsymbol{\mu}^\top$.
2. *For every matrix $A \in \mathbb{R}^{m \times d}$ and vector $\mathbf{b} \in \mathbb{R}^m$, $\mathbb{E}[A\mathbf{X} + \mathbf{b}] = A\boldsymbol{\mu} + \mathbf{b}$ and $\mathrm{Cov}(A\mathbf{X} + \mathbf{b}) = A\Sigma A^\top$.*
3. *For every $\mathbf{u} \in \mathbb{R}^d$, $\mathrm{Var}(\mathbf{u}^\top \mathbf{X}) = \mathbf{u}^\top \Sigma \mathbf{u}$. In particular, Σ is symmetric and positive semidefinite.*

Proof. (2) Each entry of $A\mathbf{X} + \mathbf{b}$ is a linear combination of the X_j plus a constant, so linearity of expectation gives $\mathbb{E}[A\mathbf{X} + \mathbf{b}] = A\boldsymbol{\mu} + \mathbf{b}$, and in the same way $\mathbb{E}[AMB] = A\,\mathbb{E}[M]\,B$ for a random matrix M and fixed matrices A and B. Hence

$$\mathrm{Cov}(A\mathbf{X} + \mathbf{b}) = \mathbb{E}\big[A(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top A^\top\big] = A\Sigma A^\top.$$

(1) is proved like the formula $\mathrm{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$. (3) is (2) with $A = \mathbf{u}^\top$, and $\mathrm{Var}(\mathbf{u}^\top \mathbf{X}) \geq 0$; Σ is symmetric since $\mathrm{Cov}(X_i, X_j) = \mathrm{Cov}(X_j, X_i)$. □

By (3), Σ fails to be positive definite exactly if $\mathrm{Var}(\mathbf{u}^\top \mathbf{X}) = 0$ for some $\mathbf{u} \neq \mathbf{0}$, i.e., if $\mathbf{u}^\top \mathbf{X}$ is constant: then $\mathbf{X}$ lies in an affine hyperplane (with probability 1). Part (3) also describes the variance in every direction: for a unit vector $\mathbf{u}$, the variance of the projection $\mathbf{u}^\top \mathbf{X}$ is $\mathbf{u}^\top \Sigma \mathbf{u}$. By the Rayleigh quotient (Lecture 2), it is largest for an eigenvector of Σ with the largest eigenvalue. This is the population version of principal component analysis, and the total variance $\sum_i \mathrm{Var}(X_i) = \mathrm{tr}\,\Sigma$ is the sum of the eigenvalues of Σ.

Example 5.3. Let $\Sigma = \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix}$. Then $\mathrm{Var}(X_1 + X_2) = (1,1)\Sigma(1,1)^\top = 4 + 3 + 2 \cdot 2 = 11$, $\mathrm{Var}(X_1 - X_2) = 4 + 3 - 4 = 3$, and $\mathrm{Corr}(X_1, X_2) = \frac{2}{\sqrt{12}} \approx 0.58$. The eigenvalues of Σ are $\frac{7 \pm \sqrt{17}}{2} \approx 5.56$ and 1.44: the variance in the direction of the first principal component is 5.56, out of the total variance $\mathrm{tr}\,\Sigma = 7$.

Definition 5.4 (Whitening and Mahalanobis distance)**.** Let Σ be symmetric positive definite, with spectral decomposition $\Sigma = Q\Lambda Q^\top$ (Lecture 2), and let $\Sigma^{-1/2} = Q\Lambda^{-1/2}Q^\top$, where $\Lambda^{-1/2} = \operatorname{diag}(\lambda_1^{-1/2}, \dots, \lambda_d^{-1/2})$. The whitened random vector is $\mathbf{Z} = \Sigma^{-1/2}(\mathbf{X} - \boldsymbol{\mu})$, and the Mahalanobis distance of $\mathbf{x}$ from $\boldsymbol{\mu}$ is

$$d_\Sigma(\mathbf{x}, \boldsymbol{\mu}) = \sqrt{(\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})} = \|\Sigma^{-1/2}(\mathbf{x} - \boldsymbol{\mu})\|.$$

The matrix $\Sigma^{-1/2}$ is symmetric with $(\Sigma^{-1/2})^2 = \Sigma^{-1}$, so by Proposition 5.2, $\operatorname{Cov}(\mathbf{Z}) = \Sigma^{-1/2}\Sigma\Sigma^{-1/2} = I$: whitening removes all correlations and scales all directions to unit variance. The Mahalanobis distance is the norm of the inner product $\langle \mathbf{x}, \mathbf{y} \rangle_{\Sigma^{-1}} = \mathbf{x}^\top \Sigma^{-1} \mathbf{y}$ (Lecture 2): it measures distances in units of standard deviations along each principal axis. Standardizing or whitening the features of a data set is a common preprocessing step; by Lecture 3, it improves the condition number and speeds up gradient descent.

Sample statistics. Given data points $\mathbf{x}_1, \dots, \mathbf{x}_N \in \mathbb{R}^d$, the sample mean and the sample covariance matrix are

$$\bar{\mathbf{x}} = \frac{1}{N} \sum_{i=1}^{N} \mathbf{x}_i \qquad \text{and} \qquad S = \frac{1}{N} \sum_{i=1}^{N} (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top = \frac{1}{N} X_c^\top X_c,$$

where X_c is the centered data matrix with rows $(\mathbf{x}_i - \bar{\mathbf{x}})^\top$. These are the mean and covariance matrix of the empirical distribution, which gives probability $\frac{1}{N}$ to each data point; principal component analysis in Lecture 2 used exactly this matrix. In Section 9, we will see why one often divides by $N - 1$ instead of N.

6 The Gaussian Distribution

The Gaussian (normal) distribution is the most important distribution in statistics and machine learning. It appears as the limit of sums of many small random effects (Section 7), it is completely described by its mean and covariance matrix, and its marginals and conditionals can be computed with linear algebra.

Definition 6.1 (Multivariate Gaussian distribution)**.** Let $\boldsymbol{\mu} \in \mathbb{R}^d$, and let Σ be a symmetric positive definite $d \times d$ matrix. A random vector $\mathbf{X}$ has the Gaussian distribution $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ if it has the density

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} \det(\Sigma)^{1/2}} \exp\Big(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \Big).$$

For $d = 1$ and $\Sigma = \sigma^2$, this is the density of Section 2. For $\boldsymbol{\mu} = \mathbf{0}$ and $\Sigma = I$, the density is $(2\pi)^{-d/2} e^{-\|\mathbf{z}\|^2/2} = \prod_i \varphi(z_i)$: a standard Gaussian vector $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$ has i.i.d. standard normal entries. All other Gaussians are obtained from it by affine maps.

Theorem 6.2. *Let $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$, let A be an invertible $d \times d$ matrix, and let $\boldsymbol{\mu} \in \mathbb{R}^d$. Then $\mathbf{X} = A\mathbf{Z} + \boldsymbol{\mu} \sim \mathcal{N}(\boldsymbol{\mu}, AA^\top)$. Consequently, the density of Definition 6.1 integrates to 1, and $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ has mean $\boldsymbol{\mu}$ and covariance matrix Σ.*

Proof. By the change of variables formula (Lecture 3), with $\mathbf{z} = A^{-1}(\mathbf{x} - \boldsymbol{\mu})$, whose Jacobian has the determinant $\det(A)^{-1}$,

$$p_{\mathbf{X}}(\mathbf{x}) = p_{\mathbf{Z}}\big(A^{-1}(\mathbf{x} - \boldsymbol{\mu})\big) \, \frac{1}{|\det A|} = \frac{1}{(2\pi)^{d/2} |\det A|} \exp\Big(-\frac{1}{2} \|A^{-1}(\mathbf{x} - \boldsymbol{\mu})\|^2 \Big).$$

With $\Sigma = AA^\top$, we have $\|A^{-1}\mathbf{y}\|^2 = \mathbf{y}^\top A^{-\top}A^{-1}\mathbf{y} = \mathbf{y}^\top\Sigma^{-1}\mathbf{y}$ and $\det\Sigma = \det(A)^2$, which gives the density of $\mathcal{N}(\boldsymbol{\mu},\Sigma)$. Every symmetric positive definite Σ can be written as $AA^\top$, for example with $A = \Sigma^{1/2} = Q\Lambda^{1/2}Q^\top$. So every Gaussian density is the density of some random vector, and integrates to 1. By Proposition 5.2, $\mathbb{E}[\mathbf{X}] = A\mathbb{E}[\mathbf{Z}] + \boldsymbol{\mu} = \boldsymbol{\mu}$ and $\mathrm{Cov}(\mathbf{X}) = A\,I\,A^\top = \Sigma$. □

This is also how Gaussian vectors are sampled: compute a factorization $\Sigma = AA^\top$ (in practice, the Cholesky factorization $\Sigma = LL^\top$ with a lower triangular L, as in `np.linalg.cholesky`), draw $\mathbf{z}$ with i.i.d. standard normal entries, and return $\boldsymbol{\mu} + L\mathbf{z}$. If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu},\Sigma)$ and B is invertible, then $B\mathbf{X} + \mathbf{b} = (BA)\mathbf{Z} + (B\boldsymbol{\mu} + \mathbf{b})$, so $B\mathbf{X} + \mathbf{b} \sim \mathcal{N}(B\boldsymbol{\mu} + \mathbf{b}, B\Sigma B^\top)$.

Remark 6.3 (The reparameterization trick). Writing a Gaussian random vector as $\boldsymbol{\mu} + \boldsymbol{\sigma} \odot \mathbf{Z}$ with a fixed source of randomness $\mathbf{Z}$ makes it a differentiable function of the parameters $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}$. Variational autoencoders use this to backpropagate (Lecture 3) through random sampling steps.

The density of $\mathcal{N}(\boldsymbol{\mu},\Sigma)$ is constant on the ellipsoids $(\mathbf{x}-\boldsymbol{\mu})^\top\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu}) = r^2$, i.e., on the sets of points with Mahalanobis distance r from $\boldsymbol{\mu}$. By the spectral theorem (Lecture 2), their axes point in the directions of the eigenvectors $\mathbf{q}_i$ of Σ, with half-lengths $r\sqrt{\lambda_i}$, i.e., r standard deviations in the direction $\mathbf{q}_i$ (Figure 2).

Theorem 6.4 (Independence of blocks)**.** *Let* $\mathbf{X} = (\mathbf{X}_a, \mathbf{X}_b) \sim \mathcal{N}(\boldsymbol{\mu},\Sigma)$*, and let* $\Sigma = \left(\begin{smallmatrix}\Sigma_{aa} & 0\\ 0 & \Sigma_{bb}\end{smallmatrix}\right)$ *be block diagonal. Then* $\mathbf{X}_a$ *and* $\mathbf{X}_b$ *are independent, with* $\mathbf{X}_a \sim \mathcal{N}(\boldsymbol{\mu}_a,\Sigma_{aa})$ *and* $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b,\Sigma_{bb})$*.*

Proof. Σ^{-1} is block diagonal with blocks Σ_{aa}^{-1} and Σ_{bb}^{-1}, and $\det\Sigma = \det\Sigma_{aa}\det\Sigma_{bb}$. Hence the quadratic form in the exponent splits into a sum of two terms, and the density is the product of the densities of $\mathcal{N}(\boldsymbol{\mu}_a,\Sigma_{aa})$ and $\mathcal{N}(\boldsymbol{\mu}_b,\Sigma_{bb})$. Integrating over $\mathbf{x}_b$ shows that the first factor is the marginal density of $\mathbf{X}_a$, and similarly for $\mathbf{X}_b$. □

So for jointly Gaussian random vectors, uncorrelated parts are independent. This is special to Gaussians, and it is the key to the following theorem, which is used constantly in Gaussian processes, Kalman filters and Bayesian linear regression.

Theorem 6.5 (Marginals and conditionals)**.** *Let* $\mathbf{X} = (\mathbf{X}_a, \mathbf{X}_b) \sim \mathcal{N}(\boldsymbol{\mu},\Sigma)$ *with*

$$\boldsymbol{\mu} = \begin{pmatrix}\boldsymbol{\mu}_a\\ \boldsymbol{\mu}_b\end{pmatrix}, \qquad \Sigma = \begin{pmatrix}\Sigma_{aa} & \Sigma_{ab}\\ \Sigma_{ba} & \Sigma_{bb}\end{pmatrix}.$$

Then $\mathbf{X}_a \sim \mathcal{N}(\boldsymbol{\mu}_a,\Sigma_{aa})$*,* $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b,\Sigma_{bb})$*, and the conditional distribution of* $\mathbf{X}_a$ *given* $\mathbf{X}_b = \mathbf{x}_b$ *is*

$$\mathcal{N}(\boldsymbol{\mu}_a + \Sigma_{ab}\Sigma_{bb}^{-1}(\mathbf{x}_b - \boldsymbol{\mu}_b),\ \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}).$$

Proof. The block Σ_{bb} is positive definite (since $\mathbf{u}^\top\Sigma_{bb}\mathbf{u} = (\mathbf{0},\mathbf{u})^\top\Sigma(\mathbf{0},\mathbf{u}) > 0$ for $\mathbf{u} \neq \mathbf{0}$), so $K = \Sigma_{ab}\Sigma_{bb}^{-1}$ is defined. Let $\mathbf{W} = \mathbf{X}_a - K\mathbf{X}_b$, i.e.,

$$\begin{pmatrix}\mathbf{W}\\ \mathbf{X}_b\end{pmatrix} = B\mathbf{X}, \qquad B = \begin{pmatrix}I & -K\\ 0 & I\end{pmatrix}.$$

B is invertible, so $(\mathbf{W},\mathbf{X}_b)$ is Gaussian with covariance matrix $B\Sigma B^\top$. Its off-diagonal block is $\mathrm{Cov}(\mathbf{W},\mathbf{X}_b) = \Sigma_{ab} - K\Sigma_{bb} = 0$, and its upper left block is

$$\mathrm{Cov}(\mathbf{W}) = \Sigma_{aa} - K\Sigma_{ba} - \Sigma_{ab}K^\top + K\Sigma_{bb}K^\top = \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}.$$

By Theorem 6.4, $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b,\Sigma_{bb})$, and $\mathbf{W}$ is independent of $\mathbf{X}_b$, with $\mathbf{W} \sim \mathcal{N}(\boldsymbol{\mu}_a - K\boldsymbol{\mu}_b, \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba})$. Since $\det B = 1$, the joint density of $\mathbf{X}$ is $p(\mathbf{x}_a,\mathbf{x}_b) = p_{\mathbf{W}}(\mathbf{x}_a - K\mathbf{x}_b)\,p_{\mathbf{X}_b}(\mathbf{x}_b)$, so the conditional density is $p(\mathbf{x}_a \mid \mathbf{x}_b) = p_{\mathbf{W}}(\mathbf{x}_a - K\mathbf{x}_b)$: given $\mathbf{X}_b = \mathbf{x}_b$, $\mathbf{X}_a = \mathbf{W} + K\mathbf{x}_b$, which has the claimed distribution. The marginal of $\mathbf{X}_a$ follows by exchanging the roles of a and b. □

The conditional mean is an affine function of $\mathbf{x}_b$, and the conditional covariance $\Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}$ (the Schur complement of Σ_{bb}) does not depend on $\mathbf{x}_b$. It is never larger than Σ_{aa}, since $\Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}$ is positive semidefinite: observing $\mathbf{X}_b$ reduces the uncertainty about $\mathbf{X}_a$.

Example 6.6 (Regression to the mean)**.** Let (X, Y) be Gaussian with means 0, variances 1, and correlation ρ. By Theorem 6.5, given $X = x$, $Y \sim \mathcal{N}(\rho x, 1 - \rho^2)$. For example, for $\rho = 0.5$ and $x = 2$, $Y \sim \mathcal{N}(1, 0.75)$. If X and Y are the heights of parents and their children (measured in standard deviations from the mean), then children of very tall parents are on average tall, but less extreme than their parents. Francis Galton observed this "regression to the mean" in 1886, and it gave linear regression its name. Note that the regression line $y = \rho x$ is not the major axis of the ellipses of constant density, which is the line $y = x$ (Figure 2).




Figure 2: Level curves of the density of $\mathcal{N}(\mathbf{0}, \Sigma)$ with $\Sigma = \left(\begin{smallmatrix} 3 & 1 \\ 1 & 3 \end{smallmatrix}\right)$, for Mahalanobis distances $r = 0.5, 1, 1.5, 2$. The eigenvectors $\mathbf{q}_1 = \frac{1}{\sqrt{2}}(1, 1)$ and $\mathbf{q}_2 = \frac{1}{\sqrt{2}}(1, -1)$ of Σ have the eigenvalues 4 and 2, so the axes have half-lengths $2r$ and $\sqrt{2}\,r$ (orange: one standard deviation in each direction). The conditional mean $\mathbb{E}[X_2 \mid X_1 = x_1] = \frac{\Sigma_{21}}{\Sigma_{11}}x_1 = \frac{1}{3}x_1$ (red) is flatter than the major axis $x_2 = x_1$ (dashed).

Corollary 6.7. *Let $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$, and let $A \in \mathbb{R}^{m \times d}$ have rank m. Then $A\mathbf{X} + \mathbf{b} \sim \mathcal{N}(A\boldsymbol{\mu} + \mathbf{b}, A\Sigma A^\top)$. In particular, if $X \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$ are independent, then $X + Y \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.*

Proof. The rows of A are linearly independent, so they can be extended to a basis of $\mathbb{R}^d$ (Lecture 1): there is a matrix C such that $B = \left(\begin{smallmatrix} A \\ C \end{smallmatrix}\right)$ is invertible. Then $B\mathbf{X} + (\mathbf{b}, \mathbf{0})$ is Gaussian, and $A\mathbf{X} + \mathbf{b}$ is its first block, which is Gaussian by Theorem 6.5, with the mean and covariance of Proposition 5.2. For the second statement, (X, Y) is Gaussian with the covariance matrix $\operatorname{diag}(\sigma_1^2, \sigma_2^2)$ (its density is the product of the two densities), and $X + Y = (1, 1)(X, Y)^\top$. □

7 Limit Theorems and Monte Carlo Methods

Machine learning relies on averages: the training loss is an average over the data, a mini-batch gradient is an average over a few examples, and the test accuracy is an average over the test set. The limit theorems explain why averages of many random quantities are reliable.

Proposition 7.1. *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and variance σ^2, and let $\bar{X}_N = \frac{1}{N}(X_1 + \cdots + X_N)$ be their sample mean. Then*

$$\mathbb{E}[\bar{X}_N] = \mu \qquad \textit{and} \qquad \operatorname{Var}(\bar{X}_N) = \frac{\sigma^2}{N}.$$

Proof. By linearity, $\mathbb{E}[\bar{X}_N] = \frac{1}{N} \cdot N\mu$, and by Proposition 4.6, $\operatorname{Var}(\bar{X}_N) = \frac{1}{N^2} \cdot N\sigma^2$. □

So the standard deviation of the sample mean, its standard error, is $\frac{\sigma}{\sqrt{N}}$: to halve the error, we need four times as much data.

Theorem 7.2 (Weak law of large numbers)**.** *In the situation of Proposition 7.1, for every $\varepsilon > 0$,*

$$P(|\bar{X}_N - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{N\varepsilon^2} \to 0 \qquad \textit{as } N \to \infty.$$

Proof. Chebyshev's inequality (Proposition 3.5) for $\bar{X}_N$, which has mean μ and variance $\frac{\sigma^2}{N}$. □

The law of large numbers says that the sample mean is very likely close to μ for large N. (The strong law of large numbers, which we do not prove, even says that $\bar{X}_N \to \mu$ with probability 1.) The central limit theorem describes the size and shape of the remaining fluctuations: they are approximately Gaussian, whatever the distribution of the X_i.

Theorem 7.3 (Central limit theorem)**.** *In the situation of Proposition 7.1, with $\sigma > 0$, for every $z \in \mathbb{R}$,*

$$P\Big(\frac{\sqrt{N}(\bar{X}_N - \mu)}{\sigma} \leq z\Big) \to \Phi(z) \qquad \textit{as } N \to \infty.$$

We do not prove the central limit theorem. Informally, $\bar{X}_N$ is approximately $\mathcal{N}(\mu, \frac{\sigma^2}{N})$, and the sum $X_1 + \cdots + X_N$ is approximately $\mathcal{N}(N\mu, N\sigma^2)$. This explains why Gaussian distributions are so common: a measurement error that is the sum of many small independent effects is approximately Gaussian.

Example 7.4. We flip a fair coin 100 times. The number of heads S is Binomial$(100, \frac{1}{2})$, with mean 50 and standard deviation $\sqrt{100 \cdot \frac{1}{4}} = 5$. By the central limit theorem, $P(S \geq 60) = P(S \geq 59.5) \approx 1 - \Phi\big(\frac{59.5-50}{5}\big) = 1 - \Phi(1.9) \approx 0.0287$, while the exact value is 0.0284. (Replacing 60 by 59.5, the "continuity correction", improves the approximation of a discrete distribution by a continuous one.) Figure 3 compares a binomial distribution with its Gaussian approximation.

Monte Carlo estimation. To compute an expectation $\mathbb{E}[g(X)]$ that we cannot compute exactly, we can draw i.i.d. samples $x_1, \dots, x_N$ of X and use the average $\frac{1}{N}\sum_i g(x_i)$. By the central limit theorem, its error is approximately Gaussian with standard deviation $\frac{\sigma_g}{\sqrt{N}}$, where σ_g is the standard deviation of $g(X)$, no matter how high-dimensional X is. For example, the fraction of N uniformly random points of the square $[-1, 1]^2$ that land in the unit disk estimates $\frac{\pi}{4}$, so 4 times this fraction estimates π, with standard error $4\sqrt{\frac{\pi}{4}(1 - \frac{\pi}{4})/N}$, which is 0.016 for $N = 10\,000$. Each additional correct digit costs 100 times more samples.




Figure 3: The pmf of the Binomial$(40, 0.3)$ distribution, the sum of 40 independent Bernoulli(0.3) variables, and the density of its Gaussian approximation $\mathcal{N}(12, 8.4)$ with the same mean and variance (red).

Concentration inequalities. The central limit theorem is only an approximation for finite N. For bounded random variables, there is an exact bound that is much stronger than Chebyshev's inequality.

Theorem 7.5 (Hoeffding's inequality)**.** *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and values in $[a, b]$. Then for every $\varepsilon > 0$,*

$$P(|\bar{X}_N - \mu| \geq \varepsilon) \leq 2 \exp\Big(- \frac{2N\varepsilon^2}{(b-a)^2} \Big).$$

We do not prove this theorem. For example, let a classifier have the (unknown) error rate μ, and let $\bar{X}_N$ be its error rate on a test set of $N = 1000$ examples drawn independently of the training data, where $X_i \in \{0, 1\}$ indicates an error. Then $P(|\bar{X}_N - \mu| \geq 0.05) \leq 2e^{-5} \approx 0.013$, while Chebyshev's inequality with $\sigma^2 \leq \frac{1}{4}$ only gives 0.1.

Remark 7.6 (Limit theorems in machine learning).

1. The average loss on the training data, $\frac{1}{N} \sum_i \ell(\boldsymbol{\theta}; \mathbf{x}_i, y_i)$, is a sample mean. It estimates the expected loss $\mathbb{E}[\ell(\boldsymbol{\theta}; \mathbf{X}, Y)]$ on new data, which is the quantity we actually care about. For a fixed $\boldsymbol{\theta}$, the limit theorems apply. But the trained parameters depend on the training data, so the training loss of the trained model is a biased, usually too optimistic, estimate (overfitting). This is why we evaluate on separate test data, and why learning theory combines Hoeffding's inequality with the union bound (Proposition 1.3) to control all candidate models simultaneously.

2. In stochastic gradient descent (Lecture 3), the mini-batch gradient of B randomly chosen examples is an unbiased estimate of the full gradient, and its covariance matrix decreases like $\frac{1}{B}$ (for sampling with replacement), by Proposition 7.1 applied to each coordinate. Larger batches give less noisy gradients, but each step costs more.

7.1 Monte Carlo sampling

Monte Carlo estimation needs samples from the distribution of interest. We have already seen inverse transform sampling (Theorem 2.7) and sampling Gaussian vectors by affine maps (Section 6), and we will see Markov chain Monte Carlo in Section 8. Two further general methods are rejection sampling, which produces exact samples, and importance sampling, which reweights samples from a different distribution.

Theorem 7.7 (Rejection sampling)**.** *Let p and q be densities, and let $M \geq 1$ be such that $p(x) \leq Mq(x)$ for all x. Repeat the following trial until a value is accepted: draw $X \sim q$ and, independently, $U \sim$ Uniform$(0,1)$, and accept X if $U \leq \frac{p(X)}{Mq(X)}$. Then the accepted value has the density p, and each trial is accepted with probability $\frac{1}{M}$.*

Proof. For every set A, $P(X \in A \text{ and accepted}) = \int_A q(x)\,\frac{p(x)}{Mq(x)}\,dx = \frac{1}{M}\int_A p(x)\,dx$. For $A = \mathbb{R}$, this gives $P(\text{accepted}) = \frac{1}{M}$, so $P(X \in A \mid \text{accepted}) = \int_A p(x)\,dx$. Since the trials are independent, the accepted value has this conditional distribution. □

Geometrically, the points $(X, UMq(X))$ are uniformly distributed under the graph of Mq, and we keep those that lie under the graph of p; their first coordinates have the density p. The method also works if p is only known up to a constant. For example, the Beta$(2,2)$ density $6x(1-x)$ is at most 1.5 on $[0,1]$, so with the uniform proposal and $M = 1.5$, two out of three trials are accepted. In high dimensions, however, M is usually enormous: for the target $\mathcal{N}(\mathbf{0}, I)$ in $\mathbb{R}^d$ and the proposal $\mathcal{N}(\mathbf{0}, 2I)$, the ratio $\frac{p}{q} = 2^{d/2}e^{-\|\mathbf{x}\|^2/4}$ forces $M \geq 2^{d/2}$, so only one in 1024 trials is accepted for $d = 20$. This is one reason why high-dimensional sampling relies on MCMC.

Proposition 7.8 (Importance sampling)**.** *Let p and q be densities such that $q(x) > 0$ for all x with $f(x)p(x) \neq 0$, and let $w = \frac{p}{q}$ (the importance weights). Then $\mathbb{E}_p[f(X)] = \mathbb{E}_q[f(Y)w(Y)]$. Hence, for i.i.d. samples $y_1, \ldots, y_N$ from q, the estimator*

$$\hat{I} = \frac{1}{N}\sum_{i=1}^{N} f(y_i)w(y_i)$$

is unbiased, with variance $\frac{1}{N}\operatorname{Var}_q(f(Y)w(Y))$.

Proof. $\int f(x)p(x)\,dx = \int f(y)\frac{p(y)}{q(y)}q(y)\,dy$; the variance follows from Proposition 7.1. □

The variance is small if q puts its mass where $|f|p$ is large; for $f \geq 0$, the ideal (but impractical) choice $q \propto fp$ even gives variance 0.

Example 7.9 (A rare event)**.** The tail probability $P(Z > 4) = 3.17 \cdot 10^{-5}$ of $Z \sim \mathcal{N}(0,1)$ is hard to estimate by plain Monte Carlo: among $N = 10\,000$ samples, we expect only 0.32 samples above 4, and the relative standard error of the estimate is about 178%. With the proposal $q = \mathcal{N}(4,1)$, about half of the samples land above 4, and they are reweighted by $w(y) = \frac{\varphi(y)}{\varphi(y-4)} = e^{8-4y}$. With the same N, the relative standard error drops to about 2%. Probabilities of rare failures are estimated in this way.

If p is only known up to a constant, one uses the self-normalized estimator $\sum_i f(y_i)w_i / \sum_i w_i$, which is slightly biased but consistent. If a few weights dominate, the estimate is unreliable; the effective sample size $\frac{(\sum_i w_i)^2}{\sum_i w_i^2}$ measures how many samples effectively contribute. In machine learning, importance weights correct for differences between distributions: in off-policy evaluation in reinforcement learning (data collected with one policy, used to evaluate another), and under covariate shift, where training examples are weighted by the ratio of the test and training densities.

8 Markov Chains

So far, we mostly considered independent random variables. Many random sequences have memory: tomorrow's weather depends on today's, the next word of a sentence depends on the previous ones,

and a user browsing the web chooses the next page depending on the current one. The simplest model with memory is a Markov chain, in which the future depends on the past only through the present.

Definition 8.1 (Markov chain)**.** A sequence of random variables $X_0, X_1, X_2, \ldots$ with values in a finite set of states $\{1, \ldots, n\}$ is a Markov chain with transition matrix $P \in \mathbb{R}^{n \times n}$ if

$$P(X_{t+1} = j \mid X_t = i, X_{t-1} = i_{t-1}, \ldots, X_0 = i_0) = P_{ij}$$

for all times t and all states $i, j, i_0, \ldots, i_{t-1}$ (whenever the condition has positive probability).

The entry P_{ij} is the probability of moving from state i to state j in one step. So P has nonnegative entries, and each row sums to 1, i.e., $P\mathbf{1} = \mathbf{1}$; such matrices are called stochastic matrices.

Example 8.2 (Weather)**.** Let state 1 be "sunny" and state 2 "rainy". After a sunny day, the next day is sunny with probability 0.9; after a rainy day, it is sunny with probability 0.5:

$$P = \begin{pmatrix} 0.9 & 0.1 \\ 0.5 & 0.5 \end{pmatrix}.$$

The chain is drawn as a graph with an arrow $i \to j$ labeled P_{ij} for each possible transition (Figure 4).

We write the distribution of X_t as a row vector $\boldsymbol{\pi}_t = \big(P(X_t = 1), \ldots, P(X_t = n)\big)$. (The same weather model appears in the section on eigenvalues in Lecture 1, with the distributions written as columns and multiplied by the transposed matrix $P^\top$.)

Proposition 8.3. *$\boldsymbol{\pi}_{t+1} = \boldsymbol{\pi}_t P$, hence $\boldsymbol{\pi}_t = \boldsymbol{\pi}_0 P^t$. More generally, $P(X_{t+k} = j \mid X_t = i) = (P^k)_{ij}$.*

Proof. By the law of total probability (Theorem 1.6), $P(X_{t+1} = j) = \sum_i P(X_t = i)\, P(X_{t+1} = j \mid X_t = i) = \sum_i (\boldsymbol{\pi}_t)_i P_{ij}$. The second statement follows by induction on k, in the same way. □

So the distribution evolves by repeated multiplication with P, and the long-term behavior of the chain is a question about the powers of a matrix, i.e., about its eigenvalues (Lecture 1). For the weather chain, starting on a sunny day, $\boldsymbol{\pi}_0 = (1, 0)$, $\boldsymbol{\pi}_1 = (0.9, 0.1)$, $\boldsymbol{\pi}_2 = (0.86, 0.14)$, $\boldsymbol{\pi}_3 = (0.844, 0.156)$, and the distribution seems to approach a limit.

Definition 8.4 (Stationary distribution)**.** A probability distribution $\boldsymbol{\pi}$ (a row vector with $\pi_i \geq 0$ and $\sum_i \pi_i = 1$) is stationary for P if $\boldsymbol{\pi} P = \boldsymbol{\pi}$.

If X_0 has a stationary distribution, then so does every X_t. Transposing, $P^\top \boldsymbol{\pi}^\top = \boldsymbol{\pi}^\top$: a stationary distribution is an eigenvector of $P^\top$ with the eigenvalue 1 (a left eigenvector of P), scaled to have entries that sum to 1.

Proposition 8.5. *Let P be a stochastic matrix. Then 1 is an eigenvalue of P and of $P^\top$, and every (complex) eigenvalue λ of P satisfies $|\lambda| \leq 1$.*

Proof. $P\mathbf{1} = \mathbf{1}$, and $P^\top$ has the same characteristic polynomial as P, since $\det(A^\top) = \det(A)$ (Lecture 1). If $P\mathbf{v} = \lambda \mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, choose i with $|v_i| = \max_j |v_j|$. Then $|\lambda||v_i| = |\sum_j P_{ij} v_j| \leq \sum_j P_{ij}|v_j| \leq \sum_j P_{ij}|v_i| = |v_i|$. □

Whether the distribution converges to a stationary distribution depends on two properties. A chain is irreducible if every state can be reached from every other state, i.e., for all i and j there is a t with $(P^t)_{ij} > 0$. It is aperiodic if for every state i, the numbers $t \geq 1$ with $(P^t)_{ii} > 0$ have greatest common divisor 1; for an irreducible chain, it suffices that some $P_{ii} > 0$.

Theorem 8.6 (Convergence of Markov chains)**.** *An irreducible Markov chain has a unique stationary distribution* $\boldsymbol{\pi}$*, and all* $\pi_i > 0$*. If the chain is also aperiodic, then* $\boldsymbol{\pi}_0 P^t \to \boldsymbol{\pi}$ *as* $t \to \infty$*, for every initial distribution* $\boldsymbol{\pi}_0$*.*

We do not prove this theorem in general; it is a consequence of the Perron–Frobenius theorem on matrices with nonnegative entries. The speed of convergence is governed by the second largest absolute value of an eigenvalue of P. Both assumptions are needed. For $P = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, the chain alternates between the two states: $(\frac{1}{2}, \frac{1}{2})$ is stationary, but $\boldsymbol{\pi}_0 = (1, 0)$ gives $\boldsymbol{\pi}_t = (1, 0), (0, 1), (1, 0), \ldots$ (the chain has period 2, and -1 is an eigenvalue). For $P = I$, the chain never moves, and every distribution is stationary. For two states, everything can be computed by hand.

Proposition 8.7 (Two-state chains)**.** *Let* $P = \begin{pmatrix} 1-a & a \\ b & 1-b \end{pmatrix}$ *with* $a+b > 0$*. Then* $\boldsymbol{\pi} = (\frac{b}{a+b}, \frac{a}{a+b})$ *is stationary, and for every initial distribution,*

$$\boldsymbol{\pi}_t - \boldsymbol{\pi} = (1 - a - b)^t \, (\boldsymbol{\pi}_0 - \boldsymbol{\pi}).$$

Proof. One checks $\boldsymbol{\pi} P = \boldsymbol{\pi}$ directly. The difference $\mathbf{d} = \boldsymbol{\pi}_0 - \boldsymbol{\pi}$ has entries that sum to 0, so $\mathbf{d} = c(1, -1)$, and $(1, -1)P = (1 - a - b, a - 1 + b) = (1 - a - b)(1, -1)$: the row vector $(1, -1)$ is a left eigenvector with the eigenvalue $1 - a - b$. Hence $\boldsymbol{\pi}_t - \boldsymbol{\pi} = (\boldsymbol{\pi}_0 - \boldsymbol{\pi})P^t = (1 - a - b)^t \, \mathbf{d}$. □

For the weather chain, $a = 0.1$ and $b = 0.5$, so the stationary distribution is $(\frac{5}{6}, \frac{1}{6})$, and the distance to it shrinks by the factor 0.4 (the second eigenvalue of P) every day. Starting on a sunny day, $P(X_t = \text{sunny}) = \frac{5}{6} + \frac{1}{6}\, 0.4^t$; starting on a rainy day, it is $\frac{5}{6} - \frac{5}{6}\, 0.4^t$ (Figure 4).




Figure 4: Left: the weather chain of Example 8.2. Right: the probability of a sunny day t days after a sunny day (orange) and after a rainy day (blue); both converge to the stationary probability $\frac{5}{6}$, and the distance shrinks by the factor 0.4 every day.

The ergodic theorem. For an irreducible chain, the fraction of time that the chain spends in state i converges to π_i with probability 1, and more generally $\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) \to \sum_i \pi_i f(i)$ for every function f. This is a law of large numbers for dependent random variables, which we state without proof. In the long run, $\frac{5}{6}$ of the days are sunny. Since successive states are correlated, these averages converge more slowly than averages of independent samples.

8.1 Detailed balance and Markov chain Monte Carlo

Finding a stationary distribution means solving a linear system. It is often easier to check a stronger, local condition.

Proposition 8.8 (Detailed balance)**.** *If a distribution $\boldsymbol{\pi}$ satisfies $\pi_i P_{ij} = \pi_j P_{ji}$ for all states i and j, then it is stationary.*

Proof. $(\boldsymbol{\pi}P)_j = \sum_i \pi_i P_{ij} = \sum_i \pi_j P_{ji} = \pi_j \sum_i P_{ji} = \pi_j$. □

Detailed balance says that in the stationary regime, the probability flow from i to j equals the flow from j to i; such chains are called reversible.

Example 8.9 (Random walks on graphs)**.** Let G be a connected undirected graph, and let a walker move from each vertex to a uniformly random neighbor, i.e., $P_{ij} = \frac{1}{\deg(i)}$ for each edge $\{i, j\}$. Then $\pi_i = \frac{\deg(i)}{2|E|}$, where $|E|$ is the number of edges, satisfies detailed balance: $\pi_i P_{ij} = \frac{1}{2|E|} = \pi_j P_{ji}$ for every edge. So in the long run, the walker visits each vertex in proportion to its degree. For the graph with the edges $\{1,2\}$, $\{1,3\}$, $\{2,3\}$, $\{3,4\}$, the degrees are $2, 2, 3, 1$, and $\boldsymbol{\pi} = (\frac{1}{4}, \frac{1}{4}, \frac{3}{8}, \frac{1}{8})$.

Remark 8.10 (PageRank). Google's original PageRank models a "random surfer" who, at each step, follows a uniformly random link on the current page with probability α (typically 0.85), and jumps to a uniformly random page with probability $1-\alpha$. All entries of the resulting transition matrix are positive, so the chain is irreducible and aperiodic, and its unique stationary distribution ranks the pages. It is computed by the power iteration $\boldsymbol{\pi} \leftarrow \boldsymbol{\pi}P$ of the programming exercises of Lecture 1.

Detailed balance is also the key to Markov chain Monte Carlo (MCMC), the most important method for sampling from complicated distributions, such as the posterior distributions of Section 12. These are often only known up to a constant: $p(\theta \mid \mathcal{D}) \propto p(\mathcal{D} \mid \theta)\, p(\theta)$, where the evidence $p(\mathcal{D})$ is an intractable integral. The idea is to construct a Markov chain whose stationary distribution is the target distribution, run it for a long time, and use its states as (dependent) samples.

Theorem 8.11 (Metropolis algorithm)**.** *Let $\boldsymbol{\pi}$ be a distribution with all $\pi_i > 0$, and let Q be a symmetric stochastic matrix ($Q_{ij} = Q_{ji}$), the proposal. From the state i, propose a state j with probability Q_{ij}, and move to j with the acceptance probability* $\min\left(1, \frac{\pi_j}{\pi_i}\right)$*; otherwise, stay at i. This defines a Markov chain that satisfies detailed balance with respect to $\boldsymbol{\pi}$, so $\boldsymbol{\pi}$ is stationary.*

Proof. For $i \neq j$, $P_{ij} = Q_{ij} \min\left(1, \frac{\pi_j}{\pi_i}\right)$, so $\pi_i P_{ij} = Q_{ij} \min(\pi_i, \pi_j)$, which is symmetric in i and j because Q is. For $i = j$, detailed balance holds trivially. □

The algorithm only uses the ratios $\frac{\pi_j}{\pi_i}$, so the unknown normalizing constant cancels. Moves to more probable states are always accepted, and moves to less probable states sometimes. For densities, the same algorithm works with a symmetric proposal such as $\theta' = \theta + \eta\,\varepsilon$ with $\varepsilon \sim \mathcal{N}(0,1)$, and the ratio of the (unnormalized) densities. If the resulting chain is irreducible and aperiodic, Theorem 8.6 and the ergodic theorem justify using averages along the chain as estimates of expectations under $\boldsymbol{\pi}$.

Example 8.12. Let the states $1, 2, 3, 4$ lie on a cycle, let Q propose each of the two neighbors with probability $\frac{1}{2}$, and let the target be $\boldsymbol{\pi} = (0.1, 0.2, 0.3, 0.4)$. The Metropolis chain has the transition matrix

$$P = \begin{pmatrix} 0 & 0.5 & 0 & 0.5 \\ 0.25 & 0.25 & 0.5 & 0 \\ 0 & 1/3 & 1/6 & 0.5 \\ 0.125 & 0 & 0.375 & 0.5 \end{pmatrix};$$

for example, from state 3, the move to state 2 is proposed with probability $\frac{1}{2}$ and accepted with probability $\frac{0.2}{0.3}$, so $P_{32} = \frac{1}{3}$. One checks that $\boldsymbol{\pi} P = \boldsymbol{\pi}$.

Remark 8.13 (Markov models in machine learning). An n-gram language model is a Markov chain on words (or on tuples of $n-1$ words), whose transition probabilities are estimated by counting in a large text, which is maximum likelihood estimation (Section 9). Hidden Markov models, in which only a noisy function of the state is observed, were long the standard model in speech recognition. Reinforcement learning is based on Markov decision processes, in which the transition probabilities depend on the actions of an agent. Diffusion models for image generation learn to reverse a Markov chain that gradually adds Gaussian noise to an image (Section 14.1).

9 Parameter Estimation

Statistics turns the questions of probability around: instead of computing probabilities from a known distribution, we infer an unknown distribution from data. We assume that the data $x_1, \dots, x_N$ are i.i.d. samples from a distribution $p(x \mid \theta)$ that belongs to a known family, with an unknown parameter θ (for example, the mean and variance of a Gaussian, or the weights of a model).

Definition 9.1 (Estimator, bias and variance)**.** An estimator is a function $\hat{\theta} = \hat{\theta}(X_1, \dots, X_N)$ of the data; as a function of random data, it is a random variable. Its bias is $\mathbb{E}[\hat{\theta}] - \theta$, and it is unbiased if the bias is 0 for all θ. Its mean squared error is $\mathrm{MSE}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \theta)^2]$.

Proposition 9.2 (Bias–variance decomposition)**.** $\mathrm{MSE}(\hat{\theta}) = \big(\mathbb{E}[\hat{\theta}] - \theta\big)^2 + \mathrm{Var}(\hat{\theta})$.

Proof. Let $m = \mathbb{E}[\hat{\theta}]$. Then $\mathbb{E}[(\hat{\theta} - \theta)^2] = \mathbb{E}[(\hat{\theta} - m)^2] + 2(m - \theta)\mathbb{E}[\hat{\theta} - m] + (m - \theta)^2$, and the middle term is 0. □

A good estimator has a small bias and a small variance, and sometimes we accept some bias for a much smaller variance. This is the idea behind regularization: ridge regression shrinks the weights towards 0, which introduces a bias but reduces the variance. For a prediction model, a similar decomposition says that the expected squared test error is the sum of the squared bias, the variance and the irreducible noise; very flexible models have a small bias but a large variance (they overfit), and very simple ones the opposite.

By Proposition 7.1, the sample mean is an unbiased estimator of μ, with variance $\frac{\sigma^2}{N}$. The natural estimator of the variance, however, is biased.

Proposition 9.3 (Sample variance)**.** *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and variance σ^2, and $N \geq 2$. Then*

$$\mathbb{E}\Big[\frac{1}{N}\sum_{i=1}^{N}(X_i - \bar{X}_N)^2\Big] = \frac{N-1}{N}\sigma^2, \qquad \textit{so} \qquad s^2 = \frac{1}{N-1}\sum_{i=1}^{N}(X_i - \bar{X}_N)^2$$

is an unbiased estimator of σ^2.

Proof. Expanding $X_i - \mu = (X_i - \bar{X}_N) + (\bar{X}_N - \mu)$ and using $\sum_i (X_i - \bar{X}_N) = 0$ gives

$$\sum_i (X_i - \mu)^2 = \sum_i (X_i - \bar{X}_N)^2 + N(\bar{X}_N - \mu)^2.$$

Taking expectations, $N\sigma^2 = \mathbb{E}\big[\sum_i (X_i - \bar{X}_N)^2\big] + N \cdot \frac{\sigma^2}{N}$. □

Intuitively, the data are on average closer to their own mean $\bar{X}_N$ than to the true mean μ, so the version with $\frac{1}{N}$ underestimates the variance. The factor $\frac{N}{N-1}$ (Bessel's correction) only matters for small N; `np.var` uses $\frac{1}{N}$ by default, and `np.var(x, ddof=1)` uses $\frac{1}{N-1}$.

9.1 The method of moments

The oldest general method of estimation chooses the parameters so that the model reproduces the moments of the data.

Definition 9.4 (Method of moments)**.** Let $p(x \mid \boldsymbol{\theta})$ be a family of distributions with k parameters. The method of moments estimate $\hat{\boldsymbol{\theta}}$ solves the k equations

$$\mathbb{E}_{\boldsymbol{\theta}}[X^j] = \frac{1}{N}\sum_{i=1}^{N} x_i^j, \qquad j = 1, \dots, k,$$

(or equivalent equations for the mean, the variance, and so on).

By the law of large numbers, the sample moments converge to the true moments, so the method of moments gives consistent estimates whenever the moment equations determine $\boldsymbol{\theta}$ in a continuous way.

Example 9.5.

1. Exponential(λ): $\mathbb{E}[X] = \frac{1}{\lambda}$ gives $\hat{\lambda} = \frac{1}{\bar{x}}$.
2. Uniform$(0, \theta)$: $\mathbb{E}[X] = \frac{\theta}{2}$ gives $\hat{\theta} = 2\bar{x}$. This estimator is unbiased, but it can be smaller than the largest observation, which is impossible for the true θ (compare with the MLE in the exercises).
3. The Gamma distribution with shape $k > 0$ and scale $s > 0$ has the density $\frac{x^{k-1}e^{-x/s}}{\Gamma(k)s^k}$ for $x > 0$, the mean ks and the variance ks^2. Matching the sample mean $\bar{x}$ and the sample variance $\hat{\sigma}^2$ gives $\hat{s} = \frac{\hat{\sigma}^2}{\bar{x}}$ and $\hat{k} = \frac{\bar{x}^2}{\hat{\sigma}^2}$; for example, $\bar{x} = 6$ and $\hat{\sigma}^2 = 12$ give $\hat{k} = 3$ and $\hat{s} = 2$. The maximum likelihood estimate of k has no closed form, and the method of moments provides a good starting point for its numerical computation (for example with Newton's method, Lecture 3).

The method of moments is simple, but its estimates usually have a larger variance than maximum likelihood estimates, which we discuss next.

9.2 Maximum likelihood estimation

The most important general method to construct estimators is to choose the parameter under which the observed data are most probable.

Definition 9.6 (Maximum likelihood)**.** The likelihood of a parameter θ for i.i.d. data $x_1, \dots, x_N$ is $L(\theta) = \prod_{i=1}^{N} p(x_i \mid \theta)$, and the log-likelihood is $\ell(\theta) = \log L(\theta) = \sum_{i=1}^{N} \log p(x_i \mid \theta)$. A maximum likelihood estimate (MLE) is a parameter $\hat{\theta}$ that maximizes L, or equivalently ℓ.

Taking logarithms turns the product into a sum, which is easier to differentiate and numerically more stable. Maximizing ℓ is the same as minimizing the negative log-likelihood $-\ell(\theta)$, which is a loss function in the sense of Lecture 3.

Example 9.7 (Bernoulli)**.** Let $x_i \in \{0,1\}$ with k ones among N observations, and $p(x \mid \theta) = \theta^x(1-\theta)^{1-x}$. Then $\ell(\theta) = k\log\theta + (N-k)\log(1-\theta)$, and $\ell'(\theta) = \frac{k}{\theta} - \frac{N-k}{1-\theta} = 0$ gives $\hat\theta = \frac{k}{N}$, the relative frequency. Since $\ell'' < 0$, this is the maximum. Note that after three heads in three flips, $\hat\theta = 1$: the MLE claims that tails are impossible. With little data, maximum likelihood overfits; we return to this example in Section 12.

Example 9.8 (Gaussian)**.** For $x_i \sim \mathcal{N}(\mu, \sigma^2)$,

$$\ell(\mu, \sigma^2) = -\frac{N}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}\sum_{i=1}^{N}(x_i - \mu)^2.$$

Setting the partial derivatives to zero gives $\hat\mu = \bar x$ and $\hat\sigma^2 = \frac{1}{N}\sum_i (x_i - \bar x)^2$: the MLE of the variance is the biased version of Proposition 9.3. For the data $2.1, 3.4, 1.9, 2.8, 3.3$, we get $\hat\mu = 2.7$, $\hat\sigma^2 = 0.372$, while $s^2 = 0.465$.

Theorem 9.9 (MLE of a multivariate Gaussian)**.** *For i.i.d. data $\mathbf{x}_1, \dots, \mathbf{x}_N$ from $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$, whose sample covariance matrix S is invertible, the maximum likelihood estimates are the sample mean $\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}}$ and the sample covariance matrix $\hat\Sigma = S = \frac{1}{N}\sum_i (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top$.*

Proof. Up to a constant, $\ell = -\frac{N}{2}\log\det\Sigma - \frac{1}{2}\sum_i (\mathbf{x}_i - \boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}_i - \boldsymbol{\mu})$. For fixed Σ, this is a concave quadratic function of $\boldsymbol{\mu}$ with gradient $\Sigma^{-1}\sum_i(\mathbf{x}_i - \boldsymbol{\mu})$ (Lecture 3), which vanishes for $\boldsymbol{\mu} = \bar{\mathbf{x}}$, whatever Σ is. With $\boldsymbol{\mu} = \bar{\mathbf{x}}$, we write ℓ in terms of the precision matrix $\Lambda = \Sigma^{-1}$. Since $\mathbf{a}^\top\Lambda\mathbf{a} = \operatorname{tr}(\Lambda\mathbf{a}\mathbf{a}^\top)$,

$$\ell = \frac{N}{2}\log\det\Lambda - \frac{1}{2}\sum_i \operatorname{tr}\left(\Lambda(\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top\right) = \frac{N}{2}\big(\log\det\Lambda - \operatorname{tr}(\Lambda S)\big).$$

By the matrix gradients of Lecture 3 ($\nabla \log\det\Lambda = \Lambda^{-\top}$ and $\nabla\operatorname{tr}(\Lambda S) = S^\top$), the gradient with respect to Λ is $\frac{N}{2}(\Lambda^{-1} - S)$, which vanishes for $\Lambda = S^{-1}$, i.e., $\Sigma = S$. (One can show that $\log\det$ is concave on the symmetric positive definite matrices, so this critical point is the maximum.) □

9.3 Consistency and efficiency

How good can an estimator be? Two properties are central: an estimator should converge to the true parameter as the amount of data grows, and its variance should be as small as possible.

Definition 9.10 (Consistency)**.** A sequence of estimators $\hat\theta_N$, computed from N data points, is consistent if for every $\varepsilon > 0$ and every true parameter θ, $P(|\hat\theta_N - \theta| \ge \varepsilon) \to 0$ as $N \to \infty$.

Proposition 9.11. *If the bias and the variance of $\hat\theta_N$ tend to 0 as $N \to \infty$, then $\hat\theta_N$ is consistent.*

Proof. By Markov's inequality applied to $(\hat\theta_N - \theta)^2$ and Proposition 9.2, $P(|\hat\theta_N - \theta| \ge \varepsilon) \le \frac{\mathrm{MSE}(\hat\theta_N)}{\varepsilon^2} = \frac{\mathrm{bias}^2 + \mathrm{Var}(\hat\theta_N)}{\varepsilon^2} \to 0$. □

For example, the sample mean and both versions of the sample variance are consistent, and so are the method of moments estimators. A lower bound for the variance comes from the curvature of the log-likelihood. For a family $p(x \mid \theta)$ with a single parameter, the score is $s(x;\theta) = \frac{\partial}{\partial\theta}\log p(x \mid \theta)$. Differentiating $\int p(x \mid \theta)\,dx = 1$ under the integral sign gives $\int \frac{\partial p}{\partial\theta}\,dx = \int s\,p\,dx = 0$, i.e., $\mathbb{E}_\theta[s(X;\theta)] = 0$. The Fisher information is

$$I(\theta) = \mathbb{E}_\theta\big[s(X;\theta)^2\big] = \operatorname{Var}_\theta\big(s(X;\theta)\big),$$

and one can show that also $I(\theta) = -\mathbb{E}_\theta\big[\frac{\partial^2}{\partial\theta^2}\log p(X \mid \theta)\big]$: it measures how sharply the log-likelihood is curved around the true parameter, i.e., how much information one observation carries about θ.

Theorem 9.12 (Cramér–Rao bound)**.** *Let $\hat\theta$ be an unbiased estimator of θ from N i.i.d. observations, and assume that we may differentiate under the integral sign (Lecture 3; this excludes, e.g., families whose support depends on θ). Then*

$$\operatorname{Var}_\theta(\hat\theta) \geq \frac{1}{N\, I(\theta)}.$$

Proof. Let $S = \sum_{i=1}^N s(X_i;\theta)$ be the score of the whole sample. It has mean 0 and, by independence, variance $N\,I(\theta)$. Write $p_N(\mathbf{x} \mid \theta) = \prod_i p(x_i \mid \theta)$, so that $\frac{\partial p_N}{\partial \theta} = p_N S$. Differentiating the unbiasedness condition $\theta = \int \hat\theta(\mathbf{x})\, p_N(\mathbf{x}\mid\theta)\, d\mathbf{x}$ gives

$$1 = \int \hat\theta(\mathbf{x})\, S(\mathbf{x})\, p_N(\mathbf{x}\mid\theta)\, d\mathbf{x} = \mathbb{E}[\hat\theta S] = \operatorname{Cov}(\hat\theta, S),$$

since $\mathbb{E}[S] = 0$. By Theorem 4.7 (the Cauchy–Schwarz inequality),

$$1 = \operatorname{Cov}(\hat\theta, S)^2 \leq \operatorname{Var}(\hat\theta)\operatorname{Var}(S) = \operatorname{Var}(\hat\theta)\, N\, I(\theta). \qquad \square$$

An unbiased estimator that attains the bound is called efficient.

Example 9.13.

1. Bernoulli(θ): $s(x;\theta) = \frac{x}{\theta} - \frac{1-x}{1-\theta}$ and $I(\theta) = \frac{1}{\theta(1-\theta)}$. The bound $\frac{\theta(1-\theta)}{N}$ is the variance of the relative frequency $\bar X_N$, which is therefore efficient.
2. $\mathcal{N}(\mu, \sigma^2)$ with known σ^2: $s = \frac{x-\mu}{\sigma^2}$ and $I(\mu) = \frac{1}{\sigma^2}$, so the sample mean, with variance $\frac{\sigma^2}{N}$, is efficient.
3. Uniform$(0,\theta)$ does not satisfy the assumptions, since its support depends on θ. Indeed, the unbiased estimator $\frac{N+1}{N}\max_i X_i$ has the variance $\frac{\theta^2}{N(N+2)}$, which decreases like $\frac{1}{N^2}$, faster than any bound of the form $\frac{c}{N}$.

Theorem 9.14 (Asymptotic normality of the MLE)**.** *Under regularity conditions, the maximum likelihood estimator is consistent, and $\sqrt{N}(\hat\theta_{MLE} - \theta)$ converges in distribution to $\mathcal{N}\big(0, \frac{1}{I(\theta)}\big)$.*

We do not prove this theorem. It says that for large N, the MLE is approximately unbiased and attains the Cramér–Rao bound: maximum likelihood is asymptotically efficient, which is the main theoretical justification for it. In practice, the standard error of the MLE is estimated from the curvature of the log-likelihood at its maximum, $\mathrm{SE} \approx \big(-\ell''(\hat\theta)\big)^{-1/2}$ (the observed information). For example, for k successes in N Bernoulli trials, $-\ell''(\hat\theta) = \frac{k}{\hat\theta^2} + \frac{N-k}{(1-\hat\theta)^2} = \frac{N}{\hat\theta(1-\hat\theta)}$, which gives the familiar $\mathrm{SE} = \sqrt{\hat\theta(1-\hat\theta)/N}$, e.g., 0.046 for 30 successes in 100 trials. With several parameters, the Fisher information becomes a matrix, and the approximate covariance matrix of the MLE is the inverse of the Hessian of $-\ell$ (Lecture 3). The Fisher information matrix also defines the inner product of the natural gradient method mentioned in Lecture 3.

9.4 Loss functions as negative log-likelihoods

Maximum likelihood explains the loss functions of Lectures 2 and 3.

Proposition 9.15 (Least squares as maximum likelihood)**.** *Assume the linear regression model $y_i = \mathbf{w}^\top \mathbf{x}_i + \varepsilon_i$, where the noise terms $\varepsilon_i \sim \mathcal{N}(0,\sigma^2)$ are i.i.d. Then the negative log-likelihood of $\mathbf{w}$ is*

$$-\ell(\mathbf{w}) = \frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 + \frac{N}{2}\log(2\pi\sigma^2),$$

where X has the rows $\mathbf{x}_i^\top$. So the MLE of $\mathbf{w}$ is the least squares solution, and the MLE of the noise variance is the mean squared residual $\frac{1}{N}\|\mathbf{y} - X\hat{\mathbf{w}}\|^2$.

Proof. Given $\mathbf{x}_i$, $y_i \sim \mathcal{N}(\mathbf{w}^\top \mathbf{x}_i, \sigma^2)$, so $-\log p(y_i \mid \mathbf{x}_i, \mathbf{w}) = \frac{(y_i - \mathbf{w}^\top \mathbf{x}_i)^2}{2\sigma^2} + \frac{1}{2}\log(2\pi\sigma^2)$; sum over i. The second statement is proved as in Example 9.8. □

Similarly, if $y_i \in \{0, 1\}$ is Bernoulli with $P(y_i = 1) = \sigma(\mathbf{w}^\top \mathbf{x}_i)$, the negative log-likelihood is $\sum_i \big(-y_i \log p_i - (1 - y_i)\log(1 - p_i)\big)$ with $p_i = \sigma(\mathbf{w}^\top \mathbf{x}_i)$: the cross-entropy loss of logistic regression in Lecture 3. For K classes with $P(y_i = k) = \mathrm{softmax}(\mathbf{z}_i)_k$, it is the softmax cross-entropy loss $-\sum_i \log \mathrm{softmax}(\mathbf{z}_i)_{y_i}$. So choosing a loss function amounts to choosing a noise model. For example, Laplace-distributed noise with density $\frac{1}{2b}e^{-|\varepsilon|/b}$ leads to the absolute loss $\sum_i |y_i - \mathbf{w}^\top \mathbf{x}_i|$, which is less sensitive to outliers than the squared loss.

9.5 Confidence intervals

An estimate is more useful with an error bar. By the central limit theorem, $\bar{X}_N$ is approximately $\mathcal{N}(\mu, \frac{\sigma^2}{N})$, so

$$P\Big(|\bar{X}_N - \mu| \le 1.96\frac{\sigma}{\sqrt{N}}\Big) \approx 0.95.$$

Replacing the unknown σ by the sample standard deviation s (which is accurate for large N), the interval $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$ is an approximate 95% confidence interval for μ. For example, for $N = 100$ data points with $\bar{x} = 5.2$ and $s = 1.5$, it is $[4.91, 5.49]$. Note that μ is fixed and the interval is random: if we repeated the experiment many times, about 95% of the intervals would contain μ. For a classifier with accuracy 0.9 on 1000 independent test examples, the standard error is $\sqrt{0.9 \cdot 0.1/1000} \approx 0.0095$, so the accuracy is known up to about ± 0.019, and a difference of one percentage point between two models on such a test set may well be due to chance.

10 Hypothesis Testing

Often we want to decide a yes-or-no question from data: is a coin fair? Does a new model have a lower error rate than the old one? Does a new website design increase sales? A hypothesis test makes such decisions while controlling the probability of a false alarm.

Definition 10.1 (Hypothesis test)**.** A test compares a null hypothesis H_0 (usually "no effect") with an alternative hypothesis H_1. It computes a test statistic T from the data and rejects H_0 if T falls into a rejection region. Rejecting a true H_0 is a type I error (a false positive), and keeping a false H_0 is a type II error (a false negative). A test has significance level α if the probability of a type I error is at most α; its power is the probability of rejecting H_0 when H_1 is true.

Definition 10.2 (p-value)**.** If large values of T speak against H_0, the p-value of an observed value t is $p = P_{H_0}(T \ge t)$: the probability, computed under H_0, of a test statistic at least as extreme as the observed one. The test that rejects H_0 if $p \le \alpha$ has significance level α.

Proposition 10.3. *If T has a continuous, strictly increasing CDF F under H_0, then the p-value $1 - F(T)$ is Uniform$(0, 1)$-distributed under H_0. In particular, $P_{H_0}(p \le \alpha) = \alpha$.*

Proof. $P(F(T) \le u) = P(T \le F^{-1}(u)) = F(F^{-1}(u)) = u$, so $F(T)$ is uniform on $(0, 1)$, as in Theorem 2.7, and so is $1 - F(T)$. □

Example 10.4 (Testing a coin)**.** A coin shows 60 heads in 100 flips. Is it fair? With H_0: $\theta = \frac{1}{2}$ and the number of heads S as the test statistic, the one-sided p-value is $P_{H_0}(S \ge 60) = 0.028$ (Example 7.4). If we had no reason to expect more heads than tails, a two-sided test is appropriate:

it counts deviations in both directions, and its p-value is $P_{H_0}(|S-50| \geq 10) = 0.057$. At the level $\alpha = 0.05$, the one-sided test rejects H_0, and the two-sided test does not. The test must be chosen before looking at the data; otherwise its significance level is not what it claims.

Tests for a mean. To test H_0: $\mu = \mu_0$ for i.i.d. data, we use the standardized statistic $Z = \frac{\bar{X}_N - \mu_0}{s/\sqrt{N}}$, which is approximately $\mathcal{N}(0,1)$ under H_0 by the central limit theorem (the z-test). The two-sided p-value is $2(1-\Phi(|z|))$. For small samples from a Gaussian distribution, Z has a Student t-distribution with $N-1$ degrees of freedom, whose quantiles are larger than 1.96 (for example, 2.78, 2.26 and 2.05 for $N = 5$, 10 and 30); this is the t-test. Tests and confidence intervals are two sides of the same coin: the two-sided test at level 0.05 rejects $\mu = \mu_0$ exactly if μ_0 lies outside the 95% confidence interval $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$.

Remark 10.5 (What a p-value is not). A p-value is not the probability that H_0 is true. That probability depends on the prior plausibility of H_0, as in the medical test of Example 1.7, and only Bayesian inference can compute it. A small p-value can also come from a tiny effect that is irrelevant in practice, if N is huge (statistical versus practical significance), so effect sizes and confidence intervals should be reported as well. Conversely, a large p-value does not show that there is no effect: the test may just have too little power.

10.1 Multiple testing

If we perform many tests, some of them will reject by chance. For m independent tests of true null hypotheses at the level 0.05, the probability of at least one false positive is $1 - 0.95^m$, which is 0.64 for $m = 20$. This happens when we compare many models on the same test set, screen thousands of genes or features, or analyze many metrics of an experiment, and it is a main source of irreproducible results ("p-hacking"). The probability of at least one false positive among all tests is called the family-wise error rate (FWER).

Proposition 10.6 (Bonferroni correction)**.** *If each of m tests rejects its null hypothesis when its p-value is at most $\frac{\alpha}{m}$, then the family-wise error rate is at most α, whatever the dependence between the tests.*

Proof. Let I_0 be the set of true null hypotheses. By the union bound (Proposition 1.3) and Proposition 10.3, $P\big(\bigcup_{i \in I_0} \{p_i \leq \frac{\alpha}{m}\}\big) \leq \sum_{i \in I_0} \frac{\alpha}{m} \leq \alpha$. □

The Bonferroni correction is very conservative when m is large. When we screen many hypotheses, it is often more useful to control the false discovery rate (FDR), the expected fraction of false positives among all rejected hypotheses. The Benjamini–Hochberg procedure sorts the p-values $p_{(1)} \leq \cdots \leq p_{(m)}$, finds the largest k with $p_{(k)} \leq \frac{k}{m} q$, and rejects the k hypotheses with the smallest p-values. One can show that for independent tests, its false discovery rate is at most q.

Example 10.7. Ten tests give the p-values

$$0.001,\ 0.008,\ 0.012,\ 0.030,\ 0.041,\ 0.10,\ 0.25,\ 0.40,\ 0.62,\ 0.90.$$

Without correction, five are below 0.05. Bonferroni with $\alpha = 0.05$ uses the threshold 0.005 and rejects only the first. Benjamini–Hochberg with $q = 0.05$ compares $p_{(k)}$ with $0.005k$: $0.001 \leq 0.005$, $0.008 \leq 0.010$ and $0.012 \leq 0.015$, but $p_{(k)} > 0.005k$ for all $k \geq 4$, so it rejects three hypotheses.

10.2 The bootstrap and permutation tests

The formulas for standard errors and p-values so far required either a simple statistic, such as a mean, or a known distribution. Two resampling methods work without such formulas, at the cost of computation.

The bootstrap. To estimate the uncertainty of a statistic $\hat{\theta} = s(x_1, \dots, x_N)$, such as a median, a correlation, or the AUC of a classifier, the bootstrap replaces the unknown distribution of the data by the empirical distribution (the plug-in principle). It draws B bootstrap samples $x_1^*, \dots, x_N^*$, each consisting of N draws with replacement from the data, and computes $\hat{\theta}_b^* = s(x_1^*, \dots, x_N^*)$ for $b = 1, \dots, B$. The standard deviation of the $\hat{\theta}_b^*$ estimates the standard error of $\hat{\theta}$, and the 2.5% and 97.5% quantiles of the $\hat{\theta}_b^*$ give an approximate 95% confidence interval (the percentile interval). For the sample mean, everything can be computed exactly: a bootstrap sample consists of N i.i.d. draws from the empirical distribution, whose variance is $\hat{\sigma}^2 = \frac{1}{N}\sum_i (x_i - \bar{x})^2$, so the bootstrap standard error of the mean is $\frac{\hat{\sigma}}{\sqrt{N}}$, the familiar formula. The bootstrap can fail for very small samples, for statistics such as the maximum, and for dependent data.

Permutation tests. Suppose we observe $x_1, \dots, x_n$ in group A and $y_1, \dots, y_m$ in group B, and want to test H_0: both groups come from the same distribution. Under H_0, the group labels carry no information: given the $n + m$ observed values, each of the $\binom{n+m}{n}$ ways to assign n of them to group A is equally likely. So the p-value of a statistic such as the difference of the group means is the fraction of all relabelings whose statistic is at least as large as the observed one. For large samples, we use a few thousand random relabelings instead of all of them. The test has exactly the claimed level, and it requires no assumptions about the distribution.

Example 10.8. Let group A have the values $12, 15, 14$ and group B the values $9, 10, 13$. The observed difference of the means is $13\frac{2}{3} - 10\frac{2}{3} = 3$. Of the $\binom{6}{3} = 20$ ways to split the six values into two groups of three, only 2 give a difference of at least 3: the observed split, and the split $\{13, 14, 15\}$ versus $\{9, 10, 12\}$, with the difference $3\frac{2}{3}$. So the one-sided p-value is $\frac{2}{20} = 0.1$: with so few observations, even a clear difference is not significant.

11 Evaluating Models and Experiments

11.1 Cross-validation

To choose between models or hyperparameters, we need to estimate how well a learning algorithm generalizes, and we often cannot afford to set aside a large test set. In k-fold cross-validation, the data are split into k folds of about equal size; for $j = 1, \dots, k$, the model is trained on all folds except the j-th one and evaluated on the j-th fold, which gives an error e_j. The cross-validation estimate is the average $\bar{e} = \frac{1}{k}\sum_j e_j$. Typical choices are $k = 5$ or 10; for $k = N$, it is leave-one-out cross-validation. The statistical properties of $\bar{e}$ are subtle.

- Bias. Each model is trained on only a fraction $\frac{k-1}{k}$ of the data, so $\bar{e}$ slightly overestimates the error of a model trained on all data; this bias is smaller for larger k.
- Variance. The fold errors are not independent, since the training sets overlap. So the naive standard error $\frac{s_e}{\sqrt{k}}$, with the standard deviation s_e of the fold errors, usually underestimates the true uncertainty; one can show that no estimator of the variance of $\bar{e}$ is unbiased for all distributions. For example, the fold errors $0.12, 0.15, 0.10, 0.14, 0.13$ have the mean 0.128, the

standard deviation 0.019 and the naive standard error 0.009; the true uncertainty is larger. Repeating cross-validation with different random splits reduces the variance that comes from the split, but not the dependence.

- Comparing two models. Comparing two models on the same folds (or the same test examples) and analyzing the differences is much more precise than comparing their separate error estimates, since $\mathrm{Var}(A-B) = \mathrm{Var}(A) + \mathrm{Var}(B) - 2\,\mathrm{Cov}(A,B)$ and the errors of two models on the same data are positively correlated. This is the idea of paired tests.

- Selection bias. If we evaluate many models or hyperparameter settings and choose the one with the smallest cross-validation error, the selected error estimate is biased downward: by Jensen's inequality, the expected minimum of noisy estimates is smaller than the minimum of their expectations. For example, if 20 models all have the true error 0.10 and their estimates have independent errors with standard deviation 0.01, the smallest estimate is about $0.10 - 1.87 \cdot 0.01 \approx 0.081$ on average (the expected maximum of 20 independent standard normal variables is about 1.87). The remedy is to estimate the error of the selected model on data that played no role in the selection: a separate test set, used once, or nested cross-validation, which repeats the whole selection procedure inside each training fold.

11.2 A/B testing and experiment design

To find out whether a change causes an improvement, for example a new recommendation model, the most reliable method is a randomized controlled experiment, called an A/B test in industry: each user is randomly assigned to version A (the control) or version B (the treatment), and the outcomes are compared. Randomization makes the two groups comparable in all respects, observed or not, on average. So a significant difference can be attributed to the treatment. Observational comparisons, such as "users who chose the new feature buy more", can be confounded by other differences between the groups.

Analyzing an A/B test. Let $\hat{p}_A$ and $\hat{p}_B$ be the conversion rates (for example, the fractions of users who buy something) of n_A and n_B users. Under H_0: $p_A = p_B$, the pooled rate $\hat{p}$ of all users estimates the common rate, and by the central limit theorem,

$$Z = \frac{\hat{p}_B - \hat{p}_A}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{n_A} + \frac{1}{n_B}\right)}}$$

is approximately $\mathcal{N}(0,1)$ (the two-proportion z-test). For example, if 200 of 5000 users convert with version A (4%) and 250 of 5000 with version B (5%), then $\hat{p} = 0.045$, $Z \approx 2.41$, and the two-sided p-value is 0.016. An approximate 95% confidence interval for the improvement $p_B - p_A$ is $0.01 \pm 1.96\sqrt{\frac{\hat{p}_A(1-\hat{p}_A)}{n_A} + \frac{\hat{p}_B(1-\hat{p}_B)}{n_B}} = [0.002, 0.018]$.

Sample size and power. How many users do we need to detect an improvement $\delta = p_B - p_A$? With n users per group, the standard error of $\hat{p}_B - \hat{p}_A$ is about $\sqrt{2\bar{p}(1-\bar{p})/n}$, where $\bar{p}$ is the average rate, and Z is approximately $\mathcal{N}(\delta/\mathrm{SE}, 1)$. The two-sided test at level α rejects if $|Z| > z_{1-\alpha/2}$, so it has power $1-\beta$ approximately when $\frac{\delta}{\mathrm{SE}} = z_{1-\alpha/2} + z_{1-\beta}$, i.e.,

$$n \approx \frac{(z_{1-\alpha/2} + z_{1-\beta})^2 \cdot 2\bar{p}(1-\bar{p})}{\delta^2}$$

per group, where $z_u = \Phi^{-1}(u)$. For $\alpha = 0.05$ and power 0.8, $z_{0.975} + z_{0.8} \approx 1.96 + 0.84$; to detect an increase from 4% to 5%, we need about 6750 users per group. Halving the effect that we want to detect quadruples the sample size.

Design principles and pitfalls.

- Fix the primary metric, the test and the sample size in advance. Peeking at the p-value repeatedly and stopping as soon as it is below 0.05 inflates the false positive rate: in a simulation with two identical versions (conversion rate 5%, up to 10 000 users per group), checking after every 200 users per group declared a significant difference in about one third of the runs. If early stopping is needed, use sequential tests designed for it.
- Many metrics or user segments mean many tests: correct for multiple testing, or treat secondary results as hypotheses for future experiments.
- Randomize at the right level. If users influence each other (for example in a social network or a marketplace), or one user is counted several times, the observations are not independent, and the standard errors above are too small.
- Reduce variance by design. Paired or blocked designs compare similar units (for example, the same users or the same queries under both versions), and adjusting for measurements taken before the experiment can reduce the variance considerably.
- Watch for novelty effects, which make new versions look better at first, and check that the groups have the planned sizes; a mismatch indicates a bug in the assignment.

The same principles apply to offline comparisons of machine learning models: whether model B is really better than model A is a hypothesis testing question, and many reported improvements are within the noise of the evaluation.

12 Bayesian Inference

Maximum likelihood treats the parameter θ as a fixed unknown number. Bayesian inference treats it as a random variable, which expresses our uncertainty about it. Before seeing data, our beliefs are described by a prior distribution $p(\theta)$; after seeing the data $\mathcal{D}$, Bayes' theorem gives the posterior distribution

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}, \qquad p(\mathcal{D}) = \int p(\mathcal{D} \mid \theta)\, p(\theta)\, d\theta.$$

In words: the posterior is proportional to the likelihood times the prior. The normalizing constant $p(\mathcal{D})$ is called the evidence; it does not depend on θ.

12.1 Coin flips and the Beta distribution

Definition 12.1 (Beta distribution)**.** For $a, b > 0$, the Beta(a, b) distribution on $[0, 1]$ has the density

$$p(\theta) = \frac{\theta^{a-1}(1-\theta)^{b-1}}{B(a,b)}, \qquad \text{where} \quad B(a,b) = \int_0^1 \theta^{a-1}(1-\theta)^{b-1}\, d\theta.$$

For positive integers, $B(a, b) = \frac{(a-1)!\,(b-1)!}{(a+b-1)!}$; for example, Beta$(1, 1)$ is the uniform distribution, and Beta$(2, 2)$ has the density $6\theta(1-\theta)$. The mean of Beta(a, b) is $\frac{a}{a+b}$, and for $a, b > 1$ its mode (the maximum of the density) is $\frac{a-1}{a+b-2}$.

Theorem 12.2 (Beta–Bernoulli conjugacy)**.** *Let the prior of θ be Beta(a, b), and let the data be N independent Bernoulli(θ) observations with k ones. Then the posterior is Beta$(a + k,\, b + N - k)$.*

Proof. $p(\theta \mid \mathcal{D}) \propto \theta^k(1-\theta)^{N-k} \cdot \theta^{a-1}(1-\theta)^{b-1} = \theta^{a+k-1}(1-\theta)^{b+N-k-1}$. The only density proportional to this function is that of Beta$(a + k, b + N - k)$. □

A prior is called conjugate to a likelihood if the posterior belongs to the same family as the prior; then updating our beliefs amounts to updating a few parameters. Here, the prior acts like a previously seen ones and b previously seen zeros.

Example 12.3. With the prior Beta$(2, 2)$, which expresses a mild belief that the coin is fair, and three heads in three flips, the posterior is Beta$(5, 2)$. Its mean is $\frac{5}{7} \approx 0.71$ and its mode is $\frac{4}{5} = 0.8$, whereas the MLE was 1 (Example 9.7). After 30 flips with 20 heads, the posterior is Beta$(22, 12)$, with mean 0.65 and standard deviation 0.08: with more data, the posterior concentrates, and the prior matters less (Figure 5).




Figure 5: Bayesian updating for the probability θ of heads: the prior Beta$(2, 2)$ (blue), the posterior Beta$(5, 2)$ after 3 heads in 3 flips (orange), and the posterior Beta$(22, 12)$ after 20 heads in 30 flips (red). With more data, the posterior concentrates around the relative frequency of heads.

To predict the next flip, Bayesian inference averages over all values of θ, weighted by the posterior:

$$P(X_{N+1} = 1 \mid \mathcal{D}) = \int_0^1 \theta\, p(\theta \mid \mathcal{D})\, d\theta = \mathbb{E}[\theta \mid \mathcal{D}] = \frac{a + k}{a + b + N}.$$

For the uniform prior ($a = b = 1$), this is Laplace's rule of succession $\frac{k+1}{N+2}$: after three heads in three flips, the probability of heads is $\frac{4}{5}$, not 1. The same "add pseudo-counts" idea (Laplace smoothing) prevents zero probabilities for unseen words in language models.

12.2 Maximum a posteriori estimation and regularization

Often we only want a single parameter value. The maximum a posteriori (MAP) estimate is the mode of the posterior,

$$\hat{\theta}_{\text{MAP}} = \arg\max_\theta p(\theta \mid \mathcal{D}) = \arg\min_\theta \big(-\log p(\mathcal{D} \mid \theta) - \log p(\theta) \big),$$

since $p(\mathcal{D})$ does not depend on θ. So MAP estimation minimizes the negative log-likelihood plus a penalty $-\log p(\theta)$ coming from the prior: a prior is a regularizer.

Proposition 12.4 (Ridge regression as MAP estimation)**.** *In the linear regression model of Proposition 9.15, with the prior* $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1} I)$*, the MAP estimate minimizes* $\|\mathbf{y} - X\mathbf{w}\|^2 + \lambda \|\mathbf{w}\|^2$ *with* $\lambda = \alpha\sigma^2$*, i.e., it is the ridge regression solution* $\hat{\mathbf{w}} = (X^\top X + \lambda I)^{-1} X^\top \mathbf{y}$*.*

Proof. Up to constants, $-\log p(\mathcal{D} \mid \mathbf{w}) - \log p(\mathbf{w}) = \frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 + \frac{\alpha}{2}\|\mathbf{w}\|^2$; multiply by $2\sigma^2$. The minimum was computed in the exercises of Lecture 3. □

A strong prior belief that the weights are small (large α) means strong regularization. Similarly, the Laplace prior $p(\mathbf{w}) \propto e^{-\|\mathbf{w}\|_1 / b}$ leads to ℓ_1 regularization (the lasso), which favors sparse weight vectors.

12.3 Bayesian linear regression

The MAP estimate discards the uncertainty that the posterior expresses. In linear regression with a Gaussian prior, the full posterior can be computed with the tools of Section 6.

Theorem 12.5 (Posterior of Bayesian linear regression)**.** *In the model of Proposition 9.15 with the prior* $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1} I)$*, the posterior is* $\mathcal{N}(\mathbf{m}, S)$ *with*

$$S = \left(\alpha I + \sigma^{-2} X^\top X\right)^{-1}, \qquad \mathbf{m} = \sigma^{-2} S X^\top \mathbf{y}.$$

For a new input $\mathbf{x}_*$*, the predictive distribution of* $y_* = \mathbf{w}^\top \mathbf{x}_* + \varepsilon_*$ *is* $\mathcal{N}(\mathbf{m}^\top \mathbf{x}_*,\ \mathbf{x}_*^\top S \mathbf{x}_* + \sigma^2)$*.*

Proof. Up to an additive constant that does not depend on $\mathbf{w}$,

$$\log p(\mathbf{w} \mid \mathcal{D}) = -\frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 - \frac{\alpha}{2}\|\mathbf{w}\|^2 = -\frac{1}{2}\mathbf{w}^\top S^{-1}\mathbf{w} + \sigma^{-2}\mathbf{y}^\top X\mathbf{w} + \text{const}.$$

Completing the square, this is $-\frac{1}{2}(\mathbf{w} - \mathbf{m})^\top S^{-1}(\mathbf{w} - \mathbf{m}) + \text{const}$, since the linear term of the latter is $\mathbf{m}^\top S^{-1}\mathbf{w} = \sigma^{-2}\mathbf{y}^\top X\mathbf{w}$. So the posterior density is proportional to the density of $\mathcal{N}(\mathbf{m}, S)$, hence equal to it. Given the data, $\mathbf{w}^\top \mathbf{x}_*$ is $\mathcal{N}(\mathbf{m}^\top \mathbf{x}_*, \mathbf{x}_*^\top S \mathbf{x}_*)$ by Corollary 6.7, and adding the independent noise $\varepsilon_* \sim \mathcal{N}(0, \sigma^2)$ adds σ^2 to the variance. □

The posterior mean $\mathbf{m} = (X^\top X + \alpha\sigma^2 I)^{-1} X^\top \mathbf{y}$ is the ridge regression solution, which is also the MAP estimate (the mode and the mean of a Gaussian coincide). But the posterior also gives error bars: the predictive variance $\mathbf{x}_*^\top S \mathbf{x}_* + \sigma^2$ consists of the uncertainty about the weights, which shrinks as we collect more data, and the noise, which does not.

Example 12.6. For the data $(1, 2), (2, 3), (3, 5)$, the model $y = wx + \varepsilon$ without intercept, $\alpha = 1$ and $\sigma^2 = 1$: $S = (1 + 1 + 4 + 9)^{-1} = \frac{1}{15}$ and $m = \frac{1}{15}(2 + 6 + 15) = \frac{23}{15} \approx 1.53$. At $x_* = 4$, the predictive distribution is $\mathcal{N}(6.13,\ \frac{16}{15} + 1) \approx \mathcal{N}(6.13, 2.07)$.

12.4 Gaussian processes

Bayesian linear regression puts a prior on the weights of a fixed set of features. A Gaussian process puts a prior directly on the unknown function.

Definition 12.7 (Gaussian process)**.** A Gaussian process with mean function m and covariance function (or kernel) k is a random function f such that for all points $\mathbf{x}_1, \ldots, \mathbf{x}_n$, the vector $(f(\mathbf{x}_1), \ldots, f(\mathbf{x}_n))$ is Gaussian with the mean vector $(m(\mathbf{x}_i))_i$ and the covariance matrix K with entries $K_{ij} = k(\mathbf{x}_i, \mathbf{x}_j)$. We write $f \sim \mathcal{GP}(m, k)$.

For this to make sense, every such matrix K must be symmetric and positive semidefinite. This holds, for example, for $k(\mathbf{x},\mathbf{x}') = \boldsymbol{\phi}(\mathbf{x})^\top \boldsymbol{\phi}(\mathbf{x}')$ with any feature map $\boldsymbol{\phi}$, since K is then a Gram matrix (compare with the remark on kernels in Lecture 2).

Example 12.8. 1. (Linear kernel.) If $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$ with $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1} I)$, then

$$\operatorname{Cov}(f(\mathbf{x}), f(\mathbf{x}')) = \mathbf{x}^\top \operatorname{Cov}(\mathbf{w}) \mathbf{x}' = \alpha^{-1} \mathbf{x}^\top \mathbf{x}'.$$

So the prior of Bayesian linear regression is the Gaussian process with the kernel $k(\mathbf{x},\mathbf{x}') = \alpha^{-1}\mathbf{x}^\top\mathbf{x}'$.

2. (Squared exponential kernel.) $k(\mathbf{x},\mathbf{x}') = \exp\big(-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2\ell^2}\big)$ gives smooth random functions, whose values at points closer than the length scale ℓ are strongly correlated. It corresponds to infinitely many features.

3. (Brownian motion.) $k(s,t) = \min(s,t)$ for $s,t \geq 0$ is the covariance function of Brownian motion, a random function that we study in Section 14.

Proposition 12.9 (Gaussian process regression)**.** *Let $f \sim \mathcal{GP}(0,k)$, and let $y_i = f(\mathbf{x}_i) + \varepsilon_i$ for $i = 1,\dots,N$, with i.i.d. noise $\varepsilon_i \sim \mathcal{N}(0,\sigma^2)$ independent of f. Let K be the kernel matrix of the training inputs, and for a new input $\mathbf{x}_*$, let $\mathbf{k}_* = (k(\mathbf{x}_1,\mathbf{x}_*),\dots,k(\mathbf{x}_N,\mathbf{x}_*))^\top$. Then, given the data,*

$$f(\mathbf{x}_*) \sim \mathcal{N}\big(\mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{y},\ k(\mathbf{x}_*,\mathbf{x}_*) - \mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{k}_*\big).$$

Proof. The vector $(\mathbf{y}, f(\mathbf{x}_*))$ is Gaussian with mean $\mathbf{0}$ and the covariance blocks $\operatorname{Cov}(\mathbf{y}) = K+\sigma^2 I$, $\operatorname{Cov}(\mathbf{y}, f(\mathbf{x}_*)) = \mathbf{k}_*$ and $\operatorname{Var}(f(\mathbf{x}_*)) = k(\mathbf{x}_*,\mathbf{x}_*)$; now apply Theorem 6.5. □

The posterior mean $\sum_i \alpha_i k(\mathbf{x}_i,\mathbf{x}_*)$ with $\boldsymbol{\alpha} = (K+\sigma^2 I)^{-1}\mathbf{y}$ is a combination of kernel functions centered at the data points; it is also the prediction of kernel ridge regression. The posterior variance does not depend on $\mathbf{y}$: it is small near the data, and it returns to the prior variance far away from them (Figure 6). For the linear kernel, the proposition reproduces the predictive distribution of Theorem 12.5.

Example 12.10. With the squared exponential kernel ($\ell = 1$), $\sigma^2 = 0.1$, and the data $(0,1)$ and $(1,0)$: $K + \sigma^2 I = \begin{pmatrix} 1.1 & 0.607 \\ 0.607 & 1.1 \end{pmatrix}$, and at $x_* = 0.5$, $\mathbf{k}_* = (0.882, 0.882)^\top$. The posterior mean is 0.517, and the posterior variance is $1 - \mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{k}_* \approx 0.087$; a new noisy observation at x_* has the predictive variance $0.087 + 0.1 = 0.187$.

The hyperparameters, such as ℓ and σ^2, are usually chosen by maximizing the marginal likelihood $\log p(\mathbf{y}) = -\frac{1}{2}\mathbf{y}^\top (K+\sigma^2 I)^{-1}\mathbf{y} - \frac{1}{2}\log\det(K+\sigma^2 I) - \frac{N}{2}\log(2\pi)$, since $\mathbf{y} \sim \mathcal{N}(\mathbf{0}, K+\sigma^2 I)$; its gradient can be computed with the matrix calculus of Lecture 3. The cost of solving with the $N \times N$ matrix $K+\sigma^2 I$ grows like N^3, so exact Gaussian processes are limited to moderate data sets. Remarkably, one can show that neural networks with random weights become Gaussian processes in the limit of infinitely wide layers.

Remark 12.11 (Beyond conjugate models). For most models, such as neural networks, the posterior has no closed form, and the evidence $p(\mathcal{D})$ is an intractable integral. Approximate methods include Markov chain Monte Carlo sampling (Section 8) and variational inference, which approximates the posterior by a simpler distribution by minimizing a Kullback–Leibler divergence (Section 13).




Figure 6: Gaussian process regression with the squared exponential kernel ($\ell = 0.7$, $\sigma^2 = 0.01$) for six data points (red): the posterior mean (blue) and a band of ± 2 posterior standard deviations. The uncertainty is small near the data and grows back to the prior standard deviation 1 between and beyond them.

13 Information Theory

Information theory measures uncertainty and the difference between distributions. Its quantities appear in machine learning as loss functions (cross-entropy), as regularizers (the KL term of variational autoencoders), and in the analysis of learning algorithms.

Definition 13.1 (Entropy, cross-entropy and KL divergence)**.** Let $p = (p_1, \dots, p_K)$ and $q = (q_1, \dots, q_K)$ be probability distributions on $\{1, \dots, K\}$. The entropy of p, the cross-entropy of q relative to p, and the Kullback–Leibler (KL) divergence of q from p are

$$H(p) = -\sum_k p_k \log p_k, \qquad H(p, q) = -\sum_k p_k \log q_k, \qquad \mathrm{KL}(p \,\|\, q) = \sum_k p_k \log \frac{p_k}{q_k},$$

with the conventions $0 \log 0 = 0$ and $\mathrm{KL}(p \,\|\, q) = \infty$ if $q_k = 0 < p_k$ for some k.

The number $-\log p_k$ measures the surprise of seeing the outcome k: rare outcomes are surprising. The entropy is the expected surprise. With $\log_2$ instead of log, entropy is measured in bits, and it is the minimal average number of bits per symbol needed to encode outcomes of p (Shannon's source coding theorem). A fair coin has entropy 1 bit, a coin with $P(\text{heads}) = 0.9$ only 0.47 bits, and a deterministic outcome 0 bits. Clearly $H(p) \geq 0$, and

$$H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q) :$$

the cross-entropy is the average number of bits if we encode outcomes of p with a code designed for q, and the KL divergence is the number of extra bits we pay for using the wrong distribution.

Theorem 13.2 (Gibbs' inequality)**.** $\mathrm{KL}(p \,\|\, q) \geq 0$*, with equality if and only if* $p = q$*.*

Proof. We may assume $q_k > 0$ whenever $p_k > 0$. Let T be the set of k with $p_k > 0$, and let R be the random variable that takes the value $\frac{q_k}{p_k}$ with probability p_k, for $k \in T$. By Jensen's inequality for the concave function log (Theorem 3.6),

$$-\mathrm{KL}(p \,\|\, q) = \sum_{k \in T} p_k \log \frac{q_k}{p_k} = \mathbb{E}[\log R] \leq \log \mathbb{E}[R] = \log \sum_{k \in T} q_k \leq \log 1 = 0.$$

If equality holds, then $\sum_{k\in T} q_k = 1$, and equality holds in Jensen's inequality, which for the strictly concave log happens only if R is constant; so $q_k = c\,p_k$ for $k \in T$ and $q_k = 0$ otherwise, and $c = 1$ since both sum to 1. □

Two consequences: the cross-entropy $H(p, q)$ is minimal for $q = p$, and for the uniform distribution u on K outcomes, $0 \le \mathrm{KL}(p \,\|\, u) = \log K - H(p)$, so $H(p) \le \log K$: the uniform distribution has maximal entropy (compare with the exercise on maximum entropy in Lecture 3). The KL divergence is not a distance in the sense of Lecture 2, since it is not symmetric. For $p = (\frac{1}{2}, \frac{1}{2})$ and $q = (0.9, 0.1)$, $\mathrm{KL}(p \,\|\, q) \approx 0.511$ but $\mathrm{KL}(q \,\|\, p) \approx 0.368$.

Theorem 13.3 (Maximum likelihood minimizes the KL divergence)**.** *Let $x_1, \dots, x_N \in \{1, \dots, K\}$ be data with the empirical distribution $\hat{p}$, i.e., $\hat{p}_k$ is the fraction of data points equal to k, and let q_θ be a model distribution. Then*

$$\frac{1}{N}\sum_{i=1}^{N} \log q_\theta(x_i) = -H(\hat{p}, q_\theta) = -H(\hat{p}) - \mathrm{KL}(\hat{p} \,\|\, q_\theta).$$

So maximizing the likelihood is the same as minimizing the cross-entropy $H(\hat{p}, q_\theta)$, and the same as minimizing $\mathrm{KL}(\hat{p} \,\|\, q_\theta)$.

Proof. Grouping the data points by their values, $\frac{1}{N}\sum_i \log q_\theta(x_i) = \sum_k \hat{p}_k \log q_\theta(k) = -H(\hat{p}, q_\theta)$. The term $H(\hat{p})$ does not depend on θ. □

This is why the loss of classifiers is called the cross-entropy loss: for each training example, it is the cross-entropy between the one-hot distribution of the true label and the predicted distribution, and training fits the model distribution to the data in the sense of the KL divergence.

Continuous distributions. For densities, the sums become integrals: the differential entropy is $h(p) = -\int p(x) \log p(x)\, dx$, and $\mathrm{KL}(p \,\|\, q) = \int p(x) \log \frac{p(x)}{q(x)}\, dx \ge 0$. The differential entropy of $\mathcal{N}(\mu, \sigma^2)$ is $\frac{1}{2}\log(2\pi e \sigma^2)$, and one can show that among all densities with variance σ^2, the Gaussian has the largest entropy.

Proposition 13.4 (KL divergence between Gaussians)**.** *For $p = \mathcal{N}(\mu_1, \sigma_1^2)$ and $q = \mathcal{N}(\mu_2, \sigma_2^2)$,*

$$\mathrm{KL}(p \,\|\, q) = \log \frac{\sigma_2}{\sigma_1} + \frac{\sigma_1^2 + (\mu_1 - \mu_2)^2}{2\sigma_2^2} - \frac{1}{2}.$$

Proof. $\log \frac{p(x)}{q(x)} = \log \frac{\sigma_2}{\sigma_1} - \frac{(x-\mu_1)^2}{2\sigma_1^2} + \frac{(x-\mu_2)^2}{2\sigma_2^2}$. Take the expectation for $X \sim p$, using $\mathbb{E}[(X-\mu_1)^2] = \sigma_1^2$ and $\mathbb{E}[(X-\mu_2)^2] = \sigma_1^2 + (\mu_1 - \mu_2)^2$. □

In a variational autoencoder, each data point is encoded as a Gaussian $\mathcal{N}(\mu, \sigma^2)$ (in each latent dimension), and the loss contains the term $\mathrm{KL}\left(\mathcal{N}(\mu, \sigma^2) \,\|\, \mathcal{N}(0, 1)\right) = \frac{1}{2}(\mu^2 + \sigma^2 - 1 - \log \sigma^2)$, which keeps the codes close to a standard Gaussian. For example, for $\mu = 1$ and $\sigma = 2$, it is $\frac{1}{2}(4 - \log 4) \approx 1.31$.

Mutual information. The mutual information of two random variables is the KL divergence between their joint distribution and the product of their marginals,

$$I(X;Y) = \mathrm{KL}\left(p(x,y) \,\|\, p_X(x) p_Y(y)\right) = \sum_{x,y} p(x,y) \log \frac{p(x,y)}{p_X(x) p_Y(y)}.$$

By Gibbs' inequality, $I(X;Y) \ge 0$, with equality if and only if X and Y are independent. Unlike the correlation, it detects every kind of dependence. For Example 4.3, $I(X;Y) \approx 0.086$ nats.

14 Stochastic Differential Equations

Markov chains (Section 8) move in discrete steps. Many random processes change continuously in time: the position of a particle in a fluid, the price of a stock, the parameters of a network during training with stochastic gradients, or the gradually noised images of a diffusion model. Stochastic differential equations (SDEs) describe such processes by adding random noise to ordinary differential equations (Section 15 of Lecture 3). The source of the noise is Brownian motion.

Definition 14.1 (Brownian motion)**.** A Brownian motion (or Wiener process) is a family of random variables $B(t)$, $t \geq 0$, with $B(0) = 0$, such that

1. for $0 \leq t_0 < t_1 < \cdots < t_k$, the increments $B(t_1) - B(t_0), \ldots, B(t_k) - B(t_{k-1})$ are independent;
2. $B(t) - B(s) \sim \mathcal{N}(0, t - s)$ for $s < t$;
3. the paths $t \mapsto B(t)$ are continuous.

That such a process exists is a theorem of Norbert Wiener, which we do not prove. On the grid $t_k = kh$, Brownian motion is simulated exactly by the random walk $B(t_{k+1}) = B(t_k) + \sqrt{h}\, Z_k$ with i.i.d. $Z_k \sim \mathcal{N}(0, 1)$ (Figure 7). Conversely, by the central limit theorem, every random walk with many small independent steps looks like a Brownian motion from far away.

Proposition 14.2. $\mathbb{E}[B(t)] = 0$, $\mathrm{Var}(B(t)) = t$, *and* $\mathrm{Cov}(B(s), B(t)) = \min(s, t)$.

Proof. The first two statements are property (2) with $s = 0$. For $s \leq t$, $\mathrm{Cov}(B(s), B(t)) = \mathrm{Cov}\left(B(s), B(s) + (B(t) - B(s))\right) = \mathrm{Var}(B(s)) + 0 = s$, since the increment $B(t) - B(s)$ is independent of $B(s) = B(s) - B(0)$. □

So Brownian motion is a Gaussian process with mean 0 and covariance function $\min(s, t)$ (Section 12.4). Its typical displacement in a time h is $\sqrt{h}$, which is much larger than h for small h: the difference quotient $\frac{B(t+h)-B(t)}{h}$ has the variance $\frac{1}{h} \to \infty$. In fact, the paths of Brownian motion are continuous but nowhere differentiable, and ordinary calculus does not apply to them.

Definition 14.3 (Stochastic differential equation)**.** An SDE $dX = \mu(X)\, dt + \sigma(X)\, dB$, with drift μ and diffusion coefficient σ, describes a process with $X(t) = X(0) + \int_0^t \mu(X(s))\, ds + \int_0^t \sigma(X(s))\, dB(s)$, where the last integral is a stochastic (Itô) integral. Its Euler–Maruyama approximation with step h is

$$X_{k+1} = X_k + \mu(X_k)\, h + \sigma(X_k)\sqrt{h}\, Z_k, \qquad Z_k \sim \mathcal{N}(0, 1) \text{ i.i.d.}$$

We do not define the stochastic integral; for our purposes, the Euler–Maruyama scheme, which converges to the solution as $h \to 0$, can serve as the definition. It is Euler's method for the ODE $x' = \mu(x)$ (Section 15.2 of Lecture 3), plus Gaussian noise whose standard deviation is proportional to $\sqrt{h}$, and it is a Markov chain with continuous states. For $\sigma = 0$, the SDE is an ODE.

Example 14.4 (The Ornstein–Uhlenbeck process)**.** The SDE $dX = -\theta X\, dt + \sigma\, dB$ with $\theta > 0$ pulls X towards 0 while the noise pushes it around. Its Euler–Maruyama approximation is $X_{k+1} = (1 - \theta h)X_k + \sigma\sqrt{h}\, Z_k$. Its mean satisfies $\mathbb{E}[X_{k+1}] = (1 - \theta h)\mathbb{E}[X_k]$, so it follows Euler's method for the ODE $x' = -\theta x$, with the solution $e^{-\theta t}X(0)$. If $X_k \sim \mathcal{N}(0, v)$ is independent of Z_k, then $X_{k+1} \sim \mathcal{N}\big(0, (1 - \theta h)^2 v + \sigma^2 h\big)$ by Corollary 6.7. So for $0 < \theta h < 2$, the chain has the stationary distribution $\mathcal{N}(0, v_h)$ with

$$v_h = \frac{\sigma^2 h}{1 - (1 - \theta h)^2} = \frac{\sigma^2}{\theta(2 - \theta h)} \to \frac{\sigma^2}{2\theta} \qquad \text{as } h \to 0.$$

For $\theta = 1$, $\sigma = \sqrt{2}$, the stationary distribution of the SDE is $\mathcal{N}(0, 1)$ (Figure 7); for $\theta = \sigma = 1$ and $h = 0.01$, $v_h \approx 0.5025$ instead of 0.5.




Figure 7: Left: three paths of Brownian motion, simulated with $h = 0.02$. Right: three paths of the Ornstein–Uhlenbeck process $dX = -X\,dt + \sqrt{2}\,dB$, started at 2.5, -2 and 0, together with the solutions $X(0)e^{-t}$ of the ODE without noise (dashed). The process forgets its starting point and fluctuates around 0 with the stationary distribution $\mathcal{N}(0, 1)$; about 95% of the time, it stays between -2 and 2.

14.1 Langevin dynamics and diffusion models

The Ornstein–Uhlenbeck process is an example of a general principle: a gradient flow with the right amount of noise samples from a probability distribution.

Theorem 14.5 (Langevin dynamics)**.** *Let $U : \mathbb{R}^d \to \mathbb{R}$ be smooth with $\int e^{-U(\mathbf{x})}\,d\mathbf{x} < \infty$. Then the SDE $d\mathbf{X} = -\nabla U(\mathbf{X})\,dt + \sqrt{2}\,d\mathbf{B}$, where $\mathbf{B}$ is a vector of d independent Brownian motions, has the stationary density $p(\mathbf{x}) \propto e^{-U(\mathbf{x})}$, and under mild conditions, the distribution of $\mathbf{X}(t)$ converges to it.*

We do not prove the convergence, but we can see why e^{-U} is stationary. Without noise, the density p_t of $\mathbf{X}(t)$ under the flow of the vector field $-\nabla U$ satisfies the continuity equation $\frac{\partial p_t}{\partial t} = -\operatorname{div}(-p_t \nabla U)$ (Proposition 16.3 of Lecture 3). The noise adds a diffusion term, and one can show that p_t satisfies the Fokker–Planck equation

$$\frac{\partial p_t}{\partial t} = \operatorname{div}(p_t \nabla U) + \Delta p_t = \operatorname{div}(p_t \nabla U + \nabla p_t),$$

with the Laplacian $\Delta p_t = \operatorname{div} \nabla p_t$ (Lecture 3). For $p \propto e^{-U}$, we have $\nabla p = -p \nabla U$, so the right side vanishes, and p does not change in time. For $U = 0$, the Fokker–Planck equation is the heat equation $\frac{\partial p_t}{\partial t} = \Delta p_t$, which describes how the density of $\sqrt{2}\,\mathbf{B}(t)$ spreads out. For $U(x) = \frac{x^2}{2s^2}$, the SDE is the Ornstein–Uhlenbeck process with $\theta = \frac{1}{s^2}$ and $\sigma = \sqrt{2}$, whose stationary variance $\frac{\sigma^2}{2\theta} = s^2$ agrees with the density $e^{-U} \propto e^{-x^2/(2s^2)}$ of $\mathcal{N}(0, s^2)$. The Euler–Maruyama scheme gives the unadjusted Langevin algorithm

$$\mathbf{x}_{k+1} = \mathbf{x}_k - h \nabla U(\mathbf{x}_k) + \sqrt{2h}\,\mathbf{z}_k, \qquad \mathbf{z}_k \sim \mathcal{N}(\mathbf{0}, I),$$

which is gradient descent on U plus Gaussian noise. To sample from a posterior $p(\boldsymbol{\theta} \mid \mathcal{D}) \propto e^{-U(\boldsymbol{\theta})}$, we take $U(\boldsymbol{\theta}) = -\log p(\mathcal{D} \mid \boldsymbol{\theta}) - \log p(\boldsymbol{\theta})$, the training loss plus the regularizer. Without the

noise, the iteration converges to the MAP estimate; with the noise, it explores the posterior. The discretization introduces a small bias: for the target $\mathcal{N}(2, 0.25)$ and $h = 0.01$, the stationary variance of the algorithm is 0.2551 instead of 0.25, by Example 14.4. The Metropolis-adjusted Langevin algorithm removes this bias by accepting or rejecting each step as in Theorem 8.11, and stochastic gradient Langevin dynamics uses mini-batch gradients to make the method feasible for large models.

Remark 14.6 (SDEs in machine learning). 1. Diffusion models for image generation use an SDE of Ornstein–Uhlenbeck type that gradually turns images into Gaussian noise. The time-reversed process is again an SDE, whose drift involves the score $\nabla \log p_t(\mathbf{x})$ of the noised data distribution at time t. A neural network learns this score, with a variant of the score matching loss of Lecture 3, and new images are generated by simulating the reverse SDE, starting from pure noise.

2. The noise of mini-batch gradients makes SGD with a small step size behave approximately like the Euler–Maruyama scheme for an SDE $d\boldsymbol{\theta} = -\nabla L(\boldsymbol{\theta})\, dt + \sqrt{\eta\, \Sigma(\boldsymbol{\theta})}\, d\mathbf{B}$, where Σ is the covariance matrix of the mini-batch gradients. This heuristic is used to study which minima SGD prefers.

15 Key Takeaways

- A probability assigns numbers in $[0, 1]$ to events, additively for disjoint events. Conditional probability $P(A \mid B) = \frac{P(A \cap B)}{P(B)}$ and Bayes' theorem $P(B \mid A) = \frac{P(A|B)P(B)}{P(A)}$ update beliefs with evidence; ignoring the prior $P(B)$ is the base rate fallacy.

- Random variables are described by a pmf or a density. The sum rule $p(x) = \sum_y p(x, y)$ and the product rule $p(x, y) = p(y \mid x)p(x)$ are the basic rules of probabilistic modeling; independence means that the joint distribution factorizes.

- The expectation is linear; $\operatorname{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$; Markov's, Chebyshev's and Jensen's inequalities relate them. For densities, $p_Y(y) = \frac{p_X(x)}{|g'(x)|}$ under $y = g(x)$, and $F^{-1}(U)$ samples from the CDF F.

- Covariance is an inner product on centered random variables, and correlation is the cosine of their angle, so $|\operatorname{Corr}| \leq 1$. Uncorrelated does not imply independent. $\mathbb{E}[Y \mid X]$ is the best mean-squared-error predictor of Y, the orthogonal projection of Y onto functions of X.

- The covariance matrix $\Sigma = \mathbb{E}[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top]$ is symmetric positive semidefinite, $\operatorname{Cov}(A\mathbf{X} + \mathbf{b}) = A\Sigma A^\top$, and $\operatorname{Var}(\mathbf{u}^\top \mathbf{X}) = \mathbf{u}^\top \Sigma \mathbf{u}$ is maximal for the top eigenvector (PCA). Whitening $\Sigma^{-1/2}(\mathbf{X} - \boldsymbol{\mu})$ gives identity covariance.

- $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ is the distribution of $\boldsymbol{\mu} + A\mathbf{Z}$ with $AA^\top = \Sigma$ and $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$; its level sets are ellipsoids along the eigenvectors of Σ. Affine maps, marginals and conditionals of Gaussians are Gaussian, with $\mathbf{X}_a \mid \mathbf{x}_b \sim \mathcal{N}(\boldsymbol{\mu}_a + \Sigma_{ab}\Sigma_{bb}^{-1}(\mathbf{x}_b - \boldsymbol{\mu}_b), \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba})$; for jointly Gaussian vectors, uncorrelated means independent.

- The sample mean of N i.i.d. variables has standard error $\frac{\sigma}{\sqrt{N}}$; it converges to μ (law of large numbers), its fluctuations are approximately Gaussian (central limit theorem), and for bounded variables Hoeffding's inequality bounds large deviations exponentially. This underlies Monte Carlo methods, test set evaluation, and stochastic gradients. Rejection sampling draws exact

samples using an envelope $p \le Mq$ (acceptance rate $\frac{1}{M}$); importance sampling reweights samples from q by $\frac{p}{q}$, which can reduce the variance dramatically for rare events.

- A Markov chain with transition matrix P evolves by $\boldsymbol{\pi}_{t+1} = \boldsymbol{\pi}_t P$. Stationary distributions are left eigenvectors of P with eigenvalue 1; an irreducible, aperiodic chain converges to its unique stationary distribution, at a speed governed by the second largest eigenvalue in absolute value. Detailed balance $\pi_i P_{ij} = \pi_j P_{ji}$ implies stationarity; the Metropolis algorithm uses it to sample from distributions that are only known up to a constant (MCMC).

- Estimators have a bias and a variance, and MSE = bias^2 + variance; if both tend to 0, the estimator is consistent. The sample variance needs the factor $\frac{1}{N-1}$ to be unbiased. No unbiased estimator has a variance below the Cramér–Rao bound $\frac{1}{N\,I(\theta)}$, where the Fisher information $I(\theta)$ is the variance of the score $\frac{\partial}{\partial\theta}\log p(x \mid \theta)$; the MLE attains it asymptotically, and its standard error is estimated from the curvature of the log-likelihood.

- Maximum likelihood maximizes $\sum_i \log p(x_i \mid \theta)$. Least squares is maximum likelihood with Gaussian noise, and the cross-entropy loss is maximum likelihood for categorical labels; the MLE of a Gaussian is the sample mean and the (biased) sample covariance. The method of moments matches sample moments instead; it is simpler but usually less efficient. Confidence intervals $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$ quantify uncertainty.

- A test rejects H_0 if the p-value, the probability under H_0 of data at least as extreme as observed, is at most α; under H_0, p-values are uniform, so the false positive rate is α. A p-value is not the probability that H_0 is true. Many tests need corrections: Bonferroni ($\frac{\alpha}{m}$) controls the family-wise error rate, Benjamini–Hochberg the false discovery rate. The bootstrap (resampling the data with replacement) estimates standard errors of any statistic, and permutation tests (relabeling the groups) give exact p-values without distributional assumptions.

- Cross-validation estimates the generalization error, but its naive standard error is too small, and the best of many cross-validated models is optimistically biased; compare models with paired differences and evaluate the final choice on untouched data. A/B tests randomize to measure causal effects; plan the sample size $n \approx (z_{1-\alpha/2} + z_{1-\beta})^2\, 2\bar{p}(1-\bar{p})/\delta^2$ per group in advance, and do not stop at the first significant peek.

- Bayesian inference updates a prior to the posterior $p(\theta \mid \mathcal{D}) \propto p(\mathcal{D} \mid \theta)p(\theta)$. Conjugate priors (Beta for Bernoulli, Gaussian for linear regression) give closed-form posteriors. MAP estimation adds the regularizer $-\log p(\theta)$: a Gaussian prior gives ridge regression, a Laplace prior the lasso. The Bayesian posterior also gives predictive uncertainty. A Gaussian process is a prior on functions given by a kernel; its posterior follows from Gaussian conditioning, with the mean $\mathbf{k}_*^\top (K + \sigma^2 I)^{-1}\mathbf{y}$ and a variance that is small near the data.

- Entropy measures uncertainty, cross-entropy $H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q)$, and $\mathrm{KL}(p \,\|\, q) \ge 0$ with equality only for $p = q$ (Gibbs). Maximum likelihood minimizes the cross-entropy and the KL divergence from the empirical distribution to the model.

- Brownian motion has independent Gaussian increments $B(t) - B(s) \sim \mathcal{N}(0, t-s)$ and continuous, nowhere differentiable paths. An SDE $dX = \mu(X)\,dt + \sigma(X)\,dB$ is simulated by Euler–Maruyama, $X_{k+1} = X_k + \mu(X_k)h + \sigma(X_k)\sqrt{h}\,Z_k$: noise of size $\sqrt{h}$, not h. Langevin dynamics $d\mathbf{X} = -\nabla U\,dt + \sqrt{2}\,d\mathbf{B}$ samples from $p \propto e^{-U}$, the stationary solution of the Fokker–Planck equation; its discretization is gradient descent plus noise, and diffusion models generate data by reversing a noising SDE.

16 Exercises

1. (The birthday problem.) Assume that birthdays are independent and uniformly distributed over 365 days. Show that the probability that among n people at least two share a birthday is $1 - \frac{365 \cdot 364 \cdots (365-n+1)}{365^n}$, and that it exceeds $\frac{1}{2}$ for $n = 23$. How many people are needed for 99%?

2. (a) Prove the inclusion–exclusion formula $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$.

   (b) Two fair coins are flipped. Let $A =$ "the first coin shows heads", $B =$ "the second coin shows heads", and $C =$ "exactly one coin shows heads". Show that any two of these events are independent, but the three events are not independent.

3. A spam filter knows that 40% of all emails are spam, that the word "free" appears in 30% of the spam emails and in 2% of the other emails. What is the probability that an email containing "free" is spam?

4. Let X be Geometric(p). Show that $\mathbb{E}[X] = \frac{1}{p}$ and that $P(X > s + t \mid X > s) = P(X > t)$ for all positive integers s and t. How many times do you expect to roll a fair die until you see a six?

5. Let U and V be independent and Uniform$(0, 1)$. Find the densities of (a) U^2, (b) $-\log U$, (c) $\max(U, V)$ (Hint: compute the CDF first.)

6. Let X be Poisson(λ). Compute $\mathbb{E}[X(X-1)]$, and use it to show that $\mathrm{Var}(X) = \lambda$.

7. (a) A nonnegative random variable has mean 2. Bound $P(X \geq 10)$ with Markov's inequality, and compare with the exact value for $X \sim$ Exponential($\frac{1}{2}$).

   (b) Use Jensen's inequality to show that $\mathbb{E}[\frac{1}{X}] \geq \frac{1}{\mathbb{E}[X]}$ for every positive random variable X, and that $\mathbb{E}[e^X] \geq e^{\mathbb{E}[X]}$.

8. Let (X, Y) have the joint pmf $p(0,0) = 0.1$, $p(0,1) = 0.2$, $p(0,2) = 0.1$, $p(1,0) = 0.2$, $p(1,1) = 0.1$, $p(1,2) = 0.3$. Compute the marginal distributions, $\mathbb{E}[Y \mid X = 0]$ and $\mathbb{E}[Y \mid X = 1]$, and check the tower property. Compute $\mathrm{Cov}(X, Y)$. Are X and Y independent?

9. Let $X \sim \mathcal{N}(0, 1)$. Show that X and X^2 are uncorrelated, but not independent. Show that $\mathrm{Corr}(X, aX + b) = \mathrm{sign}(a)$ for $a \neq 0$.

10. Let $\mathbf{X}$ have the covariance matrix $\Sigma = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$. Compute $\mathrm{Var}(X_1 + X_2 + X_3)$ and $\mathrm{Var}(X_1 - X_3)$. In which direction $\mathbf{u}$ (with $\|\mathbf{u}\| = 1$) is $\mathrm{Var}(\mathbf{u}^\top \mathbf{X})$ largest, and what is it? Find a matrix A such that $A\mathbf{X}$ has covariance matrix I.

11. Let $(X, Y) \sim \mathcal{N}\Big(\mathbf{0}, \begin{pmatrix} 4 & 2 \\ 2 & 2 \end{pmatrix}\Big)$. Find the conditional distributions of Y given $X = x$ and of X given $Y = y$, the distribution of $X - Y$, and $P(X > Y)$. Describe the level curves of the density.

12. Let $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ in $\mathbb{R}^d$. Show that the squared Mahalanobis distance $(\mathbf{X} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{X} - \boldsymbol{\mu})$ has the same distribution as $\|\mathbf{Z}\|^2$ with $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$, and that its expectation is d. What does this say about where the probability mass of a high-dimensional Gaussian lies?

13. How many times do we have to flip a fair coin so that the fraction of heads is within 0.01 of $\frac{1}{2}$ with probability at least 95%? Answer with Chebyshev's inequality, with Hoeffding's inequality, and with the central limit theorem, and compare.

14. Let $p(x) = \sqrt{2/\pi}\, e^{-x^2/2}$ for $x > 0$ be the density of $|Z|$ for $Z \sim \mathcal{N}(0,1)$, and let $q(x) = e^{-x}$ be the Exponential(1) density.

    (a) Find the smallest M with $p \le Mq$, and the acceptance rate of rejection sampling with this proposal.

    (b) Write down the importance sampling estimator of $\mathbb{E}_p[X^2]$ with samples from q, and explain why its variance is finite.

15. Consider the Markov chain with the transition matrix

$$P = \begin{pmatrix} 0.7 & 0.2 & 0.1 \\ 0.3 & 0.5 & 0.2 \\ 0.2 & 0.3 & 0.5 \end{pmatrix}.$$

    Draw its transition graph, compute $P(X_2 = 3 \mid X_0 = 1)$, and find the stationary distribution. Is the chain irreducible and aperiodic? In the long run, what fraction of the time does it spend in state 1?

16. (Random walks on graphs.)

    (a) Let the walker of Example 8.9 move on the cycle with the four vertices $1, 2, 3, 4$ (and the edges $\{1,2\}, \{2,3\}, \{3,4\}, \{4,1\}$). Find the stationary distribution and the eigenvalues of P. Show that the distribution of X_t does not converge if $X_0 = 1$.

    (b) The lazy walk stays at the current vertex with probability $\frac{1}{2}$, i.e., it has the transition matrix $\frac{1}{2}(I + P)$. Show that it has the same stationary distribution, find its eigenvalues, and explain why it converges.

17. Let $X_1, \ldots, X_N$ be i.i.d. Uniform$(0, \theta)$.

    (a) Show that the MLE of θ is $M = \max_i X_i$, and that $\mathbb{E}[M] = \frac{N}{N+1}\theta$. (Hint: $P(M \le t) = (t/\theta)^N$.)

    (b) Show that $\frac{N+1}{N}M$ and $2\bar{X}_N$ are unbiased, and compare their variances, $\frac{\theta^2}{N(N+2)}$ and $\frac{\theta^2}{3N}$. Which estimator is better?

18. Find the maximum likelihood estimates for i.i.d. data from (a) Poisson(λ), (b) Exponential(λ), (c) Geometric(p).

19. (Fisher information.)

    (a) Show that the Fisher information of Poisson(λ) is $I(\lambda) = \frac{1}{\lambda}$, and that the MLE $\bar{X}_N$ is efficient.

    (b) Show that $I(\lambda) = \frac{1}{\lambda^2}$ for Exponential(λ). The sum S of $N \ge 3$ i.i.d. Exponential(λ) variables has the Gamma density $\frac{\lambda^N s^{N-1} e^{-\lambda s}}{(N-1)!}$ for $s > 0$. Use it to show that the MLE $\frac{1}{\bar{X}_N}$ has the expectation $\frac{N\lambda}{N-1}$, that $\frac{N-1}{N\bar{X}_N}$ is unbiased, and that its variance $\frac{\lambda^2}{N-2}$ is larger than the Cramér–Rao bound. Is the MLE asymptotically efficient?

(c) Counts of $N = 50$ independent Poisson observations have the sample mean 3. Estimate the standard error of $\hat{\lambda}$ from the observed information $-\ell''(\hat{\lambda})$, and give an approximate 95% confidence interval for λ.

20. Assume the regression model $y_i = \mathbf{w}^\top \mathbf{x}_i + \varepsilon_i$ with i.i.d. Laplace noise, with density $\frac{1}{2b} e^{-|\varepsilon|/b}$. Show that the MLE of $\mathbf{w}$ minimizes $\sum_i |y_i - \mathbf{w}^\top \mathbf{x}_i|$. In the special case $\mathbf{x}_i = 1$ (estimating a single number w), show that every median of the y_i is a minimizer.

21. Find the method of moments estimators of (a) a and b for Uniform(a, b), using the mean and the variance; (b) the shape k and the scale s of a Gamma distribution, for data with the sample mean 4 and the sample variance 8.

22. A die is rolled 120 times and shows 30 sixes. Test at the level 0.05 whether the die favors sixes: state H_0 and H_1, and compute the exact one-sided p-value $P(S \geq 30)$ for $S \sim \text{Binomial}(120, \frac{1}{6})$ with a computer, and its normal approximation.

23. (a) Show that for m independent tests of true null hypotheses at the level $\frac{\alpha}{m}$, the probability of at least one rejection is $1 - (1 - \frac{\alpha}{m})^m \leq \alpha$.

    (b) In a permutation test with two groups of three observations each, what is the smallest possible one-sided p-value? What does this mean for very small experiments?

    (c) Show that the bootstrap standard error of the sample mean (for $B \to \infty$) is $\frac{\hat{\sigma}}{\sqrt{N}}$ with $\hat{\sigma}^2 = \frac{1}{N} \sum_i (x_i - \bar{x})^2$.

24. An online shop wants to test whether a new design increases the click-through rate from 10% to 11%. How many users per group are needed for a two-sided test at the level 0.05 with power 0.8? After the experiment, the two groups of 15 000 users have 1500 and 1650 clicks. Compute the z-statistic, the p-value, and a 95% confidence interval for the difference.

25. With the uniform prior Beta$(1, 1)$, a coin shows 7 heads in 10 flips. Find the posterior distribution, its mean, the MAP estimate, the MLE, and the predictive probability of heads in the next flip.

26. (Gaussian–Gaussian conjugacy.) Let $x_1, \dots, x_N$ be i.i.d. $\mathcal{N}(\mu, \sigma^2)$ with known σ^2, and let the prior be $\mu \sim \mathcal{N}(\mu_0, \tau^2)$. Show that the posterior is Gaussian with precision $\frac{1}{\tau^2} + \frac{N}{\sigma^2}$ and mean

$$\frac{\mu_0/\tau^2 + N\bar{x}/\sigma^2}{1/\tau^2 + N/\sigma^2},$$

    a weighted average of the prior mean and the sample mean. What happens as $N \to \infty$?

27. (Gaussian processes and Bayesian linear regression.)

    (a) Show that $(X^\top X + \lambda I) X^\top = X^\top (X X^\top + \lambda I)$ for every matrix X and $\lambda > 0$, and conclude that $(X^\top X + \lambda I)^{-1} X^\top = X^\top (X X^\top + \lambda I)^{-1}$.

    (b) Use (a) to show that for the linear kernel $k(\mathbf{x}, \mathbf{x}') = \alpha^{-1} \mathbf{x}^\top \mathbf{x}'$, the posterior mean of Proposition 12.9 equals $\mathbf{m}^\top \mathbf{x}_*$ from Theorem 12.5. Which of the two formulas is cheaper to evaluate when $N \gg d$, and which when $d \gg N$?

    (c) Let $f \sim \mathcal{GP}(0, k)$ with $k(s, t) = \min(s, t)$, and observe $f(1) = 1$ without noise. Find the posterior mean and variance of $f(t)$ for $0 \leq t \leq 1$ and for $t > 1$, and sketch them.

28. (a) Compute the entropy of a fair die in bits.

(b) Compute KL(Bernoulli(p) $\|$ Bernoulli(q)), and show that

$$\mathrm{KL}\left(\mathcal{N}(\mu_1,\sigma^2)\,\|\,\mathcal{N}(\mu_2,\sigma^2)\right)=\frac{(\mu_1-\mu_2)^2}{2\sigma^2}.$$

(c) Show that $I(X;Y) = H(X) + H(Y) - H(X,Y)$ for discrete random variables, where $H(X,Y)$ is the entropy of the joint distribution. Compute $I(X;Y)$ for Exercise 8.

29. (Brownian motion and the Ornstein–Uhlenbeck process.)

(a) Compute $\mathrm{Var}(B(2)-B(0.5))$, $\mathrm{Cov}(B(1),B(3))$ and $\mathrm{Corr}(B(1),B(3))$.

(b) Let $c>0$. Show that $W(t)=\frac{1}{\sqrt{c}}B(ct)$ has the same mean and covariance functions as $B(t)$. (In fact, W is again a Brownian motion: zooming into a Brownian path by the factor c in time and $\sqrt{c}$ in space gives a Brownian path.)

(c) Let $X_{k+1}=(1-\theta h)X_k+\sigma\sqrt{h}\,Z_k$ with $X_0=x_0$ and i.i.d. $Z_k\sim\mathcal{N}(0,1)$. Show that $\mathbb{E}[X_k]=(1-\theta h)^k x_0$ and $\mathrm{Var}(X_k)=v_h\big(1-(1-\theta h)^{2k}\big)$, with v_h from Example 14.4. What happens for $\theta h>2$?

17 Programming Exercises

As in the previous lectures, implement the methods yourself with NumPy, and use library routines (such as `rng.normal` or `scipy.stats`) only to check your results. Start each file with

```
import numpy as np
import matplotlib.pyplot as plt
rng = np.random.default_rng(0)
```

P1. Monte Carlo estimation. Estimate π from N uniform random points in $[-1,1]^2$, for $N=10^2,10^3,\dots,10^7$. Plot the absolute error against N on a log-log scale, together with the standard error $4\sqrt{\frac{\pi}{4}(1-\frac{\pi}{4})/N}$. What is the slope? Then estimate the volume of the unit ball in $\mathbb{R}^d$ for $d=2,5,10,20$ in the same way. What goes wrong in high dimensions?

P2. Sampling. Generate 10^5 samples from each of the following distributions, using only `rng.random()` (uniform samples), and compare histograms with the densities.

- Exponential(λ) by inverse transform sampling (Theorem 2.7).
- $\mathcal{N}(0,1)$ with the Box–Muller method: for independent $U_1,U_2\sim\mathrm{Uniform}(0,1)$, the numbers $\sqrt{-2\log U_1}\cos(2\pi U_2)$ and $\sqrt{-2\log U_1}\sin(2\pi U_2)$ are independent standard normal.
- $\mathcal{N}(\boldsymbol{\mu},\Sigma)$ in $\mathbb{R}^2$ with $\Sigma=\begin{pmatrix}3&1\\1&3\end{pmatrix}$, via $\boldsymbol{\mu}+L\mathbf{z}$ with the Cholesky factor L. Plot the samples together with the ellipses of Figure 2, and check the sample mean and covariance matrix. What fraction of the samples lies inside the ellipse with $r=1$, and with $r=2$? (Compare with $1-e^{-r^2/2}$.)

P3. The law of large numbers and the central limit theorem. For $X_i\sim$ Exponential(1), simulate 10^4 sample means $\bar{X}_N$ for $N=1,2,5,30$, and plot histograms of the standardized means $\sqrt{N}(\bar{X}_N-1)$ together with the standard normal density. Repeat with the Cauchy distribution (the ratio of two independent standard normal variables), which has no mean: plot $\bar{X}_N$ against N for one long sequence. What happens?

P4. Gaussian classifiers. Generate two classes in $\mathbb{R}^2$ from Gaussians with means $(0, 0)$ and $(2, 1)$ and the same covariance matrix $\Sigma = \begin{pmatrix} 1 & 0.5 \\ 0.5 & 1 \end{pmatrix}$, with 200 points each.

- Estimate the means and the shared covariance matrix by maximum likelihood (Theorem 9.9).
- Show (on paper) that with equal class priors, Bayes' theorem gives $P(y = 1 \mid \mathbf{x}) = \sigma(\mathbf{w}^\top \mathbf{x} + b)$ with $\mathbf{w} = \Sigma^{-1}(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0)$ and $b = -\frac{1}{2}(\boldsymbol{\mu}_1^\top \Sigma^{-1} \boldsymbol{\mu}_1 - \boldsymbol{\mu}_0^\top \Sigma^{-1} \boldsymbol{\mu}_0)$: a logistic regression model.
- Plot the data and the decision boundary, and compare with logistic regression trained by gradient descent (Lecture 3).

P5. Maximum likelihood and bias. Draw 10^5 data sets of size $N = 5$ from $\mathcal{N}(2, 9)$. Compute the average of the MLE $\hat{\sigma}^2$ and of s^2 over the data sets, and compare with $\frac{N-1}{N}\sigma^2 = 7.2$ and $\sigma^2 = 9$. Then estimate the mean squared errors of both estimators. Which one has the smaller MSE?

P6. The bootstrap. For $N = 50$ samples from Exponential(1), estimate the standard error of the sample median by resampling the data 2000 times with replacement and computing the median of each resample. Compare with the standard error of the median over many independent data sets. Use the same method to compute a 95% interval for the mean (the 2.5% and 97.5% quantiles of the bootstrapped means), and compare with $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$.

P7. Bayesian updating and Thompson sampling.

- Simulate 100 flips of a coin with $\theta = 0.3$, and plot the Beta posterior (starting from Beta$(1, 1)$) after $0, 1, 5, 20$ and 100 flips, as in Figure 5.
- A slot machine has three arms with unknown success probabilities 0.3, 0.5 and 0.6. Thompson sampling keeps a Beta posterior for each arm, and in each round draws one sample from each posterior and plays the arm with the largest sample. Simulate 1000 rounds, and plot how often each arm is played. Compare the total reward with always playing a random arm and with always playing the best arm.

P8. Bayesian linear regression. Generate 20 points $y = \sin(3x) + \varepsilon$ with x uniform in $[-1, 1]$ and $\varepsilon \sim \mathcal{N}(0, 0.1^2)$, and use the features $1, x, \ldots, x^7$. Compute the posterior of Theorem 12.5 with $\alpha = 1$ and $\sigma = 0.1$. Plot the data, the posterior mean, the band of ± 2 predictive standard deviations, and a few functions with weights sampled from the posterior. Where is the uncertainty largest? Check that the posterior mean equals the ridge solution with $\lambda = \alpha\sigma^2$.

P9. Cross-entropy and KL divergence.

- Write functions for $H(p)$, $H(p, q)$ and $\mathrm{KL}(p \,\|\, q)$, and check $H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q)$ and $\mathrm{KL} \geq 0$ for random distributions (e.g., `rng.dirichlet(np.ones(5))`).
- Let p be the mixture $\frac{1}{2}\mathcal{N}(-2, 0.5^2) + \frac{1}{2}\mathcal{N}(2, 0.5^2)$. On a grid, find the Gaussian $q = \mathcal{N}(\mu, s^2)$ that minimizes $\mathrm{KL}(p \,\|\, q)$, and the one that minimizes $\mathrm{KL}(q \,\|\, p)$ (try several starting points). Plot both. Why does the first cover both modes, while the second picks one of them?

P10. Gaussian process regression (Section 12.4), with the squared exponential kernel $k(x, x') = \exp(-\frac{(x-x')^2}{2\ell^2})$.

- Sample and plot a few functions from the prior $\mathcal{GP}(0, k)$ on a grid of 200 points in $[-3, 3]$, for $\ell = 0.1, 0.3, 1$. (Add $10^{-8} I$ to the kernel matrix before the Cholesky factorization.) Do the same for the Brownian motion kernel $\min(s, t)$ (Section 14) on $[0, 3]$.
- For 8 noisy observations $y_i = \sin(3x_i) + \varepsilon_i$ ($\varepsilon_i \sim \mathcal{N}(0, 0.1^2)$), compute the posterior mean and variance of Proposition 12.9 on the grid, and plot them with a band of ± 2 standard deviations, as in Figure 6. Use `np.linalg.solve` (or a Cholesky factorization) rather than the inverse. Sample a few functions from the posterior.
- Compute the log marginal likelihood $\log p(\mathbf{y})$ for ℓ on a grid from 0.05 to 3, and plot it. Which length scale does it select? Compare the posterior with the one for $\ell = 0.1$ and $\ell = 1$.

P11. Markov chains and MCMC.

- (A bigram language model.) Take a text of a few thousand words, split it into words, and estimate the transition probabilities $P(\text{next word} = j \mid \text{current word} = i)$ by counting. Generate random sentences from the chain. Compute the stationary distribution by power iteration, and compare it with the word frequencies in the text.
- (Metropolis sampling.) For the probability θ of heads, use the prior density $p(\theta) \propto e^{-10(\theta - 0.5)^2}$ on $[0, 1]$, which is not a Beta distribution, and the data "7 heads in 10 flips". Sample from the posterior $p(\theta \mid \mathcal{D}) \propto \theta^7 (1-\theta)^3 e^{-10(\theta-0.5)^2}$ with the Metropolis algorithm, using the proposals $\theta' = \theta + \eta\varepsilon$ with $\varepsilon \sim \mathcal{N}(0, 1)$ (and target 0 outside $[0, 1]$). Compare the histogram of 50 000 samples with the posterior density, normalized numerically on a grid. Plot the chain for $\eta = 0.01$, 0.1 and 1, and report the acceptance rates. Which step size works best, and why?

P12. Rejection and importance sampling.

- Sample from the Beta$(2, 5)$ distribution by rejection sampling with the uniform proposal. Find the best M numerically, compare the acceptance rate with $\frac{1}{M}$, and compare a histogram with the density.
- Estimate $P(Z > 4)$ and $P(Z > 6)$ for $Z \sim \mathcal{N}(0, 1)$ with plain Monte Carlo and with importance sampling from $\mathcal{N}(a, 1)$ for $a = 4$ (or 6), using $N = 10^4$ samples. Repeat 100 times to estimate the relative standard errors, and compare with the exact values $1 - \Phi(4)$ and $1 - \Phi(6)$. What happens with the proposal $\mathcal{N}(0, 1)$ shifted to $a = 8$? Compute the effective sample size $\frac{(\sum_i w_i)^2}{\sum_i w_i^2}$ in each case.

P13. Hypothesis tests by simulation.

- Simulate 10 000 data sets of size $N = 30$ from $\mathcal{N}(0, 1)$, compute the two-sided z-test p-value for H_0: $\mu = 0$ in each, and plot the histogram of the p-values. Repeat with data from $\mathcal{N}(0.5, 1)$, and estimate the power of the test at the level 0.05.
- Generate 1000 tests, of which 50 have a real effect (data from $\mathcal{N}(0.8, 1)$) and 950 do not (data from $\mathcal{N}(0, 1)$), each with $N = 20$. Count the true and false discoveries without correction, with Bonferroni, and with Benjamini–Hochberg at the level 0.05.
- Implement a permutation test for the difference of the means of two groups, and apply it to two groups of 15 samples from Exponential(1) and Exponential(0.6). Compare its p-value with the z-test.

- (Peeking.) Simulate A/B tests with two identical versions (conversion rate 5%) and up to 10 000 users per group, computing the two-proportion z-test after every 200 users per group. What fraction of the experiments ever reaches $p < 0.05$? Compare with a single test at the end.

P14. Cross-validation. Use the polynomial regression data of the programming exercises of Lecture 2 (20 points $y = \sin(3x) + \varepsilon$).

- Implement k-fold cross-validation, and choose the polynomial degree from 1 to 12 with 5-fold cross-validation. Plot the cross-validation error and the training error against the degree.
- Repeat the cross-validation with 50 different random splits. How much does the selected degree, and the cross-validation error of a fixed degree, vary? Compare the spread with the naive standard error $\frac{s_e}{\sqrt{k}}$.
- Generate a large independent test set from the same distribution. Is the cross-validation error of the selected degree optimistic? Implement nested cross-validation and compare.

P15. Stochastic differential equations.

- Simulate 10 000 Brownian paths on $[0, 1]$ with the step $h = 0.001$. Check $\mathrm{Var}(B(t)) = t$ and $\mathrm{Cov}(B(0.3), B(0.8)) = 0.3$ empirically, and plot a few paths. Zoom into one path: plot it on $[0.5, 0.5 + \delta]$ for $\delta = 0.1, 0.01, 0.001$ (simulating the path with a step much smaller than δ). Why does it never look smooth?
- Simulate the Ornstein–Uhlenbeck process $dX = -\theta X\,dt + \sigma\,dB$ with Euler–Maruyama for $\theta = 1$, $\sigma = \sqrt{2}$, starting 10 000 paths at $x_0 = 3$. Plot the mean and the variance of $X(t)$ against t, together with $x_0 e^{-t}$ and $1 - e^{-2t}$. Compare the histogram at $t = 5$ with $\mathcal{N}(0, 1)$, and check v_h from Example 14.4 for $h = 0.5$.
- (Langevin sampling.) Use the unadjusted Langevin algorithm to sample from the mixture $p = \frac{1}{2}\mathcal{N}(-2, 0.5^2) + \frac{1}{2}\mathcal{N}(2, 0.5^2)$, i.e., $U = -\log p$ (compute ∇U by hand). Compare a histogram of one long chain with p, and count how often the chain moves between the two modes. Then sample the posterior of the logistic regression model of Lecture 3 on a small two-dimensional data set, with the prior $\mathcal{N}(\mathbf{0}, I)$, and plot the samples of $\mathbf{w}$ together with the MAP estimate.