Math for Machine Learning

Lecture Notes

Arystan Shokan

Linear Algebra · Analytic Geometry
Vector Calculus and Optimization · Probability and Statistics

November 14, 2026

Contents

1 Linear Algebra **1**

Introduction . . . 1
1.1 Number Sets and Fields . . . 2
1.2 Vector Spaces . . . 3
1.3 Bases and Dimension . . . 6
1.4 Linear Transformations . . . 9
1.5 Subspaces . . . 17
1.6 Matrices . . . 20
1.7 Null Space, Column Space and Rank . . . 52
1.8 Dual Vector Spaces . . . 55
1.9 Quotient Spaces and the Isomorphism Theorems . . . 60
1.10 Eigenvalues and Eigenvectors . . . 64
1.11 Tensors . . . 73
1.12 Key Takeaways . . . 81
1.13 Exercises . . . 83
1.14 Programming Exercises . . . 87

2 Analytic Geometry **95**

Introduction . . . 95
2.1 Norms . . . 95
2.2 Inner Products . . . 97
2.3 Lengths and Distances . . . 100
2.4 Angles and Orthogonality . . . 102
2.5 Orthonormal Bases . . . 104
2.6 Orthogonal Complements . . . 107
2.7 Inner Products of Functions . . . 109
2.8 Orthogonal Projections . . . 110
2.9 Rotations . . . 115
2.10 The Spectral Theorem . . . 117
2.11 The Singular Value Decomposition . . . 122
2.12 Key Takeaways . . . 128
2.13 Exercises . . . 129
2.14 Programming Exercises . . . 132

3 Vector Calculus and Optimization **137**

Introduction . . . 137
3.1 Multivariable Functions . . . 138
3.2 Derivatives of Functions of One Variable . . . 140
3.3 Partial Derivatives and Gradients . . . 143
3.4 Jacobians: Derivatives of Vector-Valued Functions . . . 146

3.5 The Chain Rule . . . 148
3.6 Gradients with Respect to Matrices . . . 151
3.7 Backpropagation and Automatic Differentiation . . . 154
3.8 Higher Derivatives and Taylor's Theorem . . . 157
3.9 Unconstrained Optimization . . . 159
3.10 Gradient Descent and Newton's Method . . . 162
3.11 Convexity . . . 169
3.12 Constrained Optimization . . . 175
3.13 Vector Fields: Divergence and Curl . . . 180
3.14 Multiple Integrals and the Divergence Theorem . . . 184
3.15 Ordinary Differential Equations . . . 189
3.16 Flows, Volumes and Densities . . . 191
3.17 Key Takeaways . . . 193
3.18 Exercises . . . 195
3.19 Programming Exercises . . . 199

4 Probability and Statistics 205
Introduction . . . 205
4.1 Probability Spaces . . . 206
4.2 Random Variables and Their Distributions . . . 208
4.3 Expectation and Variance . . . 210
4.4 Joint Distributions and Independence . . . 212
4.5 Random Vectors and Covariance Matrices . . . 219
4.6 The Gaussian Distribution . . . 220
4.7 Limit Theorems and Monte Carlo Methods . . . 223
4.8 Markov Chains . . . 230
4.9 Parameter Estimation . . . 239
4.10 Hypothesis Testing . . . 249
4.11 Evaluating Models and Experiments . . . 255
4.12 Bayesian Inference . . . 259
4.13 Information Theory . . . 263
4.14 Latent Variable Models and the EM Algorithm . . . 264
4.15 Stochastic Differential Equations . . . 268
4.16 Key Takeaways . . . 271
4.17 Exercises . . . 274
4.18 Programming Exercises . . . 281

A Topology, Differential Topology and Functional Analysis 287
Introduction . . . 287
A.1 Metric Spaces and Open Sets . . . 288
A.2 Continuity and Topological Spaces . . . 289
A.3 Compactness . . . 290
A.4 Connectedness . . . 292
A.5 The Inverse Function Theorem . . . 293
A.6 The Implicit Function Theorem . . . 294
A.7 Submanifolds and Tangent Spaces . . . 295
A.8 Lagrange Multipliers and Optimization on Manifolds . . . 298
A.9 Critical Points and Morse Theory . . . 300
A.10 Differential Forms and Stokes' Theorem . . . 302
A.11 Topology in Machine Learning . . . 313
A.12 Banach Spaces . . . 313

A.13 Hilbert Spaces . . . 315
A.14 Compact Operators and the Spectral Theorem . . . 318
A.15 Reproducing Kernel Hilbert Spaces . . . 320
A.16 Functional Analysis in Machine Learning . . . 321
Key Takeaways . . . 322
Exercises . . . 323
Programming Exercises . . . 326

B Measure-Theoretic Probability **329**
Introduction . . . 329
B.1 Measures and Probability Spaces . . . 329
B.2 Random Variables and Their Distributions . . . 331
B.3 The Lebesgue Integral and Expectation . . . 332
B.4 Convergence Theorems . . . 334
B.5 Densities and the Radon–Nikodym Theorem . . . 335
B.6 Product Measures and Independence . . . 336
B.7 Conditional Expectation . . . 337
B.8 Martingales . . . 338
B.9 Modes of Convergence and the Strong Law . . . 339
Key Takeaways . . . 341
Exercises . . . 342
Programming Exercises . . . 343

Lecture 1

Linear Algebra

Introduction

Machine learning works with data, and data are almost always stored as lists of numbers: an image is a list of pixel intensities, a customer is a list of features such as age and income, and a word is represented by an embedding, a list of a few hundred numbers. Linear algebra is the mathematics of such lists, which we call vectors, and of the maps between them that are compatible with adding and scaling, the linear transformations. It is the language of the whole subject:

- a data set of N examples with n features each is an $N \times n$ matrix, and a linear model predicts a number from a feature vector $\mathbf{x}$ by $w_1 x_1 + \cdots + w_n x_n$; data with more indices, such as a batch of color images, are stored as tensors;
- a layer of a neural network multiplies its input vector by a weight matrix, so a network is built from matrix multiplications and simple nonlinear functions;
- fitting a linear model to data leads to a system of linear equations, and whether its solution is unique depends on the rank of the data matrix;
- eigenvalues and eigenvectors describe what happens when a linear map is applied again and again, as in Markov chains and the PageRank algorithm, and they are the basis of principal component analysis (Lecture 2).

In this lecture:

- we start from fields, the number systems in which we can add, subtract, multiply and divide, and define vector spaces over them, including spaces of functions such as the continuous and the smooth functions, with bases, dimension and subspaces;
- we study linear transformations, including differentiation and integration, and the matrices that represent them: matrix operations, row reduction for solving linear systems and computing inverses, special types of matrices, the trace and the determinant, which measures how a linear map scales areas and volumes, and block matrices, with the LU decomposition behind every linear solver and the Woodbury identity for low-rank updates;
- we relate the null space, the column space and the rank of a matrix by the rank–nullity theorem;
- we introduce dual spaces, which explain what the transpose of a matrix is, and quotient spaces with the isomorphism theorems;

- we find eigenvalues and eigenvectors, diagonalize matrices, prove the Cayley–Hamilton theorem, characterize diagonalizable matrices by their minimal polynomials, use powers of matrices to analyze Fibonacci numbers, Markov chains and power iteration, and show that the discrete Fourier transform diagonalizes convolutions, the linear maps behind convolutional networks;
- we construct the tensor product of vector spaces, define tensors as its elements, and derive their description by arrays (the tensors of deep learning libraries), multilinear forms, the trace as a contraction, tensor rank, and Einstein notation.

Lecture 2 adds geometry to vector spaces: lengths, angles and orthogonality. The later lectures use the language of this lecture throughout: derivatives are linear maps (Lecture 3), covariance matrices describe random vectors (Lecture 4), and the eigenvalues of a matrix decide whether the solutions of a linear differential equation grow or decay (Lecture 3).

Notation. Vectors of $\mathbb{R}^n$ are written in bold, $\mathbf{x} = (x_1, \ldots, x_n)$, with their entries in normal font; so are the zero vector $\mathbf{0}$ and the standard basis vectors $\mathbf{e}_1, \ldots, \mathbf{e}_n$. Elements of general vector spaces, such as polynomials and functions, are denoted by lowercase letters in normal font, such as u, v and f. Scalars are denoted by letters such as a, b and λ, and matrices by capital letters such as A. The same conventions are used in all lectures.

1.1 Number Sets and Fields

Remark 1.1.1. When we want to say that a number x is in a set S, we write $x \in S$. When we want to say that a number x is not in a set S, we write $x \notin S$.

Definition 1.1.2 (Number sets)**.** $\mathbb{Z}$ is the set of all integers. Meaning,

$$\mathbb{Z} = \{\ldots, -2, -1, 0, 1, 2, \ldots\}.$$

$\mathbb{Q}$ is the set of all rational numbers. Meaning,

$$\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z}, q \neq 0 \right\}.$$

$\mathbb{R}$ is the set of all real numbers. Meaning, $\mathbb{R}$ is obtained from $\mathbb{Q}$ by "filling in the gaps": $\mathbb{R} \supset \mathbb{Q}$, every Cauchy sequence in $\mathbb{Q}$ has a limit in $\mathbb{R}$ (completeness), and every real number is the limit of some Cauchy sequence in $\mathbb{Q}$. In other words, $\mathbb{R}$ is the completion of $\mathbb{Q}$.

$\mathbb{C}$ is the set of all complex numbers. Meaning,

$$\mathbb{C} = \{a + bi : a, b \in \mathbb{R}\},$$

where i is a number satisfying $i^2 = -1$.

Example 1.1.3 (Number sets)**.** 2 is a real number, and also an integer and a rational number.
$\sqrt{2}$ is a real number, but not a rational number.
π is a real number, but not a rational number.
e is a real number, but not a rational number.
$2 + 5i$ is a complex number, but not a real number.

Definition 1.1.4 (Field)**.** A field is a set $\mathbb{F}$ with two operations, an addition and a multiplication, which assign to all $a, b \in \mathbb{F}$ elements $a + b \in \mathbb{F}$ and $a \cdot b = ab \in \mathbb{F}$, and with two elements $0, 1 \in \mathbb{F}$, such that for all $a, b, c \in \mathbb{F}$:

1. (associativity) $(a + b) + c = a + (b + c)$ and $(ab)c = a(bc)$;

2. (commutativity) $a + b = b + a$ and $ab = ba$;
3. (identities) $a + 0 = a$ and $a \cdot 1 = a$;
4. (inverses) there is an element $-a \in \mathbb{F}$ with $a + (-a) = 0$, and if $a \neq 0$, there is an element $a^{-1} \in \mathbb{F}$ with $aa^{-1} = 1$;
5. (distributivity) $a(b + c) = ab + ac$;
6. (nontriviality) $0 \neq 1$.

The last axiom says that a field has at least two elements. The set $\{0\}$, with $0 + 0 = 0$ and $0 \cdot 0 = 0$, satisfies axioms 1 to 5, with the element 0 as the identity for both operations (so $1 = 0$ there); the inverse axiom for nonzero elements holds because there are none. Axiom 6 excludes it: $\{0\}$ is not a field. In fact, $\{0\}$ is the only set that satisfies axioms 1 to 5 with $0 = 1$: the distributive law gives $a \cdot 0 = a(0 + 0) = a \cdot 0 + a \cdot 0$, so $a \cdot 0 = 0$ after adding $-(a \cdot 0)$, and if $0 = 1$, then $a = a \cdot 1 = a \cdot 0 = 0$ for every a.

Example 1.1.5. The real numbers $\mathbb{R}$, with the usual addition and multiplication, form a field: axioms 1, 2, 3 and 5 are the familiar rules of arithmetic, the inverses of a are $-a$ and, for $a \neq 0$, $\frac{1}{a}$, and $0 \neq 1$.

Example 1.1.6. The set of integers $\mathbb{Z}$ with the usual addition and multiplication operations does not form a field, since not every non-zero integer has a multiplicative inverse in $\mathbb{Z}$. For example, the integer 2 does not have a multiplicative inverse in $\mathbb{Z}$: there is no integer b with $2 \cdot b = 1$, because the only solution of this equation, $b = \frac{1}{2}$, is not an integer.

Exercise 1.1.7. Verify that the set of complex numbers $\mathbb{C}$ and the set of rational numbers $\mathbb{Q}$ with the usual addition and multiplication operations form fields.

1.2 Vector Spaces

Definition 1.2.1 (Vector space)**.** A vector space over a field $\mathbb{F}$ is a set V, whose elements are called vectors, with a vector addition, which assigns to $u, v \in V$ a vector $u + v \in V$, and a scalar multiplication, which assigns to $a \in \mathbb{F}$ and $v \in V$ a vector $a \cdot v = av \in V$, such that for all $u, v, w \in V$ and $a, b \in \mathbb{F}$:

1. $(u + v) + w = u + (v + w)$ and $u + v = v + u$;
2. there is a vector $0 \in V$, the zero vector (also written 0_V), with $v + 0 = v$;
3. there is a vector $-v \in V$ with $v + (-v) = 0$;
4. $a(bv) = (ab)v$ and $1v = v$;
5. $a(u + v) = au + av$ and $(a + b)v = av + bv$.

Axioms 1 to 3 are the rules for the addition of a field, now for vectors. Axioms 4 and 5 connect the two operations: in $(ab)v$, the scalars are multiplied in $\mathbb{F}$ before the vector is scaled, and in $(a + b)v = av + bv$, the left side adds scalars in $\mathbb{F}$, while the right side adds vectors in V.

Example 1.2.2. Every field $\mathbb{F}$ is a vector space over itself, with the field addition as vector addition and the field multiplication as scalar multiplication.

Proof. Since the operations are those of $\mathbb{F}$, every vector space axiom is a field axiom: axioms 1 to 3 are the field axioms for the addition, $a(bv) = (ab)v$ is the associativity of the multiplication, $1v = v$ is the identity axiom, and axiom 5 is the distributive law, which gives $(a+b)v = v(a+b) = va + vb = av + bv$ together with commutativity. □

Corollary 1.2.3. $\mathbb{Q}, \mathbb{R}$, *and* $\mathbb{C}$ *are vector spaces over themselves.*

Definition 1.2.4 (Product of vector spaces)**.** Let V and W be vector spaces over the same field $\mathbb{F}$. Their product is the set

$$V \times W = \{(v, w) : v \in V,\ w \in W\}$$

of all ordered pairs (v, w) with $v \in V$ and $w \in W$. The operations of vector addition and scalar multiplication are defined component-wise:

$$\begin{aligned}(v_1, w_1) + (v_2, w_2) &= (v_1 + v_2, w_1 + w_2),\\ a \cdot (v, w) &= (a \cdot v, a \cdot w), \qquad a \in \mathbb{F}.\end{aligned}$$

Proposition 1.2.5. *If V and W are vector spaces over the same field $\mathbb{F}$, then $V \times W$ is also a vector space over $\mathbb{F}$.*

Proof. The operations act on each component separately, so each axiom for $V \times W$ is the same axiom in V for the first components and in W for the second components. For example,

$$(a + b)(v, w) = ((a + b)v, (a + b)w) = (av + bv, aw + bw) = a(v, w) + b(v, w).$$

The zero vector is $(0_V, 0_W)$, and the additive inverse of (v, w) is $(-v, -w)$. □

Remark 1.2.6. For two (or finitely many) vector spaces, the product is also known as the (external) direct sum of the vector spaces, and it is often denoted by $V \oplus W$.

Definition 1.2.7 (The space $\mathbb{R}^n$)**.** Let n be a positive integer. The set $\mathbb{R}^n$ consists of all ordered n-tuples of real numbers,

$$\mathbb{R}^n = \{(x_1, x_2, \ldots, x_n) : x_i \in \mathbb{R} \text{ for } i = 1, 2, \ldots, n\},$$

and the operations of vector addition and scalar multiplication are defined component-wise:

$$\begin{aligned}(x_1, \ldots, x_n) + (y_1, \ldots, y_n) &= (x_1 + y_1, \ldots, x_n + y_n),\\ a \cdot (x_1, \ldots, x_n) &= (ax_1, \ldots, ax_n), \qquad a \in \mathbb{R}.\end{aligned}$$

Corollary 1.2.8. *The set $\mathbb{R}^n$ is a vector space over the field $\mathbb{R}$ for every positive integer n.*

Proof. We use induction on n. For $n = 1$, $\mathbb{R}^1 = \mathbb{R}$ is a vector space over itself by Corollary 1.2.3.

Now let $n \geq 2$ and assume that $\mathbb{R}^{n-1}$ is a vector space over $\mathbb{R}$. Identify each n-tuple $(x_1, \ldots, x_n)$ with the pair $((x_1, \ldots, x_{n-1}), x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. Under this identification, the component-wise operations on $\mathbb{R}^n$ are exactly the operations of the product $\mathbb{R}^{n-1} \times \mathbb{R}$. By Proposition 1.2.5, $\mathbb{R}^{n-1} \times \mathbb{R}$ is a vector space over $\mathbb{R}$, and hence so is $\mathbb{R}^n$. □

Vectors need not be lists of numbers. Some of the most important vector spaces consist of functions.

Definition 1.2.9 (The space of functions on a set)**.** Let S be a nonempty set. We write $\mathcal{F}(S)$ for the set of all functions $f : S \to \mathbb{R}$, with the pointwise operations

$$(f + g)(s) = f(s) + g(s), \qquad (a \cdot f)(s) = a\, f(s) \qquad (s \in S,\ a \in \mathbb{R}).$$

In words: to add two functions, we add their values at every point. Two functions $f, g \in \mathcal{F}(S)$ are equal if and only if $f(s) = g(s)$ for every $s \in S$.

Proposition 1.2.10. *For every nonempty set S, $\mathcal{F}(S)$ is a vector space over $\mathbb{R}$.*

Proof. Each axiom is an equation between functions, so it suffices to check it at every point $s \in S$, where it becomes the same rule of arithmetic in $\mathbb{R}$. For example,

$$((a+b)f)(s) = (a+b)f(s) = af(s) + bf(s) = (af + bf)(s).$$

The zero vector is the zero function, $0(s) = 0$ for all s, and the additive inverse of f is the function $-f$ with $(-f)(s) = -f(s)$. □

For $S = \{1, 2, \ldots, n\}$, a function $x : S \to \mathbb{R}$ is the same thing as the list of its values $(x(1), \ldots, x(n))$, and the pointwise operations are the componentwise operations of $\mathbb{R}^n$: so $\mathbb{R}^n$ is $\mathcal{F}(\{1, \ldots, n\})$ in disguise. In the same way, the real sequences $(a_0, a_1, a_2, \ldots)$ are the functions $\{0, 1, 2, \ldots\} \to \mathbb{R}$, so they form a vector space.

In calculus and in machine learning, we mostly work with functions that are continuous or differentiable. Let $I \subseteq \mathbb{R}$ be an open interval, such as $\mathbb{R}$ itself or $(0, 1)$. We write $f^{(k)}$ for the k-th derivative of a function f, with $f^{(0)} = f$.

Definition 1.2.11 (Continuous, differentiable and smooth functions)**.** We define the following sets of functions on the open interval I.

- $C(I)$, also written $C^0(I)$, is the set of all continuous functions $f : I \to \mathbb{R}$.
- For an integer $n \geq 1$, $C^n(I)$ is the set of all functions $f : I \to \mathbb{R}$ that are n times differentiable and whose n-th derivative $f^{(n)}$ is continuous. Such functions are called n times continuously differentiable.
- $C^\infty(I)$ is the set of all functions $f : I \to \mathbb{R}$ that have derivatives of all orders, i.e., $C^\infty(I) = C^1(I) \cap C^2(I) \cap \cdots$. Such functions are called smooth.

For a closed interval $[a, b]$, we write $C[a, b]$ for the set of continuous functions $[a, b] \to \mathbb{R}$; the proof of Proposition 1.2.13 below shows that it is a vector space as well.

A differentiable function is continuous, so a function in $C^n(I)$ also has the continuous derivatives $f, f', \ldots, f^{(n-1)}$ of all lower orders, and

$$C^\infty(I) \subseteq \cdots \subseteq C^3(I) \subseteq C^2(I) \subseteq C^1(I) \subseteq C(I) \subseteq \mathcal{F}(I).$$

Example 1.2.12.

1. Polynomials, e^x, $\sin x$ and $\cos x$ are smooth on $\mathbb{R}$: their derivatives are again functions of the same kind, so they can be differentiated again and again.
2. $f(x) = |x|$ is continuous on $\mathbb{R}$, but not differentiable at 0. So $f \in C(\mathbb{R})$, but $f \notin C^1(\mathbb{R})$.
3. $g(x) = x|x|$ has the derivative $g'(x) = 2|x|$ (check this separately for $x > 0$, for $x < 0$, and, with the definition of the derivative, for $x = 0$). So $g \in C^1(\mathbb{R})$, but $g \notin C^2(\mathbb{R})$. In the same way, $x^n|x| \in C^n(\mathbb{R})$, but $x^n|x| \notin C^{n+1}(\mathbb{R})$ (Exercise 36), so all the inclusions above are strict.
4. $h(x) = \frac{1}{x}$ is smooth on the interval $(0, \infty)$, but it is not even defined at 0, so it is not an element of $C(\mathbb{R})$. Whether a function belongs to one of these sets depends on the interval.

Proposition 1.2.13. *With the pointwise operations of $\mathcal{F}(I)$, the sets $C(I)$, $C^n(I)$ (for every $n \geq 1$) and $C^\infty(I)$ are vector spaces over $\mathbb{R}$.*

Proof. We first show that the operations do not lead outside these sets. If f and g are continuous and $a \in \mathbb{R}$, then $f + g$ and af are continuous, by the rules for limits. If f and g are differentiable, then so are $f + g$ and af, with

$$(f+g)' = f' + g' \qquad \text{and} \qquad (af)' = af',$$

by the sum rule and the constant factor rule of calculus. Applying these rules k times gives $(f+g)^{(k)} = f^{(k)} + g^{(k)}$ and $(af)^{(k)} = af^{(k)}$. So if $f, g \in C^n(I)$, then $f+g$ and af are n times differentiable, and their n-th derivatives $f^{(n)} + g^{(n)}$ and $af^{(n)}$ are continuous; hence $f+g, af \in C^n(I)$. Since this holds for every n, it also holds for $C^\infty(I)$.

The axioms of associativity, commutativity and distributivity, and $1 \cdot f = f$, hold because they hold in $\mathcal{F}(I)$ (Proposition 1.2.10), and the operations are the same. The zero function is smooth, so it lies in all of these sets. Finally, the additive inverse of a function f in one of these sets is $-f = (-1) \cdot f$, which lies in the same set, since the set is closed under scalar multiplication. □

In Section 1.5, we will see that this argument is a general principle: $C(I)$, $C^n(I)$ and $C^\infty(I)$ are subspaces of $\mathcal{F}(I)$.

Remark 1.2.14. The set of all functions that are n times differentiable, without the condition that $f^{(n)}$ is continuous, is also a vector space, by the same argument. It is slightly larger than $C^n(I)$. For example, let $f(x) = x^2 \sin(1/x)$ for $x \neq 0$, and $f(0) = 0$. Then $f'(0) = \lim_{h\to 0} h \sin(1/h) = 0$, and $f'(x) = 2x \sin(1/x) - \cos(1/x)$ for $x \neq 0$, which has no limit as $x \to 0$. So f is differentiable on $\mathbb{R}$, but $f \notin C^1(\mathbb{R})$. The spaces $C^n(I)$ are used more often, because continuous derivatives are easier to work with.

Remark 1.2.15 (Function spaces in machine learning). A trained model is a function, for example from images to class probabilities, and learning means searching for a good function in a large set of candidates. Choosing the model fixes this set: linear models give the linear functions, polynomial regression (Programming Exercise P9) the polynomials of degree at most d, and neural networks with smooth activation functions such as tanh give smooth functions, while networks with the ReLU activation $\max(0, x)$ give continuous functions with kinks, like $|x|$. Kernel methods and Gaussian processes (Lecture 4) work with vector spaces of functions directly.

1.3 Bases and Dimension

Throughout this section, V is a vector space over a field $\mathbb{F}$.

Definition 1.3.1 (Linear combination and span)**.** Let $v_1, v_2, \ldots, v_k \in V$. A linear combination of $v_1, \ldots, v_k$ is a vector of the form

$$a_1 v_1 + a_2 v_2 + \cdots + a_k v_k, \qquad a_1, \ldots, a_k \in \mathbb{F}.$$

The span of $v_1, \ldots, v_k$ is the set of all their linear combinations:

$$\operatorname{span}\{v_1, \ldots, v_k\} = \{a_1 v_1 + \cdots + a_k v_k : a_1, \ldots, a_k \in \mathbb{F}\}.$$

We say that $v_1, \ldots, v_k$ span V if $\operatorname{span}\{v_1, \ldots, v_k\} = V$, i.e., if every vector in V is a linear combination of $v_1, \ldots, v_k$.

Definition 1.3.2 (Linear independence)**.** The vectors $v_1, \ldots, v_k \in V$ are linearly independent if the only way to write the zero vector as a linear combination of them is with all coefficients equal to zero:

$$a_1 v_1 + a_2 v_2 + \cdots + a_k v_k = 0 \implies a_1 = a_2 = \cdots = a_k = 0.$$

Otherwise, they are called linearly dependent.

For an infinite set $S \subseteq V$, we say that S is linearly independent if every finite subset of S is linearly independent, and that S spans V if every vector in V is a linear combination of finitely many vectors from S.

Example 1.3.3. In $\mathbb{R}^2$, the vectors $(1,2)$ and $(2,4)$ are linearly dependent, since

$$2 \cdot (1,2) + (-1) \cdot (2,4) = (0,0).$$

Their span is the line $\text{span}\{(1,2),(2,4)\} = \{(t,2t) : t \in \mathbb{R}\}$, so they do not span $\mathbb{R}^2$. For example, $(1,0)$ is not in their span.

Definition 1.3.4 (Basis)**.** A basis of a vector space V is a set of vectors in V that is linearly independent and spans V.

Proposition 1.3.5. *Let $\{v_1, v_2, \ldots, v_n\}$ be a basis of V. Then every vector in V can be expressed as a linear combination of the basis vectors in exactly one way.*

Proof. Since the basis spans V, every $v \in V$ can be written as $v = a_1v_1 + \cdots + a_nv_n$ for some $a_1, \ldots, a_n \in \mathbb{F}$. Suppose that also $v = b_1v_1 + \cdots + b_nv_n$. Subtracting the two expressions gives

$$(a_1 - b_1)v_1 + \cdots + (a_n - b_n)v_n = 0.$$

Since $v_1, \ldots, v_n$ are linearly independent, $a_i - b_i = 0$, i.e., $a_i = b_i$ for every i. So the two expressions are the same. □

The number of vectors in a basis is the same for all bases. This follows from the next lemma, which says that a spanning set always has at least as many vectors as a linearly independent set.

Lemma 1.3.6 (Exchange lemma)**.** *If $v_1, \ldots, v_m$ span V, and $u_1, \ldots, u_k \in V$ are linearly independent, then $k \leq m$.*

Proof. We replace the vectors v_i by the vectors u_j, one at a time, keeping a spanning list of m vectors. We show by induction on $j = 0, 1, \ldots, k$ that $j \leq m$ and that, after renumbering the v_i, the vectors

$$u_1, \ldots, u_j, v_{j+1}, \ldots, v_m$$

span V. For $j = 0$, this is the assumption. Suppose that it holds for some $j < k$. Then

$$u_{j+1} = a_1u_1 + \cdots + a_ju_j + b_{j+1}v_{j+1} + \cdots + b_mv_m$$

for some scalars a_i and b_i. If all b_i were 0 (in particular, if $j = m$, so that there are no b_i), then u_{j+1} would be a linear combination of $u_1, \ldots, u_j$, which is impossible since $u_1, \ldots, u_{j+1}$ are linearly independent. So $j + 1 \leq m$, and after renumbering, $b_{j+1} \neq 0$. Solving for v_{j+1},

$$v_{j+1} = \frac{1}{b_{j+1}}\Big(u_{j+1} - a_1u_1 - \cdots - a_ju_j - b_{j+2}v_{j+2} - \cdots - b_mv_m\Big)$$

is a linear combination of $u_1, \ldots, u_{j+1}, v_{j+2}, \ldots, v_m$. Every vector of V is a linear combination of $u_1, \ldots, u_j, v_{j+1}, \ldots, v_m$; substituting this expression for v_{j+1} writes it as a linear combination of $u_1, \ldots, u_{j+1}, v_{j+2}, \ldots, v_m$. So these vectors span V, which completes the induction. For $j = k$, we get $k \leq m$. □

Theorem 1.3.7. *If a vector space V has a basis consisting of n vectors, then every basis of V consists of exactly n vectors.*

Proof. Let $\{v_1, \ldots, v_n\}$ be a basis, and let S be another basis of V. Any k distinct vectors of S are linearly independent, and $v_1, \ldots, v_n$ span V, so $k \leq n$ by Lemma 1.3.6. Hence S is finite, with $m \leq n$ vectors. Exchanging the roles of the two bases (S spans V, and $v_1, \ldots, v_n$ are linearly independent) gives $n \leq m$. □

This allows us to make the following definition.

Definition 1.3.8 (Dimension)**.** A vector space V is said to be finite-dimensional if it has a finite basis. In this case, the dimension of V, denoted by $\dim V$, is the number of vectors in any basis of V; by Theorem 1.3.7, this number does not depend on which basis we choose. If a vector space does not have a finite basis, it is called infinite-dimensional.

By convention, the vector space $\{0\}$ consisting only of the zero vector has dimension 0 (its basis is the empty set).

Definition 1.3.9 (Standard basis)**.** The standard basis for the vector space $\mathbb{R}^n$ is the set of vectors $\{\mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_n\}$, where each $\mathbf{e}_i$ is a vector in $\mathbb{R}^n$ with a 1 in the i-th position and 0s in all other positions. Formally, the standard basis vectors are defined as:

$$\mathbf{e}_i = (0, \ldots, 0, 1, 0, \ldots, 0) \qquad \text{(with 1 in the } i\text{-th position)}.$$

Proposition 1.3.10. *The standard basis $\{\mathbf{e}_1, \ldots, \mathbf{e}_n\}$ is a basis of $\mathbb{R}^n$. In particular, $\mathbb{R}^n$ is finite-dimensional and $\dim \mathbb{R}^n = n$.*

Proof. Every vector $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{R}^n$ can be written as

$$\mathbf{x} = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \cdots + x_n\mathbf{e}_n,$$

so the standard basis spans $\mathbb{R}^n$. If $a_1\mathbf{e}_1 + \cdots + a_n\mathbf{e}_n = \mathbf{0}$, then $(a_1, \ldots, a_n) = (0, \ldots, 0)$, so $a_1 = \cdots = a_n = 0$. Hence the standard basis is also linearly independent. □

Example 1.3.11. Consider the vector space $\mathbb{R}^2$. Its standard basis is the set of vectors $\{(1,0), (0,1)\}$. These vectors are linearly independent and span the entire space $\mathbb{R}^2$, meaning that any vector $(x, y) \in \mathbb{R}^2$ can be expressed as a linear combination of these basis vectors:

$$(x, y) = x \cdot (1, 0) + y \cdot (0, 1).$$

The dimension of $\mathbb{R}^2$ is 2, as there are two vectors in the basis.

Example 1.3.12. Consider the vector space $\mathbb{R}^3$. Its standard basis is the set of vectors

$$\{(1,0,0), (0,1,0), (0,0,1)\}.$$

These vectors are linearly independent and span the entire space $\mathbb{R}^3$, meaning that any vector $(x, y, z) \in \mathbb{R}^3$ can be expressed as a linear combination of these basis vectors:

$$(x, y, z) = x \cdot (1, 0, 0) + y \cdot (0, 1, 0) + z \cdot (0, 0, 1).$$

The dimension of $\mathbb{R}^3$ is 3, as there are three vectors in the basis.

Example 1.3.13. The set of all polynomials with real coefficients,

$$\mathbb{R}[x] = \{a_0 + a_1x + \cdots + a_dx^d : d \geq 0,\ a_0, \ldots, a_d \in \mathbb{R}\},$$

is a vector space over $\mathbb{R}$ with the usual addition of polynomials and multiplication of a polynomial by a real number. This vector space is infinite-dimensional.

You may check for yourself that the set of monomials $\{1, x, x^2, x^3, \ldots\}$ is linearly independent and spans the space of all polynomials, so it is an (infinite) basis of $\mathbb{R}[x]$.

To see that there is no finite basis, take any finitely many polynomials $p_1, \ldots, p_k \in \mathbb{R}[x]$, and choose an integer $d \geq 0$ such that each p_i has degree at most d. Every linear combination of $p_1, \ldots, p_k$ also has degree at most d, so $x^{d+1} \notin \operatorname{span}\{p_1, \ldots, p_k\}$. Hence no finite set of polynomials spans $\mathbb{R}[x]$, and in particular $\mathbb{R}[x]$ has no finite basis.

The exchange lemma has several consequences that we will use all the time.

Corollary 1.3.14. *Let V be a vector space.*

1. *If $v_1, \dots, v_m$ span V, then some of these vectors form a basis of V. In particular, V is finite-dimensional, and $\dim V \leq m$.*
2. *If $\dim V = n$, then every set of linearly independent vectors in V has at most n elements, and every set of vectors that spans V has at least n elements.*
3. *If $\dim V = n$, then every set of linearly independent vectors $u_1, \dots, u_k$ in V can be extended to a basis $u_1, \dots, u_k, u_{k+1}, \dots, u_n$ of V.*
4. *If $\dim V = n$, then any n linearly independent vectors in V form a basis of V, and so do any n vectors that span V.*
5. *If $\dim V = n$, then every subspace U of V is finite-dimensional, with $\dim U \leq n$, and $\dim U = n$ only if $U = V$.*

Proof. (1) If $v_1, \dots, v_m$ are linearly dependent, then $a_1 v_1 + \cdots + a_m v_m = 0$ with some $a_i \neq 0$, so $v_i = -\frac{1}{a_i} \sum_{l \neq i} a_l v_l$ is a linear combination of the other vectors. Removing v_i leaves a list that still spans V: in every linear combination of $v_1, \dots, v_m$, we can replace v_i by this expression. We repeat this until the remaining vectors are linearly independent, which happens after at most m steps (the empty list, which spans $\{0\}$, is linearly independent). The remaining vectors span V and are linearly independent, so they form a basis.

(2) A basis of V consists of n linearly independent vectors that span V, so both statements follow from Lemma 1.3.6.

(3) If $u_1, \dots, u_k$ do not span V, choose $w \in V$ with $w \notin \operatorname{span}\{u_1, \dots, u_k\}$. Then $u_1, \dots, u_k, w$ are linearly independent: if $a_1 u_1 + \cdots + a_k u_k + bw = 0$, then $b = 0$ (otherwise $w = -\frac{1}{b}(a_1 u_1 + \cdots + a_k u_k)$ would lie in the span), and then all $a_i = 0$, since the u_i are linearly independent. We repeat this with the longer list. By (2), a linearly independent list in V has at most n vectors, so the process stops, and it stops with a linearly independent list that spans V, i.e., a basis. By Theorem 1.3.7, this basis has n vectors.

(4) If n linearly independent vectors did not span V, then the argument of (3) would give $n+1$ linearly independent vectors, contradicting (2). If n vectors span V, then by (1) some of them form a basis, which has n vectors, so it consists of all of them.

(5) Linearly independent vectors in U are linearly independent in V, so by (2), there are at most n of them. Choose linearly independent vectors $u_1, \dots, u_k \in U$ with k as large as possible. They span U: otherwise, the argument of (3) would give a vector $w \in U$ such that $u_1, \dots, u_k, w$ are linearly independent. So they form a basis of U, and $\dim U = k \leq n$. If $k = n$, then $u_1, \dots, u_n$ form a basis of V by (4), so $U = V$. □

1.4 Linear Transformations

Definition 1.4.1 (Linear transformation)**.** A linear transformation between two vector spaces V and W over the same field $\mathbb{F}$ is a function $T : V \to W$ that satisfies the following two properties for all $u, v \in V$ and all scalars $a \in \mathbb{F}$:

1. Additivity: $T(u + v) = T(u) + T(v)$
2. Homogeneity: $T(a \cdot v) = a \cdot T(v)$

Lemma 1.4.2. *If $T : V \to W$ is a linear transformation, then $T(0_V) = 0_W$.*

Proof. By additivity, $T(0_V) = T(0_V + 0_V) = T(0_V) + T(0_V)$. Adding $-T(0_V)$ to both sides gives $0_W = T(0_V)$. □

Example 1.4.3. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (2x + 3y, 4x - y)$. We will verify that T is a linear transformation.

First, we check additivity:

$$\begin{aligned} T((x_1, y_1) + (x_2, y_2)) &= T(x_1 + x_2, y_1 + y_2) \\ &= (2(x_1 + x_2) + 3(y_1 + y_2), 4(x_1 + x_2) - (y_1 + y_2)) \\ &= (2x_1 + 3y_1 + 2x_2 + 3y_2, 4x_1 - y_1 + 4x_2 - y_2) \\ &= (2x_1 + 3y_1, 4x_1 - y_1) + (2x_2 + 3y_2, 4x_2 - y_2) \\ &= T(x_1, y_1) + T(x_2, y_2). \end{aligned}$$

Next, we check homogeneity:

$$\begin{aligned} T(a \cdot (x, y)) &= T(ax, ay) \\ &= (2(ax) + 3(ay), 4(ax) - (ay)) \\ &= a(2x + 3y, 4x - y) \\ &= a \cdot T(x, y). \end{aligned}$$

Since both properties hold, T is indeed a linear transformation.

Remark 1.4.4 (Visualizing linear transformations of the plane). Every vector of $\mathbb{R}^2$ can be written as $(x, y) = x\mathbf{e}_1 + y\mathbf{e}_2$, so for a linear transformation $T : \mathbb{R}^2 \to \mathbb{R}^2$,

$$T(x, y) = x\,T(\mathbf{e}_1) + y\,T(\mathbf{e}_2).$$

Hence T is completely determined by the two vectors $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$. In particular, T maps the unit square $\{x\mathbf{e}_1 + y\mathbf{e}_2 : 0 \le x, y \le 1\}$ to the parallelogram $\{xT(\mathbf{e}_1) + yT(\mathbf{e}_2) : 0 \le x, y \le 1\}$ spanned by $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$. The origin stays where it is (Lemma 1.4.2), and a grid of equally spaced parallel lines is mapped to a grid of equally spaced parallel lines (or, if $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$ lie on a common line, squashed onto that line). Figure 1.1 shows this for the linear transformation of the previous example, for which $T(\mathbf{e}_1) = (2, 4)$ and $T(\mathbf{e}_2) = (3, -1)$. Figure 1.2 shows four basic examples; you may check that each of them is linear.





Figure 1.1: The linear transformation $T(x, y) = (2x + 3y, 4x - y)$ maps the unit square (left) to the parallelogram spanned by $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$ (right). The grid lines inside the square are mapped to equally spaced parallel lines inside the parallelogram.

Linear transformations are not only maps between spaces of lists of numbers: the two basic operations of calculus are linear. Let I be an open interval.


Example 1.4.5 (Differentiation)**.** The map

$$D : C^1(I) \to C(I), \qquad D(f) = f',$$



(a) stretch
$T(x, y) = (2x, y)$

(b) rotation by 90°
$T(x, y) = (-y, x)$



(c) reflection in the x-axis
$T(x, y) = (x, -y)$

(d) shear
$T(x, y) = (x + y, y)$

Figure 1.2: Four linear transformations of the plane. The dashed square and the gray letter F show the unit square before the transformation; the blue parallelogram and the orange letter F show its image. The red and green arrows are $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$. Note that in (c), the image of the letter F is mirrored.

is well defined, since f' is continuous for $f \in C^1(I)$, and it is a linear transformation: by the sum rule and the constant factor rule,

$$D(f+g) = (f+g)' = f' + g' = D(f) + D(g) \qquad \text{and} \qquad D(af) = (af)' = af' = aD(f)$$

for all $f, g \in C^1(I)$ and $a \in \mathbb{R}$. For example, $D(3x^2 - \sin x) = 6x - \cos x = 3D(x^2) - D(\sin x)$. In the same way, differentiation gives linear transformations $C^{n+1}(I) \to C^n(I)$ and $C^\infty(I) \to C^\infty(I)$, and also $\mathbb{R}[x] \to \mathbb{R}[x]$, since the derivative of a polynomial is a polynomial.

Example 1.4.6 (Integration)**.** 1. Let $a < b$. The definite integral

$$\varphi : C[a, b] \to \mathbb{R}, \qquad \varphi(f) = \int_a^b f(x)\,dx,$$

is a linear transformation, by the linearity of the integral: $\int_a^b (f+g)(x)\,dx = \int_a^b f(x)\,dx + \int_a^b g(x)\,dx$ and $\int_a^b af(x)\,dx = a\int_a^b f(x)\,dx$. Its values are numbers, so it is a linear functional in the sense of Section 1.8. For example, the average value $\frac{1}{b-a}\int_a^b f(x)\,dx$ of f on $[a, b]$ depends linearly on f.

2. Let $c \in I$. The integral with a variable upper limit,

$$J : C(I) \to C^1(I), \qquad (Jf)(x) = \int_c^x f(t)\,dt \qquad (x \in I),$$

with the usual convention $\int_c^x = -\int_x^c$ for $x < c$, is linear, again by the linearity of the integral, applied for each fixed x. By the fundamental theorem of calculus, Jf is differentiable with $(Jf)' = f$, which is continuous, so Jf indeed lies in $C^1(I)$: it is the antiderivative of f that vanishes at c.

Differentiation and integration undo each other, but only in one order:

$$D(Jf) = f \quad \text{for all } f \in C(I), \qquad \text{but} \qquad J(Df)(x) = \int_c^x f'(t)\,dt = f(x) - f(c) \quad \text{for } f \in C^1(I).$$

So $D \circ J$ is the identity map of $C(I)$, while $J \circ D$ forgets the value $f(c)$.

Definition 1.4.7 (Kernel)**.** The kernel of a linear transformation $T : V \to W$ is the set of all vectors in V that are mapped to the zero vector in W. Formally, the kernel is defined as

$$\ker(T) = \{v \in V : T(v) = 0_W\},$$

where 0_W is the zero vector in W.

Example 1.4.8. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (2x + 3y, 4x - y)$. We will find the kernel of T.

To find the kernel, we need to solve the equation $T(x, y) = (0, 0)$:

$$\begin{aligned} 2x + 3y &= 0 \\ 4x - y &= 0 \end{aligned}$$

From the second equation, we can express y in terms of x: $y = 4x$. Substituting this into the first equation gives:

$$\begin{aligned} 2x + 3(4x) &= 0 \\ 2x + 12x &= 0 \\ 14x &= 0 \\ x &= 0 \end{aligned}$$

Substituting $x = 0$ back into $y = 4x$, we find that $y = 0$. Therefore, the only solution is $(0, 0)$.

Thus, the kernel of T is

$$\ker(T) = \{(0, 0)\}.$$

Example 1.4.9. Let $T : \mathbb{R}^3 \to \mathbb{R}^2$ be defined by $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$. We will find the kernel of T.

To find the kernel, we need to solve the equation $T(x, y, z) = (0, 0)$:

$$\begin{aligned} x + 2y - z &= 0 \\ 3x - y + 4z &= 0 \end{aligned}$$

From the first equation, we can express z in terms of x and y: $z = x + 2y$. Substituting this into the second equation gives:

$$\begin{aligned} 3x - y + 4(x + 2y) &= 0 \\ 3x - y + 4x + 8y &= 0 \\ 7x + 7y &= 0 \\ x + y &= 0 \end{aligned}$$

From this equation, we can express y in terms of x: $y = -x$. Substituting this back into the expression for z, we get:

$$z = x + 2(-x) = x - 2x = -x.$$

Therefore, any vector in the kernel can be expressed as $(x, -x, -x)$ for some scalar x. Thus, the kernel of T is

$$\ker(T) = \{(x, -x, -x) : x \in \mathbb{R}\}.$$

Definition 1.4.10 (Image)**.** The image (or range) of a linear transformation $T : V \to W$ is the set of all vectors in W that can be expressed as $T(v)$ for some vector $v \in V$. Formally, the image is defined as

$$\text{im}(T) = \{w \in W : w = T(v) \text{ for some } v \in V\}.$$

Example 1.4.11. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (2x + 3y, 4x - y)$. We will find the image of T.

To find the image, we need to determine all possible outputs of T. We will show that every vector of $\mathbb{R}^2$ is an output. Let $(u, v) \in \mathbb{R}^2$ be arbitrary. We look for $x, y \in \mathbb{R}$ such that $T(x, y) = (u, v)$, i.e.,

$$\begin{aligned} u &= 2x + 3y \\ v &= 4x - y \end{aligned}$$

We can solve this system of equations for x and y in terms of u and v. From the second equation, we can express y in terms of x and v: $y = 4x - v$. Substituting this into the first equation gives:

$$\begin{aligned} u &= 2x + 3(4x - v) \\ u &= 2x + 12x - 3v \\ u &= 14x - 3v \end{aligned}$$

Solving for x, we get:

$$14x = u + 3v \implies x = \frac{u + 3v}{14}.$$

Substituting this back into the expression for y, we get:

$$y = 4\left(\frac{u + 3v}{14}\right) - v = \frac{4u + 12v - 14v}{14} = \frac{2u - v}{7}.$$

Let us check that these values work:

$$\begin{aligned} 2x + 3y &= \frac{2u + 6v}{14} + \frac{12u - 6v}{14} = u, \\ 4x - y &= \frac{4u + 12v}{14} - \frac{4u - 2v}{14} = v. \end{aligned}$$

Therefore, for any $(u, v) \in \mathbb{R}^2$, we have found $(x, y) \in \mathbb{R}^2$ such that $T(x, y) = (u, v)$. This means that the image of T is all of $\mathbb{R}^2$, i.e.,

$$\text{im}(T) = \mathbb{R}^2.$$

Example 1.4.12. Let $T : \mathbb{R}^3 \to \mathbb{R}^2$ be defined by $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$. We will find the image of T.

As before, we will show that every vector of $\mathbb{R}^2$ is an output of T. Let $(u, v) \in \mathbb{R}^2$ be arbitrary. We look for $x, y, z \in \mathbb{R}$ such that $T(x, y, z) = (u, v)$, i.e.,

$$\begin{aligned} u &= x + 2y - z \\ v &= 3x - y + 4z \end{aligned}$$

From the first equation, we can express z in terms of x, y, and u: $z = x + 2y - u$. Substituting this into the second equation gives:

$$\begin{aligned} v &= 3x - y + 4(x + 2y - u) \\ v &= 3x - y + 4x + 8y - 4u \\ v &= 7x + 7y - 4u \end{aligned}$$

Rearranging gives:

$$7x + 7y = 4u + v.$$

For any fixed (u, v), this equation describes a line in the (x, y)-plane, and every point (x, y) on this line, together with $z = x + 2y - u$, is a solution. For example, we can take $y = 0$ and $x = \frac{4u+v}{7}$, and then $z = x - u = \frac{v-3u}{7}$. Let us check:

$$T\left(\frac{4u+v}{7}, 0, \frac{v-3u}{7}\right) = \left(\frac{(4u+v)-(v-3u)}{7}, \frac{3(4u+v)+4(v-3u)}{7}\right) = (u, v).$$

Therefore, for any $(u, v) \in \mathbb{R}^2$, we have found $(x, y, z) \in \mathbb{R}^3$ such that $T(x, y, z) = (u, v)$. This means that the image of T is all of $\mathbb{R}^2$, i.e.,

$$\mathrm{im}(T) = \mathbb{R}^2.$$

Example 1.4.13. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (x + 2y, 2x + 4y)$. This map is linear, and it can be written as

$$T(x, y) = (x + 2y) \cdot (1, 2).$$

So $T(x, y) = (0, 0)$ if and only if $x + 2y = 0$, and the kernel of T is the line

$$\ker(T) = \{(-2t, t) : t \in \mathbb{R}\} = \mathrm{span}\{(-2, 1)\}.$$

Every output of T is a multiple of $(1, 2)$, and every multiple $s(1, 2) = T(s, 0)$ is an output, so the image of T is the line

$$\mathrm{im}(T) = \{(s, 2s) : s \in \mathbb{R}\} = \mathrm{span}\{(1, 2)\}.$$

More generally, all points of the line $x + 2y = c$, which is parallel to $\ker(T)$, are mapped to the same point $c\,(1, 2)$. So T squashes the whole plane onto the line $\mathrm{im}(T)$, as shown in Figure 1.3. In particular, T is neither injective nor surjective.




Figure 1.3: The linear transformation $T(x, y) = (x + 2y, 2x + 4y)$ of Example 1.4.13. Each line $x + 2y = c$ on the left is mapped to the single point $c\,(1, 2)$ on the right, shown in the same color (here $c = -2,\ 0,\ 2$). The kernel (red, $c = 0$) is mapped to the origin, and the whole plane is squashed onto the line $\mathrm{im}(T)$.

Example 1.4.14. For differentiation $D : C^1(I) \to C(I)$ (Example 1.4.5), the kernel consists of the functions with $f' = 0$ on I, which are exactly the constant functions (by the mean value theorem of calculus). Its image is all of $C(I)$: every continuous function g is the derivative of the function Jg of Example 1.4.6, since $D(Jg) = g$.

For integration $J : C(I) \to C^1(I)$, the kernel is $\{0\}$: if $Jf = 0$, then $f = D(Jf) = D(0) = 0$. Its image consists of the functions $F \in C^1(I)$ with $F(c) = 0$: every Jf vanishes at c, and every $F \in C^1(I)$ with $F(c) = 0$ satisfies $F = F - F(c) = J(DF)$.

1.4.1 Isomorphisms

Two vector spaces can look very different and still "behave in the same way". For example, a polynomial $a_0 + a_1x + a_2x^2$ is added and multiplied by scalars exactly like the list of its coefficients $(a_0, a_1, a_2) \in \mathbb{R}^3$. Isomorphisms make this idea precise.

Definition 1.4.15 (Injective, surjective, bijective)**.** A function $f : X \to Y$ between two sets is called

- injective if different inputs give different outputs, i.e., $f(x_1) = f(x_2)$ implies $x_1 = x_2$;
- surjective if every element of Y is an output, i.e., for every $y \in Y$ there exists $x \in X$ with $f(x) = y$;
- bijective if it is both injective and surjective.

A function f is bijective if and only if it has an inverse function $f^{-1} : Y \to X$, i.e., a function with $f^{-1}(f(x)) = x$ for all $x \in X$ and $f(f^{-1}(y)) = y$ for all $y \in Y$. Note that a linear transformation $T : V \to W$ is surjective if and only if $\mathrm{im}(T) = W$.

Definition 1.4.16 (Isomorphism)**.** Let V and W be vector spaces over the same field $\mathbb{F}$. An isomorphism from V to W is a bijective linear transformation $T : V \to W$. If there exists an isomorphism from V to W, we say that V and W are isomorphic, and we write $V \cong W$.

Remark 1.4.17. An isomorphism $T : V \to W$ matches every vector of V with exactly one vector of W, in a way that respects addition and scalar multiplication. So any statement about V that only involves addition and scalar multiplication can be translated into a statement about W, and vice versa. In this sense, isomorphic vector spaces are "the same vector space, written in a different way".

To check that a linear transformation is injective, it is enough to look at its kernel.

Lemma 1.4.18. *A linear transformation $T : V \to W$ is injective if and only if* $\ker(T) = \{0_V\}$. *Hence T is an isomorphism if and only if* $\ker(T) = \{0_V\}$ *and* $\mathrm{im}(T) = W$.

Proof. Suppose that T is injective, and let $v \in \ker(T)$. Then $T(v) = 0_W = T(0_V)$ by Lemma 1.4.2, so $v = 0_V$.

Conversely, suppose that $\ker(T) = \{0_V\}$, and let $u, v \in V$ with $T(u) = T(v)$. Using linearity and $(-1) \cdot v = -v$ (Exercise 1), we get

$$T(u - v) = T(u + (-1) \cdot v) = T(u) + (-1) \cdot T(v) = T(u) - T(v) = 0_W,$$

so $u - v \in \ker(T) = \{0_V\}$, i.e., $u = v$. □

Example 1.4.19. The linear transformation $T : \mathbb{R}^2 \to \mathbb{R}^2$, $T(x, y) = (2x + 3y, 4x - y)$, is an isomorphism. Indeed, we found in Example 1.4.8 that $\ker(T) = \{(0, 0)\}$, so T is injective by Lemma 1.4.18, and we found in Example 1.4.11 that $\mathrm{im}(T) = \mathbb{R}^2$, so T is surjective. Its inverse is the function we computed in Example 1.4.11:

$$T^{-1}(u, v) = \left(\frac{u + 3v}{14}, \frac{2u - v}{7}\right).$$

Example 1.4.20. The linear transformation $T : \mathbb{R}^3 \to \mathbb{R}^2$, $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$, is surjective by Example 1.4.12. However, it is not injective, since its kernel is the line $\{(x, -x, -x) : x \in \mathbb{R}\} \neq \{0\}$ (Example 1.4.9). So T is not an isomorphism.

Similarly, the linear transformation $S : \mathbb{R} \to \mathbb{R}^2$, $S(x) = (x, 0)$, is injective, since $\ker(S) = \{0\}$, but it is not surjective, since $(0, 1)$ is not an output. So S is not an isomorphism either.

Example 1.4.21. Let $\mathbb{R}[x]_{\leq 2}$ be the set of polynomials of degree at most 2, with the usual addition of polynomials and multiplication by real numbers; one can check that it is a vector space over $\mathbb{R}$. The map

$$T : \mathbb{R}[x]_{\leq 2} \to \mathbb{R}^3, \qquad T(a_0 + a_1 x + a_2 x^2) = (a_0, a_1, a_2),$$

which lists the coefficients of a polynomial, is an isomorphism. It is linear, since adding two polynomials adds their coefficients, and multiplying a polynomial by a scalar multiplies its coefficients by that scalar. It is injective, since its kernel consists only of the zero polynomial, and it is surjective, since every $(a_0, a_1, a_2) \in \mathbb{R}^3$ is the output $T(a_0 + a_1 x + a_2 x^2)$. Hence $\mathbb{R}[x]_{\leq 2} \cong \mathbb{R}^3$: as far as addition and scalar multiplication are concerned, a polynomial of degree at most 2 is the same thing as the list of its three coefficients.

Example 1.4.22. The complex numbers $\mathbb{C}$ also form a vector space over $\mathbb{R}$, if we only allow multiplication by real scalars. The map $T : \mathbb{C} \to \mathbb{R}^2$, $T(a + bi) = (a, b)$, is linear and bijective (check this!), so it is an isomorphism of vector spaces over $\mathbb{R}$, and $\mathbb{C} \cong \mathbb{R}^2$. This is the familiar picture of complex numbers as points in the plane.

Example 1.4.23. By Example 1.4.14 and Lemma 1.4.18, differentiation $D : C^1(I) \to C(I)$ is surjective but not injective, and integration $J : C(I) \to C^1(I)$ is injective but not surjective. So neither of them is an isomorphism, although $D \circ J$ is the identity. The same happens for differentiation $D : C^\infty(I) \to C^\infty(I)$, which maps a vector space to itself: it is surjective, since J maps smooth functions to smooth functions, but it is not injective. For a linear transformation $T : V \to V$ of a finite-dimensional vector space to itself, this cannot happen: by the rank–nullity theorem (Section 1.7), T is injective if and only if it is surjective. Infinite-dimensional vector spaces behave differently.

Lemma 1.4.24. *Let U, V and W be vector spaces over the same field $\mathbb{F}$.*

1. *If $T : U \to V$ and $S : V \to W$ are linear transformations, then their composition $S \circ T : U \to W$, defined by $(S \circ T)(u) = S(T(u))$, is a linear transformation. If S and T are isomorphisms, then so is $S \circ T$.*
2. *If $T : V \to W$ is an isomorphism, then its inverse function $T^{-1} : W \to V$ is also linear, and hence an isomorphism.*
3. *The identity map $\mathrm{id}_V : V \to V$, $\mathrm{id}_V(v) = v$, is an isomorphism.*

Proof. (1) For $u_1, u_2, u \in U$ and $a \in \mathbb{F}$, we have

$$(S \circ T)(u_1 + u_2) = S(T(u_1) + T(u_2)) = S(T(u_1)) + S(T(u_2)) = (S \circ T)(u_1) + (S \circ T)(u_2),$$
$$(S \circ T)(au) = S(aT(u)) = aS(T(u)) = a(S \circ T)(u).$$

If S and T are bijective, then $S \circ T$ is bijective, with inverse function $T^{-1} \circ S^{-1}$.

(2) Let $w_1, w_2 \in W$ and $a \in \mathbb{F}$, and let $v_1 = T^{-1}(w_1)$ and $v_2 = T^{-1}(w_2)$. Since $T(v_1 + v_2) = T(v_1) + T(v_2) = w_1 + w_2$, applying T^{-1} gives

$$T^{-1}(w_1 + w_2) = v_1 + v_2 = T^{-1}(w_1) + T^{-1}(w_2).$$

Similarly, $T(av_1) = aT(v_1) = aw_1$ gives $T^{-1}(aw_1) = av_1 = aT^{-1}(w_1)$. Finally, T^{-1} is bijective, with inverse function T.

(3) The identity map is clearly linear and bijective. □

In particular, $V \cong V$; if $V \cong W$, then $W \cong V$; and if $U \cong V$ and $V \cong W$, then $U \cong W$.

Proposition 1.4.25. *Let $T : V \to W$ be an isomorphism, and let $\{v_1, \ldots, v_n\}$ be a basis of V. Then $\{T(v_1), \ldots, T(v_n)\}$ is a basis of W. In particular, if V is finite-dimensional and $V \cong W$, then W is also finite-dimensional and $\dim W = \dim V$.*

Proof. Since T is injective, the vectors $T(v_1), \dots, T(v_n)$ are n different vectors.

They span W: let $w \in W$. Since T is surjective, $w = T(v)$ for some $v \in V$, and we can write $v = a_1 v_1 + \cdots + a_n v_n$. By linearity, $w = T(v) = a_1 T(v_1) + \cdots + a_n T(v_n)$.

They are linearly independent: if $a_1 T(v_1) + \cdots + a_n T(v_n) = 0_W$, then by linearity $T(a_1 v_1 + \cdots + a_n v_n) = 0_W$, so $a_1 v_1 + \cdots + a_n v_n \in \ker(T) = \{0_V\}$ by Lemma 1.4.18. Since $v_1, \dots, v_n$ are linearly independent, $a_1 = \cdots = a_n = 0$. □

Conversely, every finite-dimensional vector space is isomorphic to some $\mathbb{R}^n$. To see this, we use coordinates. For the rest of this subsection, we work over the field $\mathbb{R}$; everything works in the same way over any field $\mathbb{F}$, with $\mathbb{F}^n$ in place of $\mathbb{R}^n$.

Definition 1.4.26 (Coordinates)**.** Let V be a vector space over $\mathbb{R}$, and let $\mathcal{B} = (v_1, \dots, v_n)$ be a basis of V whose vectors are listed in a fixed order (an ordered basis). By Proposition 1.3.5, every $v \in V$ can be written in exactly one way as $v = a_1 v_1 + \cdots + a_n v_n$. The vector

$$[v]_{\mathcal{B}} = (a_1, \dots, a_n) \in \mathbb{R}^n$$

is called the coordinate vector of v with respect to $\mathcal{B}$, and the map $\varphi_{\mathcal{B}} : V \to \mathbb{R}^n$, $\varphi_{\mathcal{B}}(v) = [v]_{\mathcal{B}}$, is called the coordinate map.

Theorem 1.4.27. *Let V be a vector space over $\mathbb{R}$ with an ordered basis $\mathcal{B} = (v_1, \dots, v_n)$. Then the coordinate map $\varphi_{\mathcal{B}} : V \to \mathbb{R}^n$ is an isomorphism. In particular, every n-dimensional vector space over $\mathbb{R}$ is isomorphic to $\mathbb{R}^n$.*

Proof. Let $v = a_1 v_1 + \cdots + a_n v_n$ and $w = b_1 v_1 + \cdots + b_n v_n$, and let $c \in \mathbb{R}$. Then

$$v + w = (a_1 + b_1)v_1 + \cdots + (a_n + b_n)v_n \qquad \text{and} \qquad cv = (ca_1)v_1 + \cdots + (ca_n)v_n,$$

and by the uniqueness of these expressions, $[v + w]_{\mathcal{B}} = [v]_{\mathcal{B}} + [w]_{\mathcal{B}}$ and $[cv]_{\mathcal{B}} = c\,[v]_{\mathcal{B}}$. So $\varphi_{\mathcal{B}}$ is linear. It is bijective, because the map $\mathbb{R}^n \to V$, $(a_1, \dots, a_n) \mapsto a_1 v_1 + \cdots + a_n v_n$, is an inverse function of $\varphi_{\mathcal{B}}$, again by the uniqueness of the expressions. □

Example 1.4.28. With the ordered basis $\mathcal{B} = (1, x, x^2)$ of $\mathbb{R}[x]_{\leq 2}$, the coordinate map $\varphi_{\mathcal{B}}$ is exactly the isomorphism of Example 1.4.21. With a different ordered basis we get different coordinates: for example, with $\mathcal{B}' = (1, 1 + x, x^2)$, the polynomial $p = 2 + 3x + x^2 = (-1) \cdot 1 + 3 \cdot (1 + x) + 1 \cdot x^2$ has $[p]_{\mathcal{B}'} = (-1, 3, 1)$, while $[p]_{\mathcal{B}} = (2, 3, 1)$.

Corollary 1.4.29. *Two finite-dimensional vector spaces V and W over $\mathbb{R}$ are isomorphic if and only if $\dim V = \dim W$.*

Proof. If $V \cong W$, then $\dim V = \dim W$ by Proposition 1.4.25. Conversely, suppose that $\dim V = \dim W = n$, and choose ordered bases $\mathcal{B}$ of V and $\mathcal{C}$ of W. By Theorem 1.4.27 and Lemma 1.4.24, the map $\varphi_{\mathcal{C}}^{-1} \circ \varphi_{\mathcal{B}} : V \to W$ is an isomorphism. □

Example 1.4.30. Since $\dim \mathbb{R}^2 = 2 \neq 3 = \dim \mathbb{R}^3$, the spaces $\mathbb{R}^2$ and $\mathbb{R}^3$ are not isomorphic: there is no bijective linear transformation between them. On the other hand, any two real vector spaces of dimension 3, such as $\mathbb{R}[x]_{\leq 2}$ and $\mathbb{R}^3$, are isomorphic. Finally, the infinite-dimensional space $\mathbb{R}[x]$ is not isomorphic to any $\mathbb{R}^n$, by Proposition 1.4.25.

1.5 Subspaces

Definition 1.5.1 (Subspace)**.** A subspace of a vector space V over $\mathbb{F}$ is a subset $U \subseteq V$ that is itself a vector space under the same operations of vector addition and scalar multiplication defined on V. In practice, a subset $U \subseteq V$ is a subspace if and only if it satisfies the following three conditions:

1. The zero vector of V is in U, i.e., $0_V \in U$.

2. U is closed under vector addition: For any $u, v \in U$, we have $u + v \in U$.

3. U is closed under scalar multiplication: For any $a \in \mathbb{F}$ and any $v \in U$, we have $a \cdot v \in U$.

Remark 1.5.2. Why are these three conditions enough? Conditions 2 and 3 guarantee that adding vectors of U or multiplying them by scalars never leads outside of U. The axioms of associativity, commutativity and distributivity, and $1 \cdot v = v$, then hold in U automatically, because they already hold in V. Condition 1 provides the additive identity. Finally, the additive inverse of $v \in U$ is $-v = (-1) \cdot v$ (see Exercise 1 in Section 1.13), which lies in U by condition 3.

Example 1.5.3. Let $V = \mathbb{R}^3$ and consider the subset $U = \{(x, y, z) \in \mathbb{R}^3 : x + y + z = 0\}$. We will verify that U is a subspace of V.

First, we check that the zero vector is in U:

$$0_V = (0, 0, 0) \in U \text{ since } 0 + 0 + 0 = 0.$$

Next, we check closure under vector addition. Let $(x_1, y_1, z_1), (x_2, y_2, z_2) \in U$. Then we have:

$$x_1 + y_1 + z_1 = 0 \qquad \text{and} \qquad x_2 + y_2 + z_2 = 0.$$

Now consider their sum:

$$(x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2).$$

We need to check if this sum is in U:

$$(x_1 + x_2) + (y_1 + y_2) + (z_1 + z_2) = (x_1 + y_1 + z_1) + (x_2 + y_2 + z_2) = 0 + 0 = 0.$$

Thus, $(x_1 + x_2, y_1 + y_2, z_1 + z_2) \in U$.

Finally, we check closure under scalar multiplication. Let $a \in \mathbb{R}$ and $(x, y, z) \in U$. Then we have:

$$x + y + z = 0.$$

Now consider the scalar multiple:

$$a \cdot (x, y, z) = (ax, ay, az).$$

We need to check if this scalar multiple is in U:

$$ax + ay + az = a(x + y + z) = a \cdot 0 = 0.$$

Thus, $(ax, ay, az) \in U$.

Therefore, U is a subspace of $\mathbb{R}^3$.

The proof of Proposition 1.2.13 was exactly this criterion: $C(I)$, $C^n(I)$ and $C^\infty(I)$ contain the zero function and are closed under addition and scalar multiplication, so they are subspaces of $\mathcal{F}(I)$. Each of them is even a subspace of the next larger one in the chain

$$C^\infty(I) \subseteq \cdots \subseteq C^2(I) \subseteq C^1(I) \subseteq C(I) \subseteq \mathcal{F}(I),$$

since the three conditions do not depend on the larger space. Polynomials fit into this chain at the smooth end.

Proposition 1.5.4. *Identify every polynomial $p = a_0 + a_1 x + \cdots + a_d x^d \in \mathbb{R}[x]$ with the function $\mathbb{R} \to \mathbb{R}$, $x \mapsto p(x)$. Then $\mathbb{R}[x]$ is a subspace of $C^\infty(\mathbb{R})$.*

Proof. First, every polynomial function is smooth: the derivative of p is the polynomial $p' = a_1 + 2a_2x + \cdots + da_dx^{d-1}$, so by induction, all derivatives $p^{(k)}$ exist and are polynomials, which are continuous. Hence $\mathbb{R}[x] \subseteq C^\infty(\mathbb{R})$.

Next, the operations agree: the polynomial $p+q$, which is obtained by adding coefficients, has the values $(p+q)(x) = p(x) + q(x)$, and the polynomial cp has the values $c\,p(x)$. So adding and scaling polynomials is the same as adding and scaling the corresponding functions pointwise. It remains to check the three conditions of Definition 1.5.1:

1. The zero polynomial is the zero function, so the zero vector of $C^\infty(\mathbb{R})$ lies in $\mathbb{R}[x]$.
2. If p and q are polynomials, then $p+q$ is a polynomial.
3. If p is a polynomial and $c \in \mathbb{R}$, then cp is a polynomial.

Therefore, $\mathbb{R}[x]$ is a subspace of $C^\infty(\mathbb{R})$. □

Identifying polynomials with functions does not lose any information: if two polynomials $p \neq q$ defined the same function, then every real number would be a root of the nonzero polynomial $p-q$, but a nonzero polynomial of degree d has at most d roots. Combining Proposition 1.5.4 with the inclusions above, we obtain a chain of subspaces

$$\mathbb{R}[x]_{\leq d} \subseteq \mathbb{R}[x] \subseteq C^\infty(\mathbb{R}) \subseteq \cdots \subseteq C^1(\mathbb{R}) \subseteq C(\mathbb{R}) \subseteq \mathcal{F}(\mathbb{R}).$$

All of them from $\mathbb{R}[x]$ on contain the monomials $1, x, x^2, \ldots$, which are infinitely many linearly independent vectors, so these spaces are infinite-dimensional, since a vector space of dimension n contains at most n linearly independent vectors (Corollary 1.3.14).

Theorem 1.5.5. *Let $T : V \to W$ be a linear transformation. Then the kernel* $\ker(T)$ *is a subspace of V, and the image* $\operatorname{im}(T)$ *is a subspace of W.*

Proof. **1. Image is a subspace of W:**

1. The zero vector of W is in $\operatorname{im}(T)$ since $T(0_V) = 0_W$ by Lemma 1.4.2.
2. Let $w_1, w_2 \in \operatorname{im}(T)$. Then there exist $v_1, v_2 \in V$ such that $T(v_1) = w_1$ and $T(v_2) = w_2$. We have:
$$w_1 + w_2 = T(v_1) + T(v_2) = T(v_1 + v_2) \in \operatorname{im}(T).$$
3. Let $a \in \mathbb{F}$ and $w \in \operatorname{im}(T)$. Then there exists $v \in V$ such that $T(v) = w$. We have:
$$a \cdot w = a \cdot T(v) = T(a \cdot v) \in \operatorname{im}(T).$$

Therefore, $\operatorname{im}(T)$ is a subspace of W.

2. Kernel is a subspace of V:

1. The zero vector of V is in $\ker(T)$ since $T(0_V) = 0_W$ by Lemma 1.4.2.
2. Let $v_1, v_2 \in \ker(T)$. Then we have:
$$T(v_1 + v_2) = T(v_1) + T(v_2) = 0_W + 0_W = 0_W,$$
which implies that $v_1 + v_2 \in \ker(T)$.
3. Let $a \in \mathbb{F}$ and $v \in \ker(T)$. Then we have:
$$T(a \cdot v) = a \cdot T(v) = a \cdot 0_W = 0_W,$$
which implies that $a \cdot v \in \ker(T)$.

Therefore, $\ker(T)$ is a subspace of V. □

For example, the constant functions form a subspace of $C^1(I)$, since they are the kernel of differentiation (Example 1.4.14), and the functions $f \in C^2(\mathbb{R})$ with $f'' + f = 0$, such as sin and cos, form a subspace of $C^2(\mathbb{R})$, since they are the kernel of the linear transformation $f \mapsto f'' + f$.

1.6 Matrices

Definition 1.6.1 (Matrix)**.** A matrix is a rectangular array of numbers arranged in rows and columns. An $m \times n$ matrix has m rows and n columns, and is typically denoted by a capital letter, such as A. The entry in the i-th row and j-th column of the matrix A is denoted by A_{ij}.

Example 1.6.2. Let A be a 2×3 matrix given by

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}.$$

The entry in the first row and second column is $A_{12} = 2$, and the entry in the second row and third column is $A_{23} = 6$.

Definition 1.6.3 (Transpose)**.** The transpose of an $m \times n$ matrix A is the $n \times m$ matrix $A^\top$ with entries $(A^\top)_{ij} = A_{ji}$. In other words, the rows of A become the columns of $A^\top$.

Example 1.6.4. For the matrix A from the previous example,

$$A^\top = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix}.$$

When working with matrices, we write vectors of $\mathbb{R}^n$ as columns, i.e., as $n \times 1$ matrices. To save space, we often write such a column vector as the transpose of a row:

$$\mathbf{x} = (x_1, x_2, \ldots, x_n)^\top = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Definition 1.6.5 (Matrix–vector product)**.** Let A be an $m \times n$ matrix and $\mathbf{x} = (x_1, \ldots, x_n)^\top \in \mathbb{R}^n$. The product $A\mathbf{x}$ is the vector in $\mathbb{R}^m$ whose i-th entry is

$$(A\mathbf{x})_i = A_{i1}x_1 + A_{i2}x_2 + \cdots + A_{in}x_n, \qquad i = 1, \ldots, m.$$

Note that $A\mathbf{x}$ is only defined when the number of columns of A equals the number of entries of $\mathbf{x}$.

Remark 1.6.6. If $\mathbf{a}_1, \mathbf{a}_2, \ldots, \mathbf{a}_n \in \mathbb{R}^m$ denote the columns of A, then

$$A\mathbf{x} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n,$$

since the i-th entry of the right-hand side is $x_1A_{i1} + x_2A_{i2} + \cdots + x_nA_{in}$. In words: $A\mathbf{x}$ is the linear combination of the columns of A whose coefficients are the entries of $\mathbf{x}$.

Example 1.6.7. Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$ and $\mathbf{x} = (1, 0, -1)^\top$. Then

$$A\mathbf{x} = \begin{pmatrix} 1 \cdot 1 + 2 \cdot 0 + 3 \cdot (-1) \\ 4 \cdot 1 + 5 \cdot 0 + 6 \cdot (-1) \end{pmatrix} = \begin{pmatrix} -2 \\ -2 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ 4 \end{pmatrix} + 0 \cdot \begin{pmatrix} 2 \\ 5 \end{pmatrix} + (-1) \cdot \begin{pmatrix} 3 \\ 6 \end{pmatrix}.$$

Remark 1.6.8. For any $m \times n$ matrix A, the map $\mathbf{x} \mapsto A\mathbf{x}$ is a linear transformation $\mathbb{R}^n \to \mathbb{R}^m$. Indeed, for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $c \in \mathbb{R}$,

$$(A(\mathbf{x} + \mathbf{y}))_i = \sum_{j=1}^{n} A_{ij}(x_j + y_j) = \sum_{j=1}^{n} A_{ij}x_j + \sum_{j=1}^{n} A_{ij}y_j = (A\mathbf{x})_i + (A\mathbf{y})_i,$$

and similarly $(A(c\mathbf{x}))_i = c\,(A\mathbf{x})_i$. The next theorem shows that, conversely, every linear transformation $\mathbb{R}^n \to \mathbb{R}^m$ is of this form.

Theorem 1.6.9. *Any linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$ can be represented by an $m \times n$ matrix A such that for any vector $\mathbf{x} \in \mathbb{R}^n$, we have*

$$T(\mathbf{x}) = A\mathbf{x}.$$

The i-th column of A is the vector $T(\mathbf{e}_i)$.

Proof. Let $T : \mathbb{R}^n \to \mathbb{R}^m$ be a linear transformation. We will construct an $m \times n$ matrix A that represents T.

Let $\{\mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_n\}$ be the standard basis for $\mathbb{R}^n$, where $\mathbf{e}_i$ is the vector with a 1 in the i-th position and 0s elsewhere. For each basis vector $\mathbf{e}_i$, we compute $T(\mathbf{e}_i)$, which is a vector in $\mathbb{R}^m$. Let

$$T(\mathbf{e}_i) = \begin{pmatrix} T(\mathbf{e}_i)_1 \\ T(\mathbf{e}_i)_2 \\ \vdots \\ T(\mathbf{e}_i)_m \end{pmatrix}.$$

We define the matrix A as the matrix whose i-th column is $T(\mathbf{e}_i)$:

$$A = \begin{pmatrix} T(\mathbf{e}_1)_1 & T(\mathbf{e}_2)_1 & \cdots & T(\mathbf{e}_n)_1 \\ T(\mathbf{e}_1)_2 & T(\mathbf{e}_2)_2 & \cdots & T(\mathbf{e}_n)_2 \\ \vdots & \vdots & \ddots & \vdots \\ T(\mathbf{e}_1)_m & T(\mathbf{e}_2)_m & \cdots & T(\mathbf{e}_n)_m \end{pmatrix}.$$

Now, for any vector $\mathbf{x} = (x_1, x_2, \ldots, x_n)^\top \in \mathbb{R}^n$, we can express $\mathbf{x}$ as a linear combination of the basis vectors:

$$\mathbf{x} = x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \cdots + x_n\mathbf{e}_n.$$

Applying the linear transformation T to $\mathbf{x}$ and using additivity and homogeneity, we have:

$$T(\mathbf{x}) = T(x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \cdots + x_n\mathbf{e}_n) = x_1T(\mathbf{e}_1) + x_2T(\mathbf{e}_2) + \cdots + x_nT(\mathbf{e}_n).$$

By Remark 1.6.6, the right-hand side is exactly the linear combination of the columns of A with coefficients $x_1, \ldots, x_n$, that is,

$$T(\mathbf{x}) = A\mathbf{x}.$$

Therefore, we have shown that any linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$ can be represented by an $m \times n$ matrix A such that $T(\mathbf{x}) = A\mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^n$. □

Example 1.6.10. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (x - y, 2x + 3y)$. We will find the matrix representation of T.

First, we compute $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$, where $\mathbf{e}_1 = (1, 0)$ and $\mathbf{e}_2 = (0, 1)$ are the standard basis vectors for $\mathbb{R}^2$:

$$T(\mathbf{e}_1) = T(1, 0) = (1 - 0, 2 \cdot 1 + 3 \cdot 0) = (1, 2),$$
$$T(\mathbf{e}_2) = T(0, 1) = (0 - 1, 2 \cdot 0 + 3 \cdot 1) = (-1, 3).$$

Now we can construct the matrix A that represents T:

$$A = \begin{pmatrix} T(\mathbf{e}_1)_1 & T(\mathbf{e}_2)_1 \\ T(\mathbf{e}_1)_2 & T(\mathbf{e}_2)_2 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix}.$$

Therefore, the matrix representation of the linear transformation T is

$$A = \begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix}.$$

Example 1.6.11. Let $T : \mathbb{R}^3 \to \mathbb{R}^2$ be defined by $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$. We will find the matrix representation of T.

First, we compute $T(\mathbf{e}_1)$, $T(\mathbf{e}_2)$, and $T(\mathbf{e}_3)$, where $\mathbf{e}_1 = (1, 0, 0)$, $\mathbf{e}_2 = (0, 1, 0)$, and $\mathbf{e}_3 = (0, 0, 1)$ are the standard basis vectors for $\mathbb{R}^3$:

$$\begin{aligned} T(\mathbf{e}_1) &= T(1, 0, 0) = (1 + 2 \cdot 0 - 0, 3 \cdot 1 - 0 + 4 \cdot 0) = (1, 3), \\ T(\mathbf{e}_2) &= T(0, 1, 0) = (0 + 2 \cdot 1 - 0, 3 \cdot 0 - 1 + 4 \cdot 0) = (2, -1), \\ T(\mathbf{e}_3) &= T(0, 0, 1) = (0 + 2 \cdot 0 - 1, 3 \cdot 0 - 0 + 4 \cdot 1) = (-1, 4). \end{aligned}$$

Now we can construct the matrix A that represents T:

$$A = \begin{pmatrix} T(\mathbf{e}_1)_1 & T(\mathbf{e}_2)_1 & T(\mathbf{e}_3)_1 \\ T(\mathbf{e}_1)_2 & T(\mathbf{e}_2)_2 & T(\mathbf{e}_3)_2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & -1 \\ 3 & -1 & 4 \end{pmatrix}.$$

Therefore, the matrix representation of the linear transformation T is

$$A = \begin{pmatrix} 1 & 2 & -1 \\ 3 & -1 & 4 \end{pmatrix}.$$

1.6.1 Matrix operations

So far, we have used matrices to represent linear transformations. In this subsection, we learn how to add and multiply matrices; inverses of matrices are discussed in Subsection 1.6.3, after we have developed row reduction. We write $\mathbb{R}^{m \times n}$ for the set of all $m \times n$ matrices with real entries. For $A \in \mathbb{R}^{m \times n}$, we write $T_A : \mathbb{R}^n \to \mathbb{R}^m$ for the linear transformation $T_A(\mathbf{x}) = A\mathbf{x}$.

Matrix addition

Definition 1.6.12 (Matrix addition and scalar multiplication)**.** Let A and B be $m \times n$ matrices and $c \in \mathbb{R}$. The sum $A + B$ and the scalar multiple cA are the $m \times n$ matrices with entries

$$(A + B)_{ij} = A_{ij} + B_{ij}, \qquad (cA)_{ij} = cA_{ij}.$$

In other words, matrices are added and multiplied by scalars entry by entry. The sum $A + B$ is only defined when A and B have the same size.

Example 1.6.13. Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}, \qquad B = \begin{pmatrix} 0 & -1 & 2 \\ 3 & 1 & -2 \end{pmatrix}.$$

Then

$$A + B = \begin{pmatrix} 1 + 0 & 2 + (-1) & 3 + 2 \\ 4 + 3 & 5 + 1 & 6 + (-2) \end{pmatrix} = \begin{pmatrix} 1 & 1 & 5 \\ 7 & 6 & 4 \end{pmatrix},$$

and

$$2A = \begin{pmatrix} 2 & 4 & 6 \\ 8 & 10 & 12 \end{pmatrix}.$$

On the other hand, $A + A^\top$ is not defined, since A is a 2×3 matrix and $A^\top$ is a 3×2 matrix.

Remark 1.6.14. With these operations, $\mathbb{R}^{m \times n}$ is a vector space over $\mathbb{R}$. Indeed, an $m \times n$ matrix is just a list of mn real numbers arranged in a rectangle, and the operations are component-wise, exactly as in $\mathbb{R}^{mn}$. The zero vector of this space is the zero matrix 0, all of whose entries are 0, and the additive inverse of A is $-A = (-1)A$. We write $A - B = A + (-B)$. Moreover, $\dim \mathbb{R}^{m \times n} = mn$: a basis consists of the mn matrices that have a single entry equal to 1 and all other entries equal to 0.

Remark 1.6.15. Matrix addition corresponds to adding linear transformations: for every $\mathbf{x} \in \mathbb{R}^n$,

$$(A+B)\mathbf{x} = A\mathbf{x} + B\mathbf{x}, \qquad (cA)\mathbf{x} = c(A\mathbf{x}).$$

Indeed, $((A+B)\mathbf{x})_i = \sum_j (A_{ij}+B_{ij})x_j = \sum_j A_{ij}x_j + \sum_j B_{ij}x_j = (A\mathbf{x})_i + (B\mathbf{x})_i$, and similarly for cA.

Matrix multiplication

Let A be an $m \times n$ matrix and B an $n \times p$ matrix. Then $T_B : \mathbb{R}^p \to \mathbb{R}^n$ and $T_A : \mathbb{R}^n \to \mathbb{R}^m$, so we can first apply T_B and then T_A. The result is the composition

$$T_A \circ T_B : \mathbb{R}^p \to \mathbb{R}^m, \qquad \mathbf{x} \mapsto A(B\mathbf{x}),$$

which is again a linear transformation by Lemma 1.4.24. By Theorem 1.6.9, it is represented by an $m \times p$ matrix, whose j-th column is $A(B\mathbf{e}_j) = A\mathbf{b}_j$, where $\mathbf{b}_j = B\mathbf{e}_j$ is the j-th column of B (Remark 1.6.6). The i-th entry of $A\mathbf{b}_j$ is $\sum_k A_{ik}B_{kj}$. This motivates the following definition.

Definition 1.6.16 (Matrix multiplication)**.** Let A be an $m \times n$ matrix and B an $n \times p$ matrix. The product AB is the $m \times p$ matrix with entries

$$(AB)_{ij} = A_{i1}B_{1j} + A_{i2}B_{2j} + \cdots + A_{in}B_{nj} = \sum_{k=1}^{n} A_{ik}B_{kj}.$$

The product AB is only defined when the number of columns of A equals the number of rows of B. An $m \times n$ matrix times an $n \times p$ matrix is an $m \times p$ matrix.

Remark 1.6.17. There are two useful ways to think about the product AB:

- Entry by entry: $(AB)_{ij}$ is obtained by multiplying the entries of the i-th row of A with the corresponding entries of the j-th column of B, and adding up the results.
- Column by column: the j-th column of AB is $A\mathbf{b}_j$, where $\mathbf{b}_j$ is the j-th column of B:

$$AB = A\begin{pmatrix}\mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_p\end{pmatrix} = \begin{pmatrix}A\mathbf{b}_1 & A\mathbf{b}_2 & \cdots & A\mathbf{b}_p\end{pmatrix}.$$

  In particular, if $p = 1$, then B is a single column $\mathbf{x}$, and AB is the matrix–vector product $A\mathbf{x}$.

Example 1.6.18. Let

$$A = \begin{pmatrix}1 & 2 & 3\\ 4 & 5 & 6\end{pmatrix}, \qquad B = \begin{pmatrix}1 & 0\\ 0 & 1\\ -1 & 2\end{pmatrix}.$$

Since A is 2×3 and B is 3×2, the product AB is a 2×2 matrix. For example, its $(1,1)$ entry is obtained from the first row of A and the first column of B:

$$(AB)_{11} = 1 \cdot 1 + 2 \cdot 0 + 3 \cdot (-1) = -2.$$

Computing all four entries in the same way, we get

$$AB = \begin{pmatrix}1\cdot 1 + 2\cdot 0 + 3\cdot(-1) & 1\cdot 0 + 2\cdot 1 + 3\cdot 2\\ 4\cdot 1 + 5\cdot 0 + 6\cdot(-1) & 4\cdot 0 + 5\cdot 1 + 6\cdot 2\end{pmatrix} = \begin{pmatrix}-2 & 8\\ -2 & 17\end{pmatrix}.$$

Note that the first column of AB is $A(1,0,-1)^\top = (-2,-2)^\top$, which we computed in Example 1.6.7.

The product BA is also defined, but it is a 3×3 matrix:

$$BA = \begin{pmatrix}1 & 2 & 3\\ 4 & 5 & 6\\ 7 & 8 & 9\end{pmatrix}.$$

In particular, $AB \neq BA$.

Theorem 1.6.19. *Let A be an $m \times n$ matrix and B an $n \times p$ matrix. Then for every $\mathbf{x} \in \mathbb{R}^p$,*

$$(AB)\mathbf{x} = A(B\mathbf{x}).$$

In other words, AB is the matrix of the composition $T_A \circ T_B$: multiplying matrices corresponds to composing linear transformations.

Proof. This holds by construction, but we can also check it directly. For each $i = 1, \dots, m$,

$$\begin{aligned}((AB)\mathbf{x})_i &= \sum_{j=1}^{p} (AB)_{ij} x_j = \sum_{j=1}^{p} \sum_{k=1}^{n} A_{ik} B_{kj} x_j \\ &= \sum_{k=1}^{n} A_{ik} \left(\sum_{j=1}^{p} B_{kj} x_j \right) = \sum_{k=1}^{n} A_{ik} (B\mathbf{x})_k = (A(B\mathbf{x}))_i.\end{aligned}$$

□

Example 1.6.20. Let $S, T : \mathbb{R}^2 \to \mathbb{R}^2$ be the linear transformations $S(x, y) = (2x + 3y, 4x - y)$ from Example 1.4.8 and $T(x, y) = (x - y, 2x + 3y)$ from Example 1.6.10. Their matrices are

$$A_S = \begin{pmatrix} 2 & 3 \\ 4 & -1 \end{pmatrix}, \qquad A_T = \begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix}.$$

Computing the composition directly, we get

$$\begin{aligned}T(S(x, y)) &= T(2x + 3y, 4x - y) \\ &= \big((2x + 3y) - (4x - y),\ 2(2x + 3y) + 3(4x - y)\big) \\ &= (-2x + 4y, 16x + 3y).\end{aligned}$$

On the other hand,

$$A_T A_S = \begin{pmatrix} 1 \cdot 2 + (-1) \cdot 4 & 1 \cdot 3 + (-1) \cdot (-1) \\ 2 \cdot 2 + 3 \cdot 4 & 2 \cdot 3 + 3 \cdot (-1) \end{pmatrix} = \begin{pmatrix} -2 & 4 \\ 16 & 3 \end{pmatrix},$$

which is exactly the matrix of $T \circ S$, as Theorem 1.6.19 predicts.

Definition 1.6.21 (Identity matrix)**.** The $n \times n$ identity matrix I_n is the matrix with 1s on the diagonal and 0s everywhere else:

$$(I_n)_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j, \end{cases} \qquad I_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Its columns are the standard basis vectors $\mathbf{e}_1, \dots, \mathbf{e}_n$, so $I_n \mathbf{x} = \mathbf{x}$ for every $\mathbf{x} \in \mathbb{R}^n$, i.e., I_n represents the identity map. When the size is clear, we simply write I.

Proposition 1.6.22. *Let A, B, C be matrices and $c \in \mathbb{R}$. Whenever the sizes are such that the sums and products below are defined, we have:*

1. *Associativity: $(AB)C = A(BC)$;*
2. *Distributivity: $A(B + C) = AB + AC$ and $(A + B)C = AC + BC$;*
3. *Scalars: $c(AB) = (cA)B = A(cB)$;*
4. *Identity: $I_m A = A = A I_n$ for every $m \times n$ matrix A;*

5. *Transpose:* $(AB)^\top = B^\top A^\top$.

Proof. All five properties are checked by comparing entries. For associativity,

$$\begin{aligned}((AB)C)_{ij} &= \sum_l (AB)_{il} C_{lj} = \sum_l \sum_k A_{ik} B_{kl} C_{lj} \\ &= \sum_k A_{ik} \left(\sum_l B_{kl} C_{lj} \right) = \sum_k A_{ik} (BC)_{kj} = (A(BC))_{ij}.\end{aligned}$$

For the first distributive law, $(A(B+C))_{ij} = \sum_k A_{ik}(B_{kj} + C_{kj}) = \sum_k A_{ik}B_{kj} + \sum_k A_{ik}C_{kj} = (AB + AC)_{ij}$; the second distributive law and the rule for scalars are checked in the same way. For the identity, $(I_m A)_{ij} = \sum_k (I_m)_{ik} A_{kj} = A_{ij}$, since the only nonzero term is the one with $k = i$; similarly, $AI_n = A$. Finally, for the transpose,

$$((AB)^\top)_{ij} = (AB)_{ji} = \sum_k A_{jk} B_{ki} = \sum_k (B^\top)_{ik} (A^\top)_{kj} = (B^\top A^\top)_{ij}. \qquad \square$$

Remark 1.6.23. Matrix multiplication is not commutative: even when AB and BA are both defined and have the same size, in general $AB \neq BA$. For example,

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}.$$

This matches the fact that composing two linear transformations in a different order usually gives a different result. Another difference from numbers is that a product of two nonzero matrices can be zero:

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

As a consequence, we cannot "cancel" matrices: $AB = AC$ does not imply $B = C$ in general.

1.6.2 Row Reduction

Definition 1.6.24 (Row echelon form)**.** The leading entry (also called the pivot) of a nonzero row of a matrix is its first nonzero entry from the left. A matrix is said to be in row echelon form (REF) if it satisfies the following conditions:

1. All nonzero rows are above any rows of all zeros.
2. The leading entry of each nonzero row after the first occurs to the right of the leading entry of the previous row.
3. The leading entry of each nonzero row is 1.

Remark 1.6.25. Conditions 1 and 2 imply that all entries below a leading entry are zero. Some textbooks do not require condition 3, i.e., they allow leading entries to be any nonzero numbers. In these notes, we always scale the leading entries to be 1.

Example 1.6.26. Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 0 \end{pmatrix}.$$

This matrix is in row echelon form because:

1. The nonzero rows are above the row of all zeros.

2. The leading entry of the second row (in column 2) is to the right of the leading entry of the first row (in column 1).

3. Both leading entries are equal to 1.

Note that nonzero entries above a leading 1, such as the 2 in the first row, are allowed.

Example 1.6.27. Consider the matrix

$$B = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{pmatrix}.$$

This matrix is also in row echelon form because:

1. There are no rows of all zeros, so the first condition is trivially satisfied.

2. The leading entry of the second row (in column 2) is to the right of the leading entry of the first row (in column 1), and the leading entry of the third row (in column 3) is to the right of the leading entry of the second row.

3. All three leading entries are equal to 1.

Example 1.6.28. Consider the matrix

$$C = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{pmatrix}.$$

This matrix is not in row echelon form because the leading entry of the second row is 3, not 1, so the third condition fails. The first two conditions do hold: the row of zeros is at the bottom, and the leading entries are in columns 2 and 3, so the leading entry of the second row is to the right of the leading entry of the first row. (A first column consisting of zeros is allowed.) Multiplying the second row by $1/3$ gives a matrix in row echelon form.

Example 1.6.29. Consider the matrix

$$D = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 1 & 4 \end{pmatrix}.$$

This matrix is not in row echelon form because the nonzero row (the third row) is below a row of all zeros (the second row). For it to be in row echelon form, all nonzero rows must be above any rows of all zeros.

To transform a matrix into row echelon form, we can use elementary row operations, which include:

1. Swapping two rows.

2. Multiplying a row by a nonzero scalar.

3. Adding or subtracting a multiple of one row to another row.

Each elementary row operation can be undone by another elementary row operation: we can swap the same two rows again, multiply the same row by the inverse of the scalar, or subtract (add) the same multiple back. Using these operations, every matrix can be brought into row echelon form; this procedure is called Gaussian elimination.

Example 1.6.30. Consider the matrix

$$E = \begin{pmatrix} 2 & 4 & 6 \\ 1 & 3 & 5 \\ 0 & 0 & 0 \end{pmatrix}.$$

We will transform this matrix into row echelon form using elementary row operations.

First, we swap the first and second rows, so that the first row starts with a 1:

$$\begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \\ 0 & 0 & 0 \end{pmatrix}.$$

Next, we can eliminate the leading entry in the second row by subtracting 2 times the first row from the second row:

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & -2 & -4 \\ 0 & 0 & 0 \end{pmatrix}.$$

Finally, we can multiply the second row by $-1/2$ to make the leading entry of the second row equal to 1:

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

The resulting matrix is now in row echelon form.

Example 1.6.31. Consider the matrix

$$F = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}.$$

We will transform this matrix into row echelon form using elementary row operations.

First, we can eliminate the leading entry in the second row by subtracting 4 times the first row from the second row:

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 7 & 8 & 9 \end{pmatrix}.$$

Next, we can eliminate the leading entry in the third row by subtracting 7 times the first row from the third row:

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix}.$$

Now, we can eliminate the leading entry in the third row by subtracting 2 times the second row from the third row:

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 0 \end{pmatrix}.$$

Finally, we can multiply the second row by $-1/3$ to make the leading entry of the second row equal to 1:

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

The resulting matrix is now in row echelon form.

Sometimes it is useful to further simplify a matrix in row echelon form.

Definition 1.6.32 (Reduced row echelon form)**.** A matrix is in reduced row echelon form (RREF) if it is in row echelon form and, in addition, each leading 1 is the only nonzero entry in its column.

For example, the matrices A and B in Examples 1.6.26 and 1.6.27 are in row echelon form, but not in reduced row echelon form, because they have nonzero entries above some of their leading 1s. Every matrix A can be brought into reduced row echelon form by elementary row operations, and one can show that the result does not depend on which operations we use; we denote it by $\mathrm{RREF}(A)$.

Example 1.6.33. Consider the matrix

$$G = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix},$$

which is the row echelon form of the matrix F that we found in Example 1.6.31. We will transform this matrix into reduced row echelon form (RREF).

We can eliminate the entry above the leading 1 in the second row by subtracting 2 times the second row from the first row:

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

The resulting matrix is now in reduced row echelon form (RREF).

Example 1.6.34. Consider the matrix

$$H = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix}.$$

We will transform this matrix into reduced row echelon form (RREF).

First, we can eliminate the entry above the leading 1 in the third row by subtracting 4 times the third row from the second row:

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Next, we can eliminate the entry above the leading 1 in the second row by subtracting 2 times the second row from the first row:

$$\begin{pmatrix} 1 & 0 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Finally, we can eliminate the entry above the leading 1 in the third row by subtracting 3 times the third row from the first row:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The resulting matrix is now in reduced row echelon form (RREF).

1.6.3 The Inverse of a Matrix

Definition and basic properties

For a nonzero number a, dividing by a means multiplying by its inverse a^{-1}, the number with $a \cdot a^{-1} = a^{-1} \cdot a = 1$. For square matrices, the role of the number 1 is played by the identity matrix.

Definition 1.6.35 (Inverse matrix)**.** An $n \times n$ matrix A is called invertible if there exists an $n \times n$ matrix B such that

$$AB = BA = I_n.$$

Such a matrix B is called an inverse of A. A square matrix that is not invertible is called singular. (We only consider inverses of square matrices.)

Proposition 1.6.36. *An invertible matrix A has exactly one inverse, which we denote by A^{-1}.*

Proof. Suppose that B and C are both inverses of A. Then, using associativity,

$$B = BI_n = B(AC) = (BA)C = I_nC = C. \qquad \square$$

Example 1.6.37. Let

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}, \qquad B = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}.$$

Then

$$AB = \begin{pmatrix} 2 \cdot 1 + 1 \cdot (-1) & 2 \cdot (-1) + 1 \cdot 2 \\ 1 \cdot 1 + 1 \cdot (-1) & 1 \cdot (-1) + 1 \cdot 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

and in the same way $BA = I_2$. So A is invertible and $A^{-1} = B$.

Remark 1.6.38. If A is invertible, then for every $\mathbf{b} \in \mathbb{R}^n$ the equation $A\mathbf{x} = \mathbf{b}$ has exactly one solution, namely $\mathbf{x} = A^{-1}\mathbf{b}$. Indeed, $\mathbf{x} = A^{-1}\mathbf{b}$ is a solution, since $A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$; and any solution $\mathbf{x}$ satisfies $\mathbf{x} = I_n\mathbf{x} = (A^{-1}A)\mathbf{x} = A^{-1}(A\mathbf{x}) = A^{-1}\mathbf{b}$. In particular, the only solution of $A\mathbf{x} = \mathbf{0}$ is $\mathbf{x} = \mathbf{0}$. In terms of linear transformations, $T_{A^{-1}}$ is the inverse function of T_A: it undoes what T_A does.

Example 1.6.39. Not every nonzero square matrix is invertible. By Remark 1.6.38, if $A\mathbf{x} = \mathbf{0}$ for some $\mathbf{x} \neq \mathbf{0}$, then A is not invertible. For example, the matrix

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$$

is singular, since $A(2, -1)^\top = (2 - 2, 4 - 4)^\top = (0, 0)^\top$.

To find the inverse of a matrix A, we can solve the equation $A\mathbf{x} = \mathbf{y}$ for $\mathbf{x}$ in terms of $\mathbf{y}$.

Proposition 1.6.40. *Let A and B be $n \times n$ matrices. Suppose that for every $\mathbf{y} \in \mathbb{R}^n$, the equation $A\mathbf{x} = \mathbf{y}$ has exactly one solution, and that this solution is $\mathbf{x} = B\mathbf{y}$. Then A is invertible and $A^{-1} = B$.*

Proof. For every $\mathbf{y} \in \mathbb{R}^n$, the vector $B\mathbf{y}$ is a solution of $A\mathbf{x} = \mathbf{y}$, so $(AB)\mathbf{y} = A(B\mathbf{y}) = \mathbf{y}$. Taking $\mathbf{y} = \mathbf{e}_j$ shows that the j-th column of AB is $\mathbf{e}_j$, so $AB = I_n$.

Now let $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{y} = A\mathbf{x}$. Then both $\mathbf{x}$ and $B\mathbf{y}$ are solutions of the equation $Az = \mathbf{y}$. Since this equation has exactly one solution, $\mathbf{x} = B\mathbf{y} = (BA)\mathbf{x}$. Taking $\mathbf{x} = \mathbf{e}_j$ shows that the j-th column of BA is $\mathbf{e}_j$, so $BA = I_n$. $\square$

Example 1.6.41. Let

$$A = \begin{pmatrix} 2 & 3 \\ 4 & -1 \end{pmatrix}$$

be the matrix of the linear transformation $T(x, y) = (2x+3y, 4x-y)$. In Example 1.4.11, we solved the equation $T(x, y) = (u, v)$, i.e., $A(x, y)^\top = (u, v)^\top$. Each step of that computation was forced, so the solution is unique, and it is

$$x = \frac{u + 3v}{14}, \qquad y = \frac{2u - v}{7} = \frac{4u - 2v}{14}.$$

In matrix form,

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{14} \begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

By Proposition 1.6.40,

$$A^{-1} = \frac{1}{14} \begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix}.$$

We can double-check this:

$$AA^{-1} = \frac{1}{14} \begin{pmatrix} 2 \cdot 1 + 3 \cdot 4 & 2 \cdot 3 + 3 \cdot (-2) \\ 4 \cdot 1 + (-1) \cdot 4 & 4 \cdot 3 + (-1) \cdot (-2) \end{pmatrix} = \frac{1}{14} \begin{pmatrix} 14 & 0 \\ 0 & 14 \end{pmatrix} = I_2.$$

Proposition 1.6.42. *Let A and B be invertible $n \times n$ matrices. Then:*

1. *A^{-1} is invertible and $(A^{-1})^{-1} = A$;*
2. *AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$;*
3. *$A^\top$ is invertible and $(A^\top)^{-1} = (A^{-1})^\top$.*

Proof. (1) The equations $AA^{-1} = A^{-1}A = I_n$ say exactly that A is an inverse of A^{-1}.

(2) Using associativity,

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AI_nA^{-1} = AA^{-1} = I_n,$$

and similarly $(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}B = I_n$.

(3) Using $(AB)^\top = B^\top A^\top$ and $I_n^\top = I_n$, we get

$$A^\top (A^{-1})^\top = (A^{-1}A)^\top = I_n^\top = I_n, \qquad (A^{-1})^\top A^\top = (AA^{-1})^\top = I_n. \qquad \square$$

Note the reversed order in part 2: to undo "first T_B, then T_A", we must first undo T_A and then undo T_B. By induction, every product of invertible matrices is invertible, with $(A_1A_2 \cdots A_k)^{-1} = A_k^{-1} \cdots A_2^{-1}A_1^{-1}$.

Computing the inverse with row reduction

For larger matrices, solving $A\mathbf{x} = \mathbf{y}$ by hand as in Example 1.6.41 quickly becomes tedious. We now use row reduction to give a systematic method for deciding whether a square matrix is invertible and for computing its inverse. The key observation is that elementary row operations can be performed by matrix multiplication.

Definition 1.6.43 (Elementary matrix)**.** An elementary matrix is a matrix obtained from an identity matrix I_m by a single elementary row operation.

Example 1.6.44. The following are 3×3 elementary matrices. They are obtained from I_3 by swapping rows 1 and 2, by multiplying row 2 by a scalar $c \neq 0$, and by adding c times row 1 to row 3, respectively:

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ c & 0 & 1 \end{pmatrix}.$$

Multiplying a 3×2 matrix from the left by the last of them gives

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ c & 0 & 1 \end{pmatrix} \begin{pmatrix} p_1 & p_2 \\ q_1 & q_2 \\ r_1 & r_2 \end{pmatrix} = \begin{pmatrix} p_1 & p_2 \\ q_1 & q_2 \\ r_1 + cp_1 & r_2 + cp_2 \end{pmatrix},$$

which is exactly the result of adding c times row 1 to row 3.

Lemma 1.6.45. *Let E be the elementary matrix obtained from I_m by some elementary row operation. Then for every $m \times n$ matrix A, the product EA is the matrix obtained from A by the same row operation.*

Proof. By the definition of matrix multiplication, $(EA)_{ij} = \sum_k E_{ik}A_{kj}$, so the i-th row of EA is

$$E_{i1} \cdot (\text{row 1 of } A) + E_{i2} \cdot (\text{row 2 of } A) + \cdots + E_{im} \cdot (\text{row } m \text{ of } A).$$

On the other hand, every elementary row operation replaces each row by a linear combination of the old rows, with coefficients that depend only on the operation and not on the matrix. For example, adding c times row 1 to row 3 replaces row 3 by $c \cdot (\text{row } 1) + 1 \cdot (\text{row } 3)$ and keeps all other rows. Say that the operation replaces row i by $c_{i1} \cdot (\text{row } 1) + \cdots + c_{im} \cdot (\text{row } m)$. Applying it to I_m, whose k-th row is $\mathbf{e}_k^\top$, the new i-th row is $c_{i1}\mathbf{e}_1^\top + \cdots + c_{im}\mathbf{e}_m^\top = (c_{i1}, \ldots, c_{im})$. Since this is the i-th row of E, we get $E_{ik} = c_{ik}$ for all i, k. Therefore, the i-th row of EA is $c_{i1} \cdot (\text{row 1 of } A) + \cdots + c_{im} \cdot (\text{row } m \text{ of } A)$, which is the i-th row of the matrix obtained from A by the row operation. □

Lemma 1.6.46. *Every elementary matrix E is invertible, and E^{-1} is the elementary matrix of the row operation that undoes the operation of E.*

Proof. Let E' be the elementary matrix of the row operation that undoes the operation of E. The matrix E is obtained from I_m by the operation of E, and by Lemma 1.6.45, $E'E$ is obtained from E by undoing this operation. Hence $E'E = I_m$. In the same way, $EE' = I_m$. □

As a consequence, if a matrix R is obtained from A by row operations with elementary matrices $E_1, E_2, \ldots, E_k$ (in this order), then

$$R = BA, \qquad \text{where } B = E_k \cdots E_2 E_1$$

is invertible as a product of invertible matrices (Proposition 1.6.42).

Theorem 1.6.47. *For an $n \times n$ matrix A, the following statements are equivalent:*

1. *A is invertible;*
2. *the only solution of $A\mathbf{x} = \mathbf{0}$ is $\mathbf{x} = \mathbf{0}$;*
3. $\mathrm{RREF}(A) = I_n$.

Proof. (1 ⇒ 2) This was shown in Remark 1.6.38.

(2 ⇒ 3) Suppose that $R = \mathrm{RREF}(A) \neq I_n$; we will find a nonzero solution of $A\mathbf{x} = \mathbf{0}$. If R had n pivots, then each of its n rows would contain a pivot, and the pivot columns $p_1 < p_2 < \cdots < p_n$ would be n different numbers among $1, \ldots, n$, so $p_i = i$ for every i. Since each pivot is a 1 and is the only nonzero entry in its column, this would mean $R = I_n$. So R has fewer than n pivots, and some column j of R contains no pivot. Set $x_j = 1$, and $x_k = 0$ for every other column k without a pivot. Then the equation given by each nonzero row i of R becomes $x_{p_i} + R_{ij} = 0$, so we set $x_{p_i} = -R_{ij}$. This gives a vector $\mathbf{x} \neq \mathbf{0}$ with $R\mathbf{x} = \mathbf{0}$. Writing $R = BA$ with B invertible as above, we get

$$A\mathbf{x} = B^{-1}(BA)\mathbf{x} = B^{-1}(R\mathbf{x}) = B^{-1}\mathbf{0} = \mathbf{0}.$$

(3 ⇒ 1) Let $E_1, \ldots, E_k$ be the elementary matrices of the row operations that bring A to its reduced row echelon form $\mathrm{RREF}(A) = I_n$, and let $B = E_k \cdots E_1$. Then $BA = I_n$, and B is invertible. Multiplying from the left by B^{-1} gives $A = B^{-1}$, so A is invertible with $A^{-1} = (B^{-1})^{-1} = B$. □

Corollary 1.6.48. *Let A and B be $n \times n$ matrices with $BA = I_n$. Then A is invertible and $A^{-1} = B$. In particular, also $AB = I_n$.*

Proof. If $A\mathbf{x} = \mathbf{0}$, then $\mathbf{x} = I_n\mathbf{x} = (BA)\mathbf{x} = B(A\mathbf{x}) = B\mathbf{0} = \mathbf{0}$. By Theorem 1.6.47, A is invertible, and

$$B = BI_n = B(AA^{-1}) = (BA)A^{-1} = I_nA^{-1} = A^{-1}.$$ □

So, to check that a square matrix B is the inverse of A, it is enough to check one of the two equations $BA = I_n$ or $AB = I_n$. (In the second case, apply the corollary with the roles of A and B swapped: then B is invertible with $B^{-1} = A$, and hence $A^{-1} = B$.)

Theorem 1.6.49 (Gauss–Jordan method)**.** *Let A be an $n \times n$ matrix, and row reduce the $n \times 2n$ matrix $(A \mid I_n)$, obtained by writing I_n to the right of A.*

- *If the left block becomes I_n, then A is invertible, and the right block is A^{-1}:*

$$(A \mid I_n) \quad \longrightarrow \quad (I_n \mid A^{-1}).$$

- *If a row of zeros appears in the left block, then A is not invertible.*

Proof. Let $E_1, \ldots, E_k$ be the elementary matrices of the row operations we perform, and let $B = E_k \cdots E_1$. By Lemma 1.6.45 and the column-by-column description of matrix multiplication (Remark 1.6.17), the result is

$$B\,(A \mid I_n) = (BA \mid BI_n) = (BA \mid B).$$

In particular, the left block BA is what we get by applying the same row operations to A alone.

If $BA = I_n$, then A is invertible and $A^{-1} = B$ by Corollary 1.6.48, and B is exactly the right block.

If BA has a row of zeros, then BA is not invertible: for every $n \times n$ matrix M, the product $(BA)M$ also has a row of zeros, so it cannot be equal to I_n. If A were invertible, then BA would be invertible as a product of invertible matrices. Hence A is not invertible. □

One of the two cases always occurs: if $\mathrm{RREF}(A) \neq I_n$, then, as we saw in the proof of Theorem 1.6.47, $\mathrm{RREF}(A)$ has fewer than n pivots, and hence a row of zeros.

Example 1.6.50. Let us compute the inverse of the matrix A from Example 1.6.41 with the Gauss–Jordan method. We start with

$$\left(\begin{array}{cc|cc} 2 & 3 & 1 & 0 \\ 4 & -1 & 0 & 1 \end{array}\right).$$

Subtracting 2 times the first row from the second row gives

$$\left(\begin{array}{cc|cc} 2 & 3 & 1 & 0 \\ 0 & -7 & -2 & 1 \end{array}\right).$$

Multiplying the second row by $-1/7$ gives

$$\left(\begin{array}{cc|cc} 2 & 3 & 1 & 0 \\ 0 & 1 & \frac{2}{7} & -\frac{1}{7} \end{array}\right).$$

Subtracting 3 times the second row from the first row gives

$$\left(\begin{array}{cc|cc} 2 & 0 & \frac{1}{7} & \frac{3}{7} \\ 0 & 1 & \frac{2}{7} & -\frac{1}{7} \end{array}\right).$$

Finally, multiplying the first row by $1/2$ gives

$$\left(\begin{array}{cc|cc} 1 & 0 & \frac{1}{14} & \frac{3}{14} \\ 0 & 1 & \frac{2}{7} & -\frac{1}{7} \end{array}\right).$$

The left block is I_2, so

$$A^{-1} = \begin{pmatrix} \frac{1}{14} & \frac{3}{14} \\ \frac{2}{7} & -\frac{1}{7} \end{pmatrix} = \frac{1}{14}\begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix},$$

as we found in Example 1.6.41.

Example 1.6.51. Let us find the inverse of

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

We start with

$$\left(\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}\right).$$

Subtracting the first row from the third row gives

$$\left(\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array}\right).$$

The left block is now in row echelon form, and we continue to its reduced row echelon form.

Subtracting the third row from the second row gives

$$\left(\begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & -1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array}\right).$$

Subtracting the second row from the first row gives

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & -1 & 1 \\ 0 & 1 & 0 & 1 & 1 & -1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array}\right).$$

The left block is I_3, so

$$A^{-1} = \begin{pmatrix} 0 & -1 & 1 \\ 1 & 1 & -1 \\ -1 & 0 & 1 \end{pmatrix}.$$

By Corollary 1.6.48, to double-check the answer it is enough to verify that $AA^{-1} = I_3$; try it!

Example 1.6.52. Let us apply the Gauss–Jordan method to the singular matrix of Example 1.6.39:

$$\left(\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 2 & 4 & 0 & 1 \end{array}\right).$$

Subtracting 2 times the first row from the second row gives

$$\left(\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & 0 & -2 & 1 \end{array}\right).$$

A row of zeros has appeared in the left block, so the matrix is not invertible, as we already knew.

Remark 1.6.53. In practice, if we only want to solve $A\mathbf{x} = \mathbf{b}$ for one particular $\mathbf{b}$, we do not need A^{-1}: it is faster, and numerically more accurate, to row reduce the matrix $(A \mid \mathbf{b})$ directly. If A is invertible, the row operations that turn A into I_n turn $(A \mid \mathbf{b})$ into $(I_n \mid A^{-1}\mathbf{b})$, so the solution can be read off from the last column. This is also what numerical libraries do: for example, in NumPy one uses `np.linalg.solve(A, b)` rather than `np.linalg.inv(A) @ b`.

Invertible matrices and isomorphisms

In Section 1.4, we defined isomorphisms of vector spaces. We now show that isomorphisms correspond exactly to invertible matrices. We start with the spaces $\mathbb{R}^n$.

Theorem 1.6.54. *Let A be an $m \times n$ matrix. The linear transformation $T_A : \mathbb{R}^n \to \mathbb{R}^m$, $T_A(\mathbf{x}) = A\mathbf{x}$, is an isomorphism if and only if $m = n$ and A is invertible. In this case, $(T_A)^{-1} = T_{A^{-1}}$.*

Proof. Suppose that $m = n$ and A is invertible. Then $A^{-1}(A\mathbf{x}) = (A^{-1}A)\mathbf{x} = \mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^n$, and $A(A^{-1}y) = (AA^{-1})y = y$ for all $y \in \mathbb{R}^n$. So $T_{A^{-1}}$ is an inverse function of T_A. Hence T_A is bijective, i.e., an isomorphism, and $(T_A)^{-1} = T_{A^{-1}}$.

Conversely, suppose that T_A is an isomorphism. Then $\mathbb{R}^n \cong \mathbb{R}^m$, so $n = \dim \mathbb{R}^n = \dim \mathbb{R}^m = m$ by Proposition 1.4.25. Since T_A is injective, Lemma 1.4.18 shows that the only solution of $A\mathbf{x} = \mathbf{0}$ is $\mathbf{x} = \mathbf{0}$. By Theorem 1.6.47, A is invertible. □

To extend this to arbitrary finite-dimensional vector spaces, we represent linear transformations between them by matrices, using coordinates (Definition 1.4.26). As before, we work over $\mathbb{R}$.

Definition 1.6.55 (Matrix of a linear transformation)**.** Let V and W be finite-dimensional vector spaces over $\mathbb{R}$, with ordered bases $\mathcal{B} = (v_1, \ldots, v_n)$ of V and $\mathcal{C} = (w_1, \ldots, w_m)$ of W, and let $T : V \to W$ be a linear transformation. The matrix of T with respect to $\mathcal{B}$ and $\mathcal{C}$ is the $m \times n$ matrix $[T]_{\mathcal{C},\mathcal{B}}$ whose j-th column is the coordinate vector $[T(v_j)]_{\mathcal{C}}$, written as a column. We say that this matrix represents T.

Proposition 1.6.56. *With the notation above, let $A = [T]_{\mathcal{C},\mathcal{B}}$. Then*

$$[T(v)]_{\mathcal{C}} = A\,[v]_{\mathcal{B}} \qquad \text{for every } v \in V.$$

In other words, $\varphi_{\mathcal{C}} \circ T = T_A \circ \varphi_{\mathcal{B}}$.

Proof. Write $v = x_1v_1 + \cdots + x_nv_n$, so that $[v]_{\mathcal{B}} = (x_1, \ldots, x_n)^\top$. Using the linearity of T and of the coordinate map $\varphi_{\mathcal{C}}$ (Theorem 1.4.27), we get

$$[T(v)]_{\mathcal{C}} = [x_1T(v_1) + \cdots + x_nT(v_n)]_{\mathcal{C}} = x_1[T(v_1)]_{\mathcal{C}} + \cdots + x_n[T(v_n)]_{\mathcal{C}}.$$

Since the columns of A are $[T(v_1)]_{\mathcal{C}}, \ldots, [T(v_n)]_{\mathcal{C}}$, the right-hand side is $A[v]_{\mathcal{B}}$ by Remark 1.6.6. □

Remark 1.6.57. If $V = \mathbb{R}^n$ and $W = \mathbb{R}^m$ with their standard bases, then every vector is its own coordinate vector, and $[T]_{\mathcal{C},\mathcal{B}}$ is exactly the matrix of Theorem 1.6.9. With other bases, the same linear transformation is usually represented by a different matrix.

We can now prove the main result of this subsection.

Theorem 1.6.58. *Let V and W be finite-dimensional vector spaces over $\mathbb{R}$, with ordered bases $\mathcal{B}$ of V and $\mathcal{C}$ of W.*

1. *A linear transformation $T : V \to W$ is an isomorphism if and only if the matrix $[T]_{\mathcal{C},\mathcal{B}}$ that represents it is invertible. In this case, $[T^{-1}]_{\mathcal{B},\mathcal{C}} = ([T]_{\mathcal{C},\mathcal{B}})^{-1}$.*
2. *Consequently, V and W are isomorphic if and only if there exists a linear transformation $T : V \to W$ that is represented by an invertible matrix.*

Proof. (1) Let $A = [T]_{\mathcal{C},\mathcal{B}}$. By Proposition 1.6.56, $\varphi_{\mathcal{C}} \circ T = T_A \circ \varphi_{\mathcal{B}}$. Since the coordinate maps are isomorphisms (Theorem 1.4.27), we can compose with their inverses to get

$$T_A = \varphi_{\mathcal{C}} \circ T \circ \varphi_{\mathcal{B}}^{-1} \qquad \text{and} \qquad T = \varphi_{\mathcal{C}}^{-1} \circ T_A \circ \varphi_{\mathcal{B}}.$$

If T is an isomorphism, then so is T_A, as a composition of isomorphisms (Lemma 1.4.24), and A is invertible by Theorem 1.6.54. Conversely, if A is invertible, then T_A is an isomorphism by Theorem 1.6.54, and hence so is T, again as a composition of isomorphisms.

In this case, $T^{-1} = \varphi_{\mathcal{B}}^{-1} \circ (T_A)^{-1} \circ \varphi_{\mathcal{C}} = \varphi_{\mathcal{B}}^{-1} \circ T_{A^{-1}} \circ \varphi_{\mathcal{C}}$, so $[T^{-1}(w)]_{\mathcal{B}} = A^{-1}[w]_{\mathcal{C}}$ for every $w \in W$. Taking for w the j-th vector of $\mathcal{C}$, whose coordinate vector is e_j, shows that the j-th column of $[T^{-1}]_{\mathcal{B},\mathcal{C}}$ is the j-th column of A^{-1}.

(2) If $V \cong W$, choose an isomorphism $T : V \to W$; by part 1, the matrix $[T]_{\mathcal{C},\mathcal{B}}$ that represents it is invertible. Conversely, if some linear transformation $T : V \to W$ is represented by an invertible matrix, then T is an isomorphism by part 1, so $V \cong W$. □

Since only square matrices can be invertible, Theorem 1.6.58 gives another explanation of Corollary 1.4.29: isomorphic spaces must have the same dimension.

Example 1.6.59. The linear transformation $T(x, y) = (2x + 3y, 4x - y)$ of Example 1.4.19 is represented, with respect to the standard bases, by the matrix A of Example 1.6.41. Since A is invertible, T is an isomorphism, and T^{-1} is represented by

$$A^{-1} = \frac{1}{14}\begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix}, \qquad \text{i.e.,} \qquad T^{-1}(u, v) = \left(\frac{u + 3v}{14}, \frac{4u - 2v}{14}\right),$$

in agreement with Example 1.4.19.

Example 1.6.60. Let $\mathbb{R}[x]_{\leq 1}$ be the vector space of polynomials of degree at most 1, and consider the linear transformation

$$T : \mathbb{R}[x]_{\leq 1} \to \mathbb{R}^2, \qquad T(p) = (p(0), p(1)),$$

which evaluates a polynomial at 0 and at 1. We use the ordered basis $\mathcal{B} = (1, x)$ of $\mathbb{R}[x]_{\leq 1}$ and the standard basis $\mathcal{C} = (\mathbf{e}_1, \mathbf{e}_2)$ of $\mathbb{R}^2$. Since $T(1) = (1, 1)$ and $T(x) = (0, 1)$,

$$[T]_{\mathcal{C},\mathcal{B}} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \qquad \text{with inverse} \qquad ([T]_{\mathcal{C},\mathcal{B}})^{-1} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}.$$

(Check that the product of these two matrices is I_2.) By Theorem 1.6.58, T is an isomorphism: a polynomial of degree at most 1 is uniquely determined by its values at 0 and 1, and every pair of values occurs. The inverse is given by $[T^{-1}(u, v)]_{\mathcal{B}} = (u, v - u)^\top$, i.e.,

$$T^{-1}(u, v) = u + (v - u)x.$$

Indeed, this polynomial takes the value u at 0 and the value v at 1.

1.6.4 Special Types of Matrices

Some kinds of square matrices appear so often that they have their own names. Recall from Remark 1.6.14 that the set $\mathbb{R}^{n\times n}$ of all $n \times n$ matrices is a vector space over $\mathbb{R}$. The entries $A_{11}, A_{22}, \ldots, A_{nn}$ of an $n \times n$ matrix A form its (main) diagonal.

Matrices defined by their entries

Definition 1.6.61 (Special types of square matrices)**.** An $n \times n$ matrix A is called

- diagonal if $A_{ij} = 0$ whenever $i \neq j$, i.e., all entries outside the diagonal are zero. We write $\operatorname{diag}(d_1, \ldots, d_n)$ for the diagonal matrix with diagonal entries $d_1, \ldots, d_n$;
- upper triangular if $A_{ij} = 0$ whenever $i > j$, i.e., all entries below the diagonal are zero;
- lower triangular if $A_{ij} = 0$ whenever $i < j$, i.e., all entries above the diagonal are zero;
- symmetric if $A^\top = A$, i.e., $A_{ij} = A_{ji}$ for all i, j;
- skew-symmetric if $A^\top = -A$, i.e., $A_{ij} = -A_{ji}$ for all i, j.

We denote the sets of diagonal, upper triangular, lower triangular, symmetric and skew-symmetric $n \times n$ matrices by $\mathcal{D}_n$, $\mathcal{U}_n$, $\mathcal{L}_n$, Sym_n and Skew_n, respectively.

Example 1.6.62. Here are examples of each type for $n = 3$:

$$\underbrace{\begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 5 \end{pmatrix}}_{\text{diagonal}}, \qquad \underbrace{\begin{pmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{pmatrix}}_{\text{upper triangular}}, \qquad \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{pmatrix}}_{\text{lower triangular}},$$

$$\underbrace{\begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 3 & 5 & 6 \end{pmatrix}}_{\text{symmetric}}, \qquad \underbrace{\begin{pmatrix} 0 & 2 & -3 \\ -2 & 0 & 1 \\ 3 & -1 & 0 \end{pmatrix}}_{\text{skew-symmetric}}.$$

The diagonal matrix above is $\operatorname{diag}(2,-1,5)$. The identity matrix I_n and the zero matrix are diagonal, and every diagonal matrix is upper triangular, lower triangular and symmetric at the same time. The diagonal entries of a skew-symmetric matrix are always zero, since $A_{ii} = -A_{ii}$ forces $A_{ii} = 0$. Finally, every square matrix in row echelon form is upper triangular, since the pivot of row i lies in column i or further to the right.

Example 1.6.63. For any $m \times n$ matrix A, the $n \times n$ matrix $A^\top A$ is symmetric, since by Proposition 1.6.22,

$$(A^\top A)^\top = A^\top (A^\top)^\top = A^\top A.$$

Matrices of this form appear all the time in machine learning, for example in least squares problems and as covariance matrices.

Remark 1.6.64. Products of diagonal matrices are easy to compute: the only nonzero term in $(AB)_{ij} = \sum_k A_{ik}B_{kj}$ can occur for $k = i = j$, so

$$\operatorname{diag}(a_1,\dots,a_n)\operatorname{diag}(b_1,\dots,b_n) = \operatorname{diag}(a_1b_1,\dots,a_nb_n).$$

In particular, if all $d_i \neq 0$, then $\operatorname{diag}(d_1,\dots,d_n)$ is invertible with inverse $\operatorname{diag}(1/d_1,\dots,1/d_n)$. On the other hand, the product of two symmetric matrices need not be symmetric:

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

The following theorem shows that several of these sets are vector spaces in their own right.

Theorem 1.6.65. *The sets $\mathcal{D}_n$, $\mathcal{U}_n$ and Sym_n are subspaces of $\mathbb{R}^{n\times n}$. In particular, each of them is a vector space over $\mathbb{R}$. Moreover, $\dim \mathcal{D}_n = n$.*

Proof. We check the three conditions of Definition 1.5.1. Throughout, let $c \in \mathbb{R}$, and recall that $(A+B)_{ij} = A_{ij} + B_{ij}$ and $(cA)_{ij} = cA_{ij}$.

Diagonal matrices. The zero matrix is diagonal. If $A, B \in \mathcal{D}_n$ and $i \neq j$, then

$$(A+B)_{ij} = 0 + 0 = 0 \qquad \text{and} \qquad (cA)_{ij} = c \cdot 0 = 0,$$

so $A + B \in \mathcal{D}_n$ and $cA \in \mathcal{D}_n$.

Upper triangular matrices. The zero matrix is upper triangular. If $A, B \in \mathcal{U}_n$ and $i > j$, then $(A+B)_{ij} = 0 + 0 = 0$ and $(cA)_{ij} = c \cdot 0 = 0$, so $A + B \in \mathcal{U}_n$ and $cA \in \mathcal{U}_n$.

Symmetric matrices. First, note that the transpose respects addition and scalar multiplication: $((A+B)^\top)_{ij} = (A+B)_{ji} = A_{ji} + B_{ji} = (A^\top + B^\top)_{ij}$, so $(A+B)^\top = A^\top + B^\top$, and similarly $(cA)^\top = cA^\top$. The zero matrix is symmetric. If $A, B \in \operatorname{Sym}_n$, then

$$(A+B)^\top = A^\top + B^\top = A + B \qquad \text{and} \qquad (cA)^\top = cA^\top = cA,$$

so $A + B \in \operatorname{Sym}_n$ and $cA \in \operatorname{Sym}_n$.

Dimension of $\mathcal{D}_n$. For $1 \leq i, j \leq n$, let e_{ij} denote the $n \times n$ matrix with a 1 in position (i,j) and 0s everywhere else. Every diagonal matrix can be written as

$$\operatorname{diag}(d_1,\dots,d_n) = d_1e_{11} + d_2e_{22} + \cdots + d_ne_{nn},$$

so $e_{11},\dots,e_{nn}$ span $\mathcal{D}_n$. If $d_1e_{11} + \cdots + d_ne_{nn} = 0$, then $\operatorname{diag}(d_1,\dots,d_n)$ is the zero matrix, so $d_1 = \cdots = d_n = 0$. Hence $e_{11},\dots,e_{nn}$ are linearly independent, they form a basis of $\mathcal{D}_n$, and $\dim \mathcal{D}_n = n$. □

Exercise 1.6.66. 1. Show that $\mathcal{L}_n$ and Skew_n are subspaces of $\mathbb{R}^{n\times n}$.

2. Show that $\dim \mathcal{U}_n = \dim \mathcal{L}_n = \dim \mathrm{Sym}_n = \frac{n(n+1)}{2}$ and that $\dim \mathrm{Skew}_n = \frac{n(n-1)}{2}$. (Hint: use the matrices e_{ij}.)

3. Show that the product of two upper triangular matrices is upper triangular.

4. Show that every $n \times n$ matrix A can be written in exactly one way as $A = S + K$, where S is symmetric and K is skew-symmetric. (Hint: try $S = \frac{1}{2}(A + A^\top)$.)

Matrix groups

The next two sets of matrices are defined using multiplication rather than by conditions on the entries, and they behave very differently.

Definition 1.6.67 (General linear group)**.** The set of all invertible $n \times n$ matrices is denoted by

$$\mathrm{GL}(n, \mathbb{R}) = \{A \in \mathbb{R}^{n \times n} : A \text{ is invertible}\}.$$

Example 1.6.68. The identity matrix I_n, all elementary matrices (Lemma 1.6.46), all diagonal matrices $\mathrm{diag}(d_1, \ldots, d_n)$ with every $d_i \neq 0$, and the matrices of Example 1.6.41 belong to $\mathrm{GL}(n, \mathbb{R})$ for the appropriate n. The zero matrix and the matrix of Example 1.6.39 do not.

Proposition 1.6.69. *The set* $\mathrm{GL}(n, \mathbb{R})$ *is not a subspace of* $\mathbb{R}^{n \times n}$*. However, it has the following properties:*

1. $I_n \in \mathrm{GL}(n, \mathbb{R})$*;*
2. *if* $A, B \in \mathrm{GL}(n, \mathbb{R})$*, then* $AB \in \mathrm{GL}(n, \mathbb{R})$*;*
3. *if* $A \in \mathrm{GL}(n, \mathbb{R})$*, then* $A^{-1} \in \mathrm{GL}(n, \mathbb{R})$*.*

Proof. The zero matrix is not invertible, since $0 \cdot B = 0 \neq I_n$ for every $n \times n$ matrix B. So $0 \notin \mathrm{GL}(n, \mathbb{R})$, and the first condition of Definition 1.5.1 fails. ($\mathrm{GL}(n, \mathbb{R})$ is not closed under addition either: I_n and $-I_n$ are invertible, but $I_n + (-I_n) = 0$ is not.)

For the three properties: $I_n I_n = I_n$, so I_n is invertible and is its own inverse. Properties 2 and 3 are parts 2 and 1 of Proposition 1.6.42. □

These three properties say that $\mathrm{GL}(n, \mathbb{R})$ is a group, a structure that appears throughout mathematics.

Definition 1.6.70 (Group)**.** A group is a set G with an operation that assigns to $a, b \in G$ an element $ab \in G$, such that:

1. (associativity) $(ab)c = a(bc)$ for all $a, b, c \in G$;
2. (identity) there is an element $e \in G$ with $ea = ae = a$ for all $a \in G$;
3. (inverses) for every $a \in G$, there is an element $a^{-1} \in G$ with $aa^{-1} = a^{-1}a = e$.

The group is abelian if moreover $ab = ba$ for all $a, b \in G$. A subgroup of G is a subset that contains e and is closed under the operation and under taking inverses; it is a group itself, with the same operation.

The identity is unique: if e' is another one, then $e = ee' = e'$. So is the inverse: if b and c are inverses of a, then $b = b(ac) = (ba)c = c$. When the operation is an addition, we write $a + b$, 0 and $-a$ instead of ab, e and a^{-1}.

- $\mathbb{Z}$, $\mathbb{R}$ and every vector space are abelian groups under addition (axioms 1 to 3 of Definition 1.2.1; by commutativity, $v + 0 = v$ and $v + (-v) = 0$ also hold in the other order). The natural numbers $\{0, 1, 2, \ldots\}$ are not a group under addition, since 1 has no inverse.

- The field axioms (Definition 1.1.4) say that $(\mathbb{F}, +)$ and $(\mathbb{F}\backslash\{0\}, \cdot)$ are abelian groups, connected by the distributive law. (The product of nonzero elements is nonzero: if $ab = 0$ with $a \neq 0$, then $b = a^{-1}ab = a^{-1} \cdot 0 = 0$.) The nonzero integers are not a group under multiplication, since 2 has no inverse.
- By the proposition above, $\mathrm{GL}(n, \mathbb{R})$ is a group under matrix multiplication, with the identity $e = I_n$; it is called the general linear group. For $n \geq 2$, it is not abelian (Remark 1.6.23).

A subgroup of $\mathrm{GL}(n, \mathbb{R})$, i.e., a set of invertible matrices that contains I_n and is closed under multiplication and under taking inverses, is called a matrix group. Important matrix groups are the orthogonal matrices $\mathrm{O}(n)$ (below), the matrices with determinant 1, which form the special linear group $\mathrm{SL}(n, \mathbb{R})$ (Subsection 1.6.5), and the rotations, which form the special orthogonal group $\mathrm{SO}(n)$ (Lecture 2). Groups and vector spaces are different kinds of structures: in a vector space we add vectors and multiply them by scalars, while in a matrix group we multiply its elements with each other, and their sums usually leave the group. The matrix groups above are also smooth curved surfaces in $\mathbb{R}^{n\times n}$ on which multiplication and inversion are smooth maps: they are Lie groups, the mathematical description of continuous symmetries such as rotations (Section A.7.1).

Definition 1.6.71 (Orthogonal matrices)**.** An $n \times n$ matrix Q is called orthogonal if $Q^\top Q = I_n$. The set of all orthogonal $n \times n$ matrices is denoted by $\mathrm{O}(n)$.

By Corollary 1.6.48, every orthogonal matrix Q is invertible with $Q^{-1} = Q^\top$; in particular, also $QQ^\top = I_n$, and $\mathrm{O}(n) \subseteq \mathrm{GL}(n, \mathbb{R})$. So for orthogonal matrices, the inverse comes for free: we only need to transpose.

Example 1.6.72. For any angle θ, the rotation matrix

$$R_\theta = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}$$

rotates vectors in the plane counterclockwise by the angle θ. It is orthogonal, since

$$\begin{aligned} R_\theta^\top R_\theta &= \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \\ &= \begin{pmatrix} \cos^2\theta + \sin^2\theta & -\cos\theta\sin\theta + \sin\theta\cos\theta \\ -\sin\theta\cos\theta + \cos\theta\sin\theta & \sin^2\theta + \cos^2\theta \end{pmatrix} = I_2. \end{aligned}$$

Hence $R_\theta^{-1} = R_\theta^\top = R_{-\theta}$: to undo a rotation by the angle θ, we rotate by $-\theta$.

Example 1.6.73. A permutation matrix is a square matrix in which each row and each column contains exactly one 1, and all other entries are 0. For example,

$$P = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \qquad P \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_3 \\ x_1 \end{pmatrix},$$

so P permutes the entries of a vector. One checks that $P^\top P = I_3$, so P is orthogonal.

Geometrically, orthogonal matrices are exactly the matrices that preserve lengths and angles, such as rotations, reflections and permutations of the coordinates; this is proved in Theorem 2.4.7.

Exercise 1.6.74. Show that $\mathrm{O}(n)$ is not a subspace of $\mathbb{R}^{n\times n}$, but that it contains I_n and is closed under multiplication and under taking inverses. In other words, $\mathrm{O}(n)$ is a matrix group, called the orthogonal group.

1.6.5 Trace and Determinant

In this subsection, we introduce two numbers that can be computed from a square matrix: the trace and the determinant. The trace is very easy to compute, and it depends linearly on the matrix. The determinant is harder to compute, but it tells us whether a matrix is invertible, and it measures how the linear transformation T_A changes areas and volumes.

The trace

Definition 1.6.75 (Trace)**.** The trace of an $n \times n$ matrix A is the sum of its diagonal entries:

$$\operatorname{tr}(A) = A_{11} + A_{22} + \cdots + A_{nn} = \sum_{i=1}^{n} A_{ii}.$$

Example 1.6.76. For the upper triangular matrix from Subsection 1.6.4,

$$\operatorname{tr}\begin{pmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{pmatrix} = 1 + 4 + 6 = 11.$$

Moreover, $\operatorname{tr}(I_n) = 1 + \cdots + 1 = n$, and the trace of every skew-symmetric matrix is 0, since its diagonal entries are 0.

Theorem 1.6.77. *Let A and B be $n \times n$ matrices and $c \in \mathbb{R}$. Then:*

1. $\operatorname{tr}(A + B) = \operatorname{tr}(A) + \operatorname{tr}(B)$ *and* $\operatorname{tr}(cA) = c \operatorname{tr}(A)$*; in other words,* $\operatorname{tr} : \mathbb{R}^{n \times n} \to \mathbb{R}$ *is a linear transformation;*
2. $\operatorname{tr}(A^\top) = \operatorname{tr}(A)$*;*
3. $\operatorname{tr}(AB) = \operatorname{tr}(BA)$*. More generally, this holds for every $m \times n$ matrix A and every $n \times m$ matrix B.*

Proof. Parts 1 and 2 follow directly from the definitions, since the diagonal entries of $A + B$, cA and $A^\top$ are $A_{ii} + B_{ii}$, cA_{ii} and A_{ii}, respectively. For part 3, let A be an $m \times n$ matrix and B an $n \times m$ matrix. Then

$$\begin{aligned} \operatorname{tr}(AB) &= \sum_{i=1}^{m} (AB)_{ii} = \sum_{i=1}^{m} \sum_{k=1}^{n} A_{ik} B_{ki} \\ &= \sum_{k=1}^{n} \sum_{i=1}^{m} B_{ki} A_{ik} = \sum_{k=1}^{n} (BA)_{kk} = \operatorname{tr}(BA). \end{aligned}$$

□

Remark 1.6.78. Part 3 is surprising, since in general $AB \neq BA$. For example, if P is an invertible $n \times n$ matrix, then

$$\operatorname{tr}(P^{-1}AP) = \operatorname{tr}\left((P^{-1}A)P\right) = \operatorname{tr}\left(P(P^{-1}A)\right) = \operatorname{tr}(A).$$

However, the trace is not multiplicative: for

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \qquad B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix},$$

we have $AB = 0$, so $\operatorname{tr}(AB) = 0$, while $\operatorname{tr}(A)\operatorname{tr}(B) = 1 \cdot 1 = 1$.

Example 1.6.79. For any $m \times n$ matrix A, the diagonal entries of the $n \times n$ matrix $A^\top A$ are

$$(A^\top A)_{jj} = \sum_{i=1}^{m} (A^\top)_{ji} A_{ij} = \sum_{i=1}^{m} A_{ij}^2.$$

Hence

$$\operatorname{tr}(A^\top A) = \sum_{i=1}^{m} \sum_{j=1}^{n} A_{ij}^2$$

is the sum of the squares of all entries of A. In particular, $\operatorname{tr}(A^\top A) \geq 0$, with equality only for $A = 0$. The number $\|A\|_F = \sqrt{\operatorname{tr}(A^\top A)}$ is called the Frobenius norm of A. It is often used in machine learning to measure the size of a matrix, for example of the weight matrix of a layer in a neural network.

Corollary 1.6.80. *The set* $\{A \in \mathbb{R}^{n \times n} : \operatorname{tr}(A) = 0\}$ *of matrices with trace zero is a subspace of* $\mathbb{R}^{n \times n}$.

Proof. This set is the kernel of the linear transformation $\operatorname{tr} : \mathbb{R}^{n \times n} \to \mathbb{R}$, so it is a subspace by Theorem 1.5.5. □

Exercise 1.6.81.

1. Show that the subspace of $n \times n$ matrices with trace zero has dimension $n^2 - 1$. (Hint: use the matrices e_{ij} from the proof of Theorem 1.6.65.)
2. Show that there are no $n \times n$ matrices A and B with $AB - BA = I_n$. (Hint: take the trace of both sides.)

The determinant

The determinant of a square matrix is defined recursively: the determinant of an $n \times n$ matrix is expressed in terms of determinants of $(n-1) \times (n-1)$ matrices.

Definition 1.6.82 (Determinant)**.** Let A be an $n \times n$ matrix. For $1 \leq i, j \leq n$, let $A^{(ij)}$ denote the $(n-1) \times (n-1)$ matrix obtained from A by deleting its i-th row and its j-th column. The determinant $\det(A)$ of A is the number defined as follows:

- if $n = 1$, then $\det(A) = A_{11}$;
- if $n \geq 2$, then

$$\det(A) = \sum_{j=1}^{n} (-1)^{1+j} A_{1j} \det(A^{(1j)}).$$

Written out, $\det(A) = A_{11} \det(A^{(11)}) - A_{12} \det(A^{(12)}) + A_{13} \det(A^{(13)}) - \cdots$, with alternating signs. This formula is called the cofactor expansion along the first row, and the numbers $C_{ij} = (-1)^{i+j} \det(A^{(ij)})$ are called the cofactors of A. The determinant is also denoted by vertical bars, e.g., $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}$.

Example 1.6.83 (2×2 and 3×3 determinants)**.** For a 2×2 matrix, deleting the first row and the first or the second column leaves the 1×1 matrices (d) and (c), so

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc.$$

For a 3×3 matrix,

$$\det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & k \end{pmatrix} = a \det \begin{pmatrix} e & f \\ h & k \end{pmatrix} - b \det \begin{pmatrix} d & f \\ g & k \end{pmatrix} + c \det \begin{pmatrix} d & e \\ g & h \end{pmatrix}$$
$$= a(ek - fh) - b(dk - fg) + c(dh - eg).$$

For example, $\det \begin{pmatrix} 2 & 3 \\ 4 & -1 \end{pmatrix} = 2 \cdot (-1) - 3 \cdot 4 = -14$, and

$$\det \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 10 \end{pmatrix} = 1 \cdot (5 \cdot 10 - 6 \cdot 8) - 2 \cdot (4 \cdot 10 - 6 \cdot 7) + 3 \cdot (4 \cdot 8 - 5 \cdot 7) = 2 + 4 - 9 = -3.$$

The following theorem collects the basic properties of the determinant. We prove part 3 and the expansion along rows right after the theorem, directly from the definition. The expansion along columns and part 2 are proved after the product rule for determinants (Theorem 1.6.92), which is the main tool for them.

Theorem 1.6.84. *Let A be an $n \times n$ matrix.*

1. *(Expansion along any row or column.) For every row index i and every column index j,*

$$\det(A) = \sum_{l=1}^{n} (-1)^{i+l} A_{il} \det(A^{(il)}) = \sum_{l=1}^{n} (-1)^{l+j} A_{lj} \det(A^{(lj)}).$$

2. *(Transpose.)* $\det(A^\top) = \det(A)$.

3. *(Row operations.)*
   - *Swapping two rows of A multiplies the determinant by -1.*
   - *Multiplying a row of A by a scalar c multiplies the determinant by c.*
   - *Adding a multiple of one row to another row does not change the determinant.*

By part 2, the same rules hold for the corresponding operations on columns. The signs $(-1)^{i+j}$ in part 1 form a checkerboard pattern, starting with $+$ in the top left corner. All proofs below use induction on n and the definition, i.e., the expansion along the first row.

Lemma 1.6.85. *Let A be an $n \times n$ matrix with rows $\mathbf{r}_1, \dots, \mathbf{r}_n$.*

1. *(Linearity in each row.) Suppose that $\mathbf{r}_i = a\mathbf{u} + b\mathbf{w}$ for row vectors $\mathbf{u}, \mathbf{w} \in \mathbb{R}^{1 \times n}$ and scalars a, b. Let A_u and A_w be the matrices obtained from A by replacing its i-th row by $\mathbf{u}$ and by $\mathbf{w}$, respectively. Then $\det(A) = a \det(A_u) + b \det(A_w)$.*

2. *(Swapping neighboring rows.) If $n \geq 2$, swapping two neighboring rows $\mathbf{r}_i$ and $\mathbf{r}_{i+1}$ of A multiplies the determinant by -1.*

Proof. (1) For $n = 1$, $\det(a\mathbf{u} + b\mathbf{w}) = a\mathbf{u} + b\mathbf{w}$. Let $n \geq 2$. If $i = 1$, then the matrices $A^{(1j)}$ do not contain the first row, so they are the same for A, A_u and A_w, and

$$\det(A) = \sum_{j=1}^{n} (-1)^{1+j} (au_j + bw_j) \det(A^{(1j)}) = a \det(A_u) + b \det(A_w).$$

If $i \geq 2$, then the entries A_{1j} are the same for the three matrices, and row $i-1$ of $A^{(1j)}$ is a times $\mathbf{u}$ plus b times $\mathbf{w}$, each with its j-th entry deleted. By induction on n, $\det(A^{(1j)}) = a\det(A_u^{(1j)}) + b\det(A_w^{(1j)})$; multiply by $(-1)^{1+j}A_{1j}$ and add over j.

(2) For $n = 2$, swapping the rows of $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ gives the determinant $cb - da = -(ad - bc)$. Let $n \geq 3$. If $i \geq 2$, then the swap exchanges two neighboring rows of each matrix $A^{(1j)}$ and does not change the first row of A; by induction, each $\det(A^{(1j)})$ changes its sign, and hence so does $\det(A)$. It remains to swap the first two rows. Let M_{jk} be the matrix obtained from A by deleting the rows 1 and 2 and the columns j and k, so that $M_{jk} = M_{kj}$. In $A^{(1j)}$, the first row consists of the entries A_{2k} with $k \neq j$; the entry A_{2k} stands in column k if $k < j$ and in column $k-1$ if $k > j$, and deleting its row and column leaves M_{jk}. Expanding each $\det(A^{(1j)})$ along its first row therefore gives

$$\det(A) = \sum_{j=1}^{n} \sum_{k \neq j} \varepsilon_{jk} A_{1j} A_{2k} \det(M_{jk}), \qquad \varepsilon_{jk} = \begin{cases} (-1)^{1+j}(-1)^{1+k} = (-1)^{j+k} & \text{if } k < j, \\ (-1)^{1+j}(-1)^{k} = -(-1)^{j+k} & \text{if } k > j. \end{cases}$$

In particular, $\varepsilon_{kj} = -\varepsilon_{jk}$. The matrix B with the first two rows swapped has $B_{1j} = A_{2j}$ and $B_{2k} = A_{1k}$, and the same matrices M_{jk}, so, exchanging the names of the summation indices j and k,

$$\det(B) = \sum_{j=1}^{n} \sum_{k \neq j} \varepsilon_{jk} A_{2j} A_{1k} \det(M_{jk}) = \sum_{k=1}^{n} \sum_{j \neq k} \varepsilon_{kj} A_{1j} A_{2k} \det(M_{jk}) = -\det(A). \qquad \square$$

Proof of Theorem 1.6.84, part 3, and of the expansion along rows. Swapping the rows $i < k$ can be done by swaps of neighboring rows: move row i down to position k by $k - i$ swaps, and then move the old row k, which is now in position $k-1$, up to position i by $k - i - 1$ swaps. The total number $2(k-i) - 1$ is odd, so by Lemma 1.6.85, the determinant changes its sign. Multiplying a row by c multiplies the determinant by c, by Lemma 1.6.85 with $a = c$ and $b = 0$.

If a matrix has two equal rows, then swapping them does not change the matrix, but changes the sign of its determinant, so the determinant is 0. Now let A' be obtained from A by adding c times row k to row $i \neq k$. By Lemma 1.6.85, $\det(A') = \det(A) + c\det(A'')$, where A'' is obtained from A by replacing row i by row k. Since A'' has two equal rows, $\det(A'') = 0$, and $\det(A') = \det(A)$.

Finally, let $1 \leq i \leq n$. Moving row i to the top by $i-1$ swaps of neighboring rows gives the matrix B with the rows $\mathbf{r}_i, \mathbf{r}_1, \ldots, \mathbf{r}_{i-1}, \mathbf{r}_{i+1}, \ldots, \mathbf{r}_n$, and $\det(B) = (-1)^{i-1}\det(A)$. The other rows keep their order, so deleting the first row and the l-th column of B gives $A^{(il)}$. By the definition of $\det(B)$,

$$\det(A) = (-1)^{i-1} \sum_{l=1}^{n} (-1)^{1+l} A_{il} \det(A^{(il)}) = \sum_{l=1}^{n} (-1)^{i+l} A_{il} \det(A^{(il)}). \qquad \square$$

Corollary 1.6.86. *Let A be an $n \times n$ matrix.*

1. *If A has a row or a column of zeros, then $\det(A) = 0$.*
2. *If A has two equal rows, then $\det(A) = 0$.*
3. *If A is upper or lower triangular, then $\det(A) = A_{11}A_{22}\cdots A_{nn}$ is the product of its diagonal entries. In particular, $\det(I_n) = 1$.*
4. *$\det(cA) = c^n \det(A)$ for every $c \in \mathbb{R}$.*

Proof. (1) For a row of zeros, expand along this row: every term of the sum contains a factor 0. For a column of zeros, say column j, we use induction on n; for $n = 1$, $A = (0)$. In the expansion along the first row, $A_{1j} = 0$, and for $l \neq j$, the matrix $A^{(1l)}$ has a column of zeros, so $\det(A^{(1l)}) = 0$ by induction.

(2) Swapping the two equal rows does not change A, but by Theorem 1.6.84 it multiplies the determinant by -1. So $\det(A) = -\det(A)$, i.e., $\det(A) = 0$.

(3) Let A be lower triangular. The only possibly nonzero entry of its first row is A_{11}, so the definition gives $\det(A) = A_{11}\det(A^{(11)})$. The matrix $A^{(11)}$ is again lower triangular, with diagonal entries $A_{22}, \dots, A_{nn}$, so repeating this argument (formally, by induction on n) gives $\det(A) = A_{11}A_{22}\cdots A_{nn}$. If A is upper triangular, then the only possibly nonzero entry of its last row is A_{nn}, and the expansion along the last row gives $\det(A) = A_{nn}\det(A^{(nn)})$, where $A^{(nn)}$ is upper triangular with diagonal entries $A_{11}, \dots, A_{n-1,n-1}$; again, induction gives the claim.

(4) The matrix cA is obtained from A by multiplying each of its n rows by c. □

Theorem 1.6.84 and Corollary 1.6.86 give an efficient way to compute determinants: bring the matrix into upper triangular form by row operations, keeping track of how each operation changes the determinant, and then multiply the diagonal entries. For large n, this is much faster than the cofactor expansion, which involves about $n!$ terms.

Example 1.6.87. Let us compute the determinant of the 3×3 matrix above again, now with row operations. Subtracting 4 times the first row from the second row, and 7 times the first row from the third row, does not change the determinant:

$$\det\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 10 \end{pmatrix} = \det\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -11 \end{pmatrix}.$$

Subtracting 2 times the second row from the third row does not change it either, and we obtain an upper triangular matrix:

$$\det\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 1 \end{pmatrix} = 1 \cdot (-3) \cdot 1 = -3,$$

in agreement with the cofactor expansion. The matrix F of Example 1.6.31 differs from this matrix only in its bottom right entry, and the same row operations turn it into a matrix with a row of zeros (see Example 1.6.31). Hence $\det(F) = 0$.

Remark 1.6.88 (Geometric meaning). Let A be a 2×2 matrix. The linear transformation T_A maps the unit square, with sides $\mathbf{e}_1$ and $\mathbf{e}_2$, to the parallelogram with sides $A\mathbf{e}_1$ and $A\mathbf{e}_2$, which are the columns of A. The area of this parallelogram is $|\det(A)|$ (see Figure 1.4). More generally, T_A multiplies the area of every region by $|\det(A)|$, and for 3×3 matrices, T_A multiplies volumes by $|\det(A)|$. The sign of $\det(A)$ tells us whether T_A preserves the orientation (positive sign) or reverses it, like a mirror (negative sign). We do not prove these facts here.

This picture also makes the rules of Theorem 1.6.84 for column operations plausible. Adding a multiple of one column to the other shears the parallelogram, but does not change its base or its height, and hence not its area (see Figure 1.5). Multiplying a column by c stretches the parallelogram by the factor $|c|$ in one direction, which multiplies its area by $|c|$. Swapping the two columns reverses the orientation, which changes the sign of the determinant.

Example 1.6.89.

- $\det(\mathrm{diag}(2,3)) = 6$: stretching by a factor 2 in the x-direction and by a factor 3 in the y-direction multiplies areas by 6.
- $\det(R_\theta) = \cos^2\theta + \sin^2\theta = 1$: rotations preserve areas and orientation.
- $\det\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = -1$: the reflection in the x-axis preserves areas, but reverses orientation.




Figure 1.4: The matrix A with columns $(3,1)$ and $(1,2)$ maps the unit square (left, area 1) to the parallelogram spanned by its columns $A\mathbf{e}_1$ and $A\mathbf{e}_2$ (right). The area of this parallelogram is $|\det(A)| = |3 \cdot 2 - 1 \cdot 1| = 5$.




Figure 1.5: The parallelograms spanned by $\mathbf{u} = (2,0)$ and $\mathbf{v} = (\frac{1}{2}, \frac{3}{2})$ (blue) and by $\mathbf{u}$ and $\mathbf{v} + \mathbf{u}$ (orange) have the same base $\mathbf{u}$ and the same height (dashed line), hence the same area 3. Correspondingly, adding the first column to the second column does not change the determinant: both matrices with these columns have determinant $2 \cdot \frac{3}{2} = 3$.

- $\det \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} = 0$: this singular matrix maps the whole plane onto the line spanned by $(1,2)$, which has area 0 (see Figure 1.3).

Compare also with the gallery in Figure 1.2. The stretch (a) has determinant $\det \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} = 2$ and doubles areas. The rotation (b) and the shear (d) have determinants $\det \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = 1$ and $\det \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = 1$: they preserve areas, even though the shear changes the shape of the square. The reflection (c) has determinant -1, which is why the image of the letter F is mirrored.

Determinants, invertibility and products

We now connect determinants with the inverse of a matrix, using the elementary matrices of Subsection 1.6.3.

Lemma 1.6.90. *Let E be an $n \times n$ elementary matrix. Then $\det(E) = -1$ if E swaps two rows, $\det(E) = c$ if E multiplies a row by $c \neq 0$, and $\det(E) = 1$ if E adds a multiple of one row to another row. In all cases, $\det(E) \neq 0$, and*

$$\det(EA) = \det(E)\det(A) \qquad \text{for every } n \times n \text{ matrix } A.$$

Proof. The matrix E is obtained from I_n by the row operation, and $\det(I_n) = 1$, so the values of $\det(E)$ follow from part 3 of Theorem 1.6.84. By Lemma 1.6.45, EA is obtained from A by the same row operation, so $\det(EA)$ is $\det(A)$ multiplied by -1, by c, or by 1, respectively. In each case, this is $\det(E)\det(A)$. □

Theorem 1.6.91. *An $n \times n$ matrix A is invertible if and only if $\det(A) \neq 0$.*

Proof. Let $E_1, \dots, E_k$ be the elementary matrices of row operations that bring A to $R = \mathrm{RREF}(A)$, so that $R = E_k \cdots E_1 A$. Applying Lemma 1.6.90 k times gives

$$\det(R) = \det(E_k) \cdots \det(E_1) \det(A),$$

where all the factors $\det(E_i)$ are nonzero. So $\det(A) \neq 0$ if and only if $\det(R) \neq 0$. If A is invertible, then $R = I_n$ by Theorem 1.6.47, and $\det(R) = 1 \neq 0$. If A is not invertible, then $R \neq I_n$, so R has a row of zeros (as we saw after Theorem 1.6.49), and $\det(R) = 0$ by Corollary 1.6.86. □

Theorem 1.6.92 (Product rule)**.** *For all $n \times n$ matrices A and B, we have* $\det(AB) = \det(A)\det(B)$.

Proof. Suppose first that A is not invertible, so that $\det(A) = 0$ by Theorem 1.6.91. Then AB is not invertible either: otherwise, $X = B(AB)^{-1}$ would satisfy $AX = I_n$, and by Corollary 1.6.48, X would be invertible with $X^{-1} = A$, so A would be invertible. Hence $\det(AB) = 0 = \det(A)\det(B)$.

Now suppose that A is invertible. By the proof of Theorem 1.6.47, $A = (E_k \cdots E_1)^{-1} = E_1^{-1} \cdots E_k^{-1}$ for some elementary matrices $E_1, \dots, E_k$, and each $F_i = E_i^{-1}$ is again an elementary matrix by Lemma 1.6.46. Applying Lemma 1.6.90 k times, we get

$$\det(AB) = \det(F_1 F_2 \cdots F_k B) = \det(F_1)\det(F_2) \cdots \det(F_k)\det(B).$$

Taking $B = I_n$ gives $\det(A) = \det(F_1) \cdots \det(F_k)$, and hence $\det(AB) = \det(A)\det(B)$. □

Proof of Theorem 1.6.84, parts 1 and 2. We already proved part 3 and the expansion along rows, and the proofs of Corollary 1.6.86, Lemma 1.6.90 and Theorems 1.6.91 and 1.6.92 used only these. For part 2, note that A is invertible if and only if $A^\top$ is (Proposition 1.6.42, applied to A and to $A^\top$). If A is not invertible, then $\det(A^\top) = 0 = \det(A)$ by Theorem 1.6.91. If A is invertible, then, as in the proof of Theorem 1.6.92, $A = F_1 \cdots F_k$ with elementary matrices F_i, so $A^\top = F_k^\top \cdots F_1^\top$. The transpose of an elementary matrix is an elementary matrix with the same determinant: the matrices that swap two rows or multiply a row by c are symmetric, and the transpose of the matrix that adds c times row j to row i is the matrix that adds c times row i to row j, with determinant 1. By the product rule,

$$\det(A^\top) = \det(F_k^\top) \cdots \det(F_1^\top) = \det(F_1) \cdots \det(F_k) = \det(A).$$

For the expansion along column j, expand $\det(A^\top)$ along its row j. Since $(A^\top)_{jl} = A_{lj}$, and $(A^\top)^{(jl)} = (A^{(lj)})^\top$ has the same determinant as $A^{(lj)}$ by part 2 (for $(n-1) \times (n-1)$ matrices),

$$\det(A) = \det(A^\top) = \sum_{l=1}^{n} (-1)^{j+l} A_{lj} \det(A^{(lj)}).$$ □

Corollary 1.6.93. *Let A be an $n \times n$ matrix.*

1. *If A is invertible, then* $\det(A^{-1}) = 1/\det(A)$.
2. *If P is an invertible $n \times n$ matrix, then* $\det(P^{-1}AP) = \det(A)$.
3. *If A is orthogonal, then* $\det(A) = 1$ *or* $\det(A) = -1$.

Proof. (1) $\det(A)\det(A^{-1}) = \det(AA^{-1}) = \det(I_n) = 1$.
(2) $\det(P^{-1}AP) = \det(P^{-1})\det(A)\det(P) = \det(A)$, by part 1.
(3) $1 = \det(I_n) = \det(A^\top A) = \det(A^\top)\det(A) = \det(A)^2$. □

Remark 1.6.94. Unlike the trace, the determinant is not linear. For example, $\det(I_2 + I_2) = \det(2I_2) = 4$, while $\det(I_2) + \det(I_2) = 2$.

Example 1.6.95 (Inverse of a 2×2 matrix)**.** Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $\det(A) = ad - bc \neq 0$. Then

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Indeed,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} ad - bc & 0 \\ 0 & ad - bc \end{pmatrix} = (ad - bc) I_2,$$

and by Corollary 1.6.48, checking one product is enough. For example, for the matrix of Example 1.6.41, with determinant -14,

$$\begin{pmatrix} 2 & 3 \\ 4 & -1 \end{pmatrix}^{-1} = \frac{1}{-14} \begin{pmatrix} -1 & -3 \\ -4 & 2 \end{pmatrix} = \frac{1}{14} \begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix},$$

as we found before.

The matrix $\begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ is the transpose of the matrix of the cofactors $C_{11} = d$, $C_{12} = -c$, $C_{21} = -b$ and $C_{22} = a$. The same holds for all n.

Proposition 1.6.96 (Cofactor formula for the inverse)**.** *Let A be an $n \times n$ matrix with $n \geq 2$, and let C be the matrix of its cofactors $C_{ij} = (-1)^{i+j} \det(A^{(ij)})$. Then $AC^\top = \det(A)\, I_n$. In particular, if $\det(A) \neq 0$, then*

$$A^{-1} = \frac{1}{\det(A)}\, C^\top.$$

Proof. The (i, k) entry of $AC^\top$ is $\sum_{l=1}^n A_{il} C_{kl} = \sum_{l=1}^n (-1)^{k+l} A_{il} \det(A^{(kl)})$. For $i = k$, this is the expansion of $\det(A)$ along row i (Theorem 1.6.84). For $i \neq k$, let A' be obtained from A by replacing row k by row i. Deleting row k gives the same matrices for A and A', so $A'^{(kl)} = A^{(kl)}$, and $A'_{kl} = A_{il}$. Hence the sum is the expansion of $\det(A')$ along row k. Since A' has two equal rows, $\det(A') = 0$ by Corollary 1.6.86. So $AC^\top = \det(A)\, I_n$. If $\det(A) \neq 0$, then $A \cdot \frac{1}{\det(A)} C^\top = I_n$, and by Corollary 1.6.48, $\frac{1}{\det(A)} C^\top$ is the inverse of A. □

The cofactor formula is useful in theory; for example, it shows that the entries of A^{-1} are rational functions of the entries of A. For computations, the Gauss–Jordan method is much more efficient.

Theorem 1.6.91 says that $\mathrm{GL}(n, \mathbb{R}) = \{A \in \mathbb{R}^{n \times n} : \det(A) \neq 0\}$. Determinants also give us another matrix group.

Definition 1.6.97 (Special linear group)**.** The special linear group is the set

$$\mathrm{SL}(n, \mathbb{R}) = \{A \in \mathbb{R}^{n \times n} : \det(A) = 1\}.$$

Proposition 1.6.98. *The set $\mathrm{SL}(n, \mathbb{R})$ is contained in $\mathrm{GL}(n, \mathbb{R})$ and is not a subspace of $\mathbb{R}^{n \times n}$. However, it contains I_n and is closed under multiplication and under taking inverses, so it is a matrix group.*

Proof. Every matrix with determinant 1 is invertible by Theorem 1.6.91. The zero matrix has determinant 0, so it is not in $\mathrm{SL}(n, \mathbb{R})$, and $\mathrm{SL}(n, \mathbb{R})$ is not a subspace. We have $\det(I_n) = 1$. If $\det(A) = \det(B) = 1$, then $\det(AB) = 1 \cdot 1 = 1$ by Theorem 1.6.92, and $\det(A^{-1}) = 1/1 = 1$ by Corollary 1.6.93. □

Example 1.6.99. All rotation matrices R_θ belong to $\mathrm{SL}(2, \mathbb{R})$, and so does $\mathrm{diag}(2, \frac{1}{2})$, which stretches the plane by a factor 2 in one direction and shrinks it by a factor 2 in the other. By the geometric meaning of the determinant, the matrices in $\mathrm{SL}(2, \mathbb{R})$ are exactly the 2×2 matrices whose linear transformations preserve areas and orientation.

1.6.6 Block Matrices, the LU Decomposition and the Woodbury Identity

Large matrices often have a natural structure: the features of a data set fall into groups, a covariance matrix describes several random vectors at once, and a matrix changes by a small update. In this subsection, we compute with matrices in blocks, and we use blocks to describe Gaussian elimination as a factorization of a matrix and to invert low-rank updates.

Block matrices and the Schur complement

A matrix can be split into blocks by horizontal and vertical lines, for example

$$M = \left(\begin{array}{cc|c} 1 & 2 & 5 \\ 3 & 4 & 6 \\ \hline 7 & 8 & 9 \end{array}\right) = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \qquad A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}, \quad B = \begin{pmatrix} 5 \\ 6 \end{pmatrix}, \quad C = \begin{pmatrix} 7 & 8 \end{pmatrix}, \quad D = (9).$$

Block matrices are multiplied like 2×2 matrices, as long as the sizes of the blocks fit.

Proposition 1.6.100 (Block multiplication)**.** *Let* $M = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$ *and* $N = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$*, where* A_{ik} *is an* $m_i \times n_k$ *matrix and* B_{kj} *is an* $n_k \times p_j$ *matrix. Then*

$$MN = \begin{pmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{pmatrix}.$$

Proof. Each entry of MN is a sum $\sum_l M_{rl}N_{ls}$ over $l = 1, \ldots, n_1 + n_2$. The terms with $l \le n_1$ form the corresponding entry of $A_{i1}B_{1j}$, and the terms with $l > n_1$ form the entry of $A_{i2}B_{2j}$, where i and j are the block row of r and the block column of s. □

Lemma 1.6.101. *Let* A *be a* $k \times k$ *matrix and* D *an* $m \times m$ *matrix. Then*

$$\det \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} = \det(A)\det(D) = \det \begin{pmatrix} A & 0 \\ C & D \end{pmatrix}$$

for all matrices B *and* C *of the appropriate sizes.*

Proof. Let $M = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$. If A is not invertible, then $A\mathbf{x} = \mathbf{0}$ for some $\mathbf{x} \neq \mathbf{0}$, and M maps the nonzero vector $\begin{pmatrix} \mathbf{x} \\ \mathbf{0} \end{pmatrix}$ to $\begin{pmatrix} A\mathbf{x} \\ \mathbf{0} \end{pmatrix} = \mathbf{0}$. So M is not invertible either, and $\det(M) = 0 = \det(A)\det(D)$. If A is invertible, then by Proposition 1.6.100,

$$M = \begin{pmatrix} A & 0 \\ 0 & I_m \end{pmatrix} \begin{pmatrix} I_k & A^{-1}B \\ 0 & D \end{pmatrix}.$$

Expanding the determinant of the first factor m times along the last row (each time, the only nonzero entry of the last row is a diagonal entry 1) gives $\det(A)$, and expanding the determinant of the second factor k times along the first column gives $\det(D)$. The product rule gives $\det(M) = \det(A)\det(D)$. The second equation follows by taking transposes. □

Definition 1.6.102 (Schur complement)**.** Let $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ be a square matrix with square blocks A and D, where A is invertible. The Schur complement of A in M is the matrix $S = D - CA^{-1}B$.

Theorem 1.6.103. *In the situation of Definition 1.6.102,*

$$M = \begin{pmatrix} I & 0 \\ CA^{-1} & I \end{pmatrix} \begin{pmatrix} A & B \\ 0 & S \end{pmatrix}, \qquad \textit{so} \qquad \det(M) = \det(A)\det(S).$$

Hence M is invertible if and only if S is invertible, and in this case

$$M^{-1} = \begin{pmatrix} A^{-1} + A^{-1}BS^{-1}CA^{-1} & -A^{-1}BS^{-1} \\ -S^{-1}CA^{-1} & S^{-1} \end{pmatrix}.$$

Proof. By Proposition 1.6.100, the product of the two factors is $\begin{pmatrix} A & B \\ CA^{-1}A & CA^{-1}B + S \end{pmatrix} = M$. By Lemma 1.6.101, the first factor has determinant 1 and the second one $\det(A)\det(S)$. Since $\det(A) \neq 0$, M is invertible if and only if $\det(S) \neq 0$. If S is invertible, block multiplication shows that the inverses of the two factors are

$$\begin{pmatrix} I & 0 \\ -CA^{-1} & I \end{pmatrix} \qquad \text{and} \qquad \begin{pmatrix} A^{-1} & -A^{-1}BS^{-1} \\ 0 & S^{-1} \end{pmatrix},$$

and multiplying them in the opposite order, $M^{-1} = \begin{pmatrix} A^{-1} & -A^{-1}BS^{-1} \\ 0 & S^{-1} \end{pmatrix} \begin{pmatrix} I & 0 \\ -CA^{-1} & I \end{pmatrix}$, gives the formula. □

The first factor in Theorem 1.6.103 is a block version of an elementary matrix: multiplying M by its inverse from the left subtracts CA^{-1} times the first block row from the second one, which eliminates C, just as row reduction eliminates the entries below a pivot. The Schur complement is what remains in the lower right corner.

Example 1.6.104. For the matrix M above, $A^{-1} = \frac{1}{-2}\begin{pmatrix} 4 & -2 \\ -3 & 1 \end{pmatrix}$, so $A^{-1}B = \frac{1}{-2}\begin{pmatrix} 8 \\ -9 \end{pmatrix} = \begin{pmatrix} -4 \\ 4.5 \end{pmatrix}$, and the Schur complement is $S = 9 - (7,8)\begin{pmatrix} -4 \\ 4.5 \end{pmatrix} = 9 - 8 = 1$. Hence $\det(M) = \det(A)\cdot 1 = -2$, and the lower right entry of M^{-1} is $S^{-1} = 1$.

In Lecture 4, the Schur complement appears as the covariance matrix of a Gaussian random vector after we condition on some of its entries.

The LU decomposition

Gaussian elimination brings a matrix into upper triangular form by subtracting multiples of rows from the rows below them. We now show that this writes the matrix as a product of a lower and an upper triangular matrix. A lower triangular matrix is called unit lower triangular if all its diagonal entries are 1, and a permutation matrix is a matrix obtained from I_n by reordering its rows; then PA is A with its rows reordered in the same way.

Lemma 1.6.105. *Products of (unit) lower triangular $n \times n$ matrices are (unit) lower triangular, and so are the inverses of invertible ones. The same holds for upper triangular matrices.*

Proof. Let L and L' be lower triangular. In $(LL')_{ij} = \sum_k L_{ik}L'_{kj}$, only the terms with $j \leq k \leq i$ can be nonzero. So $(LL')_{ij} = 0$ for $i < j$, and $(LL')_{ii} = L_{ii}L'_{ii}$, which is 1 if L and L' are unit lower triangular. If L is invertible, the j-th column $\mathbf{x}$ of L^{-1} solves $L\mathbf{x} = \mathbf{e}_j$. The first $j-1$ equations $L_{11}x_1 = 0$, $L_{21}x_1 + L_{22}x_2 = 0$, …, only involve $x_1, \ldots, x_{j-1}$, and since the diagonal entries L_{ii} are nonzero ($\det(L) = L_{11}\cdots L_{nn} \neq 0$), they give $x_1 = \cdots = x_{j-1} = 0$; the j-th equation then gives $x_j = \frac{1}{L_{jj}}$. So L^{-1} is lower triangular, with diagonal entries $\frac{1}{L_{jj}}$. Upper triangular matrices are the transposes of lower triangular ones. □

Theorem 1.6.106 (LU decomposition)**.** *Let A be an $n \times n$ matrix.*

1. *If the upper left $k \times k$ submatrices of A are invertible for $k = 1, \dots, n-1$, then $A = LU$ with a unit lower triangular matrix L and an upper triangular matrix U. If A is invertible, then L and U are unique.*
2. *If A is invertible, then $PA = LU$ for a permutation matrix P, a unit lower triangular matrix L and an upper triangular matrix U.*

Proof. (1) We use induction on n; for $n = 1$, $A = (1)(A_{11})$. Let $n \geq 2$, and write

$$A = \begin{pmatrix} a & \mathbf{r}^\top \\ \mathbf{c} & A' \end{pmatrix}$$

with a number $a = A_{11}$, which is nonzero by assumption, vectors $\mathbf{c}, \mathbf{r} \in \mathbb{R}^{n-1}$ and an $(n-1)\times(n-1)$ matrix A'. By Theorem 1.6.103 with the 1×1 block A_{11},

$$A = \begin{pmatrix} 1 & 0 \\ \frac{1}{a}\mathbf{c} & I \end{pmatrix} \begin{pmatrix} a & \mathbf{r}^\top \\ 0 & S \end{pmatrix}, \qquad S = A' - \frac{1}{a}\mathbf{c}\mathbf{r}^\top.$$

The upper left $(k+1) \times (k+1)$ submatrix of A has the same form, with $\mathbf{c}$, $\mathbf{r}$ and A' cut down to their first k entries or $k \times k$ submatrix, and its Schur complement is the upper left $k \times k$ submatrix S_k of S. So $a \det(S_k)$ is the determinant of an upper left submatrix of A, which is nonzero for $k \leq n-2$. By induction, $S = L'U'$ with L' unit lower triangular and U' upper triangular, and

$$A = \begin{pmatrix} 1 & 0 \\ \frac{1}{a}\mathbf{c} & L' \end{pmatrix} \begin{pmatrix} a & \mathbf{r}^\top \\ 0 & U' \end{pmatrix},$$

since the product has the lower right block $\frac{1}{a}\mathbf{c}\mathbf{r}^\top + L'U' = \frac{1}{a}\mathbf{c}\mathbf{r}^\top + S = A'$. For uniqueness, let $L_1U_1 = L_2U_2$ with A invertible. Then U_1 and U_2 are invertible (as $\det(U_i) = \det(A) \neq 0$), and $L_2^{-1}L_1 = U_2U_1^{-1}$. By Lemma 1.6.105, the left side is unit lower triangular and the right side is upper triangular, so both are I_n, and $L_1 = L_2$, $U_1 = U_2$.

(2) Again we use induction on n. Since A is invertible, its first column is not zero. Let P_1 be the permutation matrix that swaps the first row with a row whose first entry is nonzero (or $P_1 = I_n$). Then $P_1A = \begin{pmatrix} a & \mathbf{r}^\top \\ \mathbf{c} & A' \end{pmatrix}$ with $a \neq 0$, and its Schur complement $S = A' - \frac{1}{a}\mathbf{c}\mathbf{r}^\top$ is invertible, since $\pm\det(A) = \det(P_1A) = a\det(S)$. By induction, $P'S = L'U'$, and with the permutation matrix $P = \begin{pmatrix} 1 & 0 \\ 0 & P' \end{pmatrix} P_1$,

$$PA = \begin{pmatrix} a & \mathbf{r}^\top \\ P'\mathbf{c} & P'A' \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \frac{1}{a}P'\mathbf{c} & L' \end{pmatrix} \begin{pmatrix} a & \mathbf{r}^\top \\ 0 & U' \end{pmatrix},$$

since $\frac{1}{a}P'cr^\top + L'U' = P'(\frac{1}{a}cr^\top + S) = P'A'$. □

The proof is Gaussian elimination: the first column of L below the diagonal contains the multipliers $\frac{c_i}{a}$ of the first row that are subtracted from the other rows, the Schur complement S is the rest of the matrix after this step, and the elimination continues with S. So L stores the multipliers, and U is the upper triangular matrix at the end of the elimination. Row swaps are needed when a pivot is 0; in practice, one swaps the row with the largest pivot candidate into place (partial pivoting), which keeps the multipliers at most 1 in absolute value and limits rounding errors.

Example 1.6.107. Let $A = \begin{pmatrix} 2 & 1 & 1 \\ 4 & 3 & 3 \\ 8 & 7 & 9 \end{pmatrix}$. Subtracting 2 times and 4 times the first row from the second and third rows gives the rows $(0,1,1)$ and $(0,3,5)$, and subtracting 3 times the new second row from the third row gives $(0,0,2)$. Hence

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 4 & 3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix} = LU,$$

and $\det(A) = \det(L)\det(U) = 2 \cdot 1 \cdot 2 = 4$. To solve $A\mathbf{x} = \mathbf{b}$ for $\mathbf{b} = (5,13,31)^\top$, we solve $Ly = \mathbf{b}$ from the top (forward substitution): $y_1 = 5$, $y_2 = 13 - 2\cdot 5 = 3$, $y_3 = 31 - 4\cdot 5 - 3\cdot 3 = 2$; and then $U\mathbf{x} = y$ from the bottom (back substitution): $x_3 = 1$, $x_2 = 3 - 1 = 2$, $x_1 = \frac{1}{2}(5 - 2 - 1) = 1$.

Computing the LU decomposition of an $n \times n$ matrix takes about $\frac{2}{3}n^3$ arithmetic operations (the Schur complement of an $m \times m$ matrix costs about $2m^2$ operations, and $\sum_{m \le n} 2m^2 \approx \frac{2}{3}n^3$), while forward and back substitution take about n^2 operations each. So once $A = LU$ is known, every further system $A\mathbf{x} = \mathbf{b}$ is cheap; this is how `np.linalg.solve` works, and there is no need to compute A^{-1}. The determinant is a by-product: $\det(A) = \pm u_{11} \cdots u_{nn}$, with the sign $\det(P)^{-1} = \pm 1$ of the row permutation.

Low-rank updates and the Woodbury identity

Suppose that we know A^{-1}, and A changes by a matrix of low rank, for example by $\mathbf{u}\mathbf{v}^\top$ when a new data point arrives. Then we do not need to invert the new matrix from scratch.

Theorem 1.6.108 (Woodbury identity)**.** *Let A be an invertible $n \times n$ matrix, C an invertible $k \times k$ matrix, U an $n \times k$ matrix and V a $k \times n$ matrix, such that $C^{-1} + VA^{-1}U$ is invertible. Then $A + UCV$ is invertible, and*

$$(A + UCV)^{-1} = A^{-1} - A^{-1}U\big(C^{-1} + VA^{-1}U\big)^{-1}VA^{-1}.$$

In particular, for $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $1 + \mathbf{v}^\top A^{-1}\mathbf{u} \neq 0$ (the case $k = 1$, $C = 1$), we have the Sherman–Morrison formula

$$(A + \mathbf{u}\mathbf{v}^\top)^{-1} = A^{-1} - \frac{A^{-1}\mathbf{u}\mathbf{v}^\top A^{-1}}{1 + \mathbf{v}^\top A^{-1}\mathbf{u}}.$$

Proof. Let X be the right side, and write $K = C^{-1} + VA^{-1}U$. Then

$$\begin{aligned}(A + UCV)X &= I + UCVA^{-1} - UK^{-1}VA^{-1} - UCVA^{-1}UK^{-1}VA^{-1} \\ &= I + UCVA^{-1} - UC\big(C^{-1} + VA^{-1}U\big)K^{-1}VA^{-1} = I + UCVA^{-1} - UCVA^{-1} = I,\end{aligned}$$

and by Corollary 1.6.48, X is the inverse of $A + UCV$. □

Proposition 1.6.109 (Determinants of low-rank updates)**.** *For an $n \times k$ matrix U and a $k \times n$ matrix V, $\det(I_n + UV) = \det(I_k + VU)$. In particular, for an invertible matrix A and $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$,*

$$\det(A + \mathbf{u}\mathbf{v}^\top) = \big(1 + \mathbf{v}^\top A^{-1}\mathbf{u}\big)\det(A).$$

Proof. Let $M = \begin{pmatrix} I_n & -U \\ V & I_k \end{pmatrix}$. By Theorem 1.6.103, with the Schur complement $I_k - VI_n^{-1}(-U) = I_k + VU$ of I_n, $\det(M) = \det(I_k + VU)$. On the other hand, block multiplication gives

$$\begin{pmatrix} I_n & U \\ 0 & I_k \end{pmatrix} M = \begin{pmatrix} I_n + UV & 0 \\ V & I_k \end{pmatrix},$$

and by Lemma 1.6.101 and the product rule, $\det(M) = \det(I_n + UV)$. For the second statement, $\det(A + \mathbf{u}\mathbf{v}^\top) = \det(A)\det(I_n + A^{-1}\mathbf{u}\mathbf{v}^\top) = \det(A)(1 + \mathbf{v}^\top A^{-1}\mathbf{u})$, by the first statement with $U = A^{-1}\mathbf{u}$ and $V = \mathbf{v}^\top$. □

Example 1.6.110. Let $A = \operatorname{diag}(2, 4)$ and $\mathbf{u} = \mathbf{v} = (1, 1)^\top$, so that $A + \mathbf{u}\mathbf{v}^\top = \begin{pmatrix} 3 & 1 \\ 1 & 5 \end{pmatrix}$. Here $A^{-1}\mathbf{u} = (\frac{1}{2}, \frac{1}{4})^\top$ and $1 + \mathbf{v}^\top A^{-1}\mathbf{u} = \frac{7}{4}$, so $\det(A + \mathbf{u}\mathbf{v}^\top) = \frac{7}{4} \cdot 8 = 14$, and

$$(A + \mathbf{u}\mathbf{v}^\top)^{-1} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{4} \end{pmatrix} - \frac{4}{7}\begin{pmatrix} \frac{1}{4} & \frac{1}{8} \\ \frac{1}{8} & \frac{1}{16} \end{pmatrix} = \frac{1}{14}\begin{pmatrix} 5 & -1 \\ -1 & 3 \end{pmatrix},$$

in agreement with the formula for the inverse of a 2×2 matrix.

Remark 1.6.111 (Low-rank updates in machine learning).

1. Linear regression with the data matrix X, whose rows are the data points $x_i^\top$, uses the matrix $X^\top X = \sum_i x_i x_i^\top$ (Lecture 2). A new data point $\mathbf{x}$ adds the rank-one matrix $\mathbf{x}\mathbf{x}^\top$, and the Sherman–Morrison formula updates $(X^\top X)^{-1}$ with about n^2 operations instead of n^3. This is recursive least squares, which also underlies the Kalman filter and some bandit algorithms.

2. For an $N \times d$ matrix X and $\lambda > 0$, the push-through identity

$$(X^\top X + \lambda I_d)^{-1} X^\top = X^\top (X X^\top + \lambda I_N)^{-1}$$

follows from $X^\top(XX^\top + \lambda I_N) = (X^\top X + \lambda I_d)X^\top$ (both matrices are invertible, since $\mathbf{x}^\top(X^\top X + \lambda I_d)\mathbf{x} = (X\mathbf{x})^\top(X\mathbf{x}) + \lambda\mathbf{x}^\top\mathbf{x} > 0$ for $\mathbf{x} \neq \mathbf{0}$, and similarly for $XX^\top + \lambda I_N$). In ridge regression with many more features than data points ($d \gg N$), the right side only needs the inverse of an $N \times N$ matrix. The matrix $XX^\top$ contains the inner products of the data points; this is the starting point of kernel methods and Gaussian processes (Lecture 4).

3. Covariance matrices of the form $D + WW^\top$, with a diagonal matrix D and an $n \times k$ matrix W with $k \ll n$ (as in factor analysis), can be inverted with the Woodbury identity and their determinants computed with Proposition 1.6.109, using only $k \times k$ matrices.

1.7 Null Space, Column Space and Rank

Row reduction is our main tool for computing the kernel and the image of a linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$. To use it, we first introduce the corresponding notions for matrices.

Definition 1.7.1. Let A be an $m \times n$ matrix.

The null space of A is the set of all vectors $\mathbf{x} \in \mathbb{R}^n$ such that $A\mathbf{x} = \mathbf{0}$. It is denoted as $\operatorname{null}(A)$. If $T(\mathbf{x}) = A\mathbf{x}$, then $\operatorname{null}(A) = \ker(T)$.

The column space of A is the set of all linear combinations of the columns of A. It is denoted as $\operatorname{col}(A)$. By Remark 1.6.6, the vectors $A\mathbf{x}$ are exactly the linear combinations of the columns of A, so if $T(\mathbf{x}) = A\mathbf{x}$, then $\operatorname{col}(A) = \operatorname{im}(T)$.

The row space of A is the set of all linear combinations of the rows of A. It is denoted as $\operatorname{row}(A)$. Writing the rows as columns, $\operatorname{row}(A) = \operatorname{col}(A^\top)$, which is the image of the linear transformation represented by the transpose $A^\top$.

By Theorem 1.5.5, $\operatorname{null}(A)$ is a subspace of $\mathbb{R}^n$ and $\operatorname{col}(A)$ is a subspace of $\mathbb{R}^m$. The following theorem explains how row reduction affects these spaces.

Theorem 1.7.2. *Let A be an $m \times n$ matrix, and let R be obtained from A by a sequence of elementary row operations. Then:*

1. $\operatorname{null}(A) = \operatorname{null}(R)$*, i.e., the equations $A\mathbf{x} = \mathbf{0}$ and $R\mathbf{x} = \mathbf{0}$ have exactly the same solutions;*

2. $\mathrm{row}(A) = \mathrm{row}(R)$;

3. *if* $R = \mathrm{RREF}(A)$, *then the columns of* A *in the positions of the pivot columns of* R *(the pivot columns of* A*) form a basis of* $\mathrm{col}(A)$.

Proof. It is enough to consider a single elementary row operation, since we can then repeat the argument for each operation in the sequence.

(1) Each equation of the new system is a linear combination of the equations of the old system, so every solution of the old system is also a solution of the new one. Since the operation can be undone by another elementary row operation, the converse holds as well.

(2) Similarly, every row of the new matrix is a linear combination of the rows of the old matrix, so the new row space is contained in the old one. Undoing the operation shows the reverse inclusion.

(3) Let $\mathbf{a}_1, \ldots, \mathbf{a}_n$ be the columns of A and $\mathbf{r}_1, \ldots, \mathbf{r}_n$ the columns of R. For any scalars $c_1, \ldots, c_n$, part (1) applied to $c = (c_1, \ldots, c_n)^\top$ gives

$$c_1\mathbf{a}_1 + \cdots + c_n\mathbf{a}_n = \mathbf{0} \iff A\mathbf{c} = \mathbf{0} \iff R\mathbf{c} = \mathbf{0} \iff c_1\mathbf{r}_1 + \cdots + c_n\mathbf{r}_n = \mathbf{0}.$$

So the columns of A satisfy exactly the same linear relations as the columns of R. If R has r pivots, then its pivot columns are the standard basis vectors $\mathbf{e}_1, \ldots, \mathbf{e}_r$ of $\mathbb{R}^m$, which are linearly independent. Every other column of R has zeros in its last $m - r$ entries, so it is a linear combination of $\mathbf{e}_1, \ldots, \mathbf{e}_r$. Transferring these relations back to A, the pivot columns of A are linearly independent, and every other column of A is a linear combination of them. Hence the pivot columns of A span $\mathrm{col}(A)$, and they form a basis of $\mathrm{col}(A)$. □

Remark 1.7.3. Elementary row operations can change the column space! For example,

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \xrightarrow{\text{subtract row 1 from row 2}} \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix},$$

but the column space of the first matrix is $\mathrm{span}\{(1,1)^\top\}$, while the column space of the second matrix is $\mathrm{span}\{(1,0)^\top\}$. This is why, in part 3 of Theorem 1.7.2, we take the pivot columns of the original matrix A, not of R.

Example 1.7.4. Consider again the linear transformation $T : \mathbb{R}^3 \to \mathbb{R}^2$ defined by $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$. The matrix representation of T is

$$A = \begin{pmatrix} 1 & 2 & -1 \\ 3 & -1 & 4 \end{pmatrix}.$$

We will transform this matrix into reduced row echelon form to find the null space, column space and row space of A, and hence the kernel and image of T.

First, we can eliminate the leading entry in the second row by subtracting 3 times the first row from the second row:

$$\begin{pmatrix} 1 & 2 & -1 \\ 0 & -7 & 7 \end{pmatrix}.$$

Next, we can multiply the second row by $-1/7$ to make the leading entry of the second row equal to 1:

$$\begin{pmatrix} 1 & 2 & -1 \\ 0 & 1 & -1 \end{pmatrix}.$$

Now we can eliminate the entry above the leading 1 in the second row by subtracting 2 times the second row from the first row:

$$\mathrm{RREF}(A) = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix}.$$

Null space. By Theorem 1.7.2, the equation $A\mathbf{x} = \mathbf{0}$, where $\mathbf{x} = (x_1, x_2, x_3)^\top$, has the same solutions as $\mathrm{RREF}(A)\,\mathbf{x} = \mathbf{0}$. From the RREF, we have:

$$x_1 + x_3 = 0 \implies x_1 = -x_3,$$
$$x_2 - x_3 = 0 \implies x_2 = x_3.$$

Letting $x_3 = t$ for some scalar t, we have:

$$\mathbf{x} = (-t, t, t)^\top = t(-1, 1, 1)^\top.$$

Therefore, $\ker(T) = \mathrm{null}(A) = \mathrm{span}\{(-1, 1, 1)^\top\}$. This agrees with Example 1.4.9, where we found $\ker(T) = \{(x, -x, -x) : x \in \mathbb{R}\}$ (take $x = -t$).

Column space. The pivots of $\mathrm{RREF}(A)$ are in columns 1 and 2. By Theorem 1.7.2, the first two columns of the original matrix A,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix}, \quad \begin{pmatrix} 2 \\ -1 \end{pmatrix},$$

form a basis of $\mathrm{col}(A)$. These two vectors span all of $\mathbb{R}^2$, since every $(u, v)^\top \in \mathbb{R}^2$ can be written as

$$\begin{pmatrix} u \\ v \end{pmatrix} = \frac{u + 2v}{7} \begin{pmatrix} 1 \\ 3 \end{pmatrix} + \frac{3u - v}{7} \begin{pmatrix} 2 \\ -1 \end{pmatrix}.$$

Hence $\mathrm{im}(T) = \mathrm{col}(A) = \mathbb{R}^2$, which agrees with Example 1.4.12.

Row space. By Theorem 1.7.2, $\mathrm{row}(A) = \mathrm{row}(\mathrm{RREF}(A))$, so the row space of A is spanned by the nonzero rows of the RREF:

$$\begin{pmatrix} 1 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 & -1 \end{pmatrix}.$$

These two rows are linearly independent (look at their first two entries), so they form a basis of $\mathrm{row}(A)$.

Definition 1.7.5 (Rank)**.** The rank of a matrix A is the dimension of its column space, $\mathrm{rank}(A) = \dim \mathrm{col}(A)$. If $T(\mathbf{x}) = A\mathbf{x}$, this is the dimension of the image of T.

Theorem 1.7.6. *Let A be a matrix, and let r be the number of pivots of* $\mathrm{RREF}(A)$*, which is the same as the number of nonzero rows of* $\mathrm{RREF}(A)$*. Then*

$$\mathrm{rank}(A) = r = \dim \mathrm{row}(A).$$

Proof. By part 3 of Theorem 1.7.2, the r pivot columns of A form a basis of $\mathrm{col}(A)$, so $\mathrm{rank}(A) = r$.

By part 2 of Theorem 1.7.2, the r nonzero rows of $\mathrm{RREF}(A)$ span $\mathrm{row}(A)$. They are also linearly independent: each of them has a 1 in its pivot column, where all the other rows have a 0. So they form a basis of $\mathrm{row}(A)$, and $\dim \mathrm{row}(A) = r$. □

Remark 1.7.7. In practice, $\mathrm{rank}(A)$ is the number of nonzero rows in any row echelon form of A: clearing the entries above the leading 1s, as in Examples 1.6.33 and 1.6.34, turns a row echelon form into the RREF without changing the number of nonzero rows. Also, Theorem 1.7.6 says that $\dim \mathrm{col}(A) = \dim \mathrm{row}(A)$: a matrix has as many linearly independent columns as linearly independent rows. This is far from obvious!

Example 1.7.8. Consider the matrix

$$M = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 0 \end{pmatrix}.$$

This matrix is already in row echelon form. The number of nonzero rows is 2, so the rank of the matrix M is $\mathrm{rank}(M) = 2$. Similarly, the matrix A in Example 1.7.4 has rank 2.

Theorem 1.7.9 (Rank–nullity theorem)**.** *For any $m \times n$ matrix A,*

$$\operatorname{rank}(A) + \dim \operatorname{null}(A) = n.$$

Equivalently, if $T : \mathbb{R}^n \to \mathbb{R}^m$ is a linear transformation, then $\dim \operatorname{im}(T) + \dim \ker(T) = n$.

Idea of proof. Let r be the number of pivots of $\operatorname{RREF}(A)$, so that $\operatorname{rank}(A) = r$ by Theorem 1.7.6. When we solve $\operatorname{RREF}(A)\,\mathbf{x} = \mathbf{0}$, the r variables corresponding to the pivot columns are expressed in terms of the remaining $n - r$ free variables. Setting one free variable equal to 1 and all others equal to 0 gives $n - r$ solutions. Every solution is a linear combination of these, and they are linearly independent, since each of them has a 1 in a position where all the others have a 0. So they form a basis of $\operatorname{null}(A)$, and $\dim \operatorname{null}(A) = n - r$. □

Example 1.7.10. For $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$, we found in Example 1.7.4 that $\dim \ker(T) = 1$ and $\dim \operatorname{im}(T) = 2$, and indeed $2 + 1 = 3 = n$. For $T(x, y) = (2x + 3y, 4x - y)$, we found in Examples 1.4.8 and 1.4.11 that $\ker(T) = \{(0, 0)\}$, of dimension 0, and $\operatorname{im}(T) = \mathbb{R}^2$, of dimension 2; indeed $2 + 0 = 2 = n$.

Combining the rank–nullity theorem with Theorem 1.6.47, we obtain several ways to recognize an invertible matrix.

Corollary 1.7.11. *For an $n \times n$ matrix A, the following statements are equivalent:*

1. *A is invertible;*
2. $\operatorname{null}(A) = \{\mathbf{0}\}$*;*
3. $\operatorname{rank}(A) = n$*;*
4. *for every $\mathbf{b} \in \mathbb{R}^n$, the equation $A\mathbf{x} = \mathbf{b}$ has exactly one solution;*
5. $\det(A) \neq 0$.

Proof. Statements 1 and 2 are equivalent by Theorem 1.6.47. By the rank–nullity theorem, $\operatorname{rank}(A) = n - \dim \operatorname{null}(A)$, so $\operatorname{rank}(A) = n$ if and only if $\dim \operatorname{null}(A) = 0$, i.e., $\operatorname{null}(A) = \{\mathbf{0}\}$; hence statements 2 and 3 are equivalent. Statement 1 implies statement 4 by Remark 1.6.38, and statement 4 implies statement 2 by taking $\mathbf{b} = \mathbf{0}$. Finally, statements 1 and 5 are equivalent by Theorem 1.6.91. □

Exercise 1.7.12. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $T(x, y) = (x - y, 2x + 3y)$. Find the kernel and image of T, and check that your answer agrees with the rank–nullity theorem.

1.8 Dual Vector Spaces

Many important quantities assign a single number to a vector: a coordinate, the average of the entries, the value of a polynomial at a point, the integral of a function, or the prediction $\mathbf{w}^\top \mathbf{x}$ of a linear model. When such an assignment is linear, it is called a linear functional. The linear functionals on a vector space V form a new vector space, the dual space V^*. Duality explains what the transpose of a matrix really is, gives a convenient way to compute coordinates, and provides "certificates" showing that a system of linear equations has no solution. In machine learning, the weight vector of a linear model and the derivative of a loss function are naturally elements of a dual space.

Throughout this section, V and W are vector spaces over a field $\mathbb{F}$.

1.8.1 Linear functionals and the dual space

Definition 1.8.1 (Linear functional, dual space)**.** A linear functional (or linear form) on V is a linear transformation $\varphi : V \to \mathbb{F}$. The set of all linear functionals on V is called the dual space of V and is denoted by V^*. For $\varphi, \psi \in V^*$ and $c \in \mathbb{F}$, we define $\varphi + \psi$ and $c\varphi$ pointwise:

$$(\varphi + \psi)(v) = \varphi(v) + \psi(v), \qquad (c\varphi)(v) = c\,\varphi(v) \qquad (v \in V).$$

Proposition 1.8.2. *With these operations, V^* is a vector space over $\mathbb{F}$.*

Proof. The functions $\varphi+\psi$ and $c\varphi$ are again linear; for example, $(\varphi+\psi)(u+v) = \varphi(u)+\varphi(v)+\psi(u)+\psi(v) = (\varphi+\psi)(u)+(\varphi+\psi)(v)$, and similarly for scalar multiples. The vector space axioms hold because they hold pointwise in $\mathbb{F}$, exactly as for the space $\mathcal{F}(S)$ of all functions (Proposition 1.2.10). The zero vector of V^* is the zero functional $v \mapsto 0$. □

On $\mathbb{R}^n$, the linear functionals are exactly the row vectors.

Proposition 1.8.3. *For every row vector $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^{1\times n}$, the map $\varphi_a(\mathbf{x}) = \mathbf{a}\mathbf{x} = a_1x_1 + \cdots + a_nx_n$ is a linear functional on $\mathbb{R}^n$. Conversely, every linear functional φ on $\mathbb{R}^n$ is of this form, with $\mathbf{a} = (\varphi(\mathbf{e}_1), \dots, \varphi(\mathbf{e}_n))$. The map $\mathbf{a} \mapsto \varphi_a$ is an isomorphism $\mathbb{R}^{1\times n} \to (\mathbb{R}^n)^*$.*

Proof. The first statement is the case $m = 1$ of matrix–vector multiplication, and the second is the case $m = 1$ of Theorem 1.6.9: the matrix of $\varphi : \mathbb{R}^n \to \mathbb{R}$ is the $1 \times n$ matrix with columns $\varphi(\mathbf{e}_1), \dots, \varphi(\mathbf{e}_n)$. The map $\mathbf{a} \mapsto \varphi_a$ is linear, since $\varphi_{\mathbf{a}+\mathbf{b}}(\mathbf{x}) = (\mathbf{a}+\mathbf{b})\mathbf{x} = \mathbf{a}\mathbf{x}+\mathbf{b}\mathbf{x}$ and $\varphi_{c\mathbf{a}}(\mathbf{x}) = c\,\mathbf{a}\mathbf{x}$; it is surjective by the second statement, and injective since $\varphi_a = 0$ implies $a_i = \varphi_a(\mathbf{e}_i) = 0$ for all i. □

So vectors of $\mathbb{R}^n$ are columns, and functionals on $\mathbb{R}^n$ are rows; applying a functional to a vector is the product "row times column".

Example 1.8.4. The following maps are linear functionals.

- The coordinate functional $\mathbf{x} \mapsto x_i$ on $\mathbb{R}^n$, given by the row $\mathbf{e}_i^\top$.
- The mean $\mathbf{x} \mapsto \frac{1}{n}(x_1 + \cdots + x_n)$ on $\mathbb{R}^n$, given by the row $\frac{1}{n}(1, \dots, 1)$.
- The trace $\operatorname{tr} : \mathbb{R}^{n\times n} \to \mathbb{R}$ (Theorem 1.6.77).
- The evaluation at a point a, $\operatorname{ev}_a : \mathbb{R}[x] \to \mathbb{R}$, $p \mapsto p(a)$, and the maps $p \mapsto p'(0)$ and $p \mapsto \int_0^1 p(x)\,dx$.

On the other hand, $\mathbf{x} \mapsto x_1x_2$, $\mathbf{x} \mapsto x_1 + 1$ and $A \mapsto \det(A)$ (for $n \geq 2$) are not linear, so they are not linear functionals.

Remark 1.8.5 (Linear models). A linear model in machine learning predicts a number from a feature vector $\mathbf{x} \in \mathbb{R}^n$ by $f(\mathbf{x}) = w_1x_1+\cdots+w_nx_n = \mathbf{w}^\top\mathbf{x}$. So the model is a linear functional, and learning a linear model means choosing an element of the dual space $(\mathbb{R}^n)^*$. This is one reason why it is natural to write the weights as a row $\mathbf{w}^\top$, and the data as columns.

The kernel of a nonzero linear functional on $\mathbb{R}^n$ is the null space of a nonzero $1 \times n$ matrix a, so by the rank–nullity theorem it is a subspace of dimension $n - 1$, called a hyperplane (through the origin). For example, the kernel of $\mathbf{x} \mapsto x_1 + x_2 + x_3$ is the plane $x_1 + x_2 + x_3 = 0$ in $\mathbb{R}^3$. Using coordinates, the same holds for every n-dimensional vector space.

1.8.2 The dual basis

For indices i and j, we use the Kronecker delta

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Theorem 1.8.6 (Dual basis)**.** *Let V be finite-dimensional with an ordered basis $\mathcal{B} = (b_1, \dots, b_n)$. For $i = 1, \dots, n$, let $b^i \in V^*$ be the linear functional that maps a vector to its i-th coordinate with respect to $\mathcal{B}$:*

$$b^i(a_1 b_1 + \cdots + a_n b_n) = a_i, \qquad \text{so that} \qquad b^i(b_j) = \delta_{ij}.$$

Then $(b^1, \dots, b^n)$ is an ordered basis of V^*, called the dual basis of $\mathcal{B}$. Moreover, every $\varphi \in V^*$ can be written as

$$\varphi = \varphi(b_1)\, b^1 + \varphi(b_2)\, b^2 + \cdots + \varphi(b_n)\, b^n.$$

In particular, $\dim V^* = \dim V$*, so $V^* \cong V$.*

Here the superscripts in $b^1, \dots, b^n$ are indices, not powers.

Proof. Each b^i is linear, since the coordinate map $v \mapsto [v]_{\mathcal{B}}$ is linear (Theorem 1.4.27). First, two linear functionals that agree on $b_1, \dots, b_n$ are equal: if $v = \sum_j a_j b_j$, then $\varphi(v) = \sum_j a_j \varphi(b_j)$ is determined by the values $\varphi(b_j)$. Now the functionals φ and $\sum_i \varphi(b_i) b^i$ agree on each b_j, since $\sum_i \varphi(b_i) b^i(b_j) = \sum_i \varphi(b_i)\delta_{ij} = \varphi(b_j)$; hence they are equal, and $b^1, \dots, b^n$ span V^*. If $\sum_i c_i b^i = 0$, then applying both sides to b_j gives $c_j = 0$, so $b^1, \dots, b^n$ are linearly independent. The last statement follows from Corollary 1.4.29. □

So the coordinates of a functional in the dual basis are simply its values on the basis vectors. For the standard basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ of $\mathbb{R}^n$, the dual basis consists of the coordinate functionals $\mathbf{x} \mapsto x_i$, i.e., of the rows $\mathbf{e}_1^\top, \dots, \mathbf{e}_n^\top$. For other bases of $\mathbb{R}^n$, the dual basis can be computed with an inverse matrix.

Proposition 1.8.7. *Let $\mathbf{b}_1, \dots, \mathbf{b}_n$ be a basis of $\mathbb{R}^n$, and let P be the (invertible) matrix with columns $\mathbf{b}_1, \dots, \mathbf{b}_n$. Then the dual basis functionals are given by the rows of P^{-1}: if $\mathbf{r}_i$ is the i-th row of P^{-1}, then $b^i(\mathbf{x}) = \mathbf{r}_i \mathbf{x}$.*

Proof. The (i, j) entry of $P^{-1}P = I_n$ is $\mathbf{r}_i \mathbf{b}_j = \delta_{ij}$. So the functionals $\mathbf{x} \mapsto \mathbf{r}_i \mathbf{x}$ satisfy the defining property of the dual basis. (Equivalently, $P^{-1}\mathbf{x} = [\mathbf{x}]_{\mathcal{B}}$ is the coordinate vector of $\mathbf{x}$.) □

Example 1.8.8. For the basis $\mathbf{b}_1 = (1, 1)^\top$, $\mathbf{b}_2 = (1, -1)^\top$ of $\mathbb{R}^2$, we have $P^{-1} = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$, so

$$b^1(\mathbf{x}) = \frac{x_1 + x_2}{2}, \qquad b^2(\mathbf{x}) = \frac{x_1 - x_2}{2}.$$

Indeed, $b^1(\mathbf{b}_1) = 1$, $b^1(\mathbf{b}_2) = 0$, $b^2(\mathbf{b}_1) = 0$ and $b^2(\mathbf{b}_2) = 1$.

Example 1.8.9 (Lagrange interpolation)**.** Let $V = \mathbb{R}[x]_{\leq 2}$. The map $T(p) = (p(-1), p(0), p(1))$ is an isomorphism $V \to \mathbb{R}^3$, since its matrix with respect to the basis $(1, x, x^2)$ is $\begin{pmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}$, which has determinant $2 \neq 0$ (Theorem 1.6.58). Hence the polynomials $L_a = T^{-1}(e_a)$, i.e.,

$$L_{-1}(x) = \frac{x(x-1)}{2}, \qquad L_0(x) = 1 - x^2, \qquad L_1(x) = \frac{x(x+1)}{2},$$

form a basis of V (Proposition 1.4.25); they satisfy $L_a(c) = \delta_{ac}$ for $a, c \in \{-1, 0, 1\}$. Its dual basis consists of the evaluation functionals $\mathrm{ev}_{-1}, \mathrm{ev}_0, \mathrm{ev}_1$, and Theorem 1.8.6 gives the Lagrange interpolation formula

$$p = p(-1)\, L_{-1} + p(0)\, L_0 + p(1)\, L_1 \qquad \text{for every } p \in \mathbb{R}[x]_{\le 2}.$$

(Apply T to both sides: both have the values $p(-1)$, $p(0)$, $p(1)$.) For example, the polynomial of degree at most 2 with values $1, 2, 5$ at $-1, 0, 1$ is $L_{-1} + 2L_0 + 5L_1 = 2 + 2x + x^2$.

1.8.3 The dual map and the transpose

A linear transformation $T : V \to W$ turns functionals on W into functionals on V: first apply T, then the functional.

Definition 1.8.10 (Dual map)**.** Let $T : V \to W$ be a linear transformation. The dual map (or transpose) of T is

$$T^* : W^* \to V^*, \qquad T^*(\psi) = \psi \circ T.$$

The map $T^*(\psi) = \psi \circ T$ is a linear functional on V by Lemma 1.4.24, and T^* is linear, since $(\psi_1+\psi_2)\circ T = \psi_1\circ T + \psi_2 \circ T$ and $(c\psi)\circ T = c(\psi \circ T)$. Note that T^* goes in the opposite direction: from W^* to V^*. If $S : W \to U$ is another linear transformation, then $(S\circ T)^*(\psi) = \psi \circ S \circ T = T^*(S^*(\psi))$, so

$$(S \circ T)^* = T^* \circ S^*.$$

Theorem 1.8.11. *Let V and W be finite-dimensional vector spaces over $\mathbb{R}$, with ordered bases $\mathcal{B} = (b_1, \ldots, b_n)$ and $\mathcal{C} = (c_1, \ldots, c_m)$, and let $\mathcal{B}^* = (b^1, \ldots, b^n)$ and $\mathcal{C}^* = (c^1, \ldots, c^m)$ be the dual bases. If $T : V \to W$ has the matrix $A = [T]_{\mathcal{C},\mathcal{B}}$, then the dual map has the matrix*

$$[T^*]_{\mathcal{B}^*,\mathcal{C}^*} = A^\top.$$

Proof. By definition of A, the i-th column of A is the coordinate vector of $T(b_i)$, so $c^j(T(b_i)) = A_{ji}$. By Theorem 1.8.6, the coordinates of the functional $T^*(c^j)$ in the dual basis $\mathcal{B}^*$ are its values on $b_1, \ldots, b_n$, which are

$$T^*(c^j)(b_i) = c^j\big(T(b_i)\big) = A_{ji} \qquad (i = 1, \ldots, n).$$

So the j-th column of $[T^*]_{\mathcal{B}^*,\mathcal{C}^*}$ is $(A_{j1}, \ldots, A_{jn})^\top$, which is the j-th column of $A^\top$. □

For $T = T_A : \mathbb{R}^n \to \mathbb{R}^m$ and functionals written as rows (Proposition 1.8.3), this is very concrete: the functional $\psi(\mathbf{y}) = \mathbf{c}\mathbf{y}$ on $\mathbb{R}^m$, given by a row $\mathbf{c}$, is mapped to

$$T_A^*(\psi)(\mathbf{x}) = \psi(A\mathbf{x}) = \mathbf{c}(A\mathbf{x}) = (\mathbf{c}A)\mathbf{x},$$

i.e., to the row $\mathbf{c}A$. So the dual map multiplies rows from the right by A, while T_A multiplies columns from the left; written as columns, $(\mathbf{c}A)^\top = A^\top \mathbf{c}^\top$. This explains the transpose conceptually, and the rule $(S \circ T)^* = T^* \circ S^*$ explains why $(AB)^\top = B^\top A^\top$ reverses the order.

1.8.4 Annihilators and solvability of linear systems

Definition 1.8.12 (Annihilator)**.** Let U be a subspace of V. The annihilator of U is the set of all functionals that vanish on U:

$$U^\circ = \{\varphi \in V^* : \varphi(u) = 0 \text{ for all } u \in U\}.$$

The annihilator is a subspace of V^* (check this). In $\mathbb{R}^n$, it can be computed with a null space.

Theorem 1.8.13. *Let $U = \operatorname{col}(B)$ be a k-dimensional subspace of $\mathbb{R}^n$, where B is an $n \times k$ matrix with linearly independent columns. A row $\mathbf{a}$ defines a functional in U° if and only if $\mathbf{a}B = \mathbf{0}$, i.e., $\mathbf{a}^\top \in \operatorname{null}(B^\top)$. In particular,*

$$\dim U + \dim U^\circ = n.$$

Proof. The functional $\mathbf{x} \mapsto \mathbf{a}\mathbf{x}$ vanishes on U if and only if it vanishes on the columns $\mathbf{b}_1, \dots, \mathbf{b}_k$ of B (by linearity), i.e., if and only if $\mathbf{a}\mathbf{b}_i = 0$ for all i, which means $\mathbf{a}B = \mathbf{0}$, or equivalently $B^\top \mathbf{a}^\top = \mathbf{0}$. By Proposition 1.8.3, U° is therefore isomorphic to $\operatorname{null}(B^\top)$. Since $\operatorname{rank}(B^\top) = \operatorname{rank}(B) = k$ (Theorem 1.7.6), the rank–nullity theorem gives $\dim U^\circ = n - k$. □

Example 1.8.14. For $U = \operatorname{span}\{(1,1,1)^\top\}$ in $\mathbb{R}^3$, the annihilator consists of the rows $\mathbf{a}$ with $a_1 + a_2 + a_3 = 0$, a 2-dimensional space, and indeed $1 + 2 = 3$.

Annihilators give a clean criterion for when a system of linear equations has a solution.

Theorem 1.8.15 (Fredholm alternative)**.** *Let A be an $m \times n$ matrix and $\mathbf{b} \in \mathbb{R}^m$. The system $A\mathbf{x} = \mathbf{b}$ has a solution if and only if $\mathbf{y}\mathbf{b} = 0$ for every row $\mathbf{y} \in \mathbb{R}^{1\times m}$ with $\mathbf{y}A = \mathbf{0}$.*

Proof. If $A\mathbf{x} = \mathbf{b}$ and $\mathbf{y}A = \mathbf{0}$, then $\mathbf{y}\mathbf{b} = \mathbf{y}A\mathbf{x} = 0$. Conversely, suppose that $A\mathbf{x} = \mathbf{b}$ has no solution, i.e., $\mathbf{b} \notin U = \operatorname{col}(A)$. Let $r = \dim U$, choose a basis $u_1, \dots, u_r$ of U, and let $U' = \operatorname{span}\{u_1, \dots, u_r, \mathbf{b}\}$. The vectors $u_1, \dots, u_r, \mathbf{b}$ are linearly independent (a nontrivial relation would have to involve $\mathbf{b}$, and would express $\mathbf{b}$ as an element of U), so $\dim U' = r + 1$. By Theorem 1.8.13, $\dim U'^\circ = m - r - 1 < m - r = \dim U^\circ$. Since $U \subseteq U'$, we have $U'^\circ \subseteq U^\circ$, and the dimensions show that there is a functional $\mathbf{y} \in U^\circ$ with $\mathbf{y} \notin U'^\circ$. Such a row $\mathbf{y}$ vanishes on the columns of A, i.e., $\mathbf{y}A = \mathbf{0}$, but it does not vanish on all of U', so $\mathbf{y}\mathbf{b} \neq 0$. □

So if a system has no solution, there is always a simple certificate that proves it: a combination $\mathbf{y}$ of the equations whose left-hand sides cancel ($\mathbf{y}A = \mathbf{0}$), but whose right-hand sides do not ($\mathbf{y}\mathbf{b} \neq 0$).

Example 1.8.16. The system $x_1 + 2x_2 = 1$, $2x_1 + 4x_2 = 3$ has no solution. A certificate is $\mathbf{y} = (2, -1)$: we have $\mathbf{y}A = (2,-1)\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} = (0, 0)$, but $\mathbf{y}\mathbf{b} = 2 - 3 = -1 \neq 0$. In words: twice the first equation minus the second one gives $0 = -1$.

Remark 1.8.17. Certificates of this kind are the starting point of duality theory in optimization (Farkas' lemma, the duality theorem of linear programming, Lagrange multipliers), which is used throughout machine learning, for example to derive the dual problem of support vector machines.

1.8.5 The double dual

Applying the construction twice gives the double dual $V^{**} = (V^*)^*$. Every vector $v \in V$ gives an element of V^{**}: the functional "evaluate at v".

Theorem 1.8.18. *For $v \in V$, let $\iota(v) \in V^{**}$ be defined by $\iota(v)(\varphi) = \varphi(v)$ for $\varphi \in V^*$. Then $\iota : V \to V^{**}$ is linear and injective, and if V is finite-dimensional, it is an isomorphism.*

Proof. Each $\iota(v)$ is linear in φ, by the definition of the operations in V^*, and ι is linear in v, since every φ is linear. We only prove the finite-dimensional case. Let $b_1, \dots, b_n$ be a basis of V, with dual basis $b^1, \dots, b^n$. If $\iota(v) = 0$, then $b^i(v) = 0$ for all i, so all coordinates of v are 0, and $v = 0$; hence ι is injective. For surjectivity, let $\Lambda \in V^{**}$, and put $v = \sum_i \Lambda(b^i) b_i$. Then $\iota(v)(b^j) = b^j(v) = \Lambda(b^j)$ for all j, so the linear functionals $\iota(v)$ and Λ on V^* agree on a basis of V^*, and hence $\iota(v) = \Lambda$. □

Remark 1.8.19. For finite-dimensional V, both V^* and V^{**} are isomorphic to V, since they have the same dimension. But an isomorphism $V \to V^*$ (such as $b_i \mapsto b^i$) depends on the choice of a basis, while $\iota : V \to V^{**}$ is defined without any choices; it is called the natural isomorphism. In Lecture 2, we will see that an inner product also gives a natural way to identify V with V^*: on $\mathbb{R}^n$ with the dot product, the vector $\mathbf{a}^\top$ corresponds to the functional $\mathbf{x} \mapsto \mathbf{ax}$. (For infinite-dimensional spaces, such as $\mathbb{R}[x]$, the map ι is injective but not surjective; we do not prove this.)

Remark 1.8.20 (Gradients and backpropagation). For a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$, the derivative at a point $\mathbf{x}$ is the linear functional

$$h \mapsto \frac{\partial f}{\partial x_1}(\mathbf{x})\, h_1 + \cdots + \frac{\partial f}{\partial x_n}(\mathbf{x})\, h_n,$$

which gives the first-order change of f in the direction h. So the derivative is naturally a row vector; the gradient $\nabla f(\mathbf{x})$ is the corresponding column vector, obtained by transposing. Backpropagation, the algorithm behind the training of neural networks, computes the derivative of a loss function with respect to all parameters by passing such row vectors backwards through the layers of the network. Each layer applies the dual map (the transpose) of its own derivative, which is the rule $(S \circ T)^* = T^* \circ S^*$ at work.

1.9 Quotient Spaces and the Isomorphism Theorems

Sometimes the differences between vectors in certain directions do not matter. The antiderivative of a function is only determined up to an additive constant; the outputs of the softmax function in a neural network do not change if the same number is added to all of its inputs; and the information a linear transformation keeps about a vector does not change if we add an element of its kernel. The quotient space V/U formalizes the idea of "the vector space V, where we ignore the directions in a subspace U". The isomorphism theorems then describe how quotients, kernels and images fit together, and they give the rank–nullity theorem for arbitrary linear transformations, as well as a formula for the dimension of the intersection of two subspaces.

Throughout this section, V is a vector space over a field $\mathbb{F}$, and U is a subspace of V.

1.9.1 Cosets

Definition 1.9.1 (Coset)**.** For $v \in V$, the coset of U through v is the set

$$v + U = \{v + u : u \in U\}.$$

In $\mathbb{R}^n$, cosets are the shifted copies of the subspace U (affine subspaces, see Lecture 2). For example, if U is a line through the origin in $\mathbb{R}^2$, the cosets of U are the lines parallel to U.

Lemma 1.9.2. *For all $v, w \in V$:*

1. *$v \in v + U$;*
2. *$v + U = w + U$ if and only if $v - w \in U$;*
3. *two cosets are either equal or disjoint.*

Proof. (1) $v = v + 0$ with $0 \in U$.

(2) If $v+U = w+U$, then $v \in w+U$ by part 1, so $v = w+u$ for some $u \in U$, and $v-w = u \in U$. Conversely, if $v - w = u_0 \in U$, then every element $v + u = w + (u_0 + u)$ of $v + U$ lies in $w + U$, and similarly $w + U \subseteq v + U$, since $w - v = -u_0 \in U$.

(3) If $x \in (v + U) \cap (w + U)$, then $x - v \in U$ and $x - w \in U$, so $v - w = (x - w) - (x - v) \in U$, and $v + U = w + U$ by part 2. □

So the cosets of U cut V into disjoint pieces, and every vector lies in exactly one of them. The vectors in the same coset are those that differ by an element of U; any of them is called a representative of the coset.

Example 1.9.3. Let $U = \operatorname{span}\{(1,1)^\top\}$ in $\mathbb{R}^2$. Two vectors $\mathbf{v}$ and $\mathbf{w}$ lie in the same coset if and only if $\mathbf{v} - \mathbf{w}$ is a multiple of $(1,1)^\top$, i.e., if and only if $v_1 - v_2 = w_1 - w_2$. So the cosets are the lines $\{x : x_1 - x_2 = c\}$, $c \in \mathbb{R}$, parallel to U. For instance, $(3,1)^\top + U = (2,0)^\top + U$, since $(3,1)^\top - (2,0)^\top = (1,1)^\top \in U$.

Example 1.9.4. Let U be the subspace of constant functions in the space $\mathcal{F}(\mathbb{R})$ of all functions $\mathbb{R} \to \mathbb{R}$. If F is an antiderivative of f, then the set of all antiderivatives of f is the coset $F + U = \{F + C : C \in \mathbb{R}\}$. The familiar notation $\int f(x)\,dx = F(x) + C$ describes exactly this coset.

1.9.2 The quotient space

Definition 1.9.5 (Quotient space)**.** The quotient space of V by U is the set of all cosets of U,

$$V/U = \{v + U : v \in V\},$$

with the operations

$$(v + U) + (w + U) = (v + w) + U, \qquad c\,(v + U) = cv + U \qquad (c \in \mathbb{F}).$$

The map $\pi : V \to V/U$, $\pi(v) = v + U$, is called the quotient map.

Proposition 1.9.6. *The operations on V/U are well defined, and with them V/U is a vector space over $\mathbb{F}$, whose zero vector is the coset $0 + U = U$. The quotient map π is a surjective linear transformation with $\ker(\pi) = U$.*

Proof. A coset has many representatives, so we must check that the results of the operations do not depend on the choice of representatives. Let $v + U = v' + U$ and $w + U = w' + U$, i.e., $v - v' \in U$ and $w - w' \in U$ (Lemma 1.9.2). Then $(v + w) - (v' + w') = (v - v') + (w - w') \in U$ and $cv - cv' = c(v - v') \in U$, so $(v + w) + U = (v' + w') + U$ and $cv + U = cv' + U$. The vector space axioms for V/U follow from those for V; for example, $(v + U) + (0 + U) = v + U$, and the additive inverse of $v + U$ is $(-v) + U$. The quotient map is linear by the definition of the operations, and it is clearly surjective. Finally, $\pi(v) = 0 + U$ if and only if $v - 0 \in U$, by Lemma 1.9.2. □

So in V/U we compute with vectors as usual, except that vectors which differ by an element of U are considered equal; in particular, all vectors of U become 0.

Example 1.9.7. For $U = \operatorname{span}\{(1,1)^\top\}$ as in Example 1.9.3, every coset is determined by the number $x_1 - x_2$, and the map $\mathbb{R}^2/U \to \mathbb{R}$, $\mathbf{x} + U \mapsto x_1 - x_2$, is well defined (it does not depend on the representative $\mathbf{x}$), linear and bijective. So $\mathbb{R}^2/U \cong \mathbb{R}$: collapsing a line in the plane to a point leaves a line.

To compute the dimension of a quotient space, we extend a basis of U to a basis of V.

Theorem 1.9.8. *If V is finite-dimensional, then $\dim(V/U) = \dim V - \dim U$. More precisely, if $u_1, \dots, u_k, w_1, \dots, w_m$ is a basis of V such that $u_1, \dots, u_k$ is a basis of U, then $w_1 + U, \dots, w_m + U$ is a basis of V/U.*

Proof. By Corollary 1.3.14, U is finite-dimensional, and every basis of U, which consists of linearly independent vectors of V, can be extended to such a basis of V. The cosets $w_j + U$ span V/U: if $v = \sum_i a_i u_i + \sum_j c_j w_j$, then $v + U = \sum_j c_j (w_j + U)$, since $\sum_i a_i u_i \in U$. They are linearly independent: if $\sum_j c_j (w_j + U) = 0 + U$, then $\sum_j c_j w_j \in U$, so $\sum_j c_j w_j = \sum_i a_i u_i$ for some scalars a_i. Since $u_1, \dots, u_k, w_1, \dots, w_m$ are linearly independent, all c_j (and a_i) are 0. □

Remark 1.9.9 (Complements). The proof shows that $W = \operatorname{span}\{w_1, \dots, w_m\}$ is a subspace such that every $v \in V$ can be written uniquely as $v = u + w$ with $u \in U$ and $w \in W$; such a W is called a complement of U. Every coset $v + U$ then contains exactly one vector of W, and $w \mapsto w + U$ is an isomorphism $W \to V/U$. For example, the x_1-axis is a complement of $U = \operatorname{span}\{(1,1)^\top\}$ in $\mathbb{R}^2$: the coset $\{x_1 - x_2 = c\}$ meets the x_1-axis exactly in $(c, 0)$. Complements are not unique, so this description of V/U depends on a choice. In Lecture 2, the orthogonal complement $U^\perp$ will be a canonical choice.

1.9.3 The isomorphism theorems

Theorem 1.9.10 (First isomorphism theorem)**.** *Let $T : V \to W$ be a linear transformation, and let $K = \ker(T)$. Then*

$$\bar{T} : V/K \to \operatorname{im}(T), \qquad \bar{T}(v + K) = T(v),$$

is a well-defined isomorphism. In particular, $V/\ker(T) \cong \operatorname{im}(T)$.

Proof. If $v + K = v' + K$, then $v - v' \in K$, so $T(v) - T(v') = T(v - v') = 0$; hence $\bar{T}$ is well defined. It is linear, since T is: $\bar{T}((v + K) + (w + K)) = T(v + w) = T(v) + T(w)$, and similarly for scalar multiples. It is surjective by the definition of $\operatorname{im}(T)$. It is injective: if $\bar{T}(v + K) = T(v) = 0$, then $v \in K$, so $v + K = 0 + K$ is the zero vector of V/K (Lemma 1.4.18). □

In words: a linear transformation identifies exactly those vectors that differ by an element of its kernel, and after this identification, it becomes an isomorphism onto its image. Combined with Theorem 1.9.8, the first isomorphism theorem gives the rank–nullity theorem for all linear transformations, not only for matrices.

Corollary 1.9.11 (Rank–nullity theorem)**.** *Let $T : V \to W$ be a linear transformation, where V is finite-dimensional. Then*

$$\dim \ker(T) + \dim \operatorname{im}(T) = \dim V.$$

Proof. By Theorem 1.9.10, $\operatorname{im}(T) \cong V/\ker(T)$, which has dimension $\dim V - \dim \ker(T)$ by Theorem 1.9.8; isomorphic spaces have the same dimension (Proposition 1.4.25). □

Example 1.9.12. For $T(x, y) = (x + 2y, 2x + 4y)$ (Example 1.4.13 and Figure 1.3), the cosets of $\ker(T) = \operatorname{span}\{(-2, 1)^\top\}$ are the lines $x + 2y = c$, and $\bar{T}$ maps the coset $\{x + 2y = c\}$ to the point $c\,(1, 2)$ of $\operatorname{im}(T)$. Figure 1.3 is therefore a picture of the isomorphism $\mathbb{R}^2/\ker(T) \cong \operatorname{im}(T)$: each colored line on the left is one element of the quotient space.

Example 1.9.13. The derivative $D : \mathbb{R}[x]_{\le 3} \to \mathbb{R}[x]_{\le 2}$, $p \mapsto p'$, is surjective, and its kernel is the space of constant polynomials. So $\mathbb{R}[x]_{\le 3}/\{\text{constants}\} \cong \mathbb{R}[x]_{\le 2}$: a polynomial "up to an additive constant" is the same thing as its derivative. Similarly, the trace $\operatorname{tr} : \mathbb{R}^{n\times n} \to \mathbb{R}$ is surjective, so $\mathbb{R}^{n\times n}$ modulo the matrices with trace zero is isomorphic to $\mathbb{R}$.

To compare two subspaces, we need their sum.

Definition 1.9.14 (Sum of subspaces)**.** For subspaces U and W of V, their sum is $U + W = \{u + w : u \in U,\ w \in W\}$.

One checks that $U + W$ is a subspace of V; it is the smallest subspace that contains both U and W. Also $U \cap W$ is a subspace (practice problems).

Theorem 1.9.15 (Second isomorphism theorem)**.** *Let U and W be subspaces of V. Then*

$$(U + W)/W \cong U/(U \cap W).$$

Proof. Consider $T : U \to (U + W)/W$, $T(u) = u + W$, which is linear (it is the restriction of the quotient map). It is surjective: every element of $(U+W)/W$ has the form $(u+w)+W = u+W = T(u)$, since $w \in W$. Its kernel consists of the $u \in U$ with $u \in W$, i.e., $\ker(T) = U \cap W$. Now apply the first isomorphism theorem. □

Corollary 1.9.16 (Dimension formula)**.** *If U and W are finite-dimensional subspaces of V, then*

$$\dim(U + W) + \dim(U \cap W) = \dim U + \dim W.$$

Proof. By Theorems 1.9.15 and 1.9.8, $\dim(U + W) - \dim W = \dim U - \dim(U \cap W)$. □

Example 1.9.17. Two different planes U and W through the origin in $\mathbb{R}^3$ always meet in a line: since $U + W$ is a subspace that is strictly larger than the 2-dimensional plane U, it is all of $\mathbb{R}^3$, and hence $\dim(U \cap W) = 2 + 2 - 3 = 1$. Similarly, two 4-dimensional subspaces of $\mathbb{R}^7$ always have a nonzero intersection, since $\dim(U \cap W) = 8 - \dim(U + W) \geq 8 - 7 = 1$.

Theorem 1.9.18 (Third isomorphism theorem)**.** *Let $U \subseteq W \subseteq V$ be subspaces. Then $W/U = \{w + U : w \in W\}$ is a subspace of V/U, and*

$$(V/U)/(W/U) \cong V/W.$$

Proof. W/U is the image of W under the quotient map $V \to V/U$, hence a subspace. Consider $T : V/U \to V/W$, $T(v + U) = v + W$. It is well defined: if $v + U = v' + U$, then $v - v' \in U \subseteq W$, so $v + W = v' + W$. It is linear and surjective, and $T(v + U) = 0 + W$ if and only if $v \in W$, so $\ker(T) = W/U$. Now apply the first isomorphism theorem. □

In words: ignoring first the directions in U, and then the remaining directions in W, is the same as ignoring all directions in W at once. For example, for $U = \operatorname{span}\{\mathbf{e}_3\} \subseteq W = \operatorname{span}\{\mathbf{e}_2, \mathbf{e}_3\} \subseteq V = \mathbb{R}^3$, we have $V/U \cong \mathbb{R}^2$ (keep x_1, x_2), $W/U \cong \mathbb{R}$ (keep x_2), and both sides of the theorem are isomorphic to $\mathbb{R}$ (keep only x_1).

1.9.4 Quotients and duality

Linear functionals on a quotient space are the same as functionals that vanish on the subspace.

Theorem 1.9.19. *The map $(V/U)^* \to U^\circ$, $\varphi \mapsto \varphi \circ \pi$, is an isomorphism, where $\pi : V \to V/U$ is the quotient map and U° is the annihilator of U (Section 1.8).*

Proof. For $\varphi \in (V/U)^*$, the functional $\varphi \circ \pi$ vanishes on U, since $\pi(u) = 0$ for $u \in U$; and $\varphi \mapsto \varphi \circ \pi$ is linear. It is injective: if $\varphi \circ \pi = 0$, then $\varphi = 0$, since π is surjective. It is surjective: if $\psi \in U^\circ$, then $\varphi(v + U) = \psi(v)$ is well defined (if $v - v' \in U$, then $\psi(v) - \psi(v') = \psi(v - v') = 0$) and linear, and $\varphi \circ \pi = \psi$. □

For finite-dimensional V, this is consistent with the dimensions: $\dim(V/U)^* = \dim V - \dim U = \dim U^\circ$ (Theorems 1.9.8, 1.8.6 and 1.8.13).

Remark 1.9.20 (Quotients in machine learning). The softmax function, which turns a vector $\mathbf{z} \in \mathbb{R}^n$ of scores ("logits") into probabilities,

$$\operatorname{softmax}(\mathbf{z})_i = \frac{e^{z_i}}{e^{z_1} + \cdots + e^{z_n}},$$

satisfies $\operatorname{softmax}(\mathbf{z} + c\mathbf{1}) = \operatorname{softmax}(\mathbf{z})$ for every $c \in \mathbb{R}$, where $\mathbf{1} = (1, \ldots, 1)^\top$, since the factor e^c cancels. So the softmax only depends on the coset $\mathbf{z} + \operatorname{span}\{\mathbf{1}\}$, i.e., it is really a function on the quotient space $\mathbb{R}^n / \operatorname{span}\{\mathbf{1}\}$, and logits are only meaningful up to adding a constant. This has a practical consequence: computing e^{z_i} overflows for large scores, so softmax is implemented by

first subtracting $\max_i z_i$ from all entries, which picks a different representative of the same coset (Programming Exercise P13). Choosing the representative whose entries sum to 0 is the same as centering the vector; the centered vectors form a complement of $\operatorname{span}\{\mathbf{1}\}$. Quotient spaces also appear in the construction of the function spaces L^2, where two functions are identified if their difference has integral of squares 0.

1.10 Eigenvalues and Eigenvectors

A linear transformation usually changes the direction of a vector. The vectors whose direction does not change, and which are only stretched or flipped, are called eigenvectors. They reveal the structure of a linear transformation: in a basis of eigenvectors, the transformation simply scales each coordinate. Eigenvalues and eigenvectors appear throughout machine learning, for example in principal component analysis (Lecture 2), in the PageRank algorithm, in the analysis of Markov chains, and in spectral clustering.

Throughout this section, A is a real $n \times n$ matrix.

1.10.1 Eigenvalues, eigenvectors and eigenspaces

Definition 1.10.1 (Eigenvalue and eigenvector)**.** A number $\lambda \in \mathbb{R}$ is an eigenvalue of A if there is a nonzero vector $\mathbf{v} \in \mathbb{R}^n$ with

$$A\mathbf{v} = \lambda\mathbf{v}.$$

Such a vector $\mathbf{v}$ is called an eigenvector of A for the eigenvalue λ. Eigenvalues and eigenvectors of a linear transformation $T : V \to V$ are defined in the same way, by $T(v) = \lambda v$ with $v \neq 0$.

The condition $\mathbf{v} \neq \mathbf{0}$ is essential, since $A\mathbf{0} = \lambda\mathbf{0}$ for every λ. On the other hand, $\lambda = 0$ is allowed: the eigenvectors for the eigenvalue 0 are the nonzero vectors of $\operatorname{null}(A)$.

Example 1.10.2. Let $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$. Then

$$A\begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix} = 3\begin{pmatrix} 1 \\ 1 \end{pmatrix} \qquad \text{and} \qquad A\begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 1 \cdot \begin{pmatrix} 1 \\ -1 \end{pmatrix},$$

so $(1,1)^\top$ is an eigenvector for the eigenvalue 3, and $(1,-1)^\top$ is an eigenvector for the eigenvalue 1. On the other hand, $A\mathbf{e}_1 = (2,1)^\top$ is not a multiple of $\mathbf{e}_1$, so $\mathbf{e}_1$ is not an eigenvector. Figure 1.6 shows what T_A does to the unit circle: it stretches the direction of $(1,1)^\top$ by the factor 3, and leaves the direction of $(1,-1)^\top$ unchanged.

Definition 1.10.3 (Eigenspace)**.** For $\lambda \in \mathbb{R}$, the set

$$E_\lambda = \{\mathbf{v} \in \mathbb{R}^n : A\mathbf{v} = \lambda\mathbf{v}\} = \operatorname{null}(A - \lambda I_n)$$

is called the eigenspace of A for λ. It is a subspace of $\mathbb{R}^n$ (Theorem 1.5.5), and λ is an eigenvalue if and only if $E_\lambda \neq \{0\}$; in that case, E_λ consists of the eigenvectors for λ together with 0.

Proposition 1.10.4. *A number λ is an eigenvalue of A if and only if $A - \lambda I_n$ is not invertible, if and only if $\det(A - \lambda I_n) = 0$. In particular, 0 is an eigenvalue of A if and only if A is not invertible.*

Proof. $A\mathbf{v} = \lambda\mathbf{v}$ means $(A - \lambda I_n)\mathbf{v} = \mathbf{0}$. So λ is an eigenvalue if and only if $\operatorname{null}(A - \lambda I_n) \neq \{\mathbf{0}\}$, which by Corollary 1.7.11 means that $A - \lambda I_n$ is not invertible, or equivalently, that $\det(A - \lambda I_n) = 0$. □




Figure 1.6: The matrix $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ maps the unit circle (dashed) to an ellipse (blue). The eigenvector $\mathbf{v}_1 = \frac{1}{\sqrt{2}}(1,1)$ is stretched by its eigenvalue 3, and the eigenvector $\mathbf{v}_2 = \frac{1}{\sqrt{2}}(1,-1)$ is left unchanged, since its eigenvalue is 1. The vector $\mathbf{e}_1$ is not an eigenvector: its image $A\mathbf{e}_1 = (2,1)$ points in a different direction.

1.10.2 The characteristic polynomial

Definition 1.10.5 (Characteristic polynomial)**.** The characteristic polynomial of A is

$$p_A(\lambda) = \det(A - \lambda I_n).$$

Expanding the determinant shows that $p_A(\lambda)$ is a polynomial of degree n in λ, with leading term $(-\lambda)^n$, which comes from the product $(A_{11} - \lambda)\cdots(A_{nn} - \lambda)$ of the diagonal entries. By Proposition 1.10.4, the eigenvalues of A are exactly the real roots of p_A. So, to find the eigenvalues and eigenvectors of A:

1. compute $p_A(\lambda)$ and find its real roots λ;
2. for each root λ, find the eigenspace $E_\lambda = \operatorname{null}(A - \lambda I_n)$ by row reduction.

Proposition 1.10.6. *For a 2×2 matrix A, $p_A(\lambda) = \lambda^2 - \operatorname{tr}(A)\,\lambda + \det(A)$.*

Proof. For $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we get $p_A(\lambda) = (a-\lambda)(d-\lambda) - bc = \lambda^2 - (a+d)\lambda + (ad - bc)$. □

Example 1.10.7. For $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, we get $p_A(\lambda) = \lambda^2 - 4\lambda + 3 = (\lambda - 1)(\lambda - 3)$, so the eigenvalues are 1 and 3. The eigenspaces are

$$E_3 = \operatorname{null}\begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} = \operatorname{span}\{(1,1)^\top\}, \qquad E_1 = \operatorname{null}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \operatorname{span}\{(1,-1)^\top\},$$

in agreement with Example 1.10.2.

Example 1.10.8. Let $J = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$. Expanding along the first row,

$$p_J(\lambda) = \det\begin{pmatrix} -\lambda & 1 & 1 \\ 1 & -\lambda & 1 \\ 1 & 1 & -\lambda \end{pmatrix} = -\lambda(\lambda^2 - 1) - (-\lambda - 1) + (1 + \lambda) = -\lambda^3 + 3\lambda + 2 = -(\lambda - 2)(\lambda + 1)^2.$$

So the eigenvalues are 2 and -1. Row reduction gives

$$E_2 = \operatorname{null}(J - 2I_3) = \operatorname{span}\{(1,1,1)^\top\}, \qquad E_{-1} = \operatorname{null}(J + I_3) = \operatorname{null}\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

The eigenspace E_{-1} is the plane $x + y + z = 0$, with basis $(1,-1,0)^\top$, $(1,0,-1)^\top$.

Proposition 1.10.9. *The eigenvalues of a triangular matrix are its diagonal entries.*

Proof. If A is triangular, so is $A - \lambda I_n$, and by Corollary 1.6.86, $p_A(\lambda) = (A_{11} - \lambda)\cdots(A_{nn} - \lambda)$. □

Example 1.10.10 (Eigenvectors from geometry)**.** Geometric intuition often tells us the eigenvectors without any computation. The shear and the rotations below also appear in the gallery of Figure 1.2.

- The projection onto the x-axis, $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, has the eigenvalue 1 with eigenspace the x-axis, and the eigenvalue 0 with eigenspace the y-axis.
- The reflection in the line $y = x$, $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, fixes this line (eigenvalue 1) and flips the line $y = -x$ (eigenvalue -1).
- The shear $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has $p(\lambda) = (1-\lambda)^2$, so its only eigenvalue is 1, and $E_1 = \operatorname{null}\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \operatorname{span}\{\mathbf{e}_1\}$: only the horizontal direction is preserved.
- The rotation R_θ has $p(\lambda) = \lambda^2 - 2\cos\theta\,\lambda + 1$, with discriminant $4\cos^2\theta - 4 < 0$ unless θ is a multiple of π. So a rotation by an angle other than 0 or π has no real eigenvalues: it does not preserve any direction.

Remark 1.10.11 (Complex eigenvalues)*.* By the fundamental theorem of algebra (which we state without proof), every polynomial of degree n factors into n linear factors over $\mathbb{C}$. Hence

$$p_A(\lambda) = (\lambda_1 - \lambda)(\lambda_2 - \lambda)\cdots(\lambda_n - \lambda)$$

for some $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$, which are called the complex eigenvalues of A, counted with multiplicity. For example, the rotation R_θ has the complex eigenvalues $\cos\theta \pm i\sin\theta$, with complex eigenvectors $(1, \mp i)^\top$. In these notes, "eigenvalue" means a real eigenvalue unless we say otherwise.

Theorem 1.10.12. *Let $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$ be the complex eigenvalues of A, counted with multiplicity. Then*

$$\det(A) = \lambda_1\lambda_2\cdots\lambda_n \qquad \textit{and} \qquad \operatorname{tr}(A) = \lambda_1 + \lambda_2 + \cdots + \lambda_n.$$

Proof. Setting $\lambda = 0$ in the factorization of p_A gives $\det(A) = p_A(0) = \lambda_1\cdots\lambda_n$. For the trace, we compare the coefficients of λ^{n-1}. In $(\lambda_1 - \lambda)\cdots(\lambda_n - \lambda)$, this coefficient is $(-1)^{n-1}(\lambda_1 + \cdots + \lambda_n)$. In $\det(A - \lambda I_n)$, one can check by induction on n, using the cofactor expansion, that only the product $(A_{11} - \lambda)\cdots(A_{nn} - \lambda)$ of the diagonal entries contributes to λ^{n-1}, since every other term contains at most $n-2$ diagonal entries. So the coefficient is $(-1)^{n-1}(A_{11} + \cdots + A_{nn}) = (-1)^{n-1}\operatorname{tr}(A)$. □

Example 1.10.13. For the matrix of Example 1.10.7, $1 + 3 = 4 = \operatorname{tr}(A)$ and $1 \cdot 3 = 3 = \det(A)$. For the matrix J of Example 1.10.8, $2 + (-1) + (-1) = 0 = \operatorname{tr}(J)$ and $2 \cdot (-1) \cdot (-1) = 2 = \det(J)$. For the rotation R_θ, $(\cos\theta + i\sin\theta)(\cos\theta - i\sin\theta) = 1 = \det(R_\theta)$ and the sum is $2\cos\theta = \operatorname{tr}(R_\theta)$.

Proposition 1.10.14. *Let $A\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$. Then:*

1. $A^k\mathbf{v} = \lambda^k\mathbf{v}$ *for every* $k \geq 1$*;*
2. *if* A *is invertible, then* $\lambda \neq 0$ *and* $A^{-1}\mathbf{v} = \frac{1}{\lambda}\mathbf{v}$*;*
3. $(A + cI_n)\mathbf{v} = (\lambda + c)\mathbf{v}$ *for every* $c \in \mathbb{R}$*.*

Moreover, A *and* $A^\top$ *have the same characteristic polynomial, and hence the same eigenvalues.*

Proof. (1) By induction, $A^k\mathbf{v} = A(A^{k-1}\mathbf{v}) = A(\lambda^{k-1}\mathbf{v}) = \lambda^{k-1}A\mathbf{v} = \lambda^k\mathbf{v}$. (2) $\lambda \neq 0$ by Proposition 1.10.4, and applying A^{-1} to $A\mathbf{v} = \lambda\mathbf{v}$ gives $\mathbf{v} = \lambda A^{-1}\mathbf{v}$. (3) $(A + cI_n)\mathbf{v} = A\mathbf{v} + c\mathbf{v} = (\lambda + c)\mathbf{v}$. Finally, $\det(A^\top - \lambda I_n) = \det\left((A - \lambda I_n)^\top\right) = \det(A - \lambda I_n)$. □

Note that A and $A^\top$ usually have different eigenvectors; for the shear, $E_1 = \text{span}\{\mathbf{e}_1\}$ for A, but $E_1 = \text{span}\{\mathbf{e}_2\}$ for $A^\top$.

1.10.3 Diagonalization

Theorem 1.10.15. *Eigenvectors* $\mathbf{v}_1, \ldots, \mathbf{v}_k$ *of* A *for distinct eigenvalues* $\lambda_1, \ldots, \lambda_k$ *are linearly independent.*

Proof. Suppose not, and let m be the smallest index such that $\mathbf{v}_1, \ldots, \mathbf{v}_m$ are linearly dependent. Then $m \geq 2$ (since $\mathbf{v}_1 \neq \mathbf{0}$), $\mathbf{v}_1, \ldots, \mathbf{v}_{m-1}$ are linearly independent, and $\mathbf{v}_m = c_1\mathbf{v}_1 + \cdots + c_{m-1}\mathbf{v}_{m-1}$ for some scalars c_i. Applying $A - \lambda_m I_n$ to both sides gives

$$0 = c_1(\lambda_1 - \lambda_m)\mathbf{v}_1 + \cdots + c_{m-1}(\lambda_{m-1} - \lambda_m)\mathbf{v}_{m-1}.$$

By independence, $c_i(\lambda_i - \lambda_m) = 0$, and since the eigenvalues are distinct, $c_i = 0$ for all i. But then $\mathbf{v}_m = \mathbf{0}$, a contradiction. □

Definition 1.10.16 (Diagonalizable matrix)**.** A matrix A is called diagonalizable if there are an invertible matrix P and a diagonal matrix D with

$$A = PDP^{-1}, \qquad \text{or equivalently} \qquad P^{-1}AP = D.$$

More generally, two matrices A and B are called similar if $B = P^{-1}AP$ for some invertible matrix P.

Similar matrices have the same characteristic polynomial, since by the product rule for determinants,

$$\det(P^{-1}AP - \lambda I_n) = \det\left(P^{-1}(A - \lambda I_n)P\right) = \det(P^{-1})\det(A - \lambda I_n)\det(P) = \det(A - \lambda I_n).$$

Hence they also have the same eigenvalues, trace and determinant (compare Subsection 1.6.5).

Theorem 1.10.17 (Diagonalization)**.** *An* $n \times n$ *matrix* A *is diagonalizable if and only if it has* n *linearly independent eigenvectors* $\mathbf{p}_1, \ldots, \mathbf{p}_n$*. In this case,* $A = PDP^{-1}$*, where* P *is the matrix with columns* $\mathbf{p}_1, \ldots, \mathbf{p}_n$*, and* $D = \text{diag}(\lambda_1, \ldots, \lambda_n)$ *contains the corresponding eigenvalues, in the same order.*

Proof. Let P have columns $\mathbf{p}_1, \ldots, \mathbf{p}_n$, and let $D = \text{diag}(\lambda_1, \ldots, \lambda_n)$. By the column-by-column description of matrix multiplication, the i-th column of AP is $A\mathbf{p}_i$, and the i-th column of PD is $\lambda_i\mathbf{p}_i$. So $AP = PD$ if and only if $A\mathbf{p}_i = \lambda_i\mathbf{p}_i$ for all i. Moreover, P is invertible if and only if $\text{null}(P) = \{\mathbf{0}\}$ (Corollary 1.7.11), i.e., if and only if its columns are linearly independent. Hence: if $\mathbf{p}_1, \ldots, \mathbf{p}_n$ are linearly independent eigenvectors, then $AP = PD$ with P invertible, so $A = PDP^{-1}$. Conversely, if $A = PDP^{-1}$, then $AP = PD$, so the columns of P are eigenvectors (they are nonzero, since P is invertible), and they are linearly independent. □

Corollary 1.10.18. *If an $n \times n$ matrix has n distinct real eigenvalues, then it is diagonalizable.*

Proof. Choose an eigenvector for each eigenvalue. By Theorem 1.10.15, these n eigenvectors are linearly independent. □

Remark 1.10.19. If $A = PDP^{-1}$, then the columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ of P form a basis $\mathcal{B}$ of $\mathbb{R}^n$, and $P^{-1}\mathbf{v} = [\mathbf{v}]_{\mathcal{B}}$ is the coordinate vector of $\mathbf{v}$ with respect to this basis. So $P^{-1}AP = D$ says that the matrix of T_A with respect to the basis of eigenvectors is diagonal (Subsection 1.6.3): in these coordinates, T_A simply multiplies the i-th coordinate by λ_i.

Example 1.10.20. For the matrix A of Example 1.10.7,

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}}_{P} \underbrace{\begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}}_{D} \underbrace{\frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}}_{P^{-1}}.$$

Not every matrix is diagonalizable. To see what can go wrong, we count eigenvectors.

Definition 1.10.21 (Multiplicities)**.** Let λ be an eigenvalue of A. Its algebraic multiplicity is its multiplicity as a root of p_A, i.e., the number of times the factor $(\lambda - \cdot)$ appears in p_A. Its geometric multiplicity is $\dim E_\lambda = n - \operatorname{rank}(A - \lambda I_n)$ (by the rank–nullity theorem).

Theorem 1.10.22. *For every eigenvalue, $1 \le$ geometric multiplicity $\le$ algebraic multiplicity. Moreover, A is diagonalizable if and only if p_A has n real roots (counted with multiplicity), and the geometric multiplicity of every eigenvalue equals its algebraic multiplicity.*

Proof. Let λ be an eigenvalue with geometric multiplicity g; then $g \ge 1$, since E_λ contains an eigenvector. Choose a basis $\mathbf{p}_1, \dots, \mathbf{p}_g$ of E_λ, extend it to a basis $\mathbf{p}_1, \dots, \mathbf{p}_n$ of $\mathbb{R}^n$ (Corollary 1.3.14), and let P be the invertible matrix with these columns, so that $P\mathbf{e}_i = \mathbf{p}_i$. For $i \le g$, the i-th column of $P^{-1}AP$ is $P^{-1}A\mathbf{p}_i = \lambda P^{-1}\mathbf{p}_i = \lambda \mathbf{e}_i$. Hence

$$P^{-1}AP = \begin{pmatrix} \lambda I_g & B \\ 0 & C \end{pmatrix}$$

for some $g \times (n-g)$ matrix B and some $(n-g) \times (n-g)$ matrix C. Similar matrices have the same characteristic polynomial, and expanding $\det(P^{-1}AP - tI_n)$ along the first column g times (each time, the only nonzero entry of the first column is $\lambda - t$) gives $p_A(t) = (\lambda - t)^g \det(C - tI_{n-g})$. So the factor $\lambda - t$ appears at least g times in p_A: the algebraic multiplicity is at least g.

Now suppose that $A = PDP^{-1}$ with $D = \operatorname{diag}(\lambda_1, \dots, \lambda_n)$. Then $p_A(t) = p_D(t) = (\lambda_1 - t) \cdots (\lambda_n - t)$ has n real roots. Let λ be an eigenvalue, and let m be the number of indices i with $\lambda_i = \lambda$, which is its algebraic multiplicity. Since $A - \lambda I_n = P(D - \lambda I_n)P^{-1}$, a vector $\mathbf{v}$ lies in E_λ if and only if $P^{-1}\mathbf{v} \in \operatorname{null}(D - \lambda I_n) = \operatorname{span}\{\mathbf{e}_i : \lambda_i = \lambda\}$, i.e., if and only if $\mathbf{v}$ is a linear combination of the m columns $\mathbf{p}_i$ of P with $\lambda_i = \lambda$. These columns are linearly independent, so $\dim E_\lambda = m$.

Conversely, suppose that $p_A(t) = (\mu_1 - t)^{m_1} \cdots (\mu_k - t)^{m_k}$ with distinct real numbers μ_j and $m_1 + \cdots + m_k = n$, and that $\dim E_{\mu_j} = m_j$ for all j. Choose a basis of each eigenspace E_{μ_j}; together, these are n eigenvectors. They are linearly independent: in a linear combination of them that equals 0, collect the terms from each eigenspace, so that $w_1 + \cdots + w_k = 0$ with $w_j \in E_{\mu_j}$. If some w_j were nonzero, then the nonzero ones would be eigenvectors for distinct eigenvalues whose sum is 0, which contradicts Theorem 1.10.15. So every $w_j = 0$, and since w_j is a linear combination of a basis of E_{μ_j}, all its coefficients are 0. By Theorem 1.10.17, A is diagonalizable. □

The second part says that we need "enough" eigenvectors: A is diagonalizable exactly when the eigenspaces together contain n linearly independent vectors.

Example 1.10.23.

- The matrix J of Example 1.10.8 has the eigenvalue -1 with algebraic multiplicity 2, and geometric multiplicity $3-\operatorname{rank}(J+I_3) = 3-1 = 2$. So J is diagonalizable:

$$J = PDP^{-1} \qquad \text{with} \qquad P = \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{pmatrix}, \quad D = \operatorname{diag}(2,-1,-1).$$

- The shear $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has the eigenvalue 1 with algebraic multiplicity 2, but geometric multiplicity 1. So it is not diagonalizable.
- A rotation R_θ with θ not a multiple of π has no real eigenvalues, so it is not diagonalizable with real matrices P and D. (It is diagonalizable with complex matrices.)

Remark 1.10.24. Symmetric matrices are always diagonalizable, and their eigenvectors can even be chosen to be orthonormal. This is the spectral theorem, which we prove in Lecture 2 once we have inner products.

1.10.4 The Cayley–Hamilton theorem and the minimal polynomial

Matrices can be substituted into polynomials: for $q(t) = a_0 + a_1 t + \cdots + a_m t^m$ and an $n \times n$ matrix A, let $q(A) = a_0 I_n + a_1 A + \cdots + a_m A^m$. Since powers of A commute with each other, $(qr)(A) = q(A)r(A)$ for all polynomials q and r. If $A\mathbf{v} = \lambda\mathbf{v}$, then $q(A)\mathbf{v} = q(\lambda)\mathbf{v}$, by Proposition 1.10.14.

Theorem 1.10.25 (Cayley–Hamilton)**.** *Every square matrix A satisfies $p_A(A) = 0$.*

Substituting A for λ in $\det(A - \lambda I_n)$ would give $\det(A - A) = 0$, a number rather than the zero matrix, so this is not a proof; we use the cofactor formula instead.

Proof. For $n = 1$, $p_A(t) = A_{11} - t$, and $p_A(A) = 0$. Let $n \geq 2$, and let $B(t)$ be the transposed cofactor matrix of $A - tI_n$, so that $(A - tI_n)B(t) = p_A(t)\, I_n$ for every $t \in \mathbb{R}$, by Proposition 1.6.96. Each entry of $B(t)$ is $\pm$ the determinant of an $(n-1)\times(n-1)$ matrix whose entries are polynomials of degree at most 1 in t, so it is a polynomial of degree at most $n-1$. Collecting powers of t, we can write $B(t) = B_0 + B_1 t + \cdots + B_{n-1} t^{n-1}$ with constant matrices B_j. Let $p_A(t) = c_0 + c_1 t + \cdots + c_n t^n$. Comparing the coefficients of t^j on both sides of $(A - tI_n)B(t) = p_A(t)\, I_n$ (entry by entry, two polynomials that agree for all t have the same coefficients) gives

$$AB_0 = c_0 I_n, \qquad AB_j - B_{j-1} = c_j I_n \quad (1 \leq j \leq n-1), \qquad -B_{n-1} = c_n I_n.$$

We multiply the equation for t^j by A^j from the left, and add all of them:

$$p_A(A) = AB_0 + \sum_{j=1}^{n-1} \left(A^{j+1}B_j - A^j B_{j-1}\right) - A^n B_{n-1} = 0,$$

since the sum telescopes: each term $A^{j+1}B_j$ with $0 \leq j \leq n-1$ appears once with a plus sign and once with a minus sign. □

So A^n is a linear combination of $I_n, A, \ldots, A^{n-1}$, and by induction, so is every higher power of A. If A is invertible, then $c_0 = p_A(0) = \det(A) \neq 0$, and multiplying $p_A(A) = 0$ by A^{-1} gives

$$A^{-1} = -\frac{1}{c_0}\left(c_1 I_n + c_2 A + \cdots + c_n A^{n-1}\right):$$

the inverse is a polynomial in A. This is the idea behind the Krylov subspace methods for large linear systems, such as the conjugate gradient method, which approximate $A^{-1}\mathbf{b}$ by vectors in $\operatorname{span}\{\mathbf{b}, A\mathbf{b}, \ldots, A^{k-1}\mathbf{b}\}$ and only need matrix-vector products.

Example 1.10.26. For $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, $p_A(t) = t^2 - 4t + 3$, so $A^2 = 4A - 3I_2$. Hence $A^3 = 4A^2 - 3A = 13A - 12I_2 = \begin{pmatrix} 14 & 13 \\ 13 & 14 \end{pmatrix}$, and $A(4I_2 - A) = 3I_2$ gives $A^{-1} = \frac{1}{3}(4I_2 - A) = \frac{1}{3}\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$.

The characteristic polynomial is not always the polynomial of smallest degree that A satisfies: I_n satisfies $I_n - I_n = 0$, a polynomial equation of degree 1, while $p_{I_n}(t) = (1-t)^n$.

Definition 1.10.27 (Minimal polynomial)**.** The minimal polynomial m_A of a square matrix A is a polynomial of smallest degree with leading coefficient 1 such that $m_A(A) = 0$.

Such polynomials exist, since $(-1)^n p_A$ has leading coefficient 1 and annihilates A by the Cayley–Hamilton theorem.

Proposition 1.10.28. *Let A be a square matrix.*

1. *If $q(A) = 0$ for a polynomial q, then m_A divides q. In particular, m_A divides p_A, and m_A is unique.*
2. *The real roots of m_A are exactly the eigenvalues of A.*

Proof. (1) Division with remainder gives $q = sm_A + r$ with polynomials s and r, where $r = 0$ or $\deg r < \deg m_A$. Then $r(A) = q(A) - s(A)m_A(A) = 0$, so $r = 0$ by the minimality of $\deg m_A$ (otherwise, r divided by its leading coefficient would be a smaller polynomial annihilating A). If m and m' are two minimal polynomials, each divides the other, and since both have leading coefficient 1, $m = m'$.

(2) If $A\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, then $\mathbf{0} = m_A(A)\mathbf{v} = m_A(\lambda)\mathbf{v}$, so $m_A(\lambda) = 0$. Conversely, let $m_A(\lambda) = 0$, and write $m_A(t) = (t - \lambda)s(t)$. Since $\deg s < \deg m_A$, $s(A) \neq 0$, so there is a vector $\mathbf{w}$ with $\mathbf{v} = s(A)\mathbf{w} \neq \mathbf{0}$. Then $(A - \lambda I_n)\mathbf{v} = m_A(A)\mathbf{w} = \mathbf{0}$, so $\mathbf{v}$ is an eigenvector for λ. □

Theorem 1.10.29. *A square matrix A is diagonalizable if and only if $m_A(t) = (t - \mu_1)\cdots(t - \mu_k)$ for distinct real numbers $\mu_1, \ldots, \mu_k$.*

Proof. Let $A = PDP^{-1}$, and let $\mu_1, \ldots, \mu_k$ be the distinct diagonal entries of D, the eigenvalues of A. For $q(t) = (t - \mu_1)\cdots(t - \mu_k)$, the matrix $q(D)$ is diagonal with the entries $q(\lambda_i) = 0$, so $q(A) = Pq(D)P^{-1} = 0$. By Proposition 1.10.28, m_A divides q, so m_A is a product of some of the distinct factors $t - \mu_j$; since every eigenvalue μ_j is a root of m_A, all of them appear, and $m_A = q$.

Conversely, let $m_A(t) = (t - \mu_1)\cdots(t - \mu_k)$ with distinct μ_j, and consider the Lagrange polynomials

$$\ell_j(t) = \prod_{i \neq j} \frac{t - \mu_i}{\mu_j - \mu_i}, \qquad \text{with } \ell_j(\mu_i) = \delta_{ij}.$$

The polynomial $\ell_1 + \cdots + \ell_k - 1$ has degree at most $k - 1$ and the k roots $\mu_1, \ldots, \mu_k$, so it is 0 (a nonzero polynomial of degree at most $k-1$ has at most $k-1$ roots). Hence $I_n = \ell_1(A) + \cdots + \ell_k(A)$, and every $\mathbf{v} \in \mathbb{R}^n$ is the sum of the vectors $\ell_j(A)\mathbf{v}$. Each of them lies in the eigenspace E_{μ_j}, since $(t - \mu_j)\ell_j(t)$ is a constant multiple of $m_A(t)$, so that $(A - \mu_j I_n)\ell_j(A)\mathbf{v} = \mathbf{0}$. So the eigenspaces together span $\mathbb{R}^n$. Bases of the eigenspaces together are linearly independent, as in the proof of Theorem 1.10.22, so they form a basis of $\mathbb{R}^n$ consisting of eigenvectors, and A is diagonalizable by Theorem 1.10.17. □

Example 1.10.30.

- For the shear $N = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, the only eigenvalue is 1, so m_N is $t - 1$ or $(t-1)^2$. Since $N - I_2 \neq 0$ but $(N - I_2)^2 = 0$, $m_N = (t-1)^2$ has a repeated root, and N is not diagonalizable.

- For the matrix J of Example 1.10.8, $(J-2I_3)(J+I_3)=0$, and both 2 and -1 must be roots of m_J, so $m_J=(t-2)(t+1)$, and J is diagonalizable, without computing any eigenvectors.
- If $P^2=P$, then m_P divides $t^2-t=t(t-1)$, so its roots lie in $\{0,1\}$ and are not repeated, and P is diagonalizable. Similarly, every matrix with $A^2=I_n$, such as a reflection, is diagonalizable, with eigenvalues in $\{1,-1\}$.

1.10.5 Powers of matrices and applications

Proposition 1.10.31. *If $A=PDP^{-1}$ with $D=\operatorname{diag}(\lambda_1,\dots,\lambda_n)$, then for every $k\geq 1$,*

$$A^k=PD^kP^{-1},\qquad \text{where}\quad D^k=\operatorname{diag}(\lambda_1^k,\dots,\lambda_n^k).$$

Proof. $A^k=(PDP^{-1})(PDP^{-1})\cdots(PDP^{-1})=PD(P^{-1}P)D(P^{-1}P)\cdots DP^{-1}=PD^kP^{-1}$, and D^k is computed entry by entry, since products of diagonal matrices are computed on the diagonal (Subsection 1.6.4). □

Example 1.10.32. For $A=\begin{pmatrix}2&1\\1&2\end{pmatrix}$,

$$A^k=\begin{pmatrix}1&1\\1&-1\end{pmatrix}\begin{pmatrix}3^k&0\\0&1\end{pmatrix}\frac{1}{2}\begin{pmatrix}1&1\\1&-1\end{pmatrix}=\frac{1}{2}\begin{pmatrix}3^k+1&3^k-1\\3^k-1&3^k+1\end{pmatrix}.$$

Computing A^{100} by multiplying A with itself 99 times would be much more work.

Example 1.10.33 (Fibonacci numbers)**.** The Fibonacci numbers $F_0=0$, $F_1=1$, $F_{n+1}=F_n+F_{n-1}$ satisfy

$$\begin{pmatrix}F_{n+1}\\F_n\end{pmatrix}=M\begin{pmatrix}F_n\\F_{n-1}\end{pmatrix}=\cdots=M^n\begin{pmatrix}1\\0\end{pmatrix},\qquad \text{where}\quad M=\begin{pmatrix}1&1\\1&0\end{pmatrix}.$$

The characteristic polynomial of M is $\lambda^2-\lambda-1$, with the roots

$$\varphi=\frac{1+\sqrt{5}}{2}\approx 1.618\quad\text{(the golden ratio)}\qquad\text{and}\qquad \psi=\frac{1-\sqrt{5}}{2}\approx -0.618.$$

One checks that $(\varphi,1)^\top$ and $(\psi,1)^\top$ are eigenvectors, and that $(1,0)^\top=\frac{1}{\sqrt{5}}((\varphi,1)^\top-(\psi,1)^\top)$, since $\varphi-\psi=\sqrt{5}$. By Proposition 1.10.14,

$$\begin{pmatrix}F_{n+1}\\F_n\end{pmatrix}=\frac{1}{\sqrt{5}}\left(\varphi^n\begin{pmatrix}\varphi\\1\end{pmatrix}-\psi^n\begin{pmatrix}\psi\\1\end{pmatrix}\right),\qquad\text{so}\qquad F_n=\frac{\varphi^n-\psi^n}{\sqrt{5}}.$$

This is Binet's formula. Since $|\psi|<1$, the term ψ^n quickly becomes tiny, and F_n is the integer closest to $\varphi^n/\sqrt{5}$. In particular, $F_{n+1}/F_n\to\varphi$.

Example 1.10.34 (Markov chains)**.** Suppose that the weather is either sunny or rainy, and that a sunny day is followed by a sunny day with probability 0.9, while a rainy day is followed by a sunny day with probability 0.5. If $\mathbf{x}_k=(s_k,r_k)^\top$ contains the probabilities that day k is sunny or rainy, then

$$\mathbf{x}_{k+1}=A\mathbf{x}_k,\qquad\text{where}\quad A=\begin{pmatrix}0.9&0.5\\0.1&0.5\end{pmatrix}.$$

The columns of A are probability vectors (nonnegative entries summing to 1). By Proposition 1.10.6, $p_A(\lambda)=\lambda^2-1.4\lambda+0.4=(\lambda-1)(\lambda-0.4)$, with eigenvectors $(5,1)^\top$ for $\lambda=1$ and $(1,-1)^\top$ for $\lambda=0.4$. If today is rainy, then $\mathbf{x}_0=(0,1)^\top=(\frac{5}{6},\frac{1}{6})^\top-\frac{5}{6}(1,-1)^\top$, so

$$\mathbf{x}_k=A^k\mathbf{x}_0=\begin{pmatrix}5/6\\1/6\end{pmatrix}-\frac{5}{6}(0.4)^k\begin{pmatrix}1\\-1\end{pmatrix}\longrightarrow\begin{pmatrix}5/6\\1/6\end{pmatrix}\qquad(k\to\infty).$$

In the long run, $\frac{5}{6}$ of the days are sunny, whatever the weather today. The limit $\left(\frac{5}{6}, \frac{1}{6}\right)^\top$ is the eigenvector for the eigenvalue 1, scaled to be a probability vector; it is called the stationary distribution of the Markov chain.

Remark 1.10.35 (PageRank and power iteration). The PageRank algorithm, originally used by Google to rank web pages, models a random surfer who follows a random link on each page. The importance of a page is its probability in the stationary distribution of this Markov chain, i.e., an entry of the eigenvector for the eigenvalue 1 of a huge matrix. Such eigenvectors are computed by power iteration: if A has an eigenvalue λ_1 with $|\lambda_1| > |\lambda_i|$ for all other eigenvalues, and A is diagonalizable with eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$, then writing $\mathbf{x} = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$ gives

$$A^k\mathbf{x} = \lambda_1^k\Big(c_1\mathbf{v}_1 + \sum_{i\geq 2} c_i\Big(\frac{\lambda_i}{\lambda_1}\Big)^k\mathbf{v}_i\Big),$$

so after rescaling, $A^k\mathbf{x}$ points more and more in the direction of $\mathbf{v}_1$ (if $c_1 \neq 0$). You will implement this in Programming Exercise P10.

1.10.6 Convolutions and circulant matrices

Convolutional neural networks process images and signals with filters that are applied in the same way at every position. We study the simplest version of this: signals $\mathbf{x} \in \mathbb{R}^n$ whose indices $0, 1, \dots, n-1$ are taken modulo n, so that, for example, $x_{-1} = x_{n-1}$, as if the signal were periodic. In this subsection, the entries of vectors and matrices are numbered from 0.

Definition 1.10.36 (Cyclic convolution and circulant matrices)**.** The cyclic convolution of $\mathbf{c}, \mathbf{x} \in \mathbb{R}^n$ is the vector $\mathbf{c} * \mathbf{x} \in \mathbb{R}^n$ with the entries

$$(\mathbf{c} * \mathbf{x})_j = \sum_{k=0}^{n-1} c_k x_{j-k} \qquad (j = 0, \dots, n-1),$$

where indices are taken modulo n. For fixed $\mathbf{c}$, it is linear in $\mathbf{x}$, with the matrix $\mathrm{circ}(\mathbf{c})$ whose entries are $\mathrm{circ}(\mathbf{c})_{jk} = c_{j-k}$. Such matrices are called circulant: the first column is $\mathbf{c}$, and each further column is the previous one, shifted down cyclically by one position.

For example, for $n = 4$ and $\mathbf{c} = (2, 1, 0, 1)^\top$, we get $(\mathbf{c} * \mathbf{x})_j = 2x_j + x_{j-1} + x_{j+1}$, which smooths $\mathbf{x}$ by averaging each entry with its neighbors (up to the factor 4), and

$$\mathrm{circ}(\mathbf{c}) = \begin{pmatrix} 2 & 1 & 0 & 1 \\ 1 & 2 & 1 & 0 \\ 0 & 1 & 2 & 1 \\ 1 & 0 & 1 & 2 \end{pmatrix}, \qquad \mathbf{c} * \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 8 \\ 8 \\ 12 \\ 12 \end{pmatrix}.$$

A convolutional layer learns a filter $\mathbf{c}$ with only a few nonzero entries; for images, the filters are small two-dimensional arrays, such as 3×3 filters, and the convolution is defined with two indices in the same way. (Deep learning libraries compute $\sum_k c_k x_{j+k}$ and still call it a convolution; this is the same operation with $\mathbf{c}$ reversed.)

Let $S = \mathrm{circ}(\mathbf{e}_1)$, where $\mathbf{e}_1 = (0, 1, 0, \dots, 0)^\top$ in the numbering from 0, be the cyclic shift, with $(S\mathbf{x})_j = x_{j-1}$, which moves every entry one position down and the last entry to the top.

Proposition 1.10.37. *1.* $\mathrm{circ}(\mathbf{c}) = c_0 I_n + c_1 S + c_2 S^2 + \cdots + c_{n-1} S^{n-1}$.

2. An $n \times n$ matrix A commutes with S, i.e., $AS = SA$, if and only if A is circulant.

Proof. (1) S^k shifts by k positions, $(S^k\mathbf{x})_j = x_{j-k}$, so $\big(\sum_k c_k S^k\mathbf{x}\big)_j = \sum_k c_k x_{j-k} = (\mathbf{c}*\mathbf{x})_j$.

(2) By (1), a circulant matrix is a polynomial in S, so it commutes with S. Conversely, the k-th column of AS is A applied to the k-th column $\mathbf{e}_{k+1}$ of S, so $(AS)_{jk} = A_{j,k+1}$, and SA is A with its rows shifted down, so $(SA)_{jk} = A_{j-1,k}$. Hence $AS = SA$ means that $A_{j+1,k+1} = A_{jk}$ for all j and k: the entries of A are constant along the cyclic diagonals, i.e., A_{jk} only depends on $j-k$ modulo n, and $A = \operatorname{circ}(\mathbf{c})$ with $c_m = A_{m0}$. □

Part 2 says that the linear maps that commute with shifts, i.e., that treat every position in the same way (shifting the input shifts the output), are exactly the convolutions. This is why convolutional networks use convolutions: their layers are translation equivariant, and a pattern is detected wherever it appears in an image. The eigenvectors of circulant matrices are complex. Let $\omega = e^{2\pi i/n} = \cos\frac{2\pi}{n} + i\sin\frac{2\pi}{n}$, so that $\omega^n = 1$, and let

$$\mathbf{f}_m = (1, \omega^m, \omega^{2m}, \dots, \omega^{(n-1)m})^\top \in \mathbb{C}^n \qquad (m = 0, \dots, n-1),$$

the discrete Fourier basis, which consists of the sampled oscillations with the frequencies m. Let F be the $n \times n$ matrix with the columns $\mathbf{f}_0, \dots, \mathbf{f}_{n-1}$, i.e., $F_{jm} = \omega^{jm}$, and let $\overline{F}$ be its complex conjugate. The discrete Fourier transform (DFT) of $\mathbf{x} \in \mathbb{R}^n$ is $\hat{\mathbf{x}} = \overline{F}\mathbf{x}$, with $\hat{x}_m = \sum_j x_j \omega^{-jm}$. We use complex numbers as in the remark on complex eigenvalues; the rules of matrix algebra hold over $\mathbb{C}$ as well.

Theorem 1.10.38 (The DFT diagonalizes convolutions)**.** *1. $\overline{F}F = nI_n$, so F is invertible, with $F^{-1} = \frac{1}{n}\overline{F}$.*

2. $\operatorname{circ}(\mathbf{c})\mathbf{f}_m = \hat{c}_m\mathbf{f}_m$ for every m, so $\operatorname{circ}(\mathbf{c}) = F\operatorname{diag}(\hat{c}_0, \dots, \hat{c}_{n-1})F^{-1}$: the eigenvalues of a convolution are the DFT of its filter.

3. (Convolution theorem.) $\widehat{\mathbf{c}*\mathbf{x}}_m = \hat{c}_m\hat{x}_m$ for every m.

Proof. (1) The (l,m) entry of $\overline{F}F$ is $\sum_j \omega^{-jl}\omega^{jm} = \sum_{j=0}^{n-1} r^j$ with $r = \omega^{m-l}$. For $l = m$, this is n. Otherwise $0 < |m-l| < n$, so $r \neq 1$, and the geometric sum is $\frac{r^n-1}{r-1} = 0$, since $r^n = (\omega^n)^{m-l} = 1$.

(2) Since $\omega^n = 1$, powers of ω only depend on exponents modulo n, and

$$\big(\operatorname{circ}(\mathbf{c})\mathbf{f}_m\big)_j = \sum_k c_k\omega^{(j-k)m} = \Big(\sum_k c_k\omega^{-km}\Big)\omega^{jm} = \hat{c}_m(\mathbf{f}_m)_j.$$

So $\operatorname{circ}(\mathbf{c})F = F\operatorname{diag}(\hat{c}_0, \dots, \hat{c}_{n-1})$, and we multiply by F^{-1} from the right.

(3) By (1) and (2), $\overline{F}(\mathbf{c}*\mathbf{x}) = \overline{F}F\operatorname{diag}(\hat{c}_0, \dots, \hat{c}_{n-1})\frac{1}{n}\overline{F}\mathbf{x} = \operatorname{diag}(\hat{c}_0, \dots, \hat{c}_{n-1})\hat{\mathbf{x}}$. □

All circulant matrices have the same eigenvectors $\mathbf{f}_m$, so they commute with each other. For the smoothing filter $\mathbf{c} = (2,1,0,1)^\top$ above, $\omega = i$ and $\hat{\mathbf{c}} = (4,2,0,2)$: the constant signal $\mathbf{f}_0$ is multiplied by 4, and the fastest oscillation $\mathbf{f}_2 = (1,-1,1,-1)^\top$ is mapped to $\mathbf{0}$. Smoothing damps high frequencies, and since 0 is an eigenvalue, it cannot be undone. Theorem 1.10.38 is also how convolutions are computed in practice: the fast Fourier transform (FFT) computes $\hat{\mathbf{x}}$ with $O(n\log n)$ operations instead of the $O(n^2)$ operations of a matrix-vector product, so $\mathbf{c}*\mathbf{x}$ is computed as the inverse DFT of $\hat{c}_m\hat{x}_m$.

1.11 Tensors

Data in machine learning often have more than two indices. A color image of $H \times W$ pixels has three color channels, so it is described by $3HW$ numbers x_{cij}, indexed by a channel c, a row i and a column j; a batch of N such images has four indices; and a batch of N sentences of L tokens, each represented by an embedding of dimension d, has three. Deep learning libraries such as PyTorch and

TensorFlow call such arrays tensors. In mathematics, a tensor is an element of a tensor product of vector spaces, a vector space in which vectors can be multiplied with each other, bilinearly. Arrays are the coordinates of tensors with respect to bases, just as a list of numbers is the coordinate vector of a vector (Theorem 1.4.27). In this section, we construct tensor products with the quotient spaces of Section 1.9, and we derive the properties of tensors, including their description by arrays, from this construction. We work over $\mathbb{R}$, and all vector spaces are finite-dimensional, except for the auxiliary spaces $\mathcal{F}_0(S)$ below.

1.11.1 Multilinear maps

Definition 1.11.1 (Multilinear map)**.** Let $V_1, \dots, V_k$ and W be vector spaces. A map $\tau : V_1 \times \cdots \times V_k \to W$ is multilinear (or k-linear) if it is linear in each argument when the other arguments are fixed: for each j, all $v_j, v_j' \in V_j$ and all $a \in \mathbb{R}$,

$$\tau(\dots, v_j + v_j', \dots) = \tau(\dots, v_j, \dots) + \tau(\dots, v_j', \dots),$$
$$\tau(\dots, a v_j, \dots) = a\,\tau(\dots, v_j, \dots),$$

where the dots stand for the same vectors in all terms. A multilinear map with $W = \mathbb{R}$ is called a multilinear form; for $k = 2$, we speak of bilinear maps and bilinear forms.

A multilinear map is *not* a linear map on the product space $V_1 \times \cdots \times V_k$: for a bilinear map, $\tau(2v_1, 2v_2) = 4\,\tau(v_1, v_2)$, and not $2\,\tau(v_1, v_2)$.

Example 1.11.2.

1. The dot product $(\mathbf{x}, \mathbf{y}) \mapsto \mathbf{x}^\top \mathbf{y} = x_1 y_1 + \cdots + x_n y_n$ is a bilinear form on $\mathbb{R}^n \times \mathbb{R}^n$.
2. For every $m \times n$ matrix A, the map
$$\beta_A(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top A \mathbf{y} = \sum_{i=1}^{m} \sum_{j=1}^{n} A_{ij} x_i y_j$$
is a bilinear form on $\mathbb{R}^m \times \mathbb{R}^n$: by Proposition 1.6.22, $\mathbf{x}^\top A\mathbf{y}$ is linear in $\mathbf{y}$ for fixed $\mathbf{x}$, and linear in $\mathbf{x}$ for fixed $\mathbf{y}$.
3. Matrix multiplication $(A, B) \mapsto AB$ is a bilinear map $\mathbb{R}^{m\times n} \times \mathbb{R}^{n\times p} \to \mathbb{R}^{m\times p}$, by the distributive laws.
4. The determinant of an $n \times n$ matrix is an n-linear form of its columns: by the expansion along the j-th column (Theorem 1.6.84), $\det(A) = \sum_l (-1)^{l+j} A_{lj} \det(A^{(lj)})$, where the matrices $A^{(lj)}$ do not involve the j-th column, so $\det(A)$ is a linear function of the j-th column when the other columns are fixed. But the determinant is not a linear function of the matrix (Example 1.8.4).

Bilinear and multilinear maps appear everywhere, but they are not linear, so the theory of the previous sections does not apply to them directly. The tensor product is the vector space on which they become linear.

1.11.2 The tensor product of two vector spaces

We want to multiply a vector $v \in V$ and a vector $w \in W$ to a product $v \otimes w$ that depends bilinearly on v and w, and that satisfies no other rules. To construct it, we first form all formal linear combinations of pairs (v, w), and then we impose the rules of bilinearity by a quotient.

Definition 1.11.3 (Free vector space)**.** Let S be a set. The free vector space on S is the set $\mathcal{F}_0(S)$ of all functions $f : S \to \mathbb{R}$ with $f(s) \neq 0$ for only finitely many $s \in S$, with the pointwise operations. For $s \in S$, let $\delta_s \in \mathcal{F}_0(S)$ be the function with $\delta_s(s) = 1$ and $\delta_s(t) = 0$ for $t \neq s$.

$\mathcal{F}_0(S)$ is a subspace of $\mathcal{F}(S)$ (Proposition 1.2.10): it contains the zero function, and sums and multiples of functions that vanish outside finite sets again vanish outside finite sets. Every $f \in \mathcal{F}_0(S)$ can be written in exactly one way as a finite linear combination $f = \sum_s f(s)\,\delta_s$ of the functions δ_s, since two functions are equal if they agree at every point. So $\{\delta_s : s \in S\}$ is a basis of $\mathcal{F}_0(S)$. We think of δ_s as the element s itself, and of $\mathcal{F}_0(S)$ as the space of formal linear combinations $a_1 s_1 + \cdots + a_r s_r$ of elements of S. For $S = \{1, \ldots, n\}$, it is $\mathbb{R}^n$, with $\delta_i = \mathbf{e}_i$. For $S = V \times W$, however, S is an infinite set, and $\mathcal{F}_0(V \times W)$ is a huge space, in which, for example, $\delta_{(2v,w)}$ and $2\,\delta_{(v,w)}$ are different vectors (for $v \neq 0$).

Definition 1.11.4 (Tensor product)**.** Let V and W be vector spaces, and let $R \subseteq \mathcal{F}_0(V \times W)$ be the subspace spanned by all the vectors

$$\delta_{(v+v',w)} - \delta_{(v,w)} - \delta_{(v',w)}, \qquad \delta_{(v,w+w')} - \delta_{(v,w)} - \delta_{(v,w')},$$
$$\delta_{(av,w)} - a\,\delta_{(v,w)}, \qquad \delta_{(v,aw)} - a\,\delta_{(v,w)},$$

for all $v, v' \in V$, $w, w' \in W$ and $a \in \mathbb{R}$. The tensor product of V and W is the quotient space

$$V \otimes W = \mathcal{F}_0(V \times W)/R,$$

and for $v \in V$ and $w \in W$, we write $v \otimes w = \delta_{(v,w)} + R$. The elements of $V \otimes W$ are called tensors (of order 2), and the tensors of the form $v \otimes w$ are called pure tensors.

Proposition 1.11.5. *1. The map $V \times W \to V \otimes W$, $(v, w) \mapsto v \otimes w$, is bilinear: for all $v, v' \in V$, $w, w' \in W$ and $a \in \mathbb{R}$,*

$$(v + v') \otimes w = v \otimes w + v' \otimes w, \qquad v \otimes (w + w') = v \otimes w + v \otimes w',$$
$$(av) \otimes w = a(v \otimes w) = v \otimes (aw).$$

2. Every tensor in $V \otimes W$ is a finite sum of pure tensors.

Proof. (1) Two cosets $f + R$ and $g + R$ are equal if and only if $f - g \in R$ (Section 1.9). For example, $\delta_{(v+v',w)} - (\delta_{(v,w)} + \delta_{(v',w)}) \in R$, so $(v + v') \otimes w = v \otimes w + v' \otimes w$. The other rules follow in the same way from the other generators of R.

(2) Every $f \in \mathcal{F}_0(V \times W)$ is a finite linear combination $f = \sum_l a_l\,\delta_{(v_l,w_l)}$, so $f + R = \sum_l a_l\,(v_l \otimes w_l) = \sum_l (a_l v_l) \otimes w_l$. □

For example, $0 \otimes w = (0 \cdot 0) \otimes w = 0\,(0 \otimes w) = 0$ for every $w \in W$. Not every tensor is pure; we will see below that $e_1 \otimes e_1 + e_2 \otimes e_2 \in \mathbb{R}^2 \otimes \mathbb{R}^2$ cannot be written as $v \otimes w$. The construction is designed so that the following theorem holds.

Theorem 1.11.6 (Universal property)**.** *Let $\beta : V \times W \to U$ be a bilinear map into a vector space U. Then there is exactly one linear transformation $\tilde{\beta} : V \otimes W \to U$ with*

$$\tilde{\beta}(v \otimes w) = \beta(v, w) \qquad \text{for all } v \in V \text{ and } w \in W.$$

Proof. Define $L : \mathcal{F}_0(V \times W) \to U$ by $L(f) = \sum_{(v,w)} f(v, w)\,\beta(v, w)$, where the sum runs over the finitely many pairs (v, w) with $f(v, w) \neq 0$. This map is linear, and $L(\delta_{(v,w)}) = \beta(v, w)$. It vanishes on the generators of R, since β is bilinear; for example,

$$L\big(\delta_{(v+v',w)} - \delta_{(v,w)} - \delta_{(v',w)}\big) = \beta(v + v', w) - \beta(v, w) - \beta(v', w) = 0.$$

Since $\ker(L)$ is a subspace (Theorem 1.5.5), it contains the whole span R of these vectors. Now define $\tilde{\beta}(f + R) = L(f)$. This does not depend on the choice of the representative f of the coset: if $f + R = g + R$, then $f - g \in R \subseteq \ker(L)$, so $L(f) = L(g)$. The map $\tilde{\beta}$ is linear, since L is, and $\tilde{\beta}(v \otimes w) = L(\delta_{(v,w)}) = \beta(v, w)$. It is the only such map, since a linear map on $V \otimes W$ is determined by its values on the pure tensors, which span $V \otimes W$ by Proposition 1.11.5. □

So bilinear maps $V \times W \to U$ are the same thing as linear maps $V \otimes W \to U$: the tensor product turns bilinear problems into linear ones. For example, a bilinear model $f(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top A \mathbf{y}$, which combines two inputs $\mathbf{x}$ and $\mathbf{y}$, is a linear model in the features $\mathbf{x} \otimes \mathbf{y}$; this is called bilinear pooling.

Theorem 1.11.7. *Let $b_1, \dots, b_m$ be a basis of V, and let $c_1, \dots, c_n$ be a basis of W. Then the mn pure tensors $b_i \otimes c_j$ form a basis of $V \otimes W$. In particular,*

$$\dim(V \otimes W) = \dim V \cdot \dim W,$$

and every tensor $t \in V \otimes W$ can be written in exactly one way as

$$t = \sum_{i=1}^{m} \sum_{j=1}^{n} T_{ij}\, b_i \otimes c_j.$$

For $v = \sum_i x_i b_i$ and $w = \sum_j y_j c_j$, the coefficients of $v \otimes w$ are $T_{ij} = x_i y_j$.

Proof. By bilinearity, $v \otimes w = \sum_{i,j} x_i y_j\, b_i \otimes c_j$. Since the pure tensors span $V \otimes W$, so do the tensors $b_i \otimes c_j$. To show that they are linearly independent, let $b^1, \dots, b^m$ and $c^1, \dots, c^n$ be the dual bases (Theorem 1.8.6). For fixed k and l, the map $(v, w) \mapsto b^k(v)\, c^l(w)$ is bilinear, so by Theorem 1.11.6, there is a linear functional λ_{kl} on $V \otimes W$ with $\lambda_{kl}(v \otimes w) = b^k(v)\, c^l(w)$; in particular, $\lambda_{kl}(b_i \otimes c_j) = \delta_{ki}\delta_{lj}$. If $\sum_{i,j} T_{ij}\, b_i \otimes c_j = 0$, then applying λ_{kl} gives $T_{kl} = 0$, for all k and l. □

Corollary 1.11.8. *The coordinate map with respect to the standard bases,*

$$\mathbb{R}^m \otimes \mathbb{R}^n \to \mathbb{R}^{m \times n}, \qquad \sum_{i,j} T_{ij}\, \mathbf{e}_i \otimes \mathbf{e}_j \mapsto (T_{ij})_{i,j},$$

is an isomorphism, and it maps the pure tensor $\mathbf{u} \otimes \mathbf{v}$ to the matrix $\mathbf{u}\mathbf{v}^\top$.

Proof. By Theorem 1.11.7, the tensors $\mathbf{e}_i \otimes \mathbf{e}_j$ form a basis, so the coordinate map is an isomorphism (Theorem 1.4.27, with the mn coordinates arranged as a matrix). The coordinates of $\mathbf{u} \otimes \mathbf{v}$ are $u_i v_j = (\mathbf{u}\mathbf{v}^\top)_{ij}$. □

From now on, we identify $\mathbb{R}^m \otimes \mathbb{R}^n$ with $\mathbb{R}^{m \times n}$, and the pure tensor $\mathbf{u} \otimes \mathbf{v}$ with the matrix $\mathbf{u}\mathbf{v}^\top$, which is called the outer product of $\mathbf{u}$ and $\mathbf{v}$. For example,

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix} \otimes \begin{pmatrix} 3 \\ -1 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 & -1 & 0 \\ 6 & -2 & 0 \end{pmatrix}.$$

All columns of $\mathbf{u}\mathbf{v}^\top$ are multiples of $\mathbf{u}$, so a pure tensor in $\mathbb{R}^m \otimes \mathbb{R}^n$ is a matrix of rank at most 1. Hence $\mathbf{e}_1 \otimes \mathbf{e}_1 + \mathbf{e}_2 \otimes \mathbf{e}_2 = I_2$, which has rank 2, is not a pure tensor.

1.11.3 Tensors of higher order and arrays

The same construction works for more factors. For vector spaces $V_1, \dots, V_k$, the tensor product $V_1 \otimes \cdots \otimes V_k$ is the quotient of $\mathcal{F}_0(V_1 \times \cdots \times V_k)$ by the subspace spanned by the rules of additivity and homogeneity in each of the k arguments, and $v_1 \otimes \cdots \otimes v_k$ is the coset of $\delta_{(v_1,\dots,v_k)}$. The proofs of Proposition 1.11.5 and Theorems 1.11.6 and 1.11.7 carry over word by word:

- $(v_1, \dots, v_k) \mapsto v_1 \otimes \cdots \otimes v_k$ is multilinear, and the pure tensors $v_1 \otimes \cdots \otimes v_k$ span $V_1 \otimes \cdots \otimes V_k$;
- for every multilinear map $\tau : V_1 \times \cdots \times V_k \to U$, there is exactly one linear map $\tilde{\tau} : V_1 \otimes \cdots \otimes V_k \to U$ with $\tilde{\tau}(v_1 \otimes \cdots \otimes v_k) = \tau(v_1, \dots, v_k)$;

- if $b_1^{(j)}, \dots, b_{n_j}^{(j)}$ is a basis of V_j for $j = 1, \dots, k$, then the tensors $b_{i_1}^{(1)} \otimes \cdots \otimes b_{i_k}^{(k)}$ form a basis of $V_1 \otimes \cdots \otimes V_k$, and $\dim(V_1 \otimes \cdots \otimes V_k) = n_1 n_2 \cdots n_k$.

One can also show, with the universal property, that grouping the factors does not matter: $u \otimes v \otimes w \mapsto (u \otimes v) \otimes w$ defines an isomorphism $U \otimes V \otimes W \to (U \otimes V) \otimes W$, and similarly for $U \otimes (V \otimes W)$.

Definition 1.11.9 (Tensor)**.** A tensor of order k is an element t of a tensor product $V_1 \otimes \cdots \otimes V_k$. With respect to bases of $V_1, \dots, V_k$ as above, its coordinates are the numbers $T_{i_1 \dots i_k}$ with

$$t = \sum_{i_1, \dots, i_k} T_{i_1 \dots i_k} \, b_{i_1}^{(1)} \otimes \cdots \otimes b_{i_k}^{(k)},$$

which form an array with k indices.

For $V_j = \mathbb{R}^{n_j}$ with the standard bases, we identify a tensor with the array of its coordinates, as we did for matrices. The arrays of shape $n_1 \times \cdots \times n_k$ form a vector space $\mathbb{R}^{n_1 \times \cdots \times n_k}$ with the entrywise operations (it is $\mathcal{F}(S)$ for the set S of all index tuples, Proposition 1.2.10), and the coordinate map is an isomorphism

$$\mathbb{R}^{n_1} \otimes \cdots \otimes \mathbb{R}^{n_k} \cong \mathbb{R}^{n_1 \times \cdots \times n_k}, \qquad \text{with} \qquad (u^{(1)} \otimes \cdots \otimes u^{(k)})_{i_1 \dots i_k} = u_{i_1}^{(1)} u_{i_2}^{(2)} \cdots u_{i_k}^{(k)}.$$

These arrays are the tensors of deep learning libraries. A tensor of order 1 is a vector, a tensor of order 2 is a matrix, and a number is sometimes called a tensor of order 0. A color image of $H \times W$ pixels is a tensor in $\mathbb{R}^3 \otimes \mathbb{R}^H \otimes \mathbb{R}^W$, and a batch of 64 color images of 224×224 pixels is a tensor in $\mathbb{R}^{64} \otimes \mathbb{R}^3 \otimes \mathbb{R}^{224} \otimes \mathbb{R}^{224}$, a space of dimension almost 10 million.

Example 1.11.10 (Reshaping)**.** Listing the coordinates of a tensor in a fixed order gives an isomorphism $\mathbb{R}^{n_1} \otimes \cdots \otimes \mathbb{R}^{n_k} \to \mathbb{R}^{n_1 \cdots n_k}$ (Theorem 1.4.27). The usual order is row by row, with the last index changing fastest; for example,

$$\begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \mapsto (a, b, c, d, e, f).$$

This is what `reshape` does in NumPy and PyTorch. For vectors $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$, it maps $\mathbf{u} \otimes \mathbf{v}$ to the vector $(u_1 v_1, \dots, u_1 v_n, u_2 v_1, \dots, u_m v_n)$, which NumPy computes as `np.kron(u, v)`. Grouping only some of the factors gives matrices: by the isomorphism $\mathbb{R}^{n_1} \otimes \mathbb{R}^{n_2} \otimes \mathbb{R}^{n_3} \cong \mathbb{R}^{n_1} \otimes (\mathbb{R}^{n_2} \otimes \mathbb{R}^{n_3}) \cong \mathbb{R}^{n_1} \otimes \mathbb{R}^{n_2 n_3}$, a tensor of shape $n_1 \times n_2 \times n_3$ becomes an $n_1 \times (n_2 n_3)$ matrix, its unfolding, whose i-th row lists the coordinates T_{ijk} with the first index i. Reshaping changes only the arrangement of the numbers, so it costs no computation at all.

1.11.4 Multilinear forms and linear maps as tensors

By the universal property, a multilinear form on $V_1 \times \cdots \times V_k$ is the same thing as a linear functional on $V_1 \otimes \cdots \otimes V_k$. The dual space of a tensor product is again a tensor product.

Theorem 1.11.11. *1. There is an isomorphism $V_1^* \otimes \cdots \otimes V_k^* \to (V_1 \otimes \cdots \otimes V_k)^*$ that maps $\varphi_1 \otimes \cdots \otimes \varphi_k$ to the functional with $v_1 \otimes \cdots \otimes v_k \mapsto \varphi_1(v_1)\, \varphi_2(v_2) \cdots \varphi_k(v_k)$. So the multilinear forms on $V_1 \times \cdots \times V_k$ correspond to the tensors in $V_1^* \otimes \cdots \otimes V_k^*$.*

2. Every multilinear form τ on $\mathbb{R}^{n_1} \times \cdots \times \mathbb{R}^{n_k}$ is given by an array $T \in \mathbb{R}^{n_1 \times \cdots \times n_k}$ as

$$\tau(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(k)}) = \sum_{i_1, \dots, i_k} T_{i_1 \dots i_k} \, \mathbf{x}_{i_1}^{(1)} \cdots \mathbf{x}_{i_k}^{(k)}, \qquad \textit{with} \quad T_{i_1 \dots i_k} = \tau(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k}).$$

Proof. (1) For fixed $\varphi_1, \dots, \varphi_k$, the map $(v_1, \dots, v_k) \mapsto \varphi_1(v_1) \cdots \varphi_k(v_k)$ is multilinear, so by the universal property, it defines a linear functional $\Lambda(\varphi_1, \dots, \varphi_k)$ on $V_1 \otimes \cdots \otimes V_k$. This functional depends multilinearly on $(\varphi_1, \dots, \varphi_k)$ (it suffices to check this on pure tensors, which span), so by the universal property again, we get a linear map $\tilde{\Lambda} : V_1^* \otimes \cdots \otimes V_k^* \to (V_1 \otimes \cdots \otimes V_k)^*$. Choose bases of all V_j and the dual bases. Then $\tilde{\Lambda}$ maps the basis tensor $b^{(1)i_1} \otimes \cdots \otimes b^{(k)i_k}$, made of dual basis vectors, to the functional that is 1 on $b^{(1)}_{i_1} \otimes \cdots \otimes b^{(k)}_{i_k}$ and 0 on all other basis tensors. So $\tilde{\Lambda}$ maps a basis to the dual basis of the basis of $V_1 \otimes \cdots \otimes V_k$, and it is an isomorphism.

(2) Writing each argument in the standard basis, $\mathbf{x}^{(j)} = \sum_i \mathbf{x}^{(j)}_i \mathbf{e}_i$, and using the linearity in each argument, we get

$$\tau(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(k)}) = \sum_{i_1, \dots, i_k} \mathbf{x}^{(1)}_{i_1} \cdots \mathbf{x}^{(k)}_{i_k} \tau(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k}). \qquad \square$$

For $k = 1$, part (2) is Proposition 1.8.3. For $k = 2$, it says that every bilinear form on $\mathbb{R}^m \times \mathbb{R}^n$ is $\beta_A(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top A \mathbf{y}$ for exactly one $m \times n$ matrix A, namely the one with $A_{ij} = \beta(\mathbf{e}_i, \mathbf{e}_j)$; this matrix is the coordinate array of a tensor in $(\mathbb{R}^m)^* \otimes (\mathbb{R}^n)^*$. Matrices also describe linear maps, and linear maps are tensors, too.

Proposition 1.11.12. *Let* $\mathrm{Hom}(V, W)$ *be the vector space of linear transformations* $V \to W$*, with the pointwise operations. The linear map* $\Psi : W \otimes V^* \to \mathrm{Hom}(V, W)$ *with*

$$\Psi(w \otimes \varphi)(v) = \varphi(v)\, w$$

is an isomorphism. If $b_1, \dots, b_n$ *and* $c_1, \dots, c_m$ *are bases of* V *and* W*, and* $T : V \to W$ *has the matrix* $A = [T]_{\mathcal{C},\mathcal{B}}$*, then* $T = \Psi\big(\sum_{i,j} A_{ij}\, c_i \otimes b^j\big)$*: the matrix of* T *is the coordinate array of the tensor* $\Psi^{-1}(T)$*.*

Proof. The map $(w, \varphi) \mapsto (v \mapsto \varphi(v)w)$ is bilinear, so Ψ exists by Theorem 1.11.6. For a tensor $t = \sum_{i,j} A_{ij}\, c_i \otimes b^j$, we get

$$\Psi(t)(b_l) = \sum_{i,j} A_{ij}\, b^j(b_l)\, c_i = \sum_i A_{il}\, c_i.$$

If A is the matrix of T, this is $T(b_l)$, since the l-th column of A is the coordinate vector of $T(b_l)$. Two linear maps that agree on a basis are equal, so $\Psi(t) = T$, and Ψ is surjective. If $\Psi(t) = 0$, then the same computation gives $\sum_i A_{il}\, c_i = 0$, so $A_{il} = 0$ for all i and l, and $t = 0$; so Ψ is injective. $\square$

For $V = \mathbb{R}^n$ and $W = \mathbb{R}^m$, with a functional φ given by a row $\mathbf{a}$, the map $\Psi(\mathbf{u} \otimes \varphi)$ is $\mathbf{x} \mapsto (\mathbf{ax})\, \mathbf{u} = (\mathbf{ua})\, \mathbf{x}$, multiplication by the rank-one matrix $\mathbf{ua}$. So one and the same $m \times n$ array can be the coordinate array of a tensor in $\mathbb{R}^m \otimes \mathbb{R}^n$, of a linear map in $\mathbb{R}^m \otimes (\mathbb{R}^n)^*$, or of a bilinear form in $(\mathbb{R}^m)^* \otimes (\mathbb{R}^n)^*$. The difference becomes visible when we change the basis (Subsection 1.11.7).

1.11.5 Contractions and Einstein notation

Definition 1.11.13 (Contraction)**.** The contraction on $V \otimes V^*$ is the linear functional $C : V \otimes V^* \to \mathbb{R}$ with $C(v \otimes \varphi) = \varphi(v)$. It exists by Theorem 1.11.6, since $(v, \varphi) \mapsto \varphi(v)$ is bilinear.

Proposition 1.11.14. *If a linear transformation* $T : V \to V$ *has the matrix* A *with respect to a basis of* V*, then* $\mathrm{tr}(A) = C(\Psi^{-1}(T))$*. In particular, the trace of the matrix of* T *is the same for all bases.*

Proof. By Proposition 1.11.12, $\Psi^{-1}(T) = \sum_{i,j} A_{ij}\, b_i \otimes b^j$, so $C(\Psi^{-1}(T)) = \sum_{i,j} A_{ij}\, b^j(b_i) = \sum_i A_{ii} = \mathrm{tr}(A)$. The second statement holds since C and Ψ are defined without choosing a basis. $\square$

Since similar matrices A and $P^{-1}AP$ are the matrices of the same linear transformation in two bases, this explains why $\operatorname{tr}(P^{-1}AP) = \operatorname{tr}(A)$, which also follows from $\operatorname{tr}(AB) = \operatorname{tr}(BA)$ (Theorem 1.6.77). Contractions are also defined for tensors with more factors, by pairing a factor V with a factor V^*. For example, the composition of linear maps is a contraction: for $S = \Psi(u \otimes \varphi) : V \to W$ and $T = \Psi(w \otimes \psi) : W \to U$, we have $(T \circ S)(v) = \varphi(v)\,\psi(u)\,w$, so

$$T \circ S = \Psi\big(\psi(u)\,w \otimes \varphi\big) :$$

the middle pair $u \in W$ and $\psi \in W^*$ is contracted to the number $\psi(u)$. In coordinates, this is the product of matrices, $(AB)_{ik} = \sum_j A_{ij}B_{jk}$, which sums over the pair of indices that belong to the middle factor. In general, a contraction sums over an index that appears twice. Since such sums appear everywhere, it is convenient to leave out the summation signs.

Definition 1.11.15 (Einstein summation convention)**.** In an expression with indices, every index that appears twice is summed over its whole range. The other indices are free, and the result is a tensor indexed by the free indices.

In this notation, the product of two matrices is $A_{ij}B_{jk}$, with the free indices i and k, the trace is A_{ii}, with no free index (a number), and the bilinear form of a matrix is $x_iA_{ij}y_j$. The NumPy function `np.einsum` implements this convention: the indices of the arguments are given as letters, and the letters after the arrow are the free indices, in the desired order (Table 1.1).

operation	Einstein notation	NumPy
matrix product AB	$A_{ij}B_{jk}$	`np.einsum("ij,jk->ik", A, B)`
trace $\operatorname{tr}(A)$	A_{ii}	`np.einsum("ii->", A)`
dot product $\mathbf{x}^\top\mathbf{y}$	x_iy_i	`np.einsum("i,i->", x, y)`
outer product $\mathbf{x} \otimes \mathbf{y}$	x_iy_j	`np.einsum("i,j->ij", x, y)`
transpose $A^\top$	A_{ji}	`np.einsum("ij->ji", A)`
bilinear form $\mathbf{x}^\top A\mathbf{y}$	$x_iA_{ij}y_j$	`np.einsum("i,ij,j->", x, A, y)`
batch of products A_bB_b	$A_{bij}B_{bjk}$	`np.einsum("bij,bjk->bik", A, B)`

Table 1.1: Operations on vectors, matrices and tensors in Einstein notation, and the corresponding calls of `np.einsum`. In the last row, A and B are tensors of order 3, which store a batch of matrices A_b and B_b.

Example 1.11.16 (Attention scores)**.** In a transformer, a sentence of L tokens is represented by a query matrix $Q \in \mathbb{R}^{L\times d}$ and a key matrix $K \in \mathbb{R}^{L\times d}$, and the attention scores $S_{ij} = \sum_c Q_{ic}K_{jc}$ measure how much token i attends to token j, i.e., $S = QK^\top$ (before a scaling by $\frac{1}{\sqrt{d}}$ and a softmax). For a batch of N sentences, Q and K are tensors of shape $N \times L \times d$, and the scores are $S_{bij} = Q_{bic}K_{bjc}$ in Einstein notation, or `np.einsum("bic,bjc->bij", Q, K)`. This contraction needs NL^2d multiplications, one for each choice of b, i, j and c.

In general, a contraction needs one multiplication for each combination of values of all the indices involved, free and summed. This is why the order of contractions matters: for $n \times n$ matrices A and B and a vector $\mathbf{x} \in \mathbb{R}^n$, both $(AB)\mathbf{x}$ and $A(B\mathbf{x})$ compute the contraction $A_{ij}B_{jk}x_k$, but the first needs $n^3 + n^2$ multiplications, and the second only $2n^2$.

Remark 1.11.17 (The Kronecker product)*.* For an $m \times n$ matrix A and a $p \times q$ matrix B, the tensor $A \otimes B$ of order 4 has the coordinates $A_{ij}B_{kl}$. Reshaping it into an $mp \times nq$ matrix, with the index pairs (i,k) as rows and (j,l) as columns, gives the Kronecker product of A and B (`np.kron` in NumPy), which is also written $A \otimes B$. It appears whenever a linear map acts on matrices: for $X \in \mathbb{R}^{n\times q}$, we have $(AXB^\top)_{ik} = A_{ij}X_{jl}B_{kl}$, so after flattening both X and $AXB^\top$ row by row, the linear map $X \mapsto AXB^\top$ is multiplication by the Kronecker product of A and B.

1.11.6 Tensor rank

Definition 1.11.18 (Tensor rank)**.** The rank of a nonzero tensor $t \in V_1 \otimes \cdots \otimes V_k$ is the smallest number r such that t is a sum of r pure tensors. The zero tensor has rank 0. A tensor of rank one is a nonzero pure tensor.

Proposition 1.11.19. *For tensors in $\mathbb{R}^m \otimes \mathbb{R}^n = \mathbb{R}^{m\times n}$, the tensor rank is the rank of Section 1.7: an $m \times n$ matrix A has rank r if and only if it is a sum of r, but not fewer, outer products $\mathbf{u} \otimes \mathbf{v} = \mathbf{u}\mathbf{v}^\top$.*

Proof. Let $\operatorname{rank}(A) = r$, and let $\mathbf{c}_1, \dots, \mathbf{c}_r$ be a basis of $\operatorname{col}(A)$. Each column $\mathbf{a}_j$ of A is a linear combination $\mathbf{a}_j = \sum_{l=1}^{r} R_{lj}\, \mathbf{c}_l$, so $A = CR$, where C is the $m \times r$ matrix with the columns $\mathbf{c}_l$, and R is the $r \times n$ matrix of the coefficients. If $\boldsymbol{\rho}_l$ denotes the l-th row of R, then

$$A = CR = \sum_{l=1}^{r} \mathbf{c}_l \boldsymbol{\rho}_l = \sum_{l=1}^{r} \mathbf{c}_l \otimes \boldsymbol{\rho}_l^\top,$$

a sum of r outer products. Conversely, if $A = \sum_{l=1}^{s} u_l \otimes v_l = \sum_{l=1}^{s} u_l v_l^\top$, then every column of A, $\mathbf{a}_j = \sum_l (v_l)_j\, u_l$, lies in $\operatorname{span}\{u_1, \dots, u_s\}$, so $\operatorname{rank}(A) = \dim \operatorname{col}(A) \le s$, since a subspace spanned by s vectors has dimension at most s (Corollary 1.3.14). □

For tensors of order 3 or more, the rank is no longer the rank of a matrix, but matrices still help to compute it.

Example 1.11.20. Let $t = \mathbf{e}_1 \otimes \mathbf{e}_1 \otimes \mathbf{e}_1 + \mathbf{e}_2 \otimes \mathbf{e}_2 \otimes \mathbf{e}_2 \in \mathbb{R}^2 \otimes \mathbb{R}^2 \otimes \mathbb{R}^2$, with the coordinates $T_{111} = T_{222} = 1$ and all other coordinates 0. So t has rank at most 2. It does not have rank one: the coordinates of a pure tensor $\mathbf{u} \otimes \mathbf{v} \otimes w$ are $u_i v_j w_k$, so its two slices $(T_{ij1})_{i,j}$ and $(T_{ij2})_{i,j}$ are the matrices $w_1\, \mathbf{u}\mathbf{v}^\top$ and $w_2\, \mathbf{u}\mathbf{v}^\top$, which are multiples of the same matrix. For our t, the slices are $\left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right)$ and $\left(\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}\right)$, which are not multiples of each other. Hence t has rank 2.

A pure tensor of order k in $\mathbb{R}^n \otimes \cdots \otimes \mathbb{R}^n$ is described by the kn coordinates of its factors instead of n^k numbers, so tensors of low rank can be stored and processed much more cheaply.

Remark 1.11.21 (Low-rank tensors in machine learning). For matrices, the rank is computed by row reduction, and the best approximation of a matrix by matrices of rank r is given by the singular value decomposition (Lecture 2). For tensors of order 3 and more, the rank is much harder to handle: computing it is NP-hard in general, and it can even depend on the field, since there are real $2 \times 2 \times 2$ tensors of rank 3 that have rank 2 as complex tensors. Nevertheless, approximating a tensor by a sum of a few pure tensors (the CP decomposition) is a standard tool to compress the weight tensors of neural networks, to find structure in data with several indices (such as users × products × time), and to learn mixture models.

1.11.7 Change of basis

Let $b_1, \dots, b_n$ and $b'_1, \dots, b'_n$ be two bases of V, related by $b'_j = \sum_i P_{ij}\, b_i$ with an invertible matrix P. (For $V = \mathbb{R}^n$ and $b_i = e_i$, P is the matrix with the columns b'_j.) Then $b_i = \sum_k (P^{-1})_{ki}\, b'_k$, and the dual bases are related by

$$b^i = \sum_l P_{il}\, b'^l,$$

since both sides take the value P_{ij} at b'_j. Substituting these formulas into the expansion of a tensor in the old basis gives its coordinates in the new basis:

- a vector $v = \sum_i x_i b_i \in V$ has the new coordinates $x' = P^{-1}x$;
- a linear transformation, $t = \sum_{i,j} A_{ij}\, b_i \otimes b^j \in V \otimes V^*$, has the new coordinates $A' = P^{-1}AP$, since $A'_{kl} = \sum_{i,j} (P^{-1})_{ki}\, A_{ij}\, P_{jl}$;

- a bilinear form, $t = \sum_{i,j} A_{ij}\, b^i \otimes b^j \in V^* \otimes V^*$, has the new coordinates $A' = P^\top A P$, since $A'_{kl} = \sum_{i,j} P_{ik}\, A_{ij}\, P_{jl}$;
- a tensor $t = \sum_{i,j} A_{ij}\, b_i \otimes b_j \in V \otimes V$ has the new coordinates $A' = P^{-1} A (P^{-1})^\top$.

In general, every index that belongs to a factor V transforms with P^{-1}, and every index that belongs to a factor V^* transforms with P. So the same array A transforms in different ways, depending on the tensor it describes. For example, the identity matrix describes both the identity map of V, i.e., the tensor $\sum_i b_i \otimes b^i \in V \otimes V^*$, and the dot product on $\mathbb{R}^n$, i.e., the tensor $\sum_i e^i \otimes e^i \in (\mathbb{R}^n)^* \otimes (\mathbb{R}^n)^*$. In every basis, the identity map is still described by $P^{-1} I_n P = I_n$, but the dot product is described by $P^\top P$, which is I_n only if P is orthogonal. Similarly, the trace is a property of linear transformations (Proposition 1.11.14), while the trace of the matrix of a bilinear form depends on the basis.

Remark 1.11.22 (Tensors in mathematics, physics and machine learning). A tensor of type (p, q) on V is an element of $V^{\otimes p} \otimes (V^*)^{\otimes q}$, the tensor product of p copies of V and q copies of V^*. Vectors are tensors of type $(1, 0)$, linear functionals of type $(0, 1)$, linear maps $V \to V$ of type $(1, 1)$, and bilinear forms of type $(0, 2)$. By Theorem 1.11.11 and the isomorphism $V \cong V^{**}$ (Theorem 1.8.18), a tensor of type (p, q) is the same thing as a multilinear form on $(V^*)^p \times V^q$, and in physics, tensors are often introduced as arrays of numbers that transform as above under a change of basis. In machine learning, the word "tensor" usually just means an array of coordinates in the standard bases, which are never changed, and the meaning of the indices is kept track of separately. The distinction still matters: the derivative of a loss with respect to the parameters is a linear functional (Section 1.8), so under a linear change of the parameters, it transforms like an index of type V^*, and not like the parameters themselves.

1.12 Key Takeaways

- A field is a set of "numbers" in which we can add, subtract, multiply, and divide by nonzero elements. $\mathbb{Q}$, $\mathbb{R}$ and $\mathbb{C}$ are fields; $\mathbb{Z}$ is not.
- A vector space over a field $\mathbb{F}$ is a set of vectors that can be added and multiplied by scalars from $\mathbb{F}$, subject to the vector space axioms. Examples: $\mathbb{F}$ itself, products $V \times W$, $\mathbb{R}^n$, the polynomials $\mathbb{R}[x]$, and spaces of functions with the pointwise operations: all functions $\mathcal{F}(S)$ on a set S, the continuous functions $C(I)$, the n times continuously differentiable functions $C^n(I)$, and the smooth functions $C^\infty(I)$.
- A basis is a linearly independent set that spans the space; every vector is a unique linear combination of the basis vectors. All bases of a finite-dimensional space have the same number of vectors, the dimension: $\dim \mathbb{R}^n = n$, while $\mathbb{R}[x]$ is infinite-dimensional.
- A linear transformation $T : V \to W$ preserves addition and scalar multiplication, and $T(0_V) = 0_W$. Its kernel is a subspace of V, and its image is a subspace of W. Differentiation $D : C^1(I) \to C(I)$ and integration $Jf(x) = \int_c^x f(t)\, dt$ are linear, with $D \circ J = \mathrm{id}$ but $J(Df) = f - f(c)$; on $C^\infty(I)$, D is surjective but not injective, which is impossible for a map of a finite-dimensional space to itself.
- A subspace is a subset that contains 0 and is closed under addition and scalar multiplication. The polynomials and the function spaces form a chain of subspaces $\mathbb{R}[x] \subseteq C^\infty(\mathbb{R}) \subseteq C^n(\mathbb{R}) \subseteq C(\mathbb{R}) \subseteq \mathcal{F}(\mathbb{R})$, all of them infinite-dimensional.
- An isomorphism is a bijective linear transformation, i.e., one with $\ker(T) = \{0\}$ and $\mathrm{im}(T) = W$. Via coordinates, every n-dimensional real vector space is isomorphic to $\mathbb{R}^n$, and $V \cong W$ if and only if $\dim V = \dim W$, if and only if some linear transformation $V \to W$ is represented by an invertible matrix.

- Every linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$ is multiplication by an $m \times n$ matrix A, whose i-th column is $T(\mathbf{e}_i)$.

- Matrices of the same size are added entry by entry. The product of an $m \times n$ and an $n \times p$ matrix is the $m \times p$ matrix with $(AB)_{ij} = \sum_k A_{ik}B_{kj}$; it represents the composition $T_A \circ T_B$. Matrix multiplication is associative but not commutative, and $(AB)^\top = B^\top A^\top$.

- A square matrix A is invertible if $AB = BA = I_n$ for some matrix $B = A^{-1}$; then $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$, and $(AB)^{-1} = B^{-1}A^{-1}$. The inverse is computed without determinants by row reducing $(A \mid I_n)$ to $(I_n \mid A^{-1})$.

- Diagonal, upper and lower triangular, symmetric ($A^\top = A$) and skew-symmetric ($A^\top = -A$) matrices form subspaces of $\mathbb{R}^{n\times n}$. The invertible matrices $\mathrm{GL}(n, \mathbb{R})$ and the orthogonal matrices $\mathrm{O}(n)$ (with $Q^\top Q = I_n$, so $Q^{-1} = Q^\top$) do not form subspaces, but they contain I_n and are closed under multiplication and inverses: they are matrix groups, i.e., subgroups of the group $\mathrm{GL}(n, \mathbb{R})$. A group is a set with an associative operation, an identity element and inverses; examples are $(\mathbb{R}, +)$, every vector space under addition, and $(\mathbb{F} \setminus \{0\}, \cdot)$ for a field $\mathbb{F}$.

- The trace $\mathrm{tr}(A) = \sum_i A_{ii}$ is linear and satisfies $\mathrm{tr}(AB) = \mathrm{tr}(BA)$. The determinant is defined by cofactor expansion and is best computed by row reduction to triangular form; it satisfies $\det(AB) = \det(A)\det(B)$ and $\det(A^\top) = \det(A)$, and $|\det(A)|$ is the factor by which T_A scales areas and volumes.

- Block matrices multiply like 2×2 matrices. With the Schur complement $S = D - CA^{-1}B$, $\det\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A)\det(S)$. Gaussian elimination computes the LU decomposition $A = LU$ (or $PA = LU$ with row swaps), which solves $A\mathbf{x} = \mathbf{b}$ by forward and back substitution and gives $\det(A)$ as a product of pivots. Low-rank updates are inverted with the Woodbury identity, for example $(A + \mathbf{u}\mathbf{v}^\top)^{-1} = A^{-1} - \frac{A^{-1}\mathbf{u}\mathbf{v}^\top A^{-1}}{1+\mathbf{v}^\top A^{-1}\mathbf{u}}$, and $\det(A + \mathbf{u}\mathbf{v}^\top) = (1 + \mathbf{v}^\top A^{-1}\mathbf{u})\det(A)$.

- Elementary row operations bring a matrix into (reduced) row echelon form. They do not change the null space or the row space, but they can change the column space; the pivot columns of the original matrix form a basis of its column space.

- $\mathrm{rank}(A) = \dim \mathrm{col}(A) = \dim \mathrm{row}(A)$ is the number of pivots, and $\mathrm{rank}(A) + \dim \mathrm{null}(A) = n$ for every $m \times n$ matrix A.

- For an $n \times n$ matrix A: A is invertible $\iff$ $\mathrm{null}(A) = \{\mathbf{0}\}$ $\iff$ $\mathrm{rank}(A) = n$ $\iff$ $\mathrm{RREF}(A) = I_n$ $\iff$ $\det(A) \neq 0$.

- The dual space V^* consists of the linear functionals $V \to \mathbb{F}$; on $\mathbb{R}^n$, these are the row vectors. A basis $b_1, \ldots, b_n$ has a dual basis $b^1, \ldots, b^n$ with $b^i(b_j) = \delta_{ij}$ (in $\mathbb{R}^n$: the rows of P^{-1}), and $\dim V^* = \dim V$. The dual map $T^*(\psi) = \psi \circ T$ is represented by the transpose $A^\top$. A system $Ax = b$ is solvable if and only if $yb = 0$ for all rows y with $yA = 0$.

- The quotient space V/U consists of the cosets $v + U$, where $v + U = w + U$ if and only if $v - w \in U$; it has dimension $\dim V - \dim U$. First isomorphism theorem: $V/\ker(T) \cong \mathrm{im}(T)$, which gives rank–nullity for every linear map. Second: $(U + W)/W \cong U/(U \cap W)$, so $\dim(U + W) + \dim(U \cap W) = \dim U + \dim W$. Third: $(V/U)/(W/U) \cong V/W$.

- An eigenvector $\mathbf{v} \neq \mathbf{0}$ of A satisfies $A\mathbf{v} = \lambda\mathbf{v}$; the eigenvalues are the roots of the characteristic polynomial $\det(A - \lambda I_n)$, and the eigenspace $E_\lambda = \mathrm{null}(A - \lambda I_n)$ is found by row reduction. The (complex) eigenvalues add up to $\mathrm{tr}(A)$ and multiply to $\det(A)$.

- A is diagonalizable, $A = PDP^{-1}$, exactly when it has n linearly independent eigenvectors (the columns of P); this holds, for example, when A has n distinct eigenvalues. Then $A^k = PD^kP^{-1}$, which explains Fibonacci numbers, the long-run behavior of Markov chains, and power iteration (PageRank).
- Cayley–Hamilton: $p_A(A) = 0$, so every power of A, and A^{-1} if it exists, is a polynomial in A of degree less than n. The minimal polynomial m_A divides every polynomial that annihilates A, its real roots are the eigenvalues, and A is diagonalizable if and only if m_A is a product of distinct real linear factors.
- A cyclic convolution $\mathbf{c} * \mathbf{x}$ is multiplication by the circulant matrix $\operatorname{circ}(\mathbf{c})$; the matrices that commute with the cyclic shift are exactly the circulant ones. The Fourier vectors $\mathbf{f}_m$ are eigenvectors of every circulant matrix, with the eigenvalues $\hat{c}_m$, so $\widehat{\mathbf{c} * \mathbf{x}}_m = \hat{c}_m \hat{x}_m$, and the FFT computes convolutions in $O(n \log n)$ operations.
- The tensor product $V \otimes W$ is the quotient of the free vector space on $V \times W$ by the rules of bilinearity; $(v, w) \mapsto v \otimes w$ is bilinear, and bilinear maps $V \times W \to U$ are the same as linear maps $V \otimes W \to U$ (universal property). If b_i and c_j are bases, the pure tensors $b_i \otimes c_j$ form a basis, so $\dim(V \otimes W) = \dim V \cdot \dim W$. A tensor is an element of $V_1 \otimes \cdots \otimes V_k$; its coordinates form an array, the "tensor" of deep learning libraries, and $\mathbb{R}^m \otimes \mathbb{R}^n = \mathbb{R}^{m \times n}$ with $u \otimes v = uv^\top$.
- Multilinear forms are tensors in $V_1^* \otimes \cdots \otimes V_k^*$, and linear maps $V \to W$ are tensors in $W \otimes V^*$; the contraction $v \otimes \varphi \mapsto \varphi(v)$ is the trace. Under a change of basis, indices of type V transform with P^{-1} and indices of type V^* with P: linear maps change to $P^{-1}AP$, bilinear forms to $P^\top AP$.
- The rank of a tensor is the smallest number of pure tensors with sum t; for matrices, it is the usual rank. Contractions such as $A_{ij}B_{jk}$ in Einstein notation, computed with `np.einsum`, describe matrix products, traces, batched products and attention scores.

1.13 Exercises

1. Using only the vector space axioms, prove that in every vector space V over a field $\mathbb{F}$:
   (a) $0 \cdot v = 0_V$ for every $v \in V$. (Hint: $0 \cdot v = (0 + 0) \cdot v$.)
   (b) $(-1) \cdot v = -v$ for every $v \in V$. (Hint: compute $v + (-1) \cdot v$.)
   (c) The axiom $u + v = v + u$ is redundant: it follows from the other axioms. (Hint: first show that $(-v) + v = 0$ and $0 + v = v$, using only associativity and axioms 2 and 3; for the first, add $w = (-v) + v$ to itself. Then expand $(1 + 1)(u + v)$ in two ways, and cancel u on the left and v on the right.)
2. Which of the following subsets of $\mathbb{R}^2$ are subspaces? Justify your answers.
   (a) $\{(x, y) \in \mathbb{R}^2 : y = 2x\}$;
   (b) $\{(x, y) \in \mathbb{R}^2 : y = 2x + 1\}$;
   (c) $\{(x, y) \in \mathbb{R}^2 : xy = 0\}$.
3. Show that $\{(1, 2), (3, 4)\}$ is a basis of $\mathbb{R}^2$, and write $(5, 6)$ as a linear combination of these two vectors.
4. Let $\mathbb{R}[x]_{\leq 2}$ be the set of polynomials with real coefficients of degree at most 2. Show that it is a subspace of $\mathbb{R}[x]$, and find its dimension.

5. Let $T : \mathbb{R}^3 \to \mathbb{R}^3$ be defined by $T(x, y, z) = (x + y, y + z, x + 2y + z)$. Find the matrix A of T and $\mathrm{RREF}(A)$. Then find a basis of $\mathrm{null}(A)$, a basis of $\mathrm{col}(A)$, and $\mathrm{rank}(A)$, and check that your answers agree with the rank–nullity theorem.

6. Write down explicit isomorphisms $\mathbb{R}^{2\times 2} \to \mathbb{R}^4$ and $\mathrm{Sym}_2 \to \mathbb{R}^3$, and prove that they are isomorphisms. Are $\mathbb{R}^{2\times 2}$ and Sym_2 isomorphic?

7. Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Compute AB and BA. Describe in words what multiplying a 2×2 matrix by B from the left does, and what multiplying it by B from the right does.

8. Use the Gauss–Jordan method to find the inverse of
$$A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 5 & 6 & 0 \end{pmatrix},$$
and check your answer.

9. Show that the matrix F from Example 1.6.31 is not invertible, and find a nonzero vector $\mathbf{x}$ with $F\mathbf{x} = \mathbf{0}$.

10. Let A be an $n \times n$ matrix with $A^2 = A$ (where $A^2 = AA$) and $A \neq I_n$. Show that A is not invertible.

11. Show that an upper triangular matrix is invertible if and only if all of its diagonal entries are nonzero, in two ways: using Theorem 1.6.47, and using determinants.

12. Compute the determinant of
$$A = \begin{pmatrix} 2 & 0 & 1 \\ 1 & 3 & 2 \\ 1 & 1 & 1 \end{pmatrix}$$
in two ways: by cofactor expansion, and by row reduction. Is A invertible?

13. Let A be a skew-symmetric $n \times n$ matrix, where n is odd. Show that $\det(A) = 0$. (Hint: compute $\det(A^\top)$ in two ways.)

14. Show that $\mathrm{tr}(ABC) = \mathrm{tr}(CAB)$ for all $n \times n$ matrices A, B, C, but find 2×2 matrices with $\mathrm{tr}(ABC) \neq \mathrm{tr}(ACB)$.

15. Let $A = \begin{pmatrix} 2 & 4 & -2 \\ 4 & 9 & -3 \\ -2 & -3 & 7 \end{pmatrix}$.
    (a) Find the LU decomposition of A, and use it to compute $\det(A)$ and to solve $A\mathbf{x} = (2, 8, 10)^\top$.
    (b) Let D be the diagonal matrix with the diagonal entries of U. Check that $U = DL^\top$, so that $A = LDL^\top$. Show that every invertible symmetric matrix that has an LU decomposition can be written as $A = LDL^\top$ in this way. (Hint: uniqueness.)

16. Show that $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ has no LU decomposition, and find a permutation matrix P such that PA has one.

17. Let A be an invertible $k \times k$ matrix and D an invertible $m \times m$ matrix. Show that $\begin{pmatrix} A & B \\ 0 & D \end{pmatrix}^{-1} = \begin{pmatrix} A^{-1} & -A^{-1}BD^{-1} \\ 0 & D^{-1} \end{pmatrix}$, and that $\det \begin{pmatrix} I_k & B \\ B^\top & I_m \end{pmatrix} = \det(I_m - B^\top B)$ for every $k \times m$ matrix B.

18. Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$. Show that $I_n + \mathbf{u}\mathbf{v}^\top$ is invertible if and only if $\mathbf{v}^\top \mathbf{u} \neq -1$, and find its inverse. Show that every nonzero vector $\mathbf{x}$ with $\mathbf{v}^\top \mathbf{x} = 0$ is an eigenvector with the eigenvalue 1, and that $\mathbf{u}$ (if $\mathbf{u} \neq \mathbf{0}$) is an eigenvector with the eigenvalue $1 + \mathbf{v}^\top \mathbf{u}$.

19. Find the eigenvalues and eigenspaces of $A = \begin{pmatrix} 4 & 1 \\ 2 & 3 \end{pmatrix}$. Diagonalize A, and use the diagonalization to find a formula for A^k.

20. Show that $A = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$ is not diagonalizable. Find all of its eigenvectors.

21. Find the complex eigenvalues of the rotation by 90°, $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, and check Theorem 1.10.12 for this matrix.

22. Let A be a 3×3 matrix with eigenvalues 1, 2 and 3. Compute $\operatorname{tr}(A^2)$, $\det(A)$ and $\det(A^{-1})$. (Hint: A is diagonalizable.)

23. Let A be an $n \times n$ matrix and λ an eigenvalue of A.

    (a) Show that if $A^2 = A$, then $\lambda = 0$ or $\lambda = 1$.

    (b) Show that if $A^k = 0$ for some $k \geq 1$, then $\lambda = 0$.

    (c) Show that if A is orthogonal, then $\lambda = 1$ or $\lambda = -1$.

24. A Markov chain has the transition matrix $A = \begin{pmatrix} 0.8 & 0.3 \\ 0.2 & 0.7 \end{pmatrix}$. Find its eigenvalues and its stationary distribution, and describe what happens to $A^k \mathbf{x}_0$ as $k \to \infty$ for a probability vector $\mathbf{x}_0$.

25. Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$. Use the Cayley–Hamilton theorem to write A^{-1} and A^5 in the form $aA + bI_2$, and check your answers by multiplying out.

26. Find the minimal polynomials of $\operatorname{diag}(2, 2, 3)$ and of $\begin{pmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$, and decide which of the two matrices is diagonalizable. Show that every matrix A with $A^3 = A$ is diagonalizable, and that its eigenvalues lie in $\{-1, 0, 1\}$.

27. (Circulant matrices.) Let $\mathbf{c}, \mathbf{d} \in \mathbb{R}^n$.

    (a) Show that $\operatorname{circ}(\mathbf{c}) \operatorname{circ}(\mathbf{d}) = \operatorname{circ}(\mathbf{c} * \mathbf{d})$, and conclude that $\mathbf{c} * \mathbf{d} = \mathbf{d} * \mathbf{c}$. (Hint: Proposition 1.10.37 and $S^n = I_n$.)

    (b) Find the eigenvalues of the difference filter, $(\mathbf{c} * \mathbf{x})_j = x_j - x_{j-1}$. Which signals does it map to $\mathbf{0}$?

    (c) Show that $\operatorname{circ}(\mathbf{c})$ is invertible if and only if $\hat{c}_m \neq 0$ for all m, and that its inverse is again circulant.

28. Find the dual basis of the basis $(1,0,0)^\top$, $(1,1,0)^\top$, $(1,1,1)^\top$ of $\mathbb{R}^3$, and check that $b^i(\mathbf{b}_j) = \delta_{ij}$.

29. On $V = \mathbb{R}[x]_{\leq 2}$, consider the functionals $\varphi_1(p) = p(0)$, $\varphi_2(p) = p'(0)$ and $\varphi_3(p) = \int_{-1}^{1} p(x)\,dx$. Show that $\varphi_1, \varphi_2, \varphi_3$ is a basis of V^*, and find the basis q_1, q_2, q_3 of V whose dual basis it is, i.e., with $\varphi_i(q_j) = \delta_{ij}$.

30. Let $T : \mathbb{R}^2 \to \mathbb{R}^2$, $T(x,y) = (x + 2y, 3y)$. Compute $T^*(\psi)$ for the coordinate functionals $\psi(x,y) = x$ and $\psi(x,y) = y$, and check that the matrix of T^* with respect to the dual of the standard basis is the transpose of the matrix of T.

31. Let φ and ψ be nonzero linear functionals on a finite-dimensional vector space with $\ker(\varphi) = \ker(\psi)$. Show that $\psi = c\varphi$ for some $c \neq 0$. (Hint: choose v with $\varphi(v) = 1$, and show that every vector is of the form $w + tv$ with $w \in \ker(\varphi)$.)

32. Let $U = \operatorname{span}\{(1,0,1)^\top, (0,1,1)^\top\} \subseteq \mathbb{R}^3$. Show that the cosets of U are the planes $x_1 + x_2 - x_3 = c$, and that $\mathbf{x} + U \mapsto x_1 + x_2 - x_3$ is an isomorphism $\mathbb{R}^3/U \to \mathbb{R}$. Check Theorem 1.9.8.

33. Use the first isomorphism theorem and the map $A \mapsto \frac{1}{2}(A + A^\top)$ to show that $\mathbb{R}^{n\times n}/\mathrm{Skew}_n \cong \mathrm{Sym}_n$, and check that the dimensions agree.

34. Let U and W be 3-dimensional subspaces of $\mathbb{R}^5$. Show that $U \cap W \neq \{0\}$. Which values can $\dim(U \cap W)$ take?

35. Show that $V/U = \{0\}$ if and only if $U = V$, and that $V/\{0\} \cong V$.

36. For $n \geq 0$, let $f_n(x) = x^n|x|$.

    (a) Show that for $n \geq 1$, f_n is differentiable on $\mathbb{R}$, with $f_n' = (n+1)f_{n-1}$. (Hint: $f_n(x) = x^{n+1}$ for $x \geq 0$, and $f_n(x) = -x^{n+1}$ for $x < 0$; at $x = 0$, use the definition of the derivative.)

    (b) Conclude that $f_n \in C^n(\mathbb{R})$, but $f_n \notin C^{n+1}(\mathbb{R})$, so that $C^{n+1}(\mathbb{R}) \neq C^n(\mathbb{R})$.

37. Let $D : C^\infty(\mathbb{R}) \to C^\infty(\mathbb{R})$, $Df = f'$, and $M : C^\infty(\mathbb{R}) \to C^\infty(\mathbb{R})$, $(Mf)(x) = xf(x)$.

    (a) Show that M is linear, and that $D \circ M - M \circ D = \mathrm{id}$.

    (b) Show that there are no $n \times n$ matrices A and B with $AB - BA = I_n$. (Hint: take the trace.) So the identity in (a) has no analogue for matrices.

38. Let $D : \mathbb{R}[x]_{\leq 3} \to \mathbb{R}[x]_{\leq 2}$, $Dp = p'$, and $J : \mathbb{R}[x]_{\leq 2} \to \mathbb{R}[x]_{\leq 3}$, $(Jp)(x) = \int_0^x p(t)\,dt$. Find the matrices of D and J with respect to the bases of monomials, and compute the products of these two matrices in both orders. Which product is the identity matrix? Why can the other product not be the identity?

39. Which of the following maps $\mathbb{R}^2 \times \mathbb{R}^2 \to \mathbb{R}$ are bilinear? For the bilinear ones, find the matrix A with $\beta(\mathbf{x},\mathbf{y}) = \mathbf{x}^\top A\mathbf{y}$. (a) $\beta(\mathbf{x},\mathbf{y}) = x_1y_2 - x_2y_1$; (b) $\beta(\mathbf{x},\mathbf{y}) = x_1 + y_1$; (c) $\beta(\mathbf{x},\mathbf{y}) = (x_1 + x_2)(y_1 - 3y_2)$; (d) $\beta(\mathbf{x},\mathbf{y}) = x_1x_2y_1$.

40. Let $\beta(\mathbf{x},\mathbf{y}) = \mathbf{x}^\top A\mathbf{y}$ be a bilinear form on $\mathbb{R}^n$.

    (a) Show that $\beta(\mathbf{x},\mathbf{y}) = \beta(\mathbf{y},\mathbf{x})$ for all $\mathbf{x},\mathbf{y}$ if and only if A is symmetric.

    (b) Let P be invertible. Show that the matrix $P^\top AP$ of β in the basis of the columns of P is symmetric if A is, and that $\det(P^\top AP) = \det(P)^2 \det(A)$. So the sign of the determinant of the matrix of a bilinear form does not depend on the basis, but its value does.

41. (a) Let $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$. Show that $(\mathbf{u} \otimes \mathbf{v})\,\mathbf{y} = (\mathbf{v}^\top\mathbf{y})\,\mathbf{u}$ for every $\mathbf{y} \in \mathbb{R}^n$, and that $\mathrm{im}(T_{\mathbf{u}\otimes\mathbf{v}}) = \operatorname{span}\{\mathbf{u}\}$ if $\mathbf{v} \neq \mathbf{0}$.

(b) Write the $2 \times 2 \times 2$ tensor with the entries $T_{ijk} = i + j + k$ as a sum of three rank-one tensors. (Hint: $i+j+k = i \cdot 1 \cdot 1 + 1 \cdot j \cdot 1 + 1 \cdot 1 \cdot k$.) Show that T does not have rank one.

42. Write the following in Einstein notation and as a call of `np.einsum`, and count the multiplications: (a) the vector $A\mathbf{x}$ for $A \in \mathbb{R}^{m \times n}$; (b) $\operatorname{tr}(A^\top B)$ for $A, B \in \mathbb{R}^{m \times n}$; (c) the matrix $XX^\top$ of all dot products of the rows of $X \in \mathbb{R}^{N \times d}$; (d) the number $\mathbf{x}^\top AB\mathbf{y}$ for $n \times n$ matrices A and B, computed as $(\mathbf{x}^\top A)(B\mathbf{y})$ and as $\mathbf{x}^\top (AB)\mathbf{y}$.

43. Let V and W be vector spaces.

    (a) Show that $(2v) \otimes w = v \otimes (2w)$ and $v \otimes 0 = 0$ in $V \otimes W$.

    (b) Use the universal property to construct linear maps $V \otimes W \to W \otimes V$ with $v \otimes w \mapsto w \otimes v$, and $W \otimes V \to V \otimes W$ with $w \otimes v \mapsto v \otimes w$, and show that they are inverse to each other. So $V \otimes W \cong W \otimes V$.

    (c) Show that $\mathbb{R} \otimes V \cong V$, via $a \otimes v \mapsto av$.

44. Let $b_1, \ldots, b_n$ be a basis of V, with the dual basis $b^1, \ldots, b^n$. Show that the tensor $\sum_{i=1}^n b_i \otimes b^i \in V \otimes V^*$ corresponds to the identity map of V under the isomorphism Ψ of Proposition 1.11.12. Conclude that this tensor does not depend on the choice of the basis, and that its contraction is $\dim V$.

1.14 Programming Exercises

In these exercises, you implement the main algorithms of this lecture in Python. You need Python 3 with NumPy, and Matplotlib for Exercises P6 and P12. Implement the algorithms yourself, using only loops, indexing and arithmetic on NumPy arrays; use NumPy's built-in linear algebra routines (`np.linalg`) only to check your results. Start each file with

```
import numpy as np
rng = np.random.default_rng(0)  # random number generator for test matrices
```

A word on floating-point numbers. Computers store real numbers with only about 16 significant digits, so almost every computation has tiny rounding errors; for example, `0.1 + 0.2 == 0.3` is `False` in Python. Therefore, never test whether a computed number is exactly zero: test `abs(x) < tol` for a small tolerance such as `tol = 1e-10` instead, and compare arrays with `np.allclose(X, Y)`. Also remember that Python counts from 0: the entry A_{ij} of these notes is `A[i-1, j-1]` in NumPy.

P1. Matrices of linear transformations. Write a function that takes a linear transformation $T : \mathbb{R}^n \to \mathbb{R}^m$, given as a Python function that maps NumPy vectors of length n to vectors of length m, and returns its $m \times n$ matrix, whose i-th column is $T(\mathbf{e}_i)$ (Theorem 1.6.9).

```
def matrix_of(T, n):
    """Return the matrix of the linear transformation T: R^n -> R^m."""
    ...

T = lambda v: np.array([2*v[0] + 3*v[1], 4*v[0] - v[1]])
print(matrix_of(T, 2))  # should print [[ 2.  3.] [ 4. -1.]]
```

- Also test your function on $T(x, y, z) = (x + 2y - z, 3x - y + 4z)$, and compare with the matrix computed in the examples after Theorem 1.6.9.

- Write a function `looks_linear(T, n, trials=100)` that checks additivity and homogeneity for random vectors `u = rng.normal(size=n)`, `v = rng.normal(size=n)` and random scalars `a = rng.normal()`. Test it on the T above and on the maps $(x, y) \mapsto (x + 1, y)$ and $(x, y) \mapsto (xy, y)$.
- Why can such a random test prove that a map is *not* linear, but never prove that a map *is* linear?

P2. Matrix multiplication from scratch. Implement `matmul(A, B)` with three nested loops, directly from the definition $(AB)_{ij} = \sum_k A_{ik}B_{kj}$. Raise a `ValueError` if the number of columns of `A` is not equal to the number of rows of `B`.

- Compare `matmul(A, B)` with NumPy's `A @ B` for random matrices of several shapes.
- Use your function to check numerically that $(AB)C = A(BC)$ and $(AB)^\top = B^\top A^\top$ (Proposition 1.6.22), and find random 2×2 matrices with $AB \neq BA$.
- Measure the running times (with `time.perf_counter()`) of `matmul` and of `@` for random $n \times n$ matrices with $n = 50, 100, 200$. By roughly what factor does the running time of `matmul` grow when n doubles? Explain this by counting the multiplications that are performed. (NumPy is much faster because it runs highly optimized compiled code.)

P3. Row reduction. Implement a function that computes the reduced row echelon form of a matrix and its pivot columns.

```
def rref(A, tol=1e-10):
    """Return (R, pivots): the RREF of A and the list of pivot columns."""
    R = np.array(A, dtype=float)  # work on a copy of A
    ...
```

Go through the columns from left to right, keeping track of the row r in which the next pivot will be placed. For each column c, find the entry with the largest absolute value among the rows $r, r+1, \ldots$ of that column. If it is smaller than `tol`, the column has no pivot. Otherwise, swap its row with row r, divide row r by the pivot, and subtract multiples of row r from all the other rows so that all other entries of column c become zero. (Choosing the largest available entry as the pivot is called partial pivoting; it reduces rounding errors.)

- Test your function: the matrix $\begin{pmatrix} 1 & 2 & -1 \\ 3 & -1 & 4 \end{pmatrix}$ should give $\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix}$ with pivots `[0, 1]`, and the matrix F of Example 1.6.31 should give $\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$.
- Using `rref`, write the functions `rank(A)` and `null_space_basis(A)`, which returns one basis vector for each free variable as in the proof of the rank–nullity theorem, and `col_space_basis(A)`, which returns the pivot columns of the *original* matrix (Theorem 1.7.2).
- For 20 random 5×7 matrices A, check that `A @ v` is close to zero for every vector `v` in `null_space_basis(A)`, that `rank(A) + len(null_space_basis(A)) == 7` (rank–nullity), and that `rank(A)` agrees with `np.linalg.matrix_rank(A)`.
- Random matrices almost always have the largest possible rank. Construct a 6×8 matrix of rank 4 as the product of a random 6×4 matrix and a random 4×8 matrix, and check that your functions find rank 4 and a 4-dimensional null space.

P4. The inverse with the Gauss–Jordan method. Implement `inverse(A)`: row reduce the $n \times 2n$ matrix $(A \mid I_n)$ with your `rref` (use `np.hstack` and `np.eye`) and return the right block. If the left block is not I_n, raise `np.linalg.LinAlgError("matrix is singular")` (Theorem 1.6.49).

- Check your function on the examples of Subsection 1.6.3: the inverse of $\begin{pmatrix} 2 & 3 \\ 4 & -1 \end{pmatrix}$ is $\frac{1}{14}\begin{pmatrix} 1 & 3 \\ 4 & -2 \end{pmatrix}$, the inverse of $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ is $\begin{pmatrix} 0 & -1 & 1 \\ 1 & 1 & -1 \\ -1 & 0 & 1 \end{pmatrix}$, and $\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$ is singular.
- For a random 10×10 matrix A, check that `A @ inverse(A)` and `inverse(A) @ A` are both close to I_{10}, and that `inverse(A)` is close to `np.linalg.inv(A)`.

P5. Determinants in two ways. Implement `det_cofactor(A)`, which computes the determinant recursively by cofactor expansion along the first row (Subsection 1.6.5), and `det_rowreduce(A)`, which brings A into upper triangular form by row operations, keeps track of the sign changes caused by row swaps, and multiplies the diagonal entries (Theorem 1.6.84 and Corollary 1.6.86). Hint: `np.delete(A[1:], j, axis=1)` is the matrix obtained from `A` by deleting its first row and the column with index `j`.

- Both functions should return -3 for $\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 10 \end{pmatrix}$, and 0 (up to rounding) for the matrix F of Example 1.6.31. Compare both functions with `np.linalg.det` on random matrices.
- Check the product rule $\det(AB) = \det(A)\det(B)$ and the rule $\det(A^\top) = \det(A)$ on random 5×5 matrices.
- Measure the running time of `det_cofactor` for random $n \times n$ matrices with $n = 2, 3, \ldots, 8$. By roughly what factor does it grow from $n-1$ to n, and why? Estimate how long `det_cofactor` would need for $n = 20$, and compare with the running time of `det_rowreduce` for $n = 200$.

P6. Visualizing linear transformations. Using Matplotlib, write a function that draws the unit square, a few grid lines inside it and the letter F, before and after applying a 2×2 matrix A, as in Figure 1.2.

```
def plot_transformation(A):
    """Draw the unit square, a grid and the letter F, before and after A."""
    ...
```

Store a shape as an array `P` with one point per row; then its image under A is `P @ A.T`. The letter F in the figures has the corners

$$(0.2, 0.1),\ (0.4, 0.1),\ (0.4, 0.45),\ (0.65, 0.45),\ (0.65, 0.6),$$
$$(0.4, 0.6),\ (0.4, 0.75),\ (0.8, 0.75),\ (0.8, 0.9),\ (0.2, 0.9).$$

- Reproduce the four panels of Figure 1.2, and also plot the singular matrix $\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$.
- The signed area of a polygon with corners $(x_1, y_1), \ldots, (x_k, y_k)$ is given by the shoelace formula

$$\frac{1}{2}\sum_{i=1}^{k}(x_i y_{i+1} - x_{i+1} y_i), \qquad \text{where } (x_{k+1}, y_{k+1}) = (x_1, y_1).$$

It is positive if the corners are listed counterclockwise, and negative if they are listed clockwise. Implement it, apply it to the image of the unit square with corners listed in the order $(0,0), (1,0), (1,1), (0,1)$, and check that the result equals $\det(A)$, including the sign, for all the matrices above and for the matrix $\begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}$ of Figure 1.4.

P7. Special matrices.

- Write functions `is_symmetric(A, tol=1e-10)` and `is_orthogonal(Q, tol=1e-10)`.
- For a random 3×5 matrix A and a random 5×3 matrix B, check that `A.T @ A` is symmetric, that $\operatorname{tr}(AB) = \operatorname{tr}(BA)$ (even though AB and BA have different sizes!), and that $\operatorname{tr}(A^\top A)$ equals `np.linalg.norm(A, "fro")**2`.
- Write a function that splits a square matrix into its symmetric and skew-symmetric parts (Exercise 1.6.66), and test it.
- The command `Q, R = np.linalg.qr(rng.normal(size=(4, 4)))` produces a random orthogonal matrix `Q`. Check that `Q` is orthogonal, that $\det(Q) = \pm 1$ (Corollary 1.6.93), and that Q preserves lengths: `np.linalg.norm(Q @ x)` equals `np.linalg.norm(x)` for random vectors $\mathbf{x}$, where the length of $\mathbf{x}$ is $\sqrt{x_1^2 + \cdots + x_n^2}$.

P8. Why you should not compute the inverse. The Hilbert matrix H_n is the $n \times n$ matrix with entries $(H_n)_{ij} = \frac{1}{i+j-1}$. It is invertible, but very close to being singular, which makes computations with it very sensitive to rounding errors. For $n = 4, 6, 8, 10, 12$, let `x_true = np.ones(n)` and `b = H @ x_true`, and solve $H_n \mathbf{x} = \mathbf{b}$ in three ways:

```
x1 = np.linalg.solve(H, b)
x2 = np.linalg.inv(H) @ b
x3 = inverse(H) @ b  # your function from P4
```

For each method, print the error $\max_i |x_i - 1|$ and the residual $\max_i |(H_n\mathbf{x} - \mathbf{b})_i|$. What do you observe? Which method keeps the residual smallest? What happens with your own `inverse` for larger n, and why? (Hint: think about the tolerance in `rref`.) Relate your observations to Remark 1.6.53.

P9. Mini-project: fitting a line to data. Given data points $(x_1, y_1), \ldots, (x_N, y_N)$, we look for the line $y = w_0 + w_1 x$ that fits the data best, in the sense that the sum of squared errors $\sum_i (y_i - w_0 - w_1 x_i)^2$ is as small as possible. This is the simplest example of linear regression. Let X be the $N \times 2$ matrix with rows $(1, x_i)$, and let $\mathbf{y} = (y_1, \ldots, y_N)^\top$. One can show (we do not prove this here) that the best $\mathbf{w} = (w_0, w_1)^\top$ is the solution of the normal equations

$$(X^\top X)\,\mathbf{w} = X^\top \mathbf{y}.$$

- Generate data on a line with some noise:

```
x = np.linspace(0, 1, 50)
y = 2 + 3 * x + rng.normal(0, 0.1, size=50)  # a line plus noise
X = np.column_stack([np.ones_like(x), x])
```

- Compute $\mathbf{w}$ using your own functions from P2 and P4, and check that $\mathbf{w}$ is close to $(2, 3)$. Compare with `np.linalg.lstsq(X, y, rcond=None)[0]`.
- Explain why $X^\top X$ is symmetric. Show that if the columns of X are linearly independent, then $X^\top X$ is invertible. (Hint: if $X^\top X\mathbf{w} = 0$, compute $\mathbf{w}^\top X^\top X \mathbf{w}$ and use Corollary 1.7.11.)
- Fit a parabola $y = w_0 + w_1 x + w_2 x^2$ to the data $y = 1 - 2x + 4x^2 + \text{noise}$, using the columns $1, x, x^2$. How is this related to the vector space $\mathbb{R}[x]_{\leq 2}$?

P10. Power iteration and PageRank. Implement power iteration:

```
def power_iteration(A, num_iter=100):
    """Approximate an eigenvector for the eigenvalue of largest absolute value."""
    x = rng.normal(size=A.shape[0])
```

```
    for _ in range(num_iter):
        x = A @ x
        x = x / np.abs(x).max()  # rescale to avoid overflow
    ...
```

and estimate the eigenvalue from the final vector $\mathbf{x}$, for instance as $(A\mathbf{x})_i / x_i$ for an index i with $x_i \neq 0$.

- Test it on the matrices of Examples 1.10.7 and 1.10.8, and compare with `np.linalg.eig`. How does the speed of convergence depend on the ratio $|\lambda_2/\lambda_1|$ of the two largest eigenvalues (in absolute value)?
- What happens for the rotation by $90°$, and for $\mathrm{diag}(1, -1)$? Explain.
- A tiny web has 4 pages, with links $1 \to 2$, $1 \to 3$, $2 \to 3$, $3 \to 1$, $4 \to 3$. Let L be the 4×4 matrix with $L_{ij} = \frac{1}{\text{number of links on page } j}$ if page j links to page i, and 0 otherwise. The PageRank vector is the stationary distribution of $G = 0.85\, L + \frac{0.15}{4}\, \mathbf{1}\mathbf{1}^\top$, where $\mathbf{1}\mathbf{1}^\top$ is the matrix of all ones. Compute it with power iteration, check that it is an eigenvector of G for the eigenvalue 1, and rank the pages.

P11. Diagonalization in floating-point arithmetic. The function `np.linalg.eig(A)` returns the eigenvalues and a matrix `P` whose columns are eigenvectors.

- For a random 5×5 matrix A, check that A is close to PDP^{-1}. (The eigenvalues may be complex.) Compare $PD^{20}P^{-1}$ with `np.linalg.matrix_power(A, 20)`.
- Compute the Fibonacci numbers F_n in two ways: exactly, with Python integers and the recursion, and with Binet's formula (Example 1.10.33) in floating-point arithmetic, rounding to the nearest integer. For which n does Binet's formula give a wrong answer for the first time, and why?
- Apply `np.linalg.eig` to the shear $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. What do you notice about the two eigenvectors it returns?

P12. Dual bases and interpolation. Write a function `dual_basis(P)` that returns the rows $r_1, \ldots, r_n$ of the dual basis of the columns of an invertible matrix `P` (Proposition 1.8.7).

- For a random invertible 5×5 matrix P, check that the matrix of all values $\mathbf{r}_i \mathbf{b}_j$ is the identity, and that applying the rows to a vector $\mathbf{x}$ gives its coordinates, i.e., that $\mathbf{x} = \sum_i (\mathbf{r}_i \mathbf{x})\, \mathbf{b}_i$.
- Polynomials of degree at most d are stored by their coefficient vectors in $\mathbb{R}^{d+1}$. For $d+1$ distinct nodes $t_0, \ldots, t_d$, the evaluation functionals $p \mapsto p(t_k)$ are the rows of the Vandermonde matrix V with entries $V_{kj} = t_k^j$. Explain why the coefficient vectors of the Lagrange polynomials $L_0, \ldots, L_d$ (with $L_j(t_k) = \delta_{jk}$) are the columns of V^{-1}, and compute them for the nodes $-1, 0, 1$; compare with Example 1.8.9.
- Interpolate $f(t) = \sin(3t)$ at 9 equally spaced nodes in $[-1, 1]$ with the formula $p = \sum_k f(t_k)\, L_k$, and plot f and p. Then repeat with 21 equally spaced nodes on $[-5, 5]$ for $f(t) = \frac{1}{1+t^2}$. What goes wrong near the ends of the interval? (This is Runge's phenomenon.)

P13. Softmax and cosets. Implement the softmax function in two ways:

```
def softmax_naive(z):
    return np.exp(z) / np.exp(z).sum()
```

```python
def softmax_stable(z):
    z = z - z.max()  # another representative of the same coset
    return np.exp(z) / np.exp(z).sum()
```

- For random vectors $\mathbf{z}$ and constants c, check that `softmax_stable(z + c)` equals `softmax_stable(z)`, i.e., that softmax only depends on the coset $\mathbf{z} + \operatorname{span}\{\mathbf{1}\}$.
- Compare both functions on $\mathbf{z} = (1, 2, 3)$ and on $\mathbf{z} = (1001, 1002, 1003)$. What happens with the naive version, and why does subtracting the maximum fix the problem without changing the result?
- Another canonical representative of the coset is the centered vector $\mathbf{z} - \bar{z}\mathbf{1}$, where $\bar{z}$ is the mean of the entries. Check that it gives the same softmax. Why is subtracting the maximum preferred in practice? (Hint: what is the largest entry after each shift?)

P14. Differentiation and integration as matrices. Store a polynomial $a_0 + a_1 x + \cdots + a_d x^d$ by its coefficient vector `np.array([a0, a1, ..., ad])`.

- Write functions `diff_matrix(d)`, which returns the $d \times (d+1)$ matrix of $D : \mathbb{R}[x]_{\le d} \to \mathbb{R}[x]_{\le d-1}$, $Dp = p'$, and `int_matrix(d)`, which returns the $(d+2) \times (d+1)$ matrix of $J : \mathbb{R}[x]_{\le d} \to \mathbb{R}[x]_{\le d+1}$, $(Jp)(x) = \int_0^x p(t)\,dt$, with respect to the bases of monomials.
- Check that `diff_matrix(d + 1) @ int_matrix(d)` is the identity matrix, and that `int_matrix(d - 1) @ diff_matrix(d)` is not. Which coefficient does the second product lose?
- The midpoint sum $M_N(f) = \frac{1}{N}\sum_{k=1}^{N} f(x_k)$ with $x_k = \frac{k-1/2}{N}$ approximates $\int_0^1 f(x)\,dx$. Explain why M_N is a linear functional on $C[0,1]$, and why it agrees with the integral on all polynomials of degree at most 1. Compute $M_N(f)$ for $f(x) = \sin(\pi x)$ and $N = 10, 100, 1000$, and compare with $\int_0^1 \sin(\pi x)\,dx = \frac{2}{\pi}$. By which factor does the error shrink when N grows by a factor of 10?

P15. Tensors with NumPy.

- Create the tensor

  ```
  T = np.arange(24).reshape(2, 3, 4)
  ```

  and check that `T[i, j, k]` equals `T.reshape(-1)[12*i + 4*j + k]` for all indices, as in Example 1.11.10. Compute the unfoldings `T.reshape(2, 12)` and `T.transpose(1, 0, 2).reshape(3, 8)`. What are their ranks?
- Compute all the operations of Table 1.1 with `np.einsum` for random arrays, and compare with `@`, `np.trace`, `np.outer` and `.T`. For a batch of 100 random 4×4 matrices, stored as an array of shape $100 \times 4 \times 4$, compute all their traces and all the products $A_b B_b$ with a single call of `np.einsum` each, and compare with a Python loop.
- For random matrices A (3×4), B (2×5) and X (4×5), check that `(A @ X @ B.T).reshape(-1)` equals `np.kron(A, B) @ X.reshape(-1)` (Remark 1.11.17). How many numbers does `np.kron(A, B)` store for $A, B \in \mathbb{R}^{100\times 100}$, and why is it much cheaper to compute $AXB^\top$ directly?
- Write a function that returns the outer product $\mathbf{u} \otimes \mathbf{v} \otimes \mathbf{w}$ of three vectors:

  ```python
  def rank_one(u, v, w):
      return np.einsum("i,j,k->ijk", u, v, w)
  ```

  For random vectors $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{w}$, check that the slices `T[:, :, k]` of `T = rank_one(u, v, w)` are multiples of the matrix $\mathbf{u}\mathbf{v}^\top$ (Example 1.11.20).

P16. The LU decomposition and rank-one updates.

```
def lu(A):
    # return L, U with A = L @ U (no row swaps)
    ...

def solve_lu(L, U, b):
    # solve L U x = b by forward and back substitution
    ...
```

- Implement `lu` by the elimination of Theorem 1.6.106: for each column, store the multipliers in L and replace the rest of the matrix by its Schur complement. Check it on Example 1.6.107, and check that `L @ U` equals A for random 6×6 matrices.
- Implement `solve_lu`, and compare it with `np.linalg.solve` for random systems. For $A = \begin{pmatrix} 10^{-17} & 1 \\ 1 & 1 \end{pmatrix}$ and $\mathbf{b} = (1,2)^\top$, the solution is very close to $(1,1)$. What does your solver return, and why? Swapping the two rows first (partial pivoting) fixes the problem.
- For a random 200×50 data matrix X, compute $(X^\top X)^{-1}$. Then add 100 new random rows one at a time, updating the inverse with the Sherman–Morrison formula of Theorem 1.6.108, and compare the result with $(X^\top X)^{-1}$ computed from scratch for the final matrix. Compare the running times of the two approaches.

P17. Convolutions and the FFT. Write a function `circulant(c)` that returns the matrix $\mathrm{circ}(\mathbf{c})$ of Definition 1.10.36.

- For random $\mathbf{c}, \mathbf{x} \in \mathbb{R}^8$, check that the product `circulant(c) @ x` agrees with the inverse DFT `np.fft.ifft(np.fft.fft(c) * np.fft.fft(x)).real` (NumPy's `fft` computes $\hat{\mathbf{x}} = \overline{F}\mathbf{x}$), and that the eigenvalues of `circulant(c)` are the entries of `np.fft.fft(c)`, up to their order.
- Time both ways of computing $\mathbf{c} * \mathbf{x}$ for $n = 2^{12}$ (without counting the construction of the matrix). Which is faster, and by how much?
- Blur a signal $\mathbf{x} \in \mathbb{R}^{101}$ of your choice with the filter $\mathbf{c} = \frac{1}{4}(2, 1, 0, \ldots, 0, 1)^\top$, add noise of size 10^{-3}, and try to recover $\mathbf{x}$ by dividing the DFT of the result by $\hat{\mathbf{c}}$. Plot $|\hat{c}_m|$ and explain why the recovered signal is so noisy. (Lecture 2 explains this with the condition number of $\mathrm{circ}(\mathbf{c})$.)

Lecture 2

Analytic Geometry

Introduction

In Lecture 1, we studied vector spaces and linear transformations. Vector spaces let us add vectors and multiply them by scalars, but they do not tell us how long a vector is, how far apart two vectors are, or what the angle between two vectors is. These geometric notions are essential in machine learning:

- data points are vectors, and many algorithms, such as k-nearest neighbors and clustering, rely on distances between them; graphs of similar data points lead to spectral clustering;
- similarity between data points, such as text documents or word embeddings, is often measured by the angle between vectors;
- linear regression and principal component analysis find the orthogonal projection of data onto a subspace, and the singular values of the data matrix decide how accurately this can be computed;
- regularization penalizes the size, i.e., the norm, of a weight vector.

In this lecture, we add geometry to vector spaces, using norms and inner products.

Throughout this lecture, all vector spaces are real vector spaces, i.e., vector spaces over $\mathbb{R}$. As in Lecture 1, vectors of $\mathbb{R}^n$ are columns written in bold, $\mathbf{x} = (x_1, \ldots, x_n)^\top$, while elements of general vector spaces, such as functions, are written in normal font. We identify a 1×1 matrix with its only entry. In particular, for $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$,

$$\mathbf{x}^\top \mathbf{y} = x_1 y_1 + x_2 y_2 + \cdots + x_n y_n.$$

2.1 Norms

The length of a vector is measured by a norm. There are many reasonable ways to measure length, so we first collect the properties that every notion of length should have.

Definition 2.1.1 (Norm)**.** A norm on a vector space V is a function $\|\cdot\| : V \to \mathbb{R}$, $v \mapsto \|v\|$, such that for all $u, v \in V$ and all $\lambda \in \mathbb{R}$:

1. Positive definiteness: $\|v\| \geq 0$, and $\|v\| = 0$ if and only if $v = 0$;
2. Absolute homogeneity: $\|\lambda v\| = |\lambda| \, \|v\|$;
3. Triangle inequality: $\|u + v\| \leq \|u\| + \|v\|$.

A vector space together with a norm is called a normed vector space, or normed space.

The triangle inequality says that in a triangle with sides u, v and $u+v$, the length of one side is at most the sum of the lengths of the other two sides.

Definition 2.1.2 (ℓ_1, ℓ_2 and ℓ_∞ norms)**.** For $\mathbf{x} = (x_1, \ldots, x_n)^\top \in \mathbb{R}^n$, we define

$$\begin{aligned} \|\mathbf{x}\|_1 &= |x_1| + |x_2| + \cdots + |x_n| && \text{(the } \ell_1 \text{ norm, or Manhattan norm),} \\ \|\mathbf{x}\|_2 &= \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}} && \text{(the } \ell_2 \text{ norm, or Euclidean norm),} \\ \|\mathbf{x}\|_\infty &= \max\{|x_1|, |x_2|, \ldots, |x_n|\} && \text{(the } \ell_\infty \text{ norm, or maximum norm).} \end{aligned}$$

More generally, for every real number $p \geq 1$, the ℓ_p norm is $\|\mathbf{x}\|_p = \left(|x_1|^p + \cdots + |x_n|^p\right)^{1/p}$.

The name "Manhattan norm" comes from the street grid of Manhattan: to get from $\mathbf{0}$ to $\mathbf{x}$ in $\mathbb{R}^2$ while only moving parallel to the axes, you have to walk the distance $|x_1| + |x_2|$. The Euclidean norm is the usual length of a vector, by the theorem of Pythagoras.

Example 2.1.3. For $\mathbf{x} = (3, -4, 1)^\top$, we have

$$\|\mathbf{x}\|_1 = 3 + 4 + 1 = 8, \qquad \|\mathbf{x}\|_2 = \sqrt{9 + 16 + 1} = \sqrt{26} \approx 5.10, \qquad \|\mathbf{x}\|_\infty = \max\{3, 4, 1\} = 4.$$

Proposition 2.1.4. *The ℓ_1 norm and the ℓ_∞ norm are norms on $\mathbb{R}^n$.*

Proof. Positive definiteness: both $\|\mathbf{x}\|_1$ and $\|\mathbf{x}\|_\infty$ are nonnegative, and they are 0 if and only if all $|x_i|$ are 0, i.e., $\mathbf{x} = \mathbf{0}$. Absolute homogeneity: $|\lambda x_i| = |\lambda||x_i|$, so $\|\lambda\mathbf{x}\|_1 = |\lambda|\|\mathbf{x}\|_1$ and $\|\lambda\mathbf{x}\|_\infty = |\lambda|\|\mathbf{x}\|_\infty$. Triangle inequality: for every i, $|x_i + y_i| \leq |x_i| + |y_i|$. Summing over i gives $\|\mathbf{x} + \mathbf{y}\|_1 \leq \|\mathbf{x}\|_1 + \|\mathbf{y}\|_1$, and since $|x_i| + |y_i| \leq \|\mathbf{x}\|_\infty + \|\mathbf{y}\|_\infty$ for every i, also $\|\mathbf{x} + \mathbf{y}\|_\infty \leq \|\mathbf{x}\|_\infty + \|\mathbf{y}\|_\infty$. □

The Euclidean norm is also a norm. Positive definiteness and absolute homogeneity are again easy to check, but the triangle inequality is harder; we prove it in Section 2.3 (Corollary 2.3.4).

A good way to visualize a norm is through its unit ball $\{\mathbf{x} : \|\mathbf{x}\| \leq 1\}$. Figure 2.1 shows the unit balls of the three norms in $\mathbb{R}^2$.




Figure 2.1: The unit balls $\{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\| \leq 1\}$ of the ℓ_1 norm (red diamond), the ℓ_2 norm (orange disk) and the ℓ_∞ norm (blue square). Each ball contains the previous one, in accordance with Proposition 2.1.5.

Proposition 2.1.5. *For every $\mathbf{x} \in \mathbb{R}^n$,*

$$\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1 \leq n\|\mathbf{x}\|_\infty.$$

Proof. Let $|x_k| = \|\mathbf{x}\|_\infty$ be the largest absolute value of an entry of $\mathbf{x}$. Then $\|\mathbf{x}\|_\infty^2 = x_k^2 \leq x_1^2 + \cdots + x_n^2 = \|\mathbf{x}\|_2^2$. Next,

$$\|\mathbf{x}\|_1^2 = \Big(\sum_i |x_i|\Big)^2 = \sum_i x_i^2 + \sum_{i \neq j} |x_i||x_j| \geq \sum_i x_i^2 = \|\mathbf{x}\|_2^2.$$

Finally, $\|\mathbf{x}\|_1 = \sum_i |x_i| \leq \sum_i |x_k| = n\|\mathbf{x}\|_\infty$. □

So all three norms are "comparable": if a vector is small in one of them, it is also small in the others (up to a factor that depends on n). In Example 2.3.5 we improve the last inequality.

Example 2.1.6 (Frobenius norm)**.** In Lecture 1, we saw that the space $\mathbb{R}^{m\times n}$ of $m \times n$ matrices is a vector space, and it is isomorphic to $\mathbb{R}^{mn}$ by listing the entries of a matrix. The Euclidean norm of this list of entries is the Frobenius norm

$$\|A\|_F = \sqrt{\sum_{i=1}^{m}\sum_{j=1}^{n} A_{ij}^2} = \sqrt{\operatorname{tr}(A^\top A)},$$

which is a norm on $\mathbb{R}^{m\times n}$.

Remark 2.1.7 (Norms in machine learning)*.* Norms are used to measure the size of model parameters. In regularized linear regression, we minimize the error of a model plus a penalty $\lambda\|\mathbf{w}\|$ on its weight vector $\mathbf{w}$. With the squared ℓ_2 norm (ridge regression), all weights are shrunk towards 0. With the ℓ_1 norm (the lasso), many weights become exactly 0, which gives sparse models; one reason is that the ℓ_1 unit ball in Figure 2.1 has corners on the coordinate axes. The ℓ_∞ norm appears, for example, when we bound how much an adversary may change each pixel of an image.

2.2 Inner Products

Norms measure lengths. To also measure angles, we need more structure: an inner product. The prototype is the dot product $\mathbf{x}^\top\mathbf{y}$ on $\mathbb{R}^n$.

Definition 2.2.1 (Inner product)**.** Let V be a vector space. A function $\langle\cdot,\cdot\rangle : V \times V \to \mathbb{R}$ is called

- bilinear if it is linear in each argument: for all $x, y, z \in V$ and $\lambda, \mu \in \mathbb{R}$,

$$\langle \lambda x + \mu y, z\rangle = \lambda\langle x, z\rangle + \mu\langle y, z\rangle \qquad \text{and} \qquad \langle z, \lambda x + \mu y\rangle = \lambda\langle z, x\rangle + \mu\langle z, y\rangle;$$

- symmetric if $\langle x, y\rangle = \langle y, x\rangle$ for all $x, y \in V$;
- positive definite if $\langle x, x\rangle > 0$ for all $x \in V$ with $x \neq 0$.

An inner product on V is a positive definite, symmetric, bilinear function $V \times V \to \mathbb{R}$. A vector space together with an inner product is called an inner product space.

Note that $\langle 0, x\rangle = \langle 0 \cdot 0, x\rangle = 0 \cdot \langle 0, x\rangle = 0$ by bilinearity, so in particular $\langle 0, 0\rangle = 0$.

Example 2.2.2 (Dot product)**.** The dot product $\langle \mathbf{x}, \mathbf{y}\rangle = \mathbf{x}^\top\mathbf{y} = \sum_i x_i y_i$ is an inner product on $\mathbb{R}^n$. It is bilinear and symmetric by the rules of matrix multiplication, and $\mathbf{x}^\top\mathbf{x} = x_1^2 + \cdots + x_n^2 > 0$ for $\mathbf{x} \neq \mathbf{0}$. The space $\mathbb{R}^n$ with the dot product is called the Euclidean space. Unless we say otherwise, we always use the dot product on $\mathbb{R}^n$.

Example 2.2.3. On $\mathbb{R}^2$, define

$$\langle \mathbf{x}, \mathbf{y} \rangle = x_1 y_1 - (x_1 y_2 + x_2 y_1) + 2 x_2 y_2.$$

This function is bilinear and symmetric. It is also positive definite, since

$$\langle \mathbf{x}, \mathbf{x} \rangle = x_1^2 - 2x_1 x_2 + 2x_2^2 = (x_1 - x_2)^2 + x_2^2 \geq 0,$$

with equality only if $x_2 = 0$ and $x_1 = x_2$, i.e., $\mathbf{x} = \mathbf{0}$. So it is an inner product, different from the dot product: for example, for $\mathbf{x} = (1, 1)^\top$, we get $\langle \mathbf{x}, \mathbf{x} \rangle = 1$, while $\mathbf{x}^\top \mathbf{x} = 2$.

Example 2.2.4.

- For positive weights $w_1, \ldots, w_n > 0$, the weighted dot product $\langle \mathbf{x}, \mathbf{y} \rangle = w_1 x_1 y_1 + \cdots + w_n x_n y_n$ is an inner product on $\mathbb{R}^n$. In machine learning, such weights can be used to make some features more important than others.
- The function $\langle \mathbf{x}, \mathbf{y} \rangle = x_1 y_1 - x_2 y_2$ on $\mathbb{R}^2$ is bilinear and symmetric, but it is not an inner product, since $\langle \mathbf{e}_2, \mathbf{e}_2 \rangle = -1 < 0$.
- On the space $\mathbb{R}^{m \times n}$ of matrices, $\langle A, B \rangle = \operatorname{tr}(A^\top B) = \sum_{i,j} A_{ij} B_{ij}$ is an inner product, the Frobenius inner product. It is the dot product of the lists of entries of A and B.

2.2.1 Symmetric positive definite matrices

All the inner products on $\mathbb{R}^n$ in the examples above can be written using a matrix. For instance, the inner product of Example 2.2.3 is

$$\langle \mathbf{x}, \mathbf{y} \rangle = \begin{pmatrix} x_1 & x_2 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \mathbf{x}^\top A \mathbf{y}.$$

The matrices that give inner products in this way are the following.

Definition 2.2.5 (Symmetric positive definite matrix)**.** A symmetric $n \times n$ matrix A is called positive definite if

$$\mathbf{x}^\top A \mathbf{x} > 0 \qquad \text{for all } \mathbf{x} \in \mathbb{R}^n \text{ with } \mathbf{x} \neq \mathbf{0},$$

and positive semidefinite if $\mathbf{x}^\top A \mathbf{x} \geq 0$ for all $\mathbf{x} \in \mathbb{R}^n$. A symmetric positive definite matrix is also called an SPD matrix.

Theorem 2.2.6. *1. If A is a symmetric positive definite $n \times n$ matrix, then $\langle \mathbf{x}, \mathbf{y} \rangle_A = \mathbf{x}^\top A \mathbf{y}$ is an inner product on $\mathbb{R}^n$.*

2. Conversely, let $\langle \cdot, \cdot \rangle$ be an inner product on an n-dimensional vector space V, and let $\mathcal{B} = (b_1, \ldots, b_n)$ be an ordered basis of V. Let A be the $n \times n$ matrix with entries $A_{ij} = \langle b_i, b_j \rangle$. Then A is symmetric positive definite, and

$$\langle x, y \rangle = [x]_{\mathcal{B}}^\top A \, [y]_{\mathcal{B}} \qquad \text{for all } x, y \in V,$$

where $[x]_{\mathcal{B}}$ is the coordinate vector of x with respect to $\mathcal{B}$, as in Lecture 1.

Proof. (1) The function $x^\top A y$ is bilinear by the distributivity of matrix multiplication and the rules for scalars. It is symmetric, since $y^\top A x$ is a 1×1 matrix and therefore equal to its transpose $(y^\top A x)^\top = x^\top A^\top y = x^\top A y$. It is positive definite by assumption.

(2) Write $x = \sum_i \psi_i b_i$ and $y = \sum_j \lambda_j b_j$, so that $[x]_{\mathcal{B}} = (\psi_1, \ldots, \psi_n)^\top$ and $[y]_{\mathcal{B}} = (\lambda_1, \ldots, \lambda_n)^\top$. By bilinearity,

$$\langle x, y \rangle = \Big\langle \sum_i \psi_i b_i, \sum_j \lambda_j b_j \Big\rangle = \sum_{i,j} \psi_i \langle b_i, b_j \rangle \lambda_j = [x]_{\mathcal{B}}^\top A [y]_{\mathcal{B}}.$$

The matrix A is symmetric, since $\langle b_i, b_j \rangle = \langle b_j, b_i \rangle$. If $\boldsymbol{\psi} \in \mathbb{R}^n$ is nonzero, then $x = \sum_i \psi_i b_i \neq 0$, since the b_i are linearly independent, and hence $\boldsymbol{\psi}^\top A \boldsymbol{\psi} = \langle x, x \rangle > 0$. □

The matrix A in part 2 is called the Gram matrix of the basis $\mathcal{B}$. So, once we fix a basis, inner products on an n-dimensional space are the same thing as symmetric positive definite $n \times n$ matrices. The dot product on $\mathbb{R}^n$ corresponds to $A = I_n$ and the standard basis.

Proposition 2.2.7. *Let A be a symmetric positive definite $n \times n$ matrix.*

1. *The diagonal entries of A are positive.*
2. *A is invertible.*
3. *For $n = 2$: a symmetric matrix $A = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$ is positive definite if and only if $a > 0$ and $\det(A) = ac - b^2 > 0$.*

Proof. (1) $A_{ii} = \mathbf{e}_i^\top A \mathbf{e}_i > 0$.
(2) If $A\mathbf{x} = \mathbf{0}$, then $\mathbf{x}^\top A\mathbf{x} = 0$, so $\mathbf{x} = \mathbf{0}$. Hence $\text{null}(A) = \{\mathbf{0}\}$, and A is invertible by Lecture 1.
(3) If A is positive definite, then $a > 0$ by part 1. If $a > 0$, completing the square gives

$$\mathbf{x}^\top A\mathbf{x} = ax_1^2 + 2bx_1x_2 + cx_2^2 = a\Big(x_1 + \frac{b}{a}x_2\Big)^2 + \frac{ac - b^2}{a}x_2^2.$$

If $ac - b^2 > 0$, both terms are nonnegative, and they are both zero only if $x_2 = 0$ and $x_1 = 0$; so A is positive definite. If $ac - b^2 \leq 0$, then $\mathbf{x} = (-b, a)^\top \neq \mathbf{0}$ gives $\mathbf{x}^\top A\mathbf{x} = (ac - b^2)a \leq 0$, so A is not positive definite. □

For larger matrices, there is a similar test, Sylvester's criterion: a symmetric matrix is positive definite if and only if the determinants of all its upper left $k \times k$ submatrices, for $k = 1, \ldots, n$, are positive. We prove it in Section 2.10 (Corollary 2.10.4), together with another test, using eigenvalues (Corollary 2.10.3).

Example 2.2.8. The matrix $A_1 = \begin{pmatrix} 9 & 6 \\ 6 & 5 \end{pmatrix}$ is positive definite, since $9 > 0$ and $\det(A_1) = 45 - 36 = 9 > 0$. Indeed, $\mathbf{x}^\top A_1\mathbf{x} = (3x_1 + 2x_2)^2 + x_2^2$. The matrix $A_2 = \begin{pmatrix} 9 & 6 \\ 6 & 3 \end{pmatrix}$ is not, since $\det(A_2) = 27 - 36 = -9 < 0$: for $\mathbf{x} = (2, -3)^\top$, we get $\mathbf{x}^\top A_2\mathbf{x} = -9$.

Example 2.2.9. Let B be an $m \times n$ matrix with linearly independent columns. Then $B^\top B$ is symmetric positive definite: it is symmetric (Lecture 1), and for $\mathbf{x} \neq \mathbf{0}$ we have $B\mathbf{x} \neq \mathbf{0}$, so

$$\mathbf{x}^\top B^\top B\mathbf{x} = (B\mathbf{x})^\top (B\mathbf{x}) = \|B\mathbf{x}\|_2^2 > 0.$$

Matrices of the form $B^\top B$ appear everywhere in machine learning, for example in linear regression (Section 2.8.3) and as (scaled) covariance matrices of data.

Conversely, every symmetric positive definite matrix is of the form $B^\top B$, even with a triangular B. This is the most efficient way to compute with such matrices.

Theorem 2.2.10 (Cholesky decomposition)**.** *A symmetric $n \times n$ matrix A is positive definite if and only if $A = LL^\top$ for a lower triangular matrix L with positive diagonal entries. In this case, L is unique.*

Proof. If $A = LL^\top$, then $L^\top$ is invertible (it is triangular, with nonzero diagonal entries), so $\mathbf{x}^\top A\mathbf{x} = \|L^\top\mathbf{x}\|^2 > 0$ for $\mathbf{x} \neq \mathbf{0}$. Conversely, we use induction on n; for $n = 1$, $A = (a)$ with $a > 0$, and $L = (\sqrt{a})$ is the only choice. For $n \geq 2$, write

$$A = \begin{pmatrix} a & \mathbf{b}^\top \\ \mathbf{b} & A' \end{pmatrix}, \qquad \text{with } a = A_{11} > 0 \text{ (Proposition 2.2.7)},$$

and let $S = A' - \frac{1}{a}\mathbf{b}\mathbf{b}^\top$ be the Schur complement of a (Lecture 1). For $\mathbf{y} \in \mathbb{R}^{n-1}$ with $\mathbf{y} \neq \mathbf{0}$, the vector $\mathbf{x} = (-\frac{1}{a}\mathbf{b}^\top\mathbf{y}, \mathbf{y})$ is nonzero, and expanding gives $0 < \mathbf{x}^\top A\mathbf{x} = \frac{1}{a}(\mathbf{b}^\top\mathbf{y})^2 - \frac{2}{a}(\mathbf{b}^\top\mathbf{y})^2 + \mathbf{y}^\top A'\mathbf{y} = \mathbf{y}^\top S\mathbf{y}$. So S is symmetric positive definite, and by induction, $S = L'L'^\top$ with a unique lower triangular L' with positive diagonal entries. Then

$$L = \begin{pmatrix} \sqrt{a} & \mathbf{0}^\top \\ \frac{1}{\sqrt{a}}\mathbf{b} & L' \end{pmatrix} \qquad \text{satisfies} \qquad LL^\top = \begin{pmatrix} a & \mathbf{b}^\top \\ \mathbf{b} & \frac{1}{a}\mathbf{b}\mathbf{b}^\top + L'L'^\top \end{pmatrix} = A.$$

For uniqueness, compare the blocks of $LL^\top = A$ for any such L: the upper left entry gives $L_{11}^2 = a$, so $L_{11} = \sqrt{a}$; the first column gives L_{11} times the rest of the first column of L equal to $\mathbf{b}$; and the lower right block then gives $L'L'^\top = S$ for the lower right block L' of L, which determines L' by induction. □

The proof is an algorithm: take the square root of the upper left entry, divide the rest of the first column by it, and continue with the Schur complement. It is the LU decomposition of Lecture 1 with $U = L^\top$, and it needs about half as many operations. It fails, with a nonpositive number under a square root, exactly when A is not positive definite, which is how `np.linalg.cholesky` detects such matrices. Once $A = LL^\top$ is known, $A\mathbf{x} = \mathbf{b}$ is solved by forward substitution in $L\mathbf{y} = \mathbf{b}$ and back substitution in $L^\top\mathbf{x} = \mathbf{y}$, and $\det(A) = (L_{11} \cdots L_{nn})^2$. Lecture 4 uses the Cholesky decomposition to sample Gaussian vectors and Gaussian processes.

Example 2.2.11. For $A = \begin{pmatrix} 9 & 6 \\ 6 & 5 \end{pmatrix}$, the algorithm gives $L_{11} = 3$, $L_{21} = \frac{6}{3} = 2$ and $L_{22} = \sqrt{5 - 2^2} = 1$, so

$$A = \begin{pmatrix} 3 & 0 \\ 2 & 1 \end{pmatrix}\begin{pmatrix} 3 & 2 \\ 0 & 1 \end{pmatrix}, \qquad \mathbf{x}^\top A\mathbf{x} = \|L^\top\mathbf{x}\|^2 = (3x_1 + 2x_2)^2 + x_2^2.$$

The Cholesky decomposition is a systematic way of completing the square, as in the proof of Proposition 2.2.7.

Remark 2.2.12 (Kernels). Many machine learning methods, such as support vector machines, only use the data through inner products $\langle \mathbf{x}_i, \mathbf{x}_j \rangle$ of data points. Kernel methods replace them by $k(\mathbf{x}_i, \mathbf{x}_j) = \langle \phi(\mathbf{x}_i), \phi(\mathbf{x}_j) \rangle$, the inner product of the data after a (possibly nonlinear) feature map ϕ. The matrix K with entries $K_{ij} = k(\mathbf{x}_i, \mathbf{x}_j)$ is then a Gram matrix, and it is symmetric positive semidefinite.

2.3 Lengths and Distances

2.3.1 The norm of an inner product

Every inner product gives a norm, in the same way that the dot product gives the Euclidean norm $\|\mathbf{x}\|_2 = \sqrt{\mathbf{x}^\top\mathbf{x}}$.

Definition 2.3.1 (Induced norm)**.** Let V be an inner product space. The length, or norm, of a vector $x \in V$ is

$$\|x\| = \sqrt{\langle x, x \rangle}.$$

A vector x with $\|x\| = 1$ is called a unit vector. For $x \neq 0$, the vector $\frac{x}{\|x\|}$ is a unit vector; replacing x by $\frac{x}{\|x\|}$ is called normalizing x.

Example 2.3.2. For $\mathbf{x} = (1, 1)^\top$, the dot product gives $\|\mathbf{x}\| = \sqrt{2}$, while the inner product of Example 2.2.3 gives $\|\mathbf{x}\| = \sqrt{1} = 1$. So the length of a vector depends on the choice of the inner product.

To show that $\|x\| = \sqrt{\langle x, x\rangle}$ really is a norm, we need the most important inequality of linear algebra.

Theorem 2.3.3 (Cauchy–Schwarz inequality)**.** *Let V be an inner product space. For all $x, y \in V$,*

$$|\langle x, y\rangle| \le \|x\|\,\|y\|,$$

with equality if and only if x and y are linearly dependent.

Proof. If $y = 0$, both sides are 0, and x, y are linearly dependent. So let $y \neq 0$. For every $t \in \mathbb{R}$, by bilinearity and symmetry,

$$0 \le \langle x - ty, x - ty\rangle = \|x\|^2 - 2t\langle x, y\rangle + t^2\|y\|^2.$$

Choosing $t = \frac{\langle x,y\rangle}{\|y\|^2}$ gives

$$0 \le \|x\|^2 - 2\frac{\langle x, y\rangle^2}{\|y\|^2} + \frac{\langle x, y\rangle^2}{\|y\|^2} = \|x\|^2 - \frac{\langle x, y\rangle^2}{\|y\|^2},$$

i.e., $\langle x, y\rangle^2 \le \|x\|^2\|y\|^2$. Taking square roots proves the inequality. Equality holds if and only if $\langle x - ty, x - ty\rangle = 0$ for this t, i.e., $x = ty$, which (for $y \neq 0$) means that x and y are linearly dependent. □

Corollary 2.3.4. *If V is an inner product space, then $\|x\| = \sqrt{\langle x, x\rangle}$ is a norm on V. In particular, the Euclidean norm $\|\cdot\|_2$ is a norm on $\mathbb{R}^n$.*

Proof. Positive definiteness follows from the positive definiteness of the inner product, and $\|\lambda x\| = \sqrt{\lambda^2\langle x, x\rangle} = |\lambda|\|x\|$. For the triangle inequality, by the Cauchy–Schwarz inequality,

$$\|x + y\|^2 = \|x\|^2 + 2\langle x, y\rangle + \|y\|^2 \le \|x\|^2 + 2\|x\|\|y\| + \|y\|^2 = \big(\|x\| + \|y\|\big)^2.$$ □

Example 2.3.5. Let $\mathbf{a} = (|x_1|, \dots, |x_n|)^\top$ and $\mathbf{1} = (1, \dots, 1)^\top$. Applying the Cauchy–Schwarz inequality for the dot product to $\mathbf{a}$ and $\mathbf{1}$ gives

$$\|\mathbf{x}\|_1 = \sum_{i=1}^{n} |x_i| \cdot 1 \le \sqrt{\sum_{i=1}^{n} x_i^2}\sqrt{\sum_{i=1}^{n} 1^2} = \sqrt{n}\,\|\mathbf{x}\|_2.$$

This improves the inequality $\|\mathbf{x}\|_1 \le n\|\mathbf{x}\|_\infty$ of Proposition 2.1.5, since $\sqrt{n}\|\mathbf{x}\|_2 \le n\|\mathbf{x}\|_\infty$.

Not every norm comes from an inner product. Norms that do come from an inner product can be recognized by the following identities.

Proposition 2.3.6 (Parallelogram law and polarization)**.** *Let V be an inner product space with the induced norm. For all $x, y \in V$:*

1. $\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2$ *(parallelogram law);*
2. $\langle x, y\rangle = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2)$ *(polarization identity).*

Proof. Expanding with bilinearity, $\|x \pm y\|^2 = \|x\|^2 \pm 2\langle x, y\rangle + \|y\|^2$. Adding the two identities gives part 1; subtracting them gives part 2. □

The parallelogram law says that in a parallelogram, the sum of the squares of the two diagonals equals the sum of the squares of the four sides. The polarization identity shows that the inner product can be recovered from its norm.

Example 2.3.7. The ℓ_1 norm on $\mathbb{R}^2$ does not come from any inner product: for $\mathbf{x} = \mathbf{e}_1$ and $\mathbf{y} = \mathbf{e}_2$, we get $\|\mathbf{x} + \mathbf{y}\|_1^2 + \|\mathbf{x} - \mathbf{y}\|_1^2 = 4 + 4 = 8$, but $2\|\mathbf{x}\|_1^2 + 2\|\mathbf{y}\|_1^2 = 4$, so the parallelogram law fails. Hence normed spaces are more general than inner product spaces.

2.3.2 Distances

Definition 2.3.8 (Distance)**.** Let V be an inner product space (or, more generally, a normed space). The distance between $x, y \in V$ is

$$d(x, y) = \|x - y\|.$$

For the dot product on $\mathbb{R}^n$, this is the Euclidean distance $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|_2 = \sqrt{\sum_i (x_i - y_i)^2}$.

Proposition 2.3.9. *The distance d has the following properties, for all $x, y, z \in V$:*

1. *Positive definiteness: $d(x, y) \geq 0$, and $d(x, y) = 0$ if and only if $x = y$;*
2. *Symmetry: $d(x, y) = d(y, x)$;*
3. *Triangle inequality: $d(x, z) \leq d(x, y) + d(y, z)$.*

Proof. Part 1 follows from the positive definiteness of the norm, and part 2 from $\|x-y\| = \|(-1)(y-x)\| = \|y-x\|$. Part 3 is the triangle inequality for the norm, applied to $x-z = (x-y)+(y-z)$. □

A function $d : X \times X \to \mathbb{R}$ on any set X with these three properties is called a metric, and (X, d) is called a metric space. So every normed space is a metric space.

Example 2.3.10. For $\mathbf{x} = (1, 2)^\top$ and $\mathbf{y} = (4, 6)^\top$, we have $\mathbf{x} - \mathbf{y} = (-3, -4)^\top$, so the distance between $\mathbf{x}$ and $\mathbf{y}$ is

$$\|\mathbf{x} - \mathbf{y}\|_2 = \sqrt{9 + 16} = 5, \qquad \|\mathbf{x} - \mathbf{y}\|_1 = 3 + 4 = 7, \qquad \|\mathbf{x} - \mathbf{y}\|_\infty = 4$$

for the Euclidean, Manhattan and maximum norms, respectively.

Remark 2.3.11 (Distances in machine learning). The k-nearest neighbor method classifies a new data point by a majority vote among the k training points that are closest to it. Which points are closest depends on the metric, so the choice of the norm matters. It also depends on the units of the features: if one feature is measured in meters and another in millimeters, the second one dominates the Euclidean distance. This is why features are usually standardized first, or, equivalently, a weighted inner product is used.

2.4 Angles and Orthogonality

2.4.1 Angles

By the Cauchy–Schwarz inequality, $-1 \leq \frac{\langle x,y\rangle}{\|x\|\|y\|} \leq 1$ for nonzero vectors x and y. So the following definition makes sense.

Definition 2.4.1 (Angle)**.** Let V be an inner product space, and let $x, y \in V$ be nonzero. The angle between x and y is the unique number $\omega \in [0, \pi]$ with

$$\cos\omega = \frac{\langle x, y\rangle}{\|x\|\,\|y\|}.$$

For the dot product in $\mathbb{R}^2$ or $\mathbb{R}^3$, this is the usual angle between two arrows. Indeed, expanding $\|\mathbf{x} - \mathbf{y}\|^2$ gives

$$\|\mathbf{x} - \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - 2\langle \mathbf{x}, \mathbf{y}\rangle = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - 2\|\mathbf{x}\|\|\mathbf{y}\|\cos\omega,$$

which is the law of cosines for the triangle with sides $\mathbf{x}$, $\mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ (see Figure 2.2).

Example 2.4.2. The angle ω between $\mathbf{x} = (3, 1)^\top$ and $\mathbf{y} = (1, 2)^\top$ satisfies

$$\cos\omega = \frac{3 \cdot 1 + 1 \cdot 2}{\sqrt{10}\,\sqrt{5}} = \frac{5}{\sqrt{50}} = \frac{1}{\sqrt{2}},$$

so $\omega = \frac{\pi}{4}$, i.e., $45°$.




Figure 2.2: The angle $\omega = 45°$ between $\mathbf{x} = (3, 1)$ and $\mathbf{y} = (1, 2)$ (Example 2.4.2). The triangle with sides $\mathbf{x}$, $\mathbf{y}$ and $\mathbf{x} - \mathbf{y}$ satisfies the law of cosines.

2.4.2 Orthogonality

Definition 2.4.3 (Orthogonality)**.** Two vectors x and y of an inner product space are orthogonal, written $x \perp y$, if $\langle x, y\rangle = 0$. If, in addition, $\|x\| = \|y\| = 1$, then x and y are called orthonormal.

So two nonzero vectors are orthogonal if and only if the angle between them is $\frac{\pi}{2}$. The zero vector is orthogonal to every vector.

Example 2.4.4. The vectors $\mathbf{x} = (1, 1)^\top$ and $\mathbf{y} = (-1, 1)^\top$ are orthogonal for the dot product, since $\mathbf{x}^\top\mathbf{y} = -1+1 = 0$. Now consider the inner product $\langle \mathbf{x}, \mathbf{y}\rangle = \mathbf{x}^\top A\mathbf{y}$ with $A = \operatorname{diag}(2, 1)$. Then

$$\langle \mathbf{x}, \mathbf{y}\rangle = 2 \cdot 1 \cdot (-1) + 1 \cdot 1 = -1, \qquad \|\mathbf{x}\|^2 = 2 + 1 = 3, \qquad \|\mathbf{y}\|^2 = 2 + 1 = 3,$$

so $\cos\omega = -\frac{1}{3}$ and $\omega \approx 1.91 \approx 109.5°$. So whether two vectors are orthogonal depends on the inner product.

Theorem 2.4.5 (Pythagoras)**.** *If $x \perp y$, then $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.*

Proof. $\|x + y\|^2 = \|x\|^2 + 2\langle x, y\rangle + \|y\|^2 = \|x\|^2 + \|y\|^2$. □

Remark 2.4.6 (Cosine similarity). In machine learning, the number $\cos\omega = \frac{\langle \mathbf{x},\mathbf{y}\rangle}{\|\mathbf{x}\|\|\mathbf{y}\|}$ is called the cosine similarity of $\mathbf{x}$ and $\mathbf{y}$. It is 1 if $\mathbf{x}$ and $\mathbf{y}$ point in the same direction, 0 if they are orthogonal, and -1 if they point in opposite directions. Unlike the distance, it does not change when $\mathbf{x}$ or $\mathbf{y}$ is multiplied by a positive scalar. This makes it useful for comparing text documents (represented by word counts, where long and short documents should be comparable) and embeddings, e.g., the vectors that language models assign to words and sentences.

2.4.3 Orthogonal matrices

In Lecture 1, we defined an $n \times n$ matrix Q to be orthogonal if $Q^\top Q = I_n$, and we claimed that orthogonal matrices preserve lengths and angles. We can now make this precise and prove it.

Theorem 2.4.7. *For an $n \times n$ matrix Q, the following statements are equivalent:*

1. *Q is orthogonal, i.e., $Q^\top Q = I_n$;*
2. *Q preserves the dot product: $(Q\mathbf{x})^\top(Q\mathbf{y}) = \mathbf{x}^\top\mathbf{y}$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$;*
3. *Q preserves lengths: $\|Q\mathbf{x}\|_2 = \|\mathbf{x}\|_2$ for all $\mathbf{x} \in \mathbb{R}^n$;*
4. *the columns of Q are orthonormal.*

In particular, orthogonal matrices preserve lengths, distances and angles.

Proof. $(1 \Rightarrow 2)$ $(Q\mathbf{x})^\top (Q\mathbf{y}) = \mathbf{x}^\top Q^\top Q\mathbf{y} = \mathbf{x}^\top \mathbf{y}$.

$(2 \Rightarrow 3)$ Take $\mathbf{y} = \mathbf{x}$ and take square roots.

$(3 \Rightarrow 2)$ By the polarization identity (Proposition 2.3.6) and linearity,

$$(Q\mathbf{x})^\top (Q\mathbf{y}) = \tfrac{1}{4}(\|Q\mathbf{x} + Q\mathbf{y}\|^2 - \|Q\mathbf{x} - Q\mathbf{y}\|^2) = \tfrac{1}{4}(\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2) = \mathbf{x}^\top \mathbf{y}.$$

$(2 \Rightarrow 1)$ The (i, j) entry of $Q^\top Q$ is $\mathbf{e}_i^\top Q^\top Q\mathbf{e}_j = (Q\mathbf{e}_i)^\top (Q\mathbf{e}_j) = \mathbf{e}_i^\top \mathbf{e}_j$, which is 1 if $i = j$ and 0 otherwise.

$(1 \Leftrightarrow 4)$ The (i, j) entry of $Q^\top Q$ is $\mathbf{q}_i^\top \mathbf{q}_j$, where $\mathbf{q}_1, \ldots, \mathbf{q}_n$ are the columns of Q. So $Q^\top Q = I_n$ means exactly that $\mathbf{q}_i^\top \mathbf{q}_j$ is 1 for $i = j$ and 0 for $i \neq j$.

Finally, distances are lengths of differences, $\|Q\mathbf{x} - Q\mathbf{y}\|_2 = \|Q(\mathbf{x} - \mathbf{y})\|_2 = \|\mathbf{x} - \mathbf{y}\|_2$, and angles are defined in terms of the dot product and lengths. □

A map that preserves distances is called an isometry. Rotations and reflections are the typical examples; we study rotations in Section 2.9.

2.5 Orthonormal Bases

In $\mathbb{R}^n$, the standard basis $\mathbf{e}_1, \ldots, \mathbf{e}_n$ consists of vectors of length 1 that are pairwise orthogonal. Bases with this property are especially convenient.

Definition 2.5.1 (Orthonormal basis)**.** Let V be an inner product space. Vectors $b_1, \ldots, b_k \in V$ are called orthogonal if $\langle b_i, b_j \rangle = 0$ for all $i \neq j$, and orthonormal if, in addition, $\|b_i\| = 1$ for all i. In other words, $b_1, \ldots, b_k$ are orthonormal if

$$\langle b_i, b_j \rangle = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

An orthonormal basis (ONB) of V is a basis of V consisting of orthonormal vectors.

Lemma 2.5.2. *Orthogonal nonzero vectors $b_1, \ldots, b_k$ are linearly independent.*

Proof. Suppose that $a_1 b_1 + \cdots + a_k b_k = 0$. Taking the inner product with b_j and using bilinearity gives

$$0 = \langle a_1 b_1 + \cdots + a_k b_k, b_j \rangle = a_1 \langle b_1, b_j \rangle + \cdots + a_k \langle b_k, b_j \rangle = a_j \|b_j\|^2.$$

Since $b_j \neq 0$, we get $a_j = 0$, for every j. □

The main advantage of an orthonormal basis is that coordinates are computed by inner products, without solving a system of linear equations.

Proposition 2.5.3. *Let $b_1, \ldots, b_n$ be an orthonormal basis of V. Then for all $u, v \in V$:*

1. $v = \langle v, b_1 \rangle b_1 + \langle v, b_2 \rangle b_2 + \cdots + \langle v, b_n \rangle b_n$;
2. $\langle u, v \rangle = \sum_{i=1}^n \langle u, b_i \rangle \langle v, b_i \rangle$, *and in particular* $\|v\|^2 = \sum_{i=1}^n \langle v, b_i \rangle^2$ *(Parseval's identity).*

Proof. (1) Write $v = c_1 b_1 + \cdots + c_n b_n$. Taking the inner product with b_j gives $\langle v, b_j \rangle = \sum_i c_i \langle b_i, b_j \rangle = c_j$.

(2) Write $u = \sum_i d_i b_i$ and $v = \sum_j c_j b_j$ with $d_i = \langle u, b_i \rangle$ and $c_j = \langle v, b_j \rangle$. Then $\langle u, v \rangle = \sum_{i,j} d_i c_j \langle b_i, b_j \rangle = \sum_i d_i c_i$. □

In the language of Theorem 2.2.6: the Gram matrix of an orthonormal basis is the identity matrix, so in coordinates with respect to an orthonormal basis, every inner product becomes the dot product.

Example 2.5.4. The vectors $\mathbf{b}_1 = \frac{1}{\sqrt{2}}(1,1)^\top$ and $\mathbf{b}_2 = \frac{1}{\sqrt{2}}(1,-1)^\top$ form an orthonormal basis of $\mathbb{R}^2$. For $\mathbf{v} = (3,1)^\top$, we get

$$\langle \mathbf{v}, \mathbf{b}_1 \rangle = \frac{4}{\sqrt{2}} = 2\sqrt{2}, \qquad \langle \mathbf{v}, \mathbf{b}_2 \rangle = \frac{2}{\sqrt{2}} = \sqrt{2},$$

so $\mathbf{v} = 2\sqrt{2}\,\mathbf{b}_1 + \sqrt{2}\,\mathbf{b}_2 = 2(1,1)^\top + (1,-1)^\top$. Parseval's identity checks out: $(2\sqrt{2})^2 + (\sqrt{2})^2 = 10 = 3^2 + 1^2$.

Remark 2.5.5. If $\mathbf{b}_1, \dots, \mathbf{b}_n$ is an orthonormal basis of $\mathbb{R}^n$ (with the dot product), then the matrix Q with columns $\mathbf{b}_1, \dots, \mathbf{b}_n$ is orthogonal (Theorem 2.4.7). The coordinate vector of $\mathbf{v}$ is then $Q^{-1}\mathbf{v} = Q^\top \mathbf{v} = (\mathbf{b}_1^\top \mathbf{v}, \dots, \mathbf{b}_n^\top \mathbf{v})^\top$, in agreement with Proposition 2.5.3.

2.5.1 The Gram–Schmidt process

Every basis can be turned into an orthonormal basis. The idea is to go through the basis vectors one by one, and to subtract from each vector its components in the directions of the previous ones.

Theorem 2.5.6 (Gram–Schmidt process)**.** *Let $v_1, \dots, v_k$ be linearly independent vectors in an inner product space V. Define recursively, for $j = 1, \dots, k$,*

$$u_j = v_j - \sum_{i=1}^{j-1} \langle v_j, e_i \rangle\, e_i, \qquad e_j = \frac{u_j}{\|u_j\|}.$$

Then $e_1, \dots, e_k$ are orthonormal, and $\operatorname{span}\{e_1, \dots, e_j\} = \operatorname{span}\{v_1, \dots, v_j\}$ *for every $j = 1, \dots, k$.*

Proof. We use induction on j. For $j = 1$, we have $u_1 = v_1 \neq 0$ and $e_1 = v_1/\|v_1\|$, so the claim holds. Now let $j \geq 2$, and assume that $e_1, \dots, e_{j-1}$ are orthonormal with $\operatorname{span}\{e_1, \dots, e_{j-1}\} = \operatorname{span}\{v_1, \dots, v_{j-1}\}$.

First, $u_j \neq 0$: otherwise $v_j = \sum_{i<j} \langle v_j, e_i \rangle e_i$ would lie in $\operatorname{span}\{e_1, \dots, e_{j-1}\}$, which equals $\operatorname{span}\{v_1, \dots, v_{j-1}\}$, contradicting the linear independence of $v_1, \dots, v_j$. So e_j is well defined and has length 1.

Next, for $l < j$, using that $e_1, \dots, e_{j-1}$ are orthonormal,

$$\langle u_j, e_l \rangle = \langle v_j, e_l \rangle - \sum_{i<j} \langle v_j, e_i \rangle \langle e_i, e_l \rangle = \langle v_j, e_l \rangle - \langle v_j, e_l \rangle = 0,$$

so e_j is orthogonal to $e_1, \dots, e_{j-1}$. Finally, e_j is a linear combination of v_j and $e_1, \dots, e_{j-1}$, and conversely $v_j = \|u_j\| e_j + \sum_{i<j} \langle v_j, e_i \rangle e_i$. Together with the induction hypothesis, this gives $\operatorname{span}\{e_1, \dots, e_j\} = \operatorname{span}\{v_1, \dots, v_j\}$. □

The vector $\langle v_j, e_i \rangle e_i$ is the component of v_j in the direction of e_i; in Section 2.8, we will call it the orthogonal projection of v_j onto $\operatorname{span}\{e_i\}$. In hand computations, it is often easier to postpone the normalization: since $\langle v_j, e_i \rangle e_i = \frac{\langle v_j, u_i \rangle}{\langle u_i, u_i \rangle} u_i$, the vectors

$$u_j = v_j - \sum_{i=1}^{j-1} \frac{\langle v_j, u_i \rangle}{\langle u_i, u_i \rangle}\, u_i$$

are orthogonal, and we normalize them at the end. Figure 2.3 illustrates the process for two vectors.




Figure 2.3: The Gram–Schmidt process for $\mathbf{v}_1 = (3, 1)$ and $\mathbf{v}_2 = (1, 2)$. The component of $\mathbf{v}_2$ in the direction of $\mathbf{u}_1 = \mathbf{v}_1$ is $\frac{5}{10}\mathbf{u}_1 = (1.5, 0.5)$; subtracting it gives $\mathbf{u}_2 = (-0.5, 1.5)$, which is orthogonal to $\mathbf{u}_1$. Normalizing $\mathbf{u}_1$ and $\mathbf{u}_2$ gives an orthonormal basis of $\mathbb{R}^2$.

Example 2.5.7. Let us apply the Gram–Schmidt process to the basis $\mathbf{v}_1 = (1, 1, 0)^\top$, $\mathbf{v}_2 = (1, 0, 1)^\top$, $\mathbf{v}_3 = (0, 1, 1)^\top$ of $\mathbb{R}^3$, using the unnormalized version:

$$\begin{aligned}
\mathbf{u}_1 &= \mathbf{v}_1 = (1, 1, 0)^\top, \\
\mathbf{u}_2 &= \mathbf{v}_2 - \frac{\mathbf{v}_2^\top \mathbf{u}_1}{\mathbf{u}_1^\top \mathbf{u}_1}\mathbf{u}_1 = (1, 0, 1)^\top - \frac{1}{2}(1, 1, 0)^\top = (\tfrac{1}{2}, -\tfrac{1}{2}, 1)^\top, \\
\mathbf{u}_3 &= \mathbf{v}_3 - \frac{\mathbf{v}_3^\top \mathbf{u}_1}{\mathbf{u}_1^\top \mathbf{u}_1}\mathbf{u}_1 - \frac{\mathbf{v}_3^\top \mathbf{u}_2}{\mathbf{u}_2^\top \mathbf{u}_2}\mathbf{u}_2 = (0, 1, 1)^\top - \frac{1}{2}(1, 1, 0)^\top - \frac{1/2}{3/2}(\tfrac{1}{2}, -\tfrac{1}{2}, 1)^\top = (-\tfrac{2}{3}, \tfrac{2}{3}, \tfrac{2}{3})^\top.
\end{aligned}$$

One checks that $\mathbf{u}_1^\top \mathbf{u}_2 = \mathbf{u}_1^\top \mathbf{u}_3 = \mathbf{u}_2^\top \mathbf{u}_3 = 0$. Normalizing gives the orthonormal basis

$$\mathbf{e}_1 = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \qquad \mathbf{e}_2 = \frac{1}{\sqrt{6}}\begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \qquad \mathbf{e}_3 = \frac{1}{\sqrt{3}}\begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}.$$

Corollary 2.5.8. *Every finite-dimensional inner product space has an orthonormal basis.*

Proof. Apply the Gram–Schmidt process to any basis $v_1, \ldots, v_n$. The resulting vectors $e_1, \ldots, e_n$ are orthonormal and span $\operatorname{span}\{v_1, \ldots, v_n\} = V$, and they are linearly independent by Lemma 2.5.2. □

The Gram–Schmidt process can also be written as a factorization of matrices.

Proposition 2.5.9 (QR decomposition)**.** *Let A be an $m \times n$ matrix with linearly independent columns $\mathbf{a}_1, \ldots, \mathbf{a}_n$. Then $A = QR$, where Q is an $m \times n$ matrix with orthonormal columns (i.e., $Q^\top Q = I_n$), and R is an upper triangular $n \times n$ matrix with positive diagonal entries.*

Proof. Apply the Gram–Schmidt process to $\mathbf{a}_1, \ldots, \mathbf{a}_n$ (with the dot product), and let Q be the matrix with columns $\mathbf{e}_1, \ldots, \mathbf{e}_n$. Since $\mathbf{a}_j \in \operatorname{span}\{\mathbf{e}_1, \ldots, \mathbf{e}_j\}$ and $\mathbf{e}_1, \ldots, \mathbf{e}_j$ are orthonormal, the argument of Proposition 2.5.3 gives $\mathbf{a}_j = \sum_{i=1}^{j} (\mathbf{e}_i^\top \mathbf{a}_j)\, \mathbf{e}_i$. So $A = QR$, where $R_{ij} = \mathbf{e}_i^\top \mathbf{a}_j$ for $i \le j$ and $R_{ij} = 0$ for $i > j$. The diagonal entries are $R_{jj} = \mathbf{e}_j^\top \mathbf{a}_j = \mathbf{e}_j^\top \mathbf{u}_j = \|\mathbf{u}_j\| > 0$, since $\mathbf{a}_j - \mathbf{u}_j$ is a linear combination of $\mathbf{e}_1, \ldots, \mathbf{e}_{j-1}$, which are orthogonal to $\mathbf{e}_j$. □

Example 2.5.10. For the matrix A with columns v_1, v_2, v_3 of Example 2.5.7, the matrix Q has columns $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$, and

$$R = Q^\top A = \begin{pmatrix} \sqrt{2} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{\sqrt{6}}{2} & \frac{1}{\sqrt{6}} \\ 0 & 0 & \frac{2}{\sqrt{3}} \end{pmatrix}.$$

Remark 2.5.11. In NumPy, `Q, R = np.linalg.qr(A)` computes a QR decomposition (the signs of some columns of Q and rows of R may differ from the ones above). It does not use the Gram–Schmidt process directly, since in floating-point arithmetic the vectors $\mathbf{e}_j$ computed by the Gram–Schmidt process can be far from orthogonal; see Programming Exercise P3. The QR decomposition is used, for example, to solve least squares problems in a numerically stable way.

2.6 Orthogonal Complements

Definition 2.6.1 (Orthogonal complement)**.** Let U be a subspace of an inner product space V. The orthogonal complement of U is the set

$$U^\perp = \{v \in V : \langle v, u\rangle = 0 \text{ for all } u \in U\}$$

of all vectors that are orthogonal to every vector of U.

Lemma 2.6.2. *Let U be a subspace of an inner product space V.*

1. *$U^\perp$ is a subspace of V, and $U \cap U^\perp = \{0\}$.*
2. *If $U = \operatorname{span}\{u_1, \dots, u_k\}$, then $v \in U^\perp$ if and only if $\langle v, u_i\rangle = 0$ for $i = 1, \dots, k$.*

Proof. (1) $\langle 0, u\rangle = 0$ for all u, so $0 \in U^\perp$. If $v, w \in U^\perp$ and $\lambda \in \mathbb{R}$, then $\langle v+w, u\rangle = \langle v, u\rangle + \langle w, u\rangle = 0$ and $\langle \lambda v, u\rangle = \lambda\langle v, u\rangle = 0$ for all $u \in U$. If $v \in U \cap U^\perp$, then $\langle v, v\rangle = 0$, so $v = 0$.

(2) If $\langle v, u_i\rangle = 0$ for all i, then $\langle v, \sum_i a_i u_i\rangle = \sum_i a_i \langle v, u_i\rangle = 0$ for every element $\sum_i a_i u_i$ of U. The converse is clear. □

To state the main theorem, we need the following notion.

Definition 2.6.3 (Direct sum of subspaces)**.** Let U and W be subspaces of a vector space V. We say that V is the direct sum of U and W, and write $V = U \oplus W$, if every $v \in V$ can be written in exactly one way as $v = u + w$ with $u \in U$ and $w \in W$.

Lemma 2.6.4. *If $V = U \oplus W$, where U has a basis $u_1, \dots, u_k$ and W has a basis $w_1, \dots, w_l$, then $u_1, \dots, u_k, w_1, \dots, w_l$ is a basis of V. In particular,* $\dim V = \dim U + \dim W$.

Proof. These vectors span V, since every $v = u + w$ is a linear combination of them. If $\sum_i a_i u_i + \sum_j b_j w_j = 0$, then $0 = \sum_i a_i u_i + \sum_j b_j w_j$ and $0 = 0 + 0$ are two ways to write 0 as an element of U plus an element of W. By uniqueness, $\sum_i a_i u_i = 0$ and $\sum_j b_j w_j = 0$, so all a_i and b_j are 0. □

In the following theorem, we use the fact that every subspace of a finite-dimensional vector space is itself finite-dimensional, which was proved in Lecture 1 with the exchange lemma.

Theorem 2.6.5 (Orthogonal decomposition)**.** *Let U be a subspace of a finite-dimensional inner product space V. Then:*

1. *$V = U \oplus U^\perp$;*
2. *$\dim U^\perp = \dim V - \dim U$;*
3. *$(U^\perp)^\perp = U$.*

Proof. (1) By Corollary 2.5.8, U has an orthonormal basis $e_1, \dots, e_k$. For $v \in V$, let

$$u = \sum_{i=1}^{k} \langle v, e_i\rangle e_i \in U \qquad \text{and} \qquad w = v - u.$$

Then $\langle w, e_j\rangle = \langle v, e_j\rangle - \langle v, e_j\rangle = 0$ for all j, so $w \in U^\perp$ by Lemma 2.6.2, and $v = u + w$. This decomposition is unique: if also $v = u' + w'$ with $u' \in U$ and $w' \in U^\perp$, then $u - u' = w' - w$ lies in $U \cap U^\perp = \{0\}$, so $u = u'$ and $w = w'$.

(2) This follows from part 1 and Lemma 2.6.4.

(3) Every $u \in U$ is orthogonal to every vector of $U^\perp$, so $U \subseteq (U^\perp)^\perp$. Conversely, let $v \in (U^\perp)^\perp$, and write $v = u + w$ with $u \in U$ and $w \in U^\perp$. Then $0 = \langle v, w\rangle = \langle u, w\rangle + \langle w, w\rangle = \|w\|^2$, so $w = 0$ and $v = u \in U$. □

Example 2.6.6. Let $U = \operatorname{span}\{(1,1,1)^\top\}$ in $\mathbb{R}^3$. By Lemma 2.6.2, $U^\perp = \{\mathbf{x} : x_1 + x_2 + x_3 = 0\}$ is the plane that we met as an example of a subspace in Lecture 1, and indeed $\dim U + \dim U^\perp = 1 + 2 = 3$. For $\mathbf{v} = (1,2,3)^\top$, the proof of Theorem 2.6.5 with $\mathbf{e}_1 = \frac{1}{\sqrt{3}}(1,1,1)^\top$ gives the decomposition

$$\mathbf{v} = \underbrace{\frac{6}{\sqrt{3}}\mathbf{e}_1}_{(2,2,2)^\top \in U} + \underbrace{(-1,0,1)^\top}_{\in U^\perp}.$$

2.6.1 The four fundamental subspaces

Orthogonal complements explain how the null space, the row space and the column space of a matrix from Lecture 1 fit together.

Theorem 2.6.7. *Let A be an $m \times n$ matrix. Then, with respect to the dot product,*

$$\operatorname{null}(A) = \operatorname{row}(A)^\perp \quad \text{in } \mathbb{R}^n \qquad \text{and} \qquad \operatorname{null}(A^\top) = \operatorname{col}(A)^\perp \quad \text{in } \mathbb{R}^m.$$

Proof. Let $\mathbf{r}_1, \ldots, \mathbf{r}_m$ be the rows of A, written as column vectors. The i-th entry of $A\mathbf{x}$ is $\mathbf{r}_i^\top \mathbf{x}$, so $A\mathbf{x} = \mathbf{0}$ if and only if $\mathbf{x}$ is orthogonal to $\mathbf{r}_1, \ldots, \mathbf{r}_m$, which by Lemma 2.6.2 means $\mathbf{x} \in \operatorname{row}(A)^\perp$. The second statement follows by applying the first one to $A^\top$, since $\operatorname{row}(A^\top) = \operatorname{col}(A)$. □

Combined with Theorem 2.6.5, this gives $\mathbb{R}^n = \operatorname{row}(A) \oplus \operatorname{null}(A)$ and $\dim \operatorname{row}(A) + \dim \operatorname{null}(A) = n$. Since $\dim \operatorname{row}(A) = \operatorname{rank}(A)$ (Lecture 1), this is another proof of the rank–nullity theorem.

Example 2.6.8. In Lecture 1, we found that the null space of $A = \begin{pmatrix} 1 & 2 & -1 \\ 3 & -1 & 4 \end{pmatrix}$ is spanned by $(-1,1,1)^\top$. Indeed, this vector is orthogonal to both rows: $-1 + 2 - 1 = 0$ and $-3 - 1 + 4 = 0$.

Remark 2.6.9. By Theorem 2.6.5(3), the second statement can also be written as $\operatorname{col}(A) = \operatorname{null}(A^\top)^\perp$. So the system $A\mathbf{x} = \mathbf{b}$ has a solution if and only if $\mathbf{b}$ is orthogonal to every solution $\mathbf{y}$ of $A^\top \mathbf{y} = \mathbf{0}$.

2.6.2 Hyperplanes and linear classifiers

Definition 2.6.10 (Hyperplane)**.** Let $\mathbf{w} \in \mathbb{R}^n$ be nonzero and $c \in \mathbb{R}$. The set

$$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{w}^\top \mathbf{x} = c\}$$

is called a hyperplane, and $\mathbf{w}$ is called a normal vector of H. If $c = 0$, then $H = \operatorname{span}\{\mathbf{w}\}^\perp$ is a subspace of dimension $n - 1$ (Theorem 2.6.5).

If $c \neq 0$, the hyperplane H does not contain $\mathbf{0}$, so it is not a subspace. But if $\mathbf{x}_0$ is any point of H, then $\mathbf{w}^\top \mathbf{x} = c$ if and only if $\mathbf{w}^\top(\mathbf{x} - \mathbf{x}_0) = 0$. Hence

$$H = \mathbf{x}_0 + \operatorname{span}\{\mathbf{w}\}^\perp = \{\mathbf{x}_0 + \mathbf{u} : \mathbf{u} \in \operatorname{span}\{\mathbf{w}\}^\perp\}$$

is a shifted copy of the subspace $\operatorname{span}\{\mathbf{w}\}^\perp$. In general, a set of the form $\mathbf{x}_0 + U$, where U is a subspace, is called an affine subspace. In $\mathbb{R}^2$, hyperplanes are lines, and in $\mathbb{R}^3$, they are planes.

Example 2.6.11. The plane $x + 2y + 2z = 2$ in $\mathbb{R}^3$ has the normal vector $\mathbf{w} = (1, 2, 2)^\top$ and contains the point $\mathbf{x}_0 = (2, 0, 0)^\top$. It is the shifted copy $\mathbf{x}_0 + \operatorname{span}\{\mathbf{w}\}^\perp$ of the plane $x + 2y + 2z = 0$ through the origin.

Remark 2.6.12 (Linear classifiers). A linear classifier assigns to a data point $\mathbf{x} \in \mathbb{R}^n$ the class

$$f(\mathbf{x}) = \operatorname{sign}(\mathbf{w}^\top \mathbf{x} + b) \in \{+1, -1\},$$

for a weight vector $\mathbf{w}$ and a bias $b \in \mathbb{R}$. The decision boundary $\{\mathbf{x} : \mathbf{w}^\top \mathbf{x} + b = 0\}$ is a hyperplane with normal vector $\mathbf{w}$, and the classifier predicts $+1$ on the side into which $\mathbf{w}$ points (see Figure 2.4). The perceptron, logistic regression and support vector machines all learn classifiers of this form. In Section 2.8.4, we compute the distance of a point to the decision boundary, which measures how confident the classification is.




Figure 2.4: A linear classifier in $\mathbb{R}^2$ with $\mathbf{w} = (1, 1)$ and $b = -2$. The decision boundary is the line $x_1 + x_2 = 2$ with normal vector $\mathbf{w}$. Points on the side into which $\mathbf{w}$ points (blue disks) are classified as $+1$, the others (orange squares) as -1. The dashed segment shows the distance of the point $(2.5, 1.5)$ to the boundary, which is $\sqrt{2}$ (Example 2.8.15).

2.7 Inner Products of Functions

Inner products are not limited to $\mathbb{R}^n$. Functions can be added and multiplied by scalars (Lecture 1), and an integral gives a natural inner product on them.

Proposition 2.7.1. *On the vector space $C[a, b]$ of continuous functions $f : [a, b] \to \mathbb{R}$ (Lecture 1),*

$$\langle f, g \rangle = \int_a^b f(x) g(x)\, dx$$

is an inner product.

Proof. Bilinearity and symmetry follow from the linearity of the integral. For positive definiteness, $\langle f, f \rangle = \int_a^b f(x)^2\, dx \geq 0$. If $f \neq 0$, then $f(x_0) \neq 0$ for some x_0, and by continuity, $f(x)^2 \geq \frac{1}{2} f(x_0)^2$ on some interval of positive length around x_0; hence $\langle f, f \rangle > 0$. □

The continuity assumption matters: a function that is 1 at a single point and 0 everywhere else is nonzero, but the integral of its square is 0.

Example 2.7.2. On $C[-\pi, \pi]$, the functions sin and cos are orthogonal, since $\sin(x)\cos(x)$ is an odd function and therefore

$$\langle \sin, \cos \rangle = \int_{-\pi}^{\pi} \sin(x) \cos(x)\, dx = 0.$$

Moreover, $\|\sin\|^2 = \int_{-\pi}^{\pi} \sin^2(x)\,dx = \pi$. More generally, the functions 1, $\cos(x)$, $\sin(x)$, $\cos(2x)$, $\sin(2x)$, $\ldots$ are pairwise orthogonal on $[-\pi, \pi]$. Writing a function in terms of them is the idea behind Fourier series, which decompose a signal into frequencies.

Example 2.7.3 (Legendre polynomials)**.** Let us apply the Gram–Schmidt process (unnormalized version) to the polynomials $1, x, x^2, x^3$ in $C[-1, 1]$. We need the integrals $\int_{-1}^{1} x^k\,dx$, which are 0 for odd k and $\frac{2}{k+1}$ for even k:

$$
\begin{aligned}
u_0 &= 1,\\
u_1 &= x - \frac{\langle x, 1\rangle}{\langle 1, 1\rangle}\cdot 1 = x,\\
u_2 &= x^2 - \frac{\langle x^2, 1\rangle}{\langle 1, 1\rangle}\cdot 1 - \frac{\langle x^2, x\rangle}{\langle x, x\rangle}\,x = x^2 - \frac{2/3}{2} = x^2 - \frac{1}{3},\\
u_3 &= x^3 - \frac{\langle x^3, 1\rangle}{\langle 1, 1\rangle} - \frac{\langle x^3, x\rangle}{\langle x, x\rangle}\,x - \frac{\langle x^3, u_2\rangle}{\langle u_2, u_2\rangle}\,u_2 = x^3 - \frac{2/5}{2/3}\,x = x^3 - \frac{3}{5}x.
\end{aligned}
$$

These orthogonal polynomials, shown in Figure 2.5, are (up to scaling) the Legendre polynomials. They are used, for example, to approximate functions by polynomials (Exercise 13).




Figure 2.5: The orthogonal polynomials of Example 2.7.3 on $[-1, 1]$. Any two of them are orthogonal with respect to $\langle f, g\rangle = \int_{-1}^{1} f(x)g(x)\,dx$.

Remark 2.7.4 (Random variables). Inner products also appear in probability. For random variables X and Y with mean 0, the covariance $\operatorname{Cov}(X, Y) = \mathbb{E}[XY]$ behaves like an inner product: it is bilinear, symmetric, and $\operatorname{Cov}(X, X) = \operatorname{Var}(X) \geq 0$. The corresponding norm is the standard deviation, and the correlation coefficient $\frac{\operatorname{Cov}(X,Y)}{\sqrt{\operatorname{Var}(X)\operatorname{Var}(Y)}}$ is the cosine of the angle between X and Y. In particular, uncorrelated random variables are orthogonal, and the Cauchy–Schwarz inequality shows that correlations lie between -1 and 1.

2.8 Orthogonal Projections

Data in machine learning is often high-dimensional, and it is useful to replace each data point by a point of a lower-dimensional subspace while losing as little information as possible. The best way to do this, in a precise sense, is the orthogonal projection. Orthogonal projections are behind linear regression (Section 2.8.3), principal component analysis (Section 2.11.6) and the Gram–Schmidt process.

Definition 2.8.1 (Orthogonal projection)**.** Let U be a subspace of a finite-dimensional inner product space V. By Theorem 2.6.5, every $v \in V$ can be written uniquely as $v = u + w$ with $u \in U$ and $w \in U^{\perp}$. The vector $\pi_U(v) = u$ is called the orthogonal projection of v onto U. In other words, $\pi_U(v)$ is the unique vector with

$$\pi_U(v) \in U \qquad \text{and} \qquad v - \pi_U(v) \in U^{\perp}.$$

Proposition 2.8.2. *Let U be a subspace of a finite-dimensional inner product space V.*

1. *If $e_1, \dots, e_k$ is an orthonormal basis of U, then $\pi_U(v) = \sum_{i=1}^{k} \langle v, e_i \rangle e_i$.*
2. *$\pi_U : V \to V$ is a linear transformation with image U and kernel $U^\perp$.*
3. *$\pi_U \circ \pi_U = \pi_U$.*

Proof. (1) This is the formula from the proof of Theorem 2.6.5. (2) Linearity follows from the formula in part 1. The image is contained in U, and $\pi_U(u) = u$ for $u \in U$ (take $w = 0$), so the image is U. Moreover, $\pi_U(v) = 0$ if and only if $v = 0 + v$ with $v \in U^\perp$. (3) Since $\pi_U(v) \in U$, we get $\pi_U(\pi_U(v)) = \pi_U(v)$. □

A linear transformation π with $\pi \circ \pi = \pi$ is called a projection; for matrices, this is the condition $P^2 = P$ (compare the idempotent matrices in the practice problems of Lecture 1).

2.8.1 Projection onto a line

Proposition 2.8.3. *Let $\mathbf{b} \neq \mathbf{0}$ and $U = \operatorname{span}\{\mathbf{b}\}$. Then*

$$\pi_U(\mathbf{x}) = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\langle \mathbf{b}, \mathbf{b} \rangle} \mathbf{b}.$$

For the dot product on $\mathbb{R}^n$, this is $\pi_U(\mathbf{x}) = P\mathbf{x}$ with the projection matrix

$$P = \frac{\mathbf{b}\mathbf{b}^\top}{\mathbf{b}^\top \mathbf{b}},$$

which is symmetric and satisfies $P^2 = P$.

Proof. We look for $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$ with $\mathbf{x} - \lambda \mathbf{b} \perp \mathbf{b}$, i.e., $\langle \mathbf{x}, \mathbf{b} \rangle - \lambda \langle \mathbf{b}, \mathbf{b} \rangle = 0$. This gives $\lambda = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\langle \mathbf{b}, \mathbf{b} \rangle}$. For the dot product, $\lambda \mathbf{b} = \mathbf{b}\lambda = \frac{\mathbf{b}(\mathbf{b}^\top \mathbf{x})}{\mathbf{b}^\top \mathbf{b}} = P\mathbf{x}$. The matrix $\mathbf{b}\mathbf{b}^\top$ is symmetric, and $P^2 = \frac{\mathbf{b}(\mathbf{b}^\top \mathbf{b})\mathbf{b}^\top}{(\mathbf{b}^\top \mathbf{b})^2} = P$. □

Note that $\mathbf{b}\mathbf{b}^\top$ is an $n \times n$ matrix, while $\mathbf{b}^\top \mathbf{b}$ is a number. The length of the projection is $\|\pi_U(\mathbf{x})\| = \frac{|\langle \mathbf{x}, \mathbf{b} \rangle|}{\|\mathbf{b}\|} = \|\mathbf{x}\| \, |\cos \omega|$, where ω is the angle between $\mathbf{x}$ and $\mathbf{b}$, as in elementary geometry (Figure 2.6).




Figure 2.6: The orthogonal projection of $\mathbf{x} = (1, 2)$ onto the line spanned by $\mathbf{b} = (2, 1)$ is $\pi_U(\mathbf{x}) = \frac{4}{5}\mathbf{b} = (1.6, 0.8)$. The difference $\mathbf{x} - \pi_U(\mathbf{x}) = (-0.6, 1.2)$ is orthogonal to $\mathbf{b}$.

Example 2.8.4. Let $\mathbf{b} = (1, 2, 2)^\top$ and $\mathbf{x} = (1, 1, 1)^\top$. Then $\mathbf{b}^\top \mathbf{b} = 9$ and $\mathbf{b}^\top \mathbf{x} = 5$, so

$$P = \frac{1}{9} \begin{pmatrix} 1 & 2 & 2 \\ 2 & 4 & 4 \\ 2 & 4 & 4 \end{pmatrix}, \qquad \pi_U(\mathbf{x}) = P\mathbf{x} = \frac{5}{9} \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}.$$

Indeed, $\mathbf{x} - \pi_U(\mathbf{x}) = \frac{1}{9}(4, -1, -1)^\top$ is orthogonal to $\mathbf{b}$, since $4 - 2 - 2 = 0$.

2.8.2 Projection onto a subspace

Now let $U = \operatorname{span}\{\mathbf{b}_1, \ldots, \mathbf{b}_m\}$ be an m-dimensional subspace of $\mathbb{R}^n$, with the dot product, and let B be the $n \times m$ matrix with columns $\mathbf{b}_1, \ldots, \mathbf{b}_m$, so that $U = \operatorname{col}(B)$.

Theorem 2.8.5 (Normal equations)**.** *Let B be an $n \times m$ matrix with linearly independent columns, and $U = \operatorname{col}(B)$. Then for every $\mathbf{x} \in \mathbb{R}^n$,*

$$\pi_U(\mathbf{x}) = B\boldsymbol{\lambda}, \qquad \text{where } \boldsymbol{\lambda} \in \mathbb{R}^m \text{ is the unique solution of } B^\top B\boldsymbol{\lambda} = B^\top \mathbf{x}.$$

In other words, $\boldsymbol{\lambda} = (B^\top B)^{-1} B^\top \mathbf{x}$, and $\pi_U(\mathbf{x}) = P\mathbf{x}$ with the projection matrix

$$P = B(B^\top B)^{-1} B^\top.$$

Proof. Every vector of $U = \operatorname{col}(B)$ has the form $B\boldsymbol{\lambda}$. By Theorem 2.6.7, $U^\perp = \operatorname{col}(B)^\perp = \operatorname{null}(B^\top)$. So $B\boldsymbol{\lambda} = \pi_U(\mathbf{x})$ if and only if $\mathbf{x} - B\boldsymbol{\lambda} \in \operatorname{null}(B^\top)$, i.e., $B^\top(\mathbf{x} - B\boldsymbol{\lambda}) = \mathbf{0}$, i.e., $B^\top B\boldsymbol{\lambda} = B^\top \mathbf{x}$. The matrix $B^\top B$ is symmetric positive definite by Example 2.2.9, hence invertible by Proposition 2.2.7, so this equation has exactly one solution $\boldsymbol{\lambda} = (B^\top B)^{-1} B^\top \mathbf{x}$. □

The equations $B^\top B\boldsymbol{\lambda} = B^\top \mathbf{x}$ are called the normal equations, since they say that $\mathbf{x} - B\boldsymbol{\lambda}$ is orthogonal ("normal") to U. If the columns of B are orthonormal, then $B^\top B = I_m$, and the formulas simplify to $\boldsymbol{\lambda} = B^\top \mathbf{x}$ and $P = BB^\top$, in agreement with Proposition 2.8.2. For $m = 1$, we recover Proposition 2.8.3.

Example 2.8.6. Let $U = \operatorname{span}\{(1,1,1)^\top, (0,1,2)^\top\}$ and $\mathbf{x} = (6,0,0)^\top$. With $B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{pmatrix}$, we get

$$B^\top B = \begin{pmatrix} 3 & 3 \\ 3 & 5 \end{pmatrix}, \qquad B^\top \mathbf{x} = \begin{pmatrix} 6 \\ 0 \end{pmatrix}, \qquad \boldsymbol{\lambda} = \begin{pmatrix} 5 \\ -3 \end{pmatrix},$$

where $\boldsymbol{\lambda}$ solves the normal equations. Hence

$$\pi_U(\mathbf{x}) = 5\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - 3\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 2 \\ -1 \end{pmatrix}, \qquad \mathbf{x} - \pi_U(\mathbf{x}) = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix},$$

and indeed $\mathbf{x} - \pi_U(\mathbf{x})$ is orthogonal to both columns of B. The projection matrix is

$$P = B(B^\top B)^{-1} B^\top = \frac{1}{6}\begin{pmatrix} 5 & 2 & -1 \\ 2 & 2 & 2 \\ -1 & 2 & 5 \end{pmatrix}.$$

Proposition 2.8.7. *The projection matrix $P = B(B^\top B)^{-1} B^\top$ is symmetric, and $P^2 = P$.*

Proof. $P^2 = B(B^\top B)^{-1}(B^\top B)(B^\top B)^{-1} B^\top = B(B^\top B)^{-1} B^\top = P$. Since $B^\top B$ is symmetric, so is its inverse (Lecture 1: $(A^{-1})^\top = (A^\top)^{-1}$), and therefore $P^\top = B\big((B^\top B)^{-1}\big)^\top B^\top = P$. □

The most important property of the orthogonal projection is that it gives the closest point.

Theorem 2.8.8 (Best approximation)**.** *Let U be a subspace of a finite-dimensional inner product space V, and let $v \in V$. Then $\pi_U(v)$ is the point of U that is closest to v:*

$$\|v - \pi_U(v)\| < \|v - u\| \qquad \text{for every } u \in U \text{ with } u \neq \pi_U(v).$$

Proof. Write $v - u = (v - \pi_U(v)) + (\pi_U(v) - u)$. The first summand lies in $U^\perp$, and the second one in U, so they are orthogonal. By the theorem of Pythagoras,

$$\|v - u\|^2 = \|v - \pi_U(v)\|^2 + \|\pi_U(v) - u\|^2 > \|v - \pi_U(v)\|^2,$$

since $\pi_U(v) - u \neq 0$. □

The number $\|v - \pi_U(v)\|$ is called the distance from v to U. In Example 2.8.6, the distance from $\mathbf{x}$ to the plane U is $\|(1, -2, 1)^\top\| = \sqrt{6}$.

Remark 2.8.9. The Gram–Schmidt process (Theorem 2.5.6) can now be described in one line: $u_j = v_j - \pi_{U_{j-1}}(v_j)$, where $U_{j-1} = \operatorname{span}\{v_1, \ldots, v_{j-1}\}$. That is, we subtract from v_j its projection onto the span of the previous vectors.

2.8.3 Least squares and linear regression

Suppose we are given data points $(x_1, y_1), \ldots, (x_N, y_N)$ with $x_i, y_i \in \mathbb{R}$, and we want to find the line $y = w_0 + w_1 x$ that fits the data best. In general, no line passes through all points, so we look for the line that minimizes the sum of squared errors

$$\sum_{i=1}^{N} \big(y_i - (w_0 + w_1 x_i)\big)^2 = \|\mathbf{y} - A\mathbf{w}\|^2, \qquad \text{where} \quad A = \begin{pmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_N \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} y_1 \\ \vdots \\ y_N \end{pmatrix}, \quad \mathbf{w} = \begin{pmatrix} w_0 \\ w_1 \end{pmatrix}.$$

The matrix A is called the design matrix. This is a special case of the following problem.

Theorem 2.8.10 (Least squares)**.** *Let A be an $N \times n$ matrix with linearly independent columns, and let $\mathbf{y} \in \mathbb{R}^N$. Then there is exactly one $\mathbf{w}^* \in \mathbb{R}^n$ that minimizes $\|\mathbf{y} - A\mathbf{w}\|$, namely the solution of the normal equations*

$$A^\top A\, \mathbf{w}^* = A^\top \mathbf{y}, \qquad \textit{i.e.,} \qquad \mathbf{w}^* = (A^\top A)^{-1} A^\top \mathbf{y}.$$

Moreover, $A\mathbf{w}^* = \pi_{\mathrm{col}(A)}(\mathbf{y})$.

Proof. As $\mathbf{w}$ ranges over $\mathbb{R}^n$, the vector $A\mathbf{w}$ ranges over $\mathrm{col}(A)$. By Theorem 2.8.8, $\|\mathbf{y} - A\mathbf{w}\|$ is minimal exactly when $A\mathbf{w} = \pi_{\mathrm{col}(A)}(\mathbf{y})$, and by Theorem 2.8.5 (with $B = A$ and $\boldsymbol{\lambda} = \mathbf{w}$), this happens exactly for $\mathbf{w} = (A^\top A)^{-1} A^\top \mathbf{y}$. □

Example 2.8.11. Let us fit a line to the data points $(-1, 0)$, $(0, 1)$ and $(1, 3)$. The design matrix and the normal equations are

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}, \qquad A^\top A = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}, \quad A^\top \mathbf{y} = \begin{pmatrix} 4 \\ 3 \end{pmatrix},$$

so $\mathbf{w}^* = (\frac{4}{3}, \frac{3}{2})^\top$, and the best line is $y = \frac{4}{3} + \frac{3}{2}x$ (Figure 2.7). The fitted values are $A\mathbf{w}^* = (-\frac{1}{6}, \frac{4}{3}, \frac{17}{6})^\top$, the residuals are $\mathbf{y} - A\mathbf{w}^* = (\frac{1}{6}, -\frac{1}{3}, \frac{1}{6})^\top$, and the minimal sum of squared errors is $\frac{1}{36} + \frac{1}{9} + \frac{1}{36} = \frac{1}{6}$. As predicted by the theory, the residual vector is orthogonal to both columns of A: its entries sum to 0, and $-\frac{1}{6} + 0 + \frac{1}{6} = 0$.

Remark 2.8.12 (Linear regression in machine learning). The same method works for any number of features: each row of the design matrix A contains the features of one data point (and a 1 for the intercept w_0), and the least squares solution gives the weights of the linear model. Using the columns $1, x, x^2, \ldots, x^d$, we can also fit polynomials of degree d (polynomial regression). In practice, the normal equations are not solved by computing $(A^\top A)^{-1}$, but with a QR decomposition or with `np.linalg.lstsq`, which are more accurate (compare Lecture 1).




Figure 2.7: The least squares line for the data points $(-1, 0)$, $(0, 1)$, $(1, 3)$ (Example 2.8.11). The red segments are the residuals; the line minimizes the sum of their squared lengths.

2.8.4 Projections onto affine subspaces and distances to hyperplanes

Let $L = \mathbf{x}_0 + U$ be an affine subspace, i.e., a subspace U shifted by a vector $\mathbf{x}_0$. To project onto L, we shift everything by $-\mathbf{x}_0$, project onto U, and shift back:

$$\pi_L(\mathbf{x}) = \mathbf{x}_0 + \pi_U(\mathbf{x} - \mathbf{x}_0).$$

Then $\mathbf{x} - \pi_L(\mathbf{x}) = (\mathbf{x} - \mathbf{x}_0) - \pi_U(\mathbf{x} - \mathbf{x}_0)$ lies in $U^\perp$, and by Theorem 2.8.8, $\pi_L(\mathbf{x})$ is the point of L closest to $\mathbf{x}$.

Theorem 2.8.13 (Distance to a hyperplane)**.** *Let $H = \{\mathbf{z} \in \mathbb{R}^n : \mathbf{w}^\top \mathbf{z} = c\}$ be a hyperplane, where $\mathbf{w} \neq \mathbf{0}$. For every $\mathbf{x} \in \mathbb{R}^n$, the point of H closest to $\mathbf{x}$ is*

$$\pi_H(\mathbf{x}) = \mathbf{x} - \frac{\mathbf{w}^\top \mathbf{x} - c}{\|\mathbf{w}\|^2}\,\mathbf{w}, \qquad \textit{and the distance from } \mathbf{x} \textit{ to } H \textit{ is} \qquad \frac{|\mathbf{w}^\top \mathbf{x} - c|}{\|\mathbf{w}\|}.$$

Proof. Let $\mathbf{x}_0 \in H$, so that $H = \mathbf{x}_0 + W$ with $W = \operatorname{span}\{\mathbf{w}\}^\perp$, and $W^\perp = \operatorname{span}\{\mathbf{w}\}$ by Theorem 2.6.5(3). Since $\mathbf{v} = \pi_W(\mathbf{v}) + \pi_{W^\perp}(\mathbf{v})$ for every $\mathbf{v}$, we get

$$\mathbf{x} - \pi_H(\mathbf{x}) = (\mathbf{x} - \mathbf{x}_0) - \pi_W(\mathbf{x} - \mathbf{x}_0) = \pi_{\operatorname{span}\{\mathbf{w}\}}(\mathbf{x} - \mathbf{x}_0) = \frac{\mathbf{w}^\top(\mathbf{x} - \mathbf{x}_0)}{\mathbf{w}^\top \mathbf{w}}\,\mathbf{w} = \frac{\mathbf{w}^\top \mathbf{x} - c}{\|\mathbf{w}\|^2}\,\mathbf{w},$$

using Proposition 2.8.3 and $\mathbf{w}^\top \mathbf{x}_0 = c$. Taking the norm gives the distance. □

The number $\frac{\mathbf{w}^\top \mathbf{x} - c}{\|\mathbf{w}\|}$, without the absolute value, is called the signed distance: it is positive on the side of H into which $\mathbf{w}$ points, and negative on the other side.

Example 2.8.14. The distance from the point $\mathbf{x} = (1, 2, 3)^\top$ to the plane H given by $x + 2y + 2z = 2$ is

$$\frac{|1 + 4 + 6 - 2|}{\sqrt{1 + 4 + 4}} = \frac{9}{3} = 3,$$

and the closest point of H is $\pi_H(\mathbf{x}) = (1, 2, 3)^\top - \frac{9}{9}(1, 2, 2)^\top = (0, 0, 1)^\top$.

Example 2.8.15. For the linear classifier of Figure 2.4, with $\mathbf{w} = (1,1)^\top$ and $b = -2$, the decision boundary is the hyperplane $\mathbf{w}^\top \mathbf{x} = 2$. The signed distance of the point $\mathbf{x} = (2.5, 1.5)^\top$ to it is $\frac{\mathbf{w}^\top \mathbf{x}+b}{\|\mathbf{w}\|} = \frac{4-2}{\sqrt{2}} = \sqrt{2}$, and the closest point on the boundary is $(2.5, 1.5)^\top - \frac{2}{2}(1,1)^\top = (1.5, 0.5)^\top$.

Remark 2.8.16 (Margins). For a linear classifier, the distance of a data point to the decision boundary is a measure of confidence: points far from the boundary are classified with a large margin. Support vector machines choose the hyperplane that maximizes the smallest distance from the training points to the boundary, which by Theorem 2.8.13 means maximizing $\min_i \frac{|\mathbf{w}^\top \mathbf{x}_i + b|}{\|\mathbf{w}\|}$.

2.9 Rotations

Definition 2.9.1 (Rotation)**.** A rotation of $\mathbb{R}^n$ is an orthogonal $n \times n$ matrix R with $\det(R) = 1$. The set of rotations is denoted by $\mathrm{SO}(n)$, the special orthogonal group.

By Theorem 2.4.7, rotations preserve lengths, distances and angles, and the condition $\det(R) = 1$ means that they preserve the orientation (Lecture 1). Orthogonal matrices with determinant -1, such as reflections, reverse the orientation. Since products and inverses of orthogonal matrices are orthogonal, and $\det(R_1 R_2) = \det(R_1)\det(R_2)$, products and inverses of rotations are again rotations; so $\mathrm{SO}(n)$ is a matrix group, i.e., a subgroup of $\mathrm{GL}(n, \mathbb{R})$ (Lecture 1). The inverse of a rotation R is simply $R^\top$. Like $\mathrm{O}(n)$, $\mathrm{SO}(n)$ is a Lie group of dimension $\frac{n(n-1)}{2}$ (Section A.7.1): a rotation of $\mathbb{R}^3$, for example, is described by three angles (Proposition 2.9.2).

2.9.1 Rotations in the plane

In Lecture 1, we met the rotation matrices

$$R_\theta = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix},$$

which rotate the plane counterclockwise by the angle θ. Every rotation R of $\mathbb{R}^2$ is of this form: its first column $(a, b)^\top$ is a unit vector, so $(a, b) = (\cos\theta, \sin\theta)$ for some angle θ; its second column is a unit vector orthogonal to the first, so it is $\pm(-\sin\theta, \cos\theta)^\top$; and $\det(R) = 1$ forces the sign $+$. Since $R_\alpha R_\beta = R_{\alpha+\beta} = R_\beta R_\alpha$, rotations of the plane commute with each other.

2.9.2 Rotations in three dimensions

The rotations of $\mathbb{R}^3$ by the angle θ about the three coordinate axes are

$$R_1(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta \\ 0 & \sin\theta & \cos\theta \end{pmatrix}, \quad R_2(\theta) = \begin{pmatrix} \cos\theta & 0 & \sin\theta \\ 0 & 1 & 0 \\ -\sin\theta & 0 & \cos\theta \end{pmatrix}, \quad R_3(\theta) = \begin{pmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

For example, $R_3(\theta)$ fixes the x_3-axis and rotates the x_1x_2-plane by θ. Each of them is orthogonal with determinant 1. Every rotation of $\mathbb{R}^3$ is a product of such rotations, and it is a rotation about a line through the origin, its axis.

Proposition 2.9.2 (Rotations of $\mathbb{R}^3$)**.** *Let $R \in \mathrm{SO}(3)$.*

1. *(Euler angles.) There are angles α, β and γ with $R = R_3(\alpha)R_1(\beta)R_3(\gamma)$.*
2. *(Axis.) There are an orthogonal matrix Q with $\det(Q) = 1$ and an angle θ with $R = QR_3(\theta)Q^\top$. So R fixes the line spanned by the third column $\mathbf{q}_3$ of Q, and it rotates the plane orthogonal to $\mathbf{q}_3$ by the angle θ.*

Proof. We first show that a rotation S with $S\mathbf{e}_3 = \mathbf{e}_3$ is of the form $R_3(\theta)$. Since $S^\top = S^{-1}$, also $S^\top \mathbf{e}_3 = \mathbf{e}_3$, so the third column and the third row of S are $\mathbf{e}_3$ and $\mathbf{e}_3^\top$:

$$S = \begin{pmatrix} S' & 0 \\ 0 & 1 \end{pmatrix}$$

with a 2×2 matrix S'. The columns of S' are orthonormal, since those of S are, and $\det(S') = \det(S) = 1$ (expand along the last row). So S' is a rotation of the plane, $S' = R_\theta$, and $S = R_3(\theta)$.

(1) The vector $\mathbf{w} = R\mathbf{e}_3$ is a unit vector. Choose $\beta \in [0, \pi]$ with $\cos\beta = w_3$. Then $w_1^2 + w_2^2 = 1 - w_3^2 = \sin^2\beta$, so we can choose α with $(w_1, w_2) = (\sin\alpha\sin\beta, -\cos\alpha\sin\beta)$ (any α if $\sin\beta = 0$). A direct computation gives

$$R_3(\alpha)R_1(\beta)\mathbf{e}_3 = R_3(\alpha)(0, -\sin\beta, \cos\beta)^\top = (\sin\alpha\sin\beta, -\cos\alpha\sin\beta, \cos\beta)^\top = \mathbf{w}.$$

Hence $S = \big(R_3(\alpha)R_1(\beta)\big)^{-1}R$ is a rotation with $S\mathbf{e}_3 = \mathbf{e}_3$, so $S = R_3(\gamma)$ for some γ, and $R = R_3(\alpha)R_1(\beta)R_3(\gamma)$.

(2) Since $\det(R) = 1$ and $R^\top R = I_3$,

$$\det(R - I_3) = \det\left(R(I_3 - R^\top)\right) = \det(R)\det\left((I_3 - R)^\top\right) = \det(I_3 - R) = (-1)^3\det(R - I_3).$$

So $\det(R - I_3) = 0$, and 1 is an eigenvalue of R (Lecture 1). Let $\mathbf{q}_3$ be a unit vector with $R\mathbf{q}_3 = \mathbf{q}_3$, and let $\mathbf{q}_1, \mathbf{q}_2$ be an orthonormal basis of $\operatorname{span}\{\mathbf{q}_3\}^\perp$ (Theorem 2.6.5 and Corollary 2.5.8). The matrix Q with the columns $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$ is orthogonal; if $\det(Q) = -1$, we replace $\mathbf{q}_1$ by $-\mathbf{q}_1$. Then $S = Q^\top RQ$ is a rotation with $S\mathbf{e}_3 = Q^\top R\mathbf{q}_3 = Q^\top\mathbf{q}_3 = \mathbf{e}_3$, so $S = R_3(\theta)$, and $R = QR_3(\theta)Q^\top$. The rotation $R_3(\theta)$ fixes $\mathbf{e}_3$ and rotates the plane spanned by $\mathbf{e}_1$ and $\mathbf{e}_2$, and Q maps $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ to $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$. □

In particular, a rotation of $\mathbb{R}^3$ is determined by its axis and its angle. The Euler angles are used, for example, to describe the orientation of a camera or of a robot arm.

Example 2.9.3. Unlike in the plane, rotations of $\mathbb{R}^3$ do not commute in general. For $\theta = \frac{\pi}{2}$,

$$R_1(\tfrac{\pi}{2})R_3(\tfrac{\pi}{2}) = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 1 & 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} = R_3(\tfrac{\pi}{2})R_1(\tfrac{\pi}{2}).$$

For instance, the first product maps $\mathbf{e}_1$ to $\mathbf{e}_3$, while the second maps $\mathbf{e}_1$ to $\mathbf{e}_2$. You can check this with a book: rotate it first about one axis and then about another, and compare with the opposite order.

2.9.3 Rotations in n dimensions

Definition 2.9.4 (Givens rotation)**.** For $1 \leq i < j \leq n$ and an angle θ, the Givens rotation $G_{ij}(\theta)$ is the $n \times n$ matrix that agrees with the identity matrix, except for the four entries

$$G_{ii} = \cos\theta, \qquad G_{ij} = -\sin\theta, \qquad G_{ji} = \sin\theta, \qquad G_{jj} = \cos\theta.$$

The Givens rotation $G_{ij}(\theta)$ rotates the plane spanned by $\mathbf{e}_i$ and $\mathbf{e}_j$ by the angle θ, and fixes all other standard basis vectors. Its columns are orthonormal, and its determinant is $\cos^2\theta + \sin^2\theta = 1$ (expand along the other rows), so it is a rotation. Givens rotations are used in numerical linear algebra: a well-chosen Givens rotation makes a chosen entry of a matrix zero, and a sequence of them computes a QR decomposition.

Remark 2.9.5 (Rotations in machine learning). Since rotations preserve dot products and distances, any method that only depends on distances or dot products between data points, such as k-nearest neighbors with the Euclidean distance, gives the same results after the data is rotated. In contrast, methods such as decision trees, which look at individual coordinates, may change. Rotated copies of images are also used for data augmentation, to make image classifiers robust to rotations.

2.10 The Spectral Theorem

In Lecture 1, we saw that a matrix is diagonalizable if it has a basis of eigenvectors, and that some matrices, such as the shear or a rotation, are not diagonalizable. Symmetric matrices are much better behaved: they always have an orthonormal basis of eigenvectors. Symmetric matrices are everywhere in machine learning: covariance matrices, Gram and kernel matrices, and the matrices $A^\top A$ of least squares problems are all symmetric.

To state the theorem, we use complex eigenvalues (Lecture 1). For a vector $\mathbf{v} \in \mathbb{C}^n$, we write $\bar{\mathbf{v}}$ for the vector of complex conjugates of its entries. Then $\bar{\mathbf{v}}^\top \mathbf{v} = \sum_i \bar{v}_i v_i = \sum_i |v_i|^2 > 0$ for $\mathbf{v} \neq \mathbf{0}$. A complex number λ is a root of p_A if and only if $A\mathbf{v} = \lambda\mathbf{v}$ for some nonzero $\mathbf{v} \in \mathbb{C}^n$, by the same argument as for real eigenvalues.

Theorem 2.10.1 (Spectral theorem)**.** *Let A be a symmetric real $n \times n$ matrix. Then:*

1. *all complex eigenvalues of A are real;*
2. *eigenvectors of A for distinct eigenvalues are orthogonal;*
3. *there is an orthonormal basis $\mathbf{q}_1, \ldots, \mathbf{q}_n$ of $\mathbb{R}^n$ consisting of eigenvectors of A. Equivalently,*

$$A = Q\Lambda Q^\top,$$

where Q is the orthogonal matrix with columns $\mathbf{q}_1, \ldots, \mathbf{q}_n$, and $\Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$ contains the corresponding eigenvalues.

Proof. (1) Let $A\mathbf{v} = \lambda\mathbf{v}$ with $\lambda \in \mathbb{C}$ and $\mathbf{v} \in \mathbb{C}^n$, $\mathbf{v} \neq \mathbf{0}$. The number $z = \bar{\mathbf{v}}^\top A\mathbf{v}$ is real: since A is real and symmetric, its complex conjugate is $\bar{z} = \mathbf{v}^\top A\bar{\mathbf{v}} = (\mathbf{v}^\top A\bar{\mathbf{v}})^\top = \bar{\mathbf{v}}^\top A^\top \mathbf{v} = z$. Since $z = \bar{\mathbf{v}}^\top(\lambda\mathbf{v}) = \lambda\,\bar{\mathbf{v}}^\top\mathbf{v}$, we get $\lambda = z/(\bar{\mathbf{v}}^\top\mathbf{v}) \in \mathbb{R}$.

(2) Let $A\mathbf{v} = \lambda\mathbf{v}$ and $A\mathbf{w} = \mu\mathbf{w}$ with $\lambda \neq \mu$. Then

$$\lambda\,\mathbf{v}^\top\mathbf{w} = (A\mathbf{v})^\top\mathbf{w} = \mathbf{v}^\top A^\top\mathbf{w} = \mathbf{v}^\top A\mathbf{w} = \mu\,\mathbf{v}^\top\mathbf{w},$$

so $(\lambda - \mu)\mathbf{v}^\top\mathbf{w} = 0$, and hence $\mathbf{v}^\top\mathbf{w} = 0$.

(3) We use induction on n; the case $n = 1$ is clear. By part 1 and the fundamental theorem of algebra, p_A has a real root λ_1. Let $\mathbf{q}_1$ be a unit eigenvector for λ_1 (it can be chosen real, since $A - \lambda_1 I_n$ is a real matrix with a nonzero null space). Let $W = \operatorname{span}\{\mathbf{q}_1\}^\perp$, and let $\mathbf{w}_2, \ldots, \mathbf{w}_n$ be an orthonormal basis of W (Theorem 2.6.5 and Corollary 2.5.8). The subspace W is mapped into itself by A: if $\mathbf{x} \in W$, then $\mathbf{q}_1^\top(A\mathbf{x}) = (A\mathbf{q}_1)^\top\mathbf{x} = \lambda_1\mathbf{q}_1^\top\mathbf{x} = 0$.

Let M be the $n \times (n-1)$ matrix with columns $\mathbf{w}_2, \ldots, \mathbf{w}_n$, so that $M^\top M = I_{n-1}$, and let $B = M^\top AM$, which is a symmetric $(n-1) \times (n-1)$ matrix. Since $A\mathbf{w}_i \in W$, Proposition 2.5.3 gives $A\mathbf{w}_i = \sum_k (\mathbf{w}_k^\top A\mathbf{w}_i)\mathbf{w}_k$, i.e., $AM = MB$. By induction, B has an orthonormal basis of eigenvectors $\mathbf{y}_2, \ldots, \mathbf{y}_n \in \mathbb{R}^{n-1}$, with $B\mathbf{y}_j = \lambda_j\mathbf{y}_j$. Then the vectors $\mathbf{q}_j = M\mathbf{y}_j$ $(j = 2, \ldots, n)$ are orthonormal, since $\mathbf{q}_i^\top\mathbf{q}_j = \mathbf{y}_i^\top M^\top M\mathbf{y}_j = \mathbf{y}_i^\top\mathbf{y}_j$; they lie in W, so they are orthogonal to $\mathbf{q}_1$; and they are eigenvectors, since $A\mathbf{q}_j = AM\mathbf{y}_j = MB\mathbf{y}_j = \lambda_j\mathbf{q}_j$. So $\mathbf{q}_1, \ldots, \mathbf{q}_n$ is an orthonormal basis of eigenvectors.

Finally, by Theorem 2.4.7, Q is orthogonal, so $Q^{-1} = Q^\top$, and by the diagonalization theorem of Lecture 1, $A = Q\Lambda Q^{-1} = Q\Lambda Q^\top$. □

The factorization $A = Q\Lambda Q^\top$ is called the eigendecomposition, or spectral decomposition, of A. Multiplying out, it says that

$$A = \lambda_1\mathbf{q}_1\mathbf{q}_1^\top + \lambda_2\mathbf{q}_2\mathbf{q}_2^\top + \cdots + \lambda_n\mathbf{q}_n\mathbf{q}_n^\top,$$

where each $\mathbf{q}_i\mathbf{q}_i^\top$ is the matrix of the orthogonal projection onto the line $\operatorname{span}\{\mathbf{q}_i\}$ (Proposition 2.8.3). So a symmetric matrix projects a vector onto n orthogonal directions, scales the projections by the eigenvalues, and adds them up.

Example 2.10.2. Let $A = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} = J + 2I_3$, where $J = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$ is the matrix from the section on eigenvalues in Lecture 1. By the computations there, the eigenvalues of A are $2+2=4$, with eigenspace $\operatorname{span}\{(1,1,1)^\top\}$, and $-1+2=1$, with eigenspace the plane $x+y+z=0$. As predicted by the spectral theorem, the two eigenspaces are orthogonal. To get an orthonormal basis of the plane, we apply the Gram–Schmidt process to its basis $(1,-1,0)^\top$, $(1,0,-1)^\top$, which gives $\frac{1}{\sqrt{2}}(1,-1,0)^\top$ and $\frac{1}{\sqrt{6}}(1,1,-2)^\top$. Hence $A = Q\Lambda Q^\top$ with

$$Q = \begin{pmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & -\frac{2}{\sqrt{6}} \end{pmatrix}, \qquad \Lambda = \operatorname{diag}(4,1,1).$$

Corollary 2.10.3. *A symmetric matrix is positive definite if and only if all its eigenvalues are positive, and positive semidefinite if and only if all its eigenvalues are nonnegative.*

Proof. Write $A = Q\Lambda Q^\top$ and $\mathbf{y} = Q^\top \mathbf{x}$. Then

$$\mathbf{x}^\top A \mathbf{x} = \mathbf{y}^\top \Lambda \mathbf{y} = \lambda_1 y_1^2 + \cdots + \lambda_n y_n^2.$$

If all $\lambda_i > 0$, this is positive for $\mathbf{x} \neq \mathbf{0}$ (then $\mathbf{y} \neq \mathbf{0}$, since $Q^\top$ is invertible). Conversely, $\mathbf{q}_i^\top A \mathbf{q}_i = \lambda_i$, so if A is positive definite, all $\lambda_i > 0$. The semidefinite case is the same, with $\geq$ instead of $>$. □

Corollary 2.10.4 (Sylvester's criterion)**.** *Let A be a symmetric $n \times n$ matrix, and let A_k be its upper left $k \times k$ submatrix, with the entries A_{ij} for $i, j \leq k$. Then A is positive definite if and only if* $\det(A_k) > 0$ *for* $k = 1, \ldots, n$.

Proof. For $\mathbf{x} \in \mathbb{R}^k$, let $\tilde{\mathbf{x}} \in \mathbb{R}^n$ be $\mathbf{x}$ followed by $n-k$ zeros. Then $\mathbf{x}^\top A_k \mathbf{x} = \tilde{\mathbf{x}}^\top A \tilde{\mathbf{x}}$. So if A is positive definite, then so is every A_k. By Corollary 2.10.3, the eigenvalues of A_k are positive, and $\det(A_k)$, which is their product (Lecture 1), is positive.

For the converse, we use induction on n; for $n = 1$, $A = (A_{11})$ with $A_{11} > 0$. Let $n \geq 2$, and assume that all $\det(A_k) > 0$. By induction, A_{n-1} is positive definite. The eigenvalues $\lambda_1, \ldots, \lambda_n$ of A have the positive product $\det(A)$, so none of them is 0, and the number of negative ones is even. Suppose that A is not positive definite. Then at least two eigenvalues are negative, say $\lambda_i, \lambda_j < 0$ with $i \neq j$, with orthonormal eigenvectors $\mathbf{q}_i$ and $\mathbf{q}_j$. The equation $a(\mathbf{q}_i)_n + b(\mathbf{q}_j)_n = 0$ has a solution $(a,b) \neq (0,0)$, and then $\mathbf{x} = a\mathbf{q}_i + b\mathbf{q}_j$ is nonzero (as $\mathbf{q}_i$ and $\mathbf{q}_j$ are linearly independent), and its last entry is 0. On the one hand, $\mathbf{x}^\top A \mathbf{x} = \lambda_i a^2 + \lambda_j b^2 < 0$. On the other hand, $\mathbf{x} = \tilde{\mathbf{x}}'$ for some nonzero $\mathbf{x}' \in \mathbb{R}^{n-1}$, so $\mathbf{x}^\top A \mathbf{x} = \mathbf{x}'^\top A_{n-1} \mathbf{x}' > 0$. This is a contradiction. □

For example, the matrix $A = J + 2I_3$ of Example 2.10.2 has $\det(A_1) = 2$, $\det(A_2) = 3$ and $\det(A_3) = 4$, so it is positive definite, in agreement with its eigenvalues $4, 1, 1$.

Example 2.10.5. The matrix $\begin{pmatrix} 9 & 6 \\ 6 & 5 \end{pmatrix}$ from the subsection on symmetric positive definite matrices has trace 14 and determinant 9, so its eigenvalues $7 \pm 2\sqrt{10}$ are both positive, and it is positive definite. The matrix $\begin{pmatrix} 9 & 6 \\ 6 & 3 \end{pmatrix}$ has determinant $-9 < 0$, so its eigenvalues have opposite signs (their product is negative), and it is not positive definite.

Positive semidefinite matrices have square roots, just like nonnegative numbers.

Proposition 2.10.6 (Square root)**.** *Let $A = Q\Lambda Q^\top$ be symmetric positive semidefinite, with $\Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$. Then $A^{1/2} = Q\Lambda^{1/2}Q^\top$, where $\Lambda^{1/2} = \operatorname{diag}(\sqrt{\lambda_1}, \ldots, \sqrt{\lambda_n})$, is the only symmetric positive semidefinite matrix B with $B^2 = A$. If A is positive definite, then so is $A^{1/2}$, and $A^{-1/2} = (A^{1/2})^{-1} = Q\Lambda^{-1/2}Q^\top$.*

Proof. The matrix $Q\Lambda^{1/2}Q^\top$ is symmetric, its eigenvalues $\sqrt{\lambda_i}$ are nonnegative, and $(Q\Lambda^{1/2}Q^\top)^2 = Q\Lambda^{1/2}Q^\top Q\Lambda^{1/2}Q^\top = Q\Lambda Q^\top = A$. For uniqueness, let B be symmetric positive semidefinite with $B^2 = A$, and write $B = PMP^\top$ with $M = \operatorname{diag}(\mu_1, \dots, \mu_n)$, all $\mu_i \geq 0$. Then $A = PM^2P^\top$, so the columns $\mathbf{p}_i$ of P are eigenvectors of A with the eigenvalues μ_i^2. For an eigenvalue λ of A, the eigenspace E_λ of A is spanned by the $\mathbf{p}_i$ with $\mu_i^2 = \lambda$ (Lecture 1, proof of the theorem on multiplicities), i.e., with $\mu_i = \sqrt{\lambda}$, since $\mu_i \geq 0$. So $B\mathbf{v} = \sqrt{\lambda}\,\mathbf{v}$ for every $\mathbf{v} \in E_\lambda$. The eigenspaces of A together span $\mathbb{R}^n$, so B is determined by A. If all $\lambda_i > 0$, then all $\sqrt{\lambda_i} > 0$, and $Q\Lambda^{-1/2}Q^\top$ is the inverse of $Q\Lambda^{1/2}Q^\top$. □

For example, $\begin{pmatrix} 5 & 4 \\ 4 & 5 \end{pmatrix}$ has the eigenvalues 9 and 1, with eigenvectors $(1,1)^\top$ and $(1,-1)^\top$, and its square root is $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, with eigenvalues 3 and 1. If Σ is the covariance matrix of data, multiplying the centered data by $\Sigma^{-1/2}$ gives data with the covariance matrix I (whitening; see Lecture 4). Both $A = A^{1/2}A^{1/2}$ and the Cholesky decomposition $A = LL^\top$ write A as $BB^\top$; they differ by an orthogonal matrix, since $L^{-1}A^{1/2}$ is orthogonal (Exercise 19).

Proposition 2.10.7 (Rayleigh quotient)**.** *Let A be symmetric with eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$ and orthonormal eigenvectors $\mathbf{q}_1, \dots, \mathbf{q}_n$. Then for every $\mathbf{x} \neq \mathbf{0}$,*

$$\lambda_n \leq \frac{\mathbf{x}^\top A\mathbf{x}}{\mathbf{x}^\top\mathbf{x}} \leq \lambda_1,$$

with equality on the right for $\mathbf{x} = \mathbf{q}_1$, and on the left for $\mathbf{x} = \mathbf{q}_n$.

Proof. With $\mathbf{y} = Q^\top\mathbf{x}$, we have $\mathbf{x}^\top\mathbf{x} = \mathbf{y}^\top\mathbf{y}$ (Theorem 2.4.7) and $\mathbf{x}^\top A\mathbf{x} = \sum_i \lambda_i y_i^2$, which lies between $\lambda_n \sum_i y_i^2$ and $\lambda_1 \sum_i y_i^2$. For $\mathbf{x} = \mathbf{q}_i$, the quotient is $\mathbf{q}_i^\top A\mathbf{q}_i = \lambda_i$. □

So the largest eigenvalue of a symmetric matrix is the maximum of $\mathbf{x}^\top A\mathbf{x}$ over all unit vectors $\mathbf{x}$, and it is attained at an eigenvector. This is the key to principal component analysis. More generally, the sum of the k largest eigenvalues is the maximum of $\sum_j \mathbf{b}_j^\top A\mathbf{b}_j$ over k orthonormal vectors $\mathbf{b}_1, \dots, \mathbf{b}_k$.

Proposition 2.10.8 (Ky Fan)**.** *Let A be symmetric with eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$ and orthonormal eigenvectors $\mathbf{q}_1, \dots, \mathbf{q}_n$, and let $1 \leq k \leq n$. For every $n \times k$ matrix B with orthonormal columns $\mathbf{b}_1, \dots, \mathbf{b}_k$,*

$$\operatorname{tr}(B^\top AB) = \sum_{j=1}^{k} \mathbf{b}_j^\top A\mathbf{b}_j \leq \lambda_1 + \cdots + \lambda_k,$$

with equality for $\mathbf{b}_j = \mathbf{q}_j$, $j = 1, \dots, k$.

Proof. Writing $\mathbf{b}_j = \sum_i (\mathbf{q}_i^\top\mathbf{b}_j)\mathbf{q}_i$ (Proposition 2.5.3) gives $\mathbf{b}_j^\top A\mathbf{b}_j = \sum_i \lambda_i (\mathbf{q}_i^\top\mathbf{b}_j)^2$, so

$$\operatorname{tr}(B^\top AB) = \sum_{i=1}^{n} \lambda_i c_i, \qquad \text{where } c_i = \sum_{j=1}^{k} (\mathbf{q}_i^\top\mathbf{b}_j)^2 = \|B^\top\mathbf{q}_i\|^2.$$

The numbers c_i satisfy $\sum_i c_i = \sum_j \sum_i (\mathbf{q}_i^\top\mathbf{b}_j)^2 = \sum_j \|\mathbf{b}_j\|^2 = k$ and $0 \leq c_i \leq 1$: extending $\mathbf{b}_1, \dots, \mathbf{b}_k$ to an orthonormal basis $\mathbf{b}_1, \dots, \mathbf{b}_n$ of $\mathbb{R}^n$ gives $c_i \leq \sum_{j=1}^n (\mathbf{b}_j^\top\mathbf{q}_i)^2 = \|\mathbf{q}_i\|^2 = 1$. Since $\lambda_i \geq \lambda_k$ and $c_i - 1 \leq 0$ for $i \leq k$, and $\lambda_i \leq \lambda_k$ and $c_i \geq 0$ for $i > k$,

$$\sum_{i=1}^{n} \lambda_i c_i - \sum_{i=1}^{k} \lambda_i = \sum_{i \leq k} \lambda_i (c_i - 1) + \sum_{i > k} \lambda_i c_i \leq \lambda_k \sum_{i \leq k} (c_i - 1) + \lambda_k \sum_{i > k} c_i = \lambda_k \Big(\sum_{i=1}^{n} c_i - k \Big) = 0.$$

For $\mathbf{b}_j = \mathbf{q}_j$, $\operatorname{tr}(B^\top AB) = \sum_{j \leq k} \mathbf{q}_j^\top A\mathbf{q}_j = \lambda_1 + \cdots + \lambda_k$. □

The Rayleigh quotient $R_A(\mathbf{x}) = \frac{\mathbf{x}^\top A\mathbf{x}}{\mathbf{x}^\top\mathbf{x}}$ also describes the eigenvalues between λ_1 and λ_n.

Theorem 2.10.9 (Courant–Fischer)**.** *Let A be symmetric with eigenvalues $\lambda_1 \geq \cdots \geq \lambda_n$. For $k = 1, \ldots, n$,*

$$\lambda_k = \max_{\dim U = k} \min_{\mathbf{x} \in U,\, \mathbf{x} \neq \mathbf{0}} R_A(\mathbf{x}),$$

where the maximum is over all k-dimensional subspaces U of $\mathbb{R}^n$, and it is attained for $U = \operatorname{span}\{\mathbf{q}_1, \ldots, \mathbf{q}_k\}$.

Proof. Write $\mathbf{x} = \sum_i y_i \mathbf{q}_i$, so that $R_A(\mathbf{x}) = \sum_i \lambda_i y_i^2 / \sum_i y_i^2$. For $U = \operatorname{span}\{\mathbf{q}_1, \ldots, \mathbf{q}_k\}$, every nonzero $\mathbf{x} \in U$ has $y_i = 0$ for $i > k$, so $R_A(\mathbf{x}) \geq \lambda_k$, with equality for $\mathbf{x} = \mathbf{q}_k$; so the minimum over this U is λ_k. Now let U be any k-dimensional subspace. The subspace $W = \operatorname{span}\{\mathbf{q}_k, \ldots, \mathbf{q}_n\}$ has dimension $n - k + 1$, and $\dim U + \dim W > n$, so U and W contain a common nonzero vector $\mathbf{x}$, as in the proof of Theorem 2.11.7. It has $y_i = 0$ for $i < k$, so $R_A(\mathbf{x}) \leq \lambda_k$. Hence the minimum over U is at most λ_k. □

Corollary 2.10.10 (Weyl)**.** *Let A and E be symmetric, and let all eigenvalues of E lie in $[-\varepsilon, \varepsilon]$. Then the k-th largest eigenvalues of A and $A + E$ differ by at most ε: $|\lambda_k(A + E) - \lambda_k(A)| \leq \varepsilon$ for every k.*

Proof. By Proposition 2.10.7, $-\varepsilon \leq R_E(\mathbf{x}) \leq \varepsilon$, so $R_A(\mathbf{x}) - \varepsilon \leq R_{A+E}(\mathbf{x}) = R_A(\mathbf{x}) + R_E(\mathbf{x}) \leq R_A(\mathbf{x}) + \varepsilon$ for all $\mathbf{x} \neq \mathbf{0}$. Taking the minimum over U and then the maximum over all k-dimensional U preserves these inequalities, and Theorem 2.10.9 gives $\lambda_k(A) - \varepsilon \leq \lambda_k(A + E) \leq \lambda_k(A) + \varepsilon$. □

So the eigenvalues of a symmetric matrix depend continuously, and even with Lipschitz constant 1, on its entries. For example, the eigenvalues of a covariance matrix that is estimated from data are close to the true ones when the estimate is close to the true matrix, which justifies PCA on estimated covariance matrices. For nonsymmetric matrices, this fails badly: $\begin{pmatrix} 0 & 1 \\ \delta & 0 \end{pmatrix}$ has the eigenvalues $\pm\sqrt{\delta}$, which change by 10^{-3} when the entry δ changes from 0 to 10^{-6}.

Remark 2.10.11 (Geometry of symmetric positive definite matrices). If A is symmetric positive definite, the set $\{\mathbf{x} : \mathbf{x}^\top A\mathbf{x} = 1\}$ is an ellipse (in $\mathbb{R}^2$) or an ellipsoid (in higher dimensions), whose axes point in the directions of the eigenvectors $\mathbf{q}_i$ and have lengths $\frac{1}{\sqrt{\lambda_i}}$, since in the coordinates $\mathbf{y} = Q^\top\mathbf{x}$ the equation becomes $\lambda_1 y_1^2 + \cdots + \lambda_n y_n^2 = 1$. In machine learning, such ellipsoids describe the shape of Gaussian distributions (with A the inverse covariance matrix) and the shape of the loss surface of least squares problems near a minimum; when the eigenvalues have very different sizes, gradient descent zig-zags and converges slowly.

2.10.1 The graph Laplacian and spectral clustering

Many data sets are graphs: users connected by friendships, web pages by links, molecules by bonds, or data points connected to their nearest neighbors. A weighted graph on the vertices $1, \ldots, n$ is described by a symmetric $n \times n$ matrix W of weights $w_{ij} = w_{ji} \geq 0$, with $w_{ii} = 0$; there is an edge between i and j if $w_{ij} > 0$. For example, $w_{ij} = 1$ for the pairs of friends, or $w_{ij} = e^{-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2s^2)}$ for data points $\mathbf{x}_i$ that are close to each other. A symmetric matrix describes the graph, and the spectral theorem describes its shape.

Definition 2.10.12 (Graph Laplacian)**.** The degree of the vertex i is $d_i = \sum_j w_{ij}$. The Laplacian of the graph is the matrix $L = D - W$, where $D = \operatorname{diag}(d_1, \ldots, d_n)$.

Proposition 2.10.13. *For every $\mathbf{x} \in \mathbb{R}^n$,*

$$\mathbf{x}^\top L\mathbf{x} = \sum_{i<j} w_{ij}(x_i - x_j)^2.$$

Hence L is symmetric and positive semidefinite, and $L\mathbf{1} = \mathbf{0}$ for the vector $\mathbf{1} = (1, \ldots, 1)^\top$.

Proof. $\mathbf{x}^\top L\mathbf{x} = \sum_i d_i x_i^2 - \sum_{i,j} w_{ij} x_i x_j = \frac{1}{2}\sum_{i,j} w_{ij}(x_i^2 + x_j^2 - 2x_i x_j) = \frac{1}{2}\sum_{i,j} w_{ij}(x_i - x_j)^2$, using $\sum_i d_i x_i^2 = \sum_{i,j} w_{ij} x_i^2 = \sum_{i,j} w_{ij} x_j^2$; the terms with $i = j$ vanish, and those with $i > j$ repeat those with $i < j$. The rows of L sum to $d_i - \sum_j w_{ij} = 0$. □

So $\mathbf{x}^\top L\mathbf{x}$ is small if $\mathbf{x}$ varies little along the edges with large weights: it measures how smooth a function $\mathbf{x}$ on the vertices is. The vertices i and j are in the same connected component of the graph if there is a path $i = i_0, i_1, \dots, i_l = j$ with $w_{i_{s-1}i_s} > 0$ for every step.

Theorem 2.10.14. *Let L be the Laplacian of a graph with the connected components $C_1, \dots, C_k$. Then* $\operatorname{null}(L)$ *is spanned by the indicator vectors $\mathbf{1}_{C_1}, \dots, \mathbf{1}_{C_k}$ (with the entries* 1 *on C_l and* 0 *elsewhere). So the multiplicity of the eigenvalue* 0 *is the number of connected components; in particular, the graph is connected if and only if the second smallest eigenvalue λ_2 of L is positive.*

Proof. Write $L = Q\Lambda Q^\top$ with eigenvalues $\lambda_i \geq 0$ (Corollary 2.10.3), and let $\mathbf{y} = Q^\top\mathbf{x}$. If $\mathbf{x}^\top L\mathbf{x} = \sum_i \lambda_i y_i^2 = 0$, then $y_i = 0$ whenever $\lambda_i > 0$, so $L\mathbf{x} = Q\Lambda\mathbf{y} = \mathbf{0}$; the converse is clear. So $L\mathbf{x} = \mathbf{0}$ if and only if $\mathbf{x}^\top L\mathbf{x} = 0$, i.e., by Proposition 2.10.13, if and only if $x_i = x_j$ whenever $w_{ij} > 0$. Following paths, this means that $\mathbf{x}$ is constant on each connected component, i.e., $\mathbf{x} = \sum_l c_l \mathbf{1}_{C_l}$. The vectors $\mathbf{1}_{C_l}$ are nonzero and orthogonal, hence linearly independent (Lemma 2.5.2), so $\dim \operatorname{null}(L) = k$, which is the multiplicity of the eigenvalue 0 in the spectral decomposition. □

Spectral clustering. Clustering means splitting the vertices into groups with many edges inside the groups and few edges between them. For a set A of vertices and its complement $\bar{A}$, the weight of the edges between them is $\operatorname{cut}(A) = \sum_{i\in A,\, j\in\bar{A}} w_{ij}$. Minimizing the cut alone tends to split off single vertices, so one balances it by the sizes of the parts:

$$\operatorname{RatioCut}(A) = \operatorname{cut}(A)\Big(\frac{1}{|A|} + \frac{1}{|\bar{A}|}\Big).$$

Finding the best A is a hard combinatorial problem, since there are $2^{n-1} - 1$ ways to split the vertices into two nonempty parts. The Laplacian turns it into an eigenvalue problem.

Theorem 2.10.15. *Let $0 = \lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n$ be the eigenvalues of the Laplacian of a graph. For every set of vertices A with $A \neq \emptyset$ and $\bar{A} \neq \emptyset$, define $\mathbf{f} \in \mathbb{R}^n$ by $f_i = \sqrt{|\bar{A}|/|A|}$ for $i \in A$ and $f_i = -\sqrt{|A|/|\bar{A}|}$ for $i \in \bar{A}$. Then*

$$\mathbf{f}^\top\mathbf{1} = 0, \qquad \|\mathbf{f}\|^2 = n, \qquad \mathbf{f}^\top L\mathbf{f} = n\operatorname{RatioCut}(A), \qquad \textit{and hence} \qquad \operatorname{RatioCut}(A) \geq \lambda_2.$$

Proof. $\mathbf{f}^\top\mathbf{1} = |A|\sqrt{|\bar{A}|/|A|} - |\bar{A}|\sqrt{|A|/|\bar{A}|} = \sqrt{|A||\bar{A}|} - \sqrt{|A||\bar{A}|} = 0$, and $\|\mathbf{f}\|^2 = |\bar{A}| + |A| = n$. In Proposition 2.10.13, only the edges between A and $\bar{A}$ contribute, each with

$$(f_i - f_j)^2 = \frac{|\bar{A}|}{|A|} + \frac{|A|}{|\bar{A}|} + 2 = \frac{(|A| + |\bar{A}|)^2}{|A||\bar{A}|} = n\Big(\frac{1}{|A|} + \frac{1}{|\bar{A}|}\Big),$$

which gives $\mathbf{f}^\top L\mathbf{f} = n\operatorname{RatioCut}(A)$. Let $\mathbf{q}_1 = \frac{1}{\sqrt{n}}\mathbf{1}, \mathbf{q}_2, \dots, \mathbf{q}_n$ be orthonormal eigenvectors of L. Since $\mathbf{f}$ is orthogonal to $\mathbf{q}_1$, it is $\mathbf{f} = \sum_{i\geq 2} c_i\mathbf{q}_i$, and $\mathbf{f}^\top L\mathbf{f} = \sum_{i\geq 2} \lambda_i c_i^2 \geq \lambda_2\|\mathbf{f}\|^2 = n\lambda_2$. □

The proof also shows that λ_2 is the minimum of $\frac{\mathbf{x}^\top L\mathbf{x}}{\mathbf{x}^\top\mathbf{x}}$ over all $\mathbf{x} \neq \mathbf{0}$ with $\mathbf{x}^\top\mathbf{1} = 0$, attained at an eigenvector $\mathbf{q}_2$, which is called the Fiedler vector. Minimizing over the special vectors $\mathbf{f}$ of the theorem is hard, but minimizing over all vectors orthogonal to $\mathbf{1}$ is an eigenvalue problem. This suggests spectral clustering: compute the Fiedler vector $\mathbf{q}_2$, and put the vertices with $(\mathbf{q}_2)_i \geq 0$ into one cluster and the others into the second one. For k clusters, one represents each vertex i by the i-th row of the $n \times k$ matrix of the eigenvectors $\mathbf{q}_1, \dots, \mathbf{q}_k$ and clusters these points with k-means (Section 4.14).

Example 2.10.16. Take two triangles on the vertices $\{1,2,3\}$ and $\{4,5,6\}$, joined by the edge between 3 and 4, all with weight 1. One checks that $L\mathbf{x} = \lambda\mathbf{x}$ for $\mathbf{x} = (1,1,t,-t,-1,-1)^\top$ if $1-t=\lambda$ (the equation for vertex 1) and $4t-2=\lambda t$ (vertex 3), i.e., $\lambda^2-5\lambda+2=0$. The smaller root $\lambda_2 = \frac{5-\sqrt{17}}{2} \approx 0.44$ is the second smallest eigenvalue (the others are 0, 3, 3, 3 and $\frac{5+\sqrt{17}}{2}$), with $t = \frac{\sqrt{17}-3}{2} \approx 0.56$. The signs of the Fiedler vector separate the two triangles, and for $A=\{1,2,3\}$, $\mathrm{RatioCut}(A) = 1\cdot(\frac{1}{3}+\frac{1}{3}) = \frac{2}{3} \geq \lambda_2$, as predicted.

Remark 2.10.17 (Graphs in machine learning). The quadratic form $\mathbf{x}^\top L\mathbf{x}$ is used as a smoothness penalty in semi-supervised learning: labels known on a few vertices are extended to the whole graph by minimizing $\mathbf{x}^\top L\mathbf{x}$ subject to the known values, so that neighbors get similar labels. Laplacian eigenmaps embed data points by the eigenvectors $\mathbf{q}_2, \mathbf{q}_3, \ldots$, and graph neural networks update the features of each vertex by averaging over its neighbors with matrices such as $D^{-1/2}WD^{-1/2}$, which are closely related to the Laplacian.

2.11 The Singular Value Decomposition

The eigendecomposition only exists for square matrices, and in general only for diagonalizable ones. The singular value decomposition (SVD) exists for every matrix, square or not. It is one of the most useful factorizations in applied mathematics and machine learning: it describes the geometry of every linear transformation, gives the best low-rank approximations of a matrix, and is the basis of principal component analysis.

2.11.1 Existence of the SVD

Theorem 2.11.1 (Singular value decomposition)**.** *Let A be a real $m\times n$ matrix. Then there are an orthogonal $m\times m$ matrix U, an orthogonal $n\times n$ matrix V, and an $m\times n$ matrix Σ with*

$$\Sigma_{11} = \sigma_1 \geq \Sigma_{22} = \sigma_2 \geq \cdots \geq \Sigma_{rr} = \sigma_r > 0$$

and all other entries 0*, such that*

$$A = U\Sigma V^\top.$$

The number r of nonzero diagonal entries is the rank of A. The numbers $\sigma_1,\ldots,\sigma_r$ are called the singular values of A; the columns $\mathbf{u}_1,\ldots,\mathbf{u}_m$ of U and $\mathbf{v}_1,\ldots,\mathbf{v}_n$ of V are called the left and right singular vectors of A.

Proof. The $n\times n$ matrix $A^\top A$ is symmetric, and positive semidefinite, since $\mathbf{x}^\top A^\top A\mathbf{x} = \|A\mathbf{x}\|^2 \geq 0$. By the spectral theorem and Corollary 2.10.3, there is an orthonormal basis $\mathbf{v}_1,\ldots,\mathbf{v}_n$ of $\mathbb{R}^n$ of eigenvectors of $A^\top A$, with eigenvalues $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$ (we order them by size). Note that

$$\|A\mathbf{v}_i\|^2 = \mathbf{v}_i^\top A^\top A\mathbf{v}_i = \lambda_i\,\mathbf{v}_i^\top\mathbf{v}_i = \lambda_i.$$

Let r be the number of positive eigenvalues, and let $\sigma_i = \sqrt{\lambda_i}$ and $\mathbf{u}_i = \frac{1}{\sigma_i}A\mathbf{v}_i$ for $i=1,\ldots,r$. These vectors are orthonormal:

$$\mathbf{u}_i^\top\mathbf{u}_j = \frac{\mathbf{v}_i^\top A^\top A\mathbf{v}_j}{\sigma_i\sigma_j} = \frac{\lambda_j\,\mathbf{v}_i^\top\mathbf{v}_j}{\sigma_i\sigma_j} = \begin{cases} 1 & \text{if } i=j,\\ 0 & \text{if } i\neq j.\end{cases}$$

Let $\mathbf{u}_{r+1},\ldots,\mathbf{u}_m$ be an orthonormal basis of $\mathrm{span}\{\mathbf{u}_1,\ldots,\mathbf{u}_r\}^\perp$ (Theorem 2.6.5 and Corollary 2.5.8), so that $\mathbf{u}_1,\ldots,\mathbf{u}_m$ is an orthonormal basis of $\mathbb{R}^m$, and let U and V be the orthogonal matrices with these columns. For $i\leq r$, we have $A\mathbf{v}_i = \sigma_i\mathbf{u}_i$ by definition, and for $i>r$, we have $\|A\mathbf{v}_i\|^2 = \lambda_i = 0$, so $A\mathbf{v}_i = \mathbf{0}$. Column by column, this says that $AV = U\Sigma$, and hence $A = U\Sigma V^{-1} = U\Sigma V^\top$.

Finally, every vector $A\mathbf{x} = \sum_i (\mathbf{v}_i^\top \mathbf{x}) A\mathbf{v}_i = \sum_{i \le r} \sigma_i (\mathbf{v}_i^\top \mathbf{x}) \mathbf{u}_i$ lies in $\operatorname{span}\{\mathbf{u}_1, \dots, \mathbf{u}_r\}$, and conversely $\mathbf{u}_i = A(\frac{1}{\sigma_i}\mathbf{v}_i) \in \operatorname{col}(A)$. So $\operatorname{col}(A) = \operatorname{span}\{\mathbf{u}_1, \dots, \mathbf{u}_r\}$, and since $\mathbf{u}_1, \dots, \mathbf{u}_r$ are linearly independent (Lemma 2.5.2), $\operatorname{rank}(A) = r$. □

The proof shows how to compute the SVD by hand:

1. compute the eigenvalues $\lambda_1 \ge \cdots \ge \lambda_n$ and orthonormal eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^\top A$;
2. set $\sigma_i = \sqrt{\lambda_i}$ and $\mathbf{u}_i = \frac{1}{\sigma_i} A\mathbf{v}_i$ for the positive eigenvalues;
3. complete $\mathbf{u}_1, \dots, \mathbf{u}_r$ to an orthonormal basis of $\mathbb{R}^m$.

Similarly, $AA^\top = U\Sigma\Sigma^\top U^\top$, so the left singular vectors are eigenvectors of $AA^\top$, with the same nonzero eigenvalues σ_i^2.

Example 2.11.2. Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{pmatrix}$. Then $A^\top A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, which we diagonalized in Lecture 1: its eigenvalues are 3 and 1, with orthonormal eigenvectors $\mathbf{v}_1 = \frac{1}{\sqrt{2}}(1,1)^\top$ and $\mathbf{v}_2 = \frac{1}{\sqrt{2}}(1,-1)^\top$. So $\sigma_1 = \sqrt{3}$ and $\sigma_2 = 1$, and

$$\mathbf{u}_1 = \frac{1}{\sqrt{3}} A\mathbf{v}_1 = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}, \qquad \mathbf{u}_2 = \frac{1}{1} A\mathbf{v}_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}.$$

A unit vector orthogonal to both is $\mathbf{u}_3 = \frac{1}{\sqrt{3}}(1,-1,-1)^\top$. Hence

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{pmatrix} = \underbrace{\begin{pmatrix} \frac{2}{\sqrt{6}} & 0 & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \end{pmatrix}}_{U} \underbrace{\begin{pmatrix} \sqrt{3} & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}}_{\Sigma} \underbrace{\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}}_{V^\top}.$$

2.11.2 The geometry of the SVD

Since U and V are orthogonal, the SVD says that every linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ is a composition of three simple steps: an orthogonal transformation $V^\top$ (a rotation, possibly combined with a reflection), a scaling Σ of the coordinate axes by the singular values (which may also change the dimension), and another orthogonal transformation U. In particular, A maps the unit sphere of $\mathbb{R}^n$ to an ellipsoid (possibly flattened) in $\mathbb{R}^m$, whose semi-axes are $\sigma_1\mathbf{u}_1, \dots, \sigma_r\mathbf{u}_r$: the right singular vector v_i is mapped to $\sigma_i \mathbf{u}_i$.

Example 2.11.3 (The SVD of a shear)**.** Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ be the shear from the gallery of linear transformations in Lecture 1. Then $A^\top A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ has trace 3 and determinant 1, so its eigenvalues are $\frac{3 \pm \sqrt{5}}{2} = \varphi^2$ and φ^{-2}, where $\varphi = \frac{1+\sqrt{5}}{2}$ is the golden ratio, which we met in Lecture 1 in Binet's formula for the Fibonacci numbers. So the singular values of the shear are

$$\sigma_1 = \varphi \approx 1.618 \qquad \text{and} \qquad \sigma_2 = \frac{1}{\varphi} \approx 0.618.$$

Using $\varphi^2 = \varphi + 1$, one checks that, with $c = \sqrt{1+\varphi^2}$,

$$\mathbf{v}_1 = \frac{1}{c} \begin{pmatrix} 1 \\ \varphi \end{pmatrix}, \quad \mathbf{v}_2 = \frac{1}{c} \begin{pmatrix} -\varphi \\ 1 \end{pmatrix}, \quad \mathbf{u}_1 = \frac{1}{c} \begin{pmatrix} \varphi \\ 1 \end{pmatrix}, \quad \mathbf{u}_2 = \frac{1}{c} \begin{pmatrix} -1 \\ \varphi \end{pmatrix}.$$

Both U and V are rotations, by the angles 31.7° and 58.3°. So the shear is the same as: rotate by $-58.3°$, stretch the x-axis by φ and shrink the y-axis by $\frac{1}{\varphi}$, and rotate by 31.7° (Figure 2.8).




Figure 2.8: The SVD of the shear $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ (Example 2.11.3). The rotation $V^\top$ moves the right singular vectors $\mathbf{v}_1, \mathbf{v}_2$ to $\mathbf{e}_1, \mathbf{e}_2$; the scaling Σ stretches the axes by $\sigma_1 = \varphi$ and $\sigma_2 = 1/\varphi$; and the rotation U moves the axes to the left singular vectors. Altogether, the unit circle is mapped to an ellipse with semi-axes $\sigma_1\mathbf{u}_1 = A\mathbf{v}_1$ and $\sigma_2\mathbf{u}_2 = A\mathbf{v}_2$.

Proposition 2.11.4. *Let $A = U\Sigma V^\top$ be an SVD of a matrix A of rank r. Then*

$$A = \sigma_1\mathbf{u}_1\mathbf{v}_1^\top + \sigma_2\mathbf{u}_2\mathbf{v}_2^\top + \cdots + \sigma_r\mathbf{u}_r\mathbf{v}_r^\top,$$

and the four fundamental subspaces of A are

$$\begin{aligned} \operatorname{col}(A) &= \operatorname{span}\{\mathbf{u}_1, \ldots, \mathbf{u}_r\}, & \operatorname{null}(A^\top) &= \operatorname{span}\{\mathbf{u}_{r+1}, \ldots, \mathbf{u}_m\}, \\ \operatorname{row}(A) &= \operatorname{span}\{\mathbf{v}_1, \ldots, \mathbf{v}_r\}, & \operatorname{null}(A) &= \operatorname{span}\{\mathbf{v}_{r+1}, \ldots, \mathbf{v}_n\}. \end{aligned}$$

Proof. For every $\mathbf{x}$, we saw in the proof of Theorem 2.11.1 that $A\mathbf{x} = \sum_{i\le r} \sigma_i\mathbf{u}_i(\mathbf{v}_i^\top\mathbf{x})$, which is the first formula applied to $\mathbf{x}$. The formula for $\operatorname{col}(A)$ was also shown there. If $\mathbf{x} = \sum_i c_i\mathbf{v}_i$, then $A\mathbf{x} = \sum_{i\le r}\sigma_i c_i\mathbf{u}_i$, which is $\mathbf{0}$ if and only if $c_1 = \cdots = c_r = 0$; this gives $\operatorname{null}(A)$. In an orthonormal basis, the orthogonal complement of the span of some basis vectors is the span of the other ones, so the remaining two formulas follow from Theorem 2.6.7 and Theorem 2.6.5(3). □

The first formula writes A as a sum of r matrices of rank 1. It also gives the reduced SVD $A = U_r\Sigma_r V_r^\top$, where U_r and V_r consist of the first r columns of U and V, and $\Sigma_r = \operatorname{diag}(\sigma_1, \ldots, \sigma_r)$; NumPy computes it with `np.linalg.svd(A, full_matrices=False)`.

2.11.3 Matrix norms and low-rank approximation

Definition 2.11.5 (Spectral norm)**.** The spectral norm (or operator norm) of an $m \times n$ matrix A is

$$\|A\|_2 = \max_{\mathbf{x}\neq\mathbf{0}} \frac{\|A\mathbf{x}\|}{\|\mathbf{x}\|},$$

the largest factor by which A stretches a vector.

Proposition 2.11.6. *Let A have singular values $\sigma_1 \ge \cdots \ge \sigma_r > 0$. Then*

$$\|A\|_2 = \sigma_1 \qquad \textit{and} \qquad \|A\|_F = \sqrt{\sigma_1^2 + \cdots + \sigma_r^2}.$$

Proof. $\frac{\|A\mathbf{x}\|^2}{\|\mathbf{x}\|^2} = \frac{\mathbf{x}^\top A^\top A\mathbf{x}}{\mathbf{x}^\top\mathbf{x}}$ is the Rayleigh quotient of $A^\top A$, whose largest eigenvalue is σ_1^2; so the maximum is σ_1 by Proposition 2.10.7. For the Frobenius norm, $\|A\|_F^2 = \operatorname{tr}(A^\top A)$ is the sum of the eigenvalues of $A^\top A$ (by Lecture 1, the trace is the sum of the eigenvalues), which is $\sigma_1^2 + \cdots + \sigma_r^2$. □

The rank-1 expansion suggests approximating A by keeping only the terms with the largest singular values. This is the best possible approximation of a given rank.

Theorem 2.11.7 (Eckart–Young)**.** *Let A have the SVD $A = \sum_{i=1}^{r} \sigma_i \mathbf{u}_i \mathbf{v}_i^\top$, and for $k < r$ let*

$$A_k = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^\top + \cdots + \sigma_k \mathbf{u}_k \mathbf{v}_k^\top.$$

Then $\operatorname{rank}(A_k) = k$*, and for every matrix B of rank at most k,*

$$\|A - A_k\|_2 = \sigma_{k+1} \le \|A - B\|_2 \qquad \text{and} \qquad \|A - A_k\|_F = \sqrt{\sigma_{k+1}^2 + \cdots + \sigma_r^2} \le \|A - B\|_F.$$

Proof. Since $A_k \mathbf{x} = \sum_{i \le k} \sigma_i (\mathbf{v}_i^\top \mathbf{x}) \mathbf{u}_i$ for every $\mathbf{x}$, and $A_k \mathbf{v}_i = \sigma_i \mathbf{u}_i$ for $i \le k$, the column space of A_k is $\operatorname{span}\{\mathbf{u}_1, \ldots, \mathbf{u}_k\}$, so $\operatorname{rank}(A_k) = k$. Moreover, $A - A_k = \sum_{i>k} \sigma_i \mathbf{u}_i \mathbf{v}_i^\top$ is an SVD of $A - A_k$, with the singular values $\sigma_{k+1}, \ldots, \sigma_r$, so the formulas for $\|A - A_k\|_2$ and $\|A - A_k\|_F$ follow from Proposition 2.11.6. Now let $\operatorname{rank}(B) \le k$, and let A be $m \times n$.

Spectral norm. By the rank–nullity theorem, $\dim \operatorname{null}(B) \ge n - k$. The subspaces $\operatorname{null}(B)$ and $\operatorname{span}\{\mathbf{v}_1, \ldots, \mathbf{v}_{k+1}\}$ of $\mathbb{R}^n$ have dimensions adding up to more than n, so they have a nonzero vector in common: together, a basis of the first and a basis of the second contain more than n vectors, so some nontrivial linear combination of them is 0, and moving the terms of the second basis to the other side gives a vector $\mathbf{x}$ that lies in both subspaces; it is nonzero, since each basis is linearly independent. We may assume $\|\mathbf{x}\| = 1$. Write $\mathbf{x} = \sum_{i \le k+1} c_i \mathbf{v}_i$ with $\sum_i c_i^2 = 1$. Then $B\mathbf{x} = \mathbf{0}$ and $A\mathbf{x} = \sum_{i \le k+1} \sigma_i c_i \mathbf{u}_i$, so

$$\|A - B\|_2^2 \ge \|(A - B)\mathbf{x}\|^2 = \|A\mathbf{x}\|^2 = \sum_{i=1}^{k+1} \sigma_i^2 c_i^2 \ge \sigma_{k+1}^2 \sum_{i=1}^{k+1} c_i^2 = \sigma_{k+1}^2.$$

Frobenius norm. Choose an orthonormal basis of $\operatorname{col}(B)$, which has at most k vectors, and add vectors of an orthonormal basis of $\operatorname{col}(B)^\perp$ (Theorem 2.6.5) until there are k vectors; this is possible since $k < r \le m$. Let W be their span, and C the $m \times k$ matrix with these columns, so that $C^\top C = I_k$, and $P = CC^\top$ is the matrix of the orthogonal projection onto $W \supseteq \operatorname{col}(B)$ (Proposition 2.8.7). For each column $\mathbf{a}_j$ of A and $\mathbf{b}_j$ of B, we have $\mathbf{a}_j - \mathbf{b}_j = (P\mathbf{a}_j - \mathbf{b}_j) + (\mathbf{a}_j - P\mathbf{a}_j)$ with $P\mathbf{a}_j - \mathbf{b}_j \in W$ and $\mathbf{a}_j - P\mathbf{a}_j \in W^\perp$, so by Pythagoras,

$$\|\mathbf{a}_j - \mathbf{b}_j\|^2 \ge \|\mathbf{a}_j - P\mathbf{a}_j\|^2 = \|\mathbf{a}_j\|^2 - \|P\mathbf{a}_j\|^2 = \|\mathbf{a}_j\|^2 - \|C^\top \mathbf{a}_j\|^2.$$

Adding over j gives $\|A - B\|_F^2 \ge \|A\|_F^2 - \|C^\top A\|_F^2 = \|A\|_F^2 - \operatorname{tr}(C^\top A A^\top C)$. The symmetric matrix $AA^\top = U\Sigma\Sigma^\top U^\top$ has the eigenvalues $\sigma_1^2 \ge \cdots \ge \sigma_r^2$, followed by zeros, so by Proposition 2.10.8, $\operatorname{tr}(C^\top A A^\top C) \le \sigma_1^2 + \cdots + \sigma_k^2$. Since $\|A\|_F^2 = \sigma_1^2 + \cdots + \sigma_r^2$, we get $\|A - B\|_F^2 \ge \sigma_{k+1}^2 + \cdots + \sigma_r^2$. □

Example 2.11.8. Let $A = \begin{pmatrix} 3 & 0 \\ 4 & 5 \end{pmatrix}$. Then $A^\top A = \begin{pmatrix} 25 & 20 \\ 20 & 25 \end{pmatrix}$ has the eigenvalues 45 and 5, with eigenvectors $\mathbf{v}_1 = \frac{1}{\sqrt{2}}(1, 1)^\top$ and $\mathbf{v}_2 = \frac{1}{\sqrt{2}}(1, -1)^\top$. So $\sigma_1 = 3\sqrt{5}$, $\sigma_2 = \sqrt{5}$, and

$$\mathbf{u}_1 = \frac{1}{3\sqrt{5}} A\mathbf{v}_1 = \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \qquad \mathbf{u}_2 = \frac{1}{\sqrt{5}} A\mathbf{v}_2 = \frac{1}{\sqrt{10}} \begin{pmatrix} 3 \\ -1 \end{pmatrix}.$$

The best rank-1 approximation of A is

$$A_1 = \sigma_1 \mathbf{u}_1 \mathbf{v}_1^\top = 3\sqrt{5} \cdot \frac{1}{\sqrt{20}} \begin{pmatrix} 1 \\ 3 \end{pmatrix} \begin{pmatrix} 1 & 1 \end{pmatrix} = \begin{pmatrix} 1.5 & 1.5 \\ 4.5 & 4.5 \end{pmatrix},$$

with error $\|A - A_1\|_F = \|A - A_1\|_2 = \sigma_2 = \sqrt{5}$. Also, $\|A\|_2 = 3\sqrt{5} \approx 6.71$.

Remark 2.11.9 (Low-rank approximation in machine learning). A grayscale image with $m \times n$ pixels is an $m \times n$ matrix. Storing its rank-k approximation A_k requires only the $k(m+n+1)$ numbers σ_i, $\mathbf{u}_i$, $\mathbf{v}_i$, instead of mn; for natural images, the singular values decay quickly, and a small k already gives a good approximation (Programming Exercise P10). The same idea is behind recommender systems, which approximate the matrix of user ratings by a low-rank matrix, latent semantic analysis of text, and the compression of large neural networks.

2.11.4 The pseudo-inverse

Definition 2.11.10 (Pseudo-inverse)**.** Let $A = U\Sigma V^\top$ be an SVD of an $m \times n$ matrix A. The pseudo-inverse of A is the $n \times m$ matrix

$$A^+ = V\Sigma^+ U^\top,$$

where Σ^+ is the $n \times m$ matrix with $(\Sigma^+)_{ii} = \frac{1}{\sigma_i}$ for $i \le r$ and all other entries 0.

If A is invertible, then $A^+ = A^{-1}$, since $A^{-1} = (U\Sigma V^\top)^{-1} = V\Sigma^{-1}U^\top$.

Proposition 2.11.11. *If A has linearly independent columns, then $A^+ = (A^\top A)^{-1}A^\top$, and $A^+\mathbf{y}$ is the least squares solution of $A\mathbf{w} \approx \mathbf{y}$ (Theorem 2.8.10).*

Proof. In this case, $r = n$, and $A^\top A = V\Sigma^\top U^\top U\Sigma V^\top = V\,\mathrm{diag}(\sigma_1^2, \dots, \sigma_n^2)V^\top$. Hence

$$(A^\top A)^{-1}A^\top = V\,\mathrm{diag}(\sigma_1^{-2}, \dots, \sigma_n^{-2})V^\top V\Sigma^\top U^\top = V\,\mathrm{diag}(\sigma_1^{-2}, \dots, \sigma_n^{-2})\Sigma^\top U^\top = V\Sigma^+ U^\top.$$

The second statement is Theorem 2.8.10. □

In general, when the columns of A are dependent, the least squares problem has infinitely many solutions, and $A^+\mathbf{y}$ is the one with the smallest norm.

Proposition 2.11.12. *Let A be an $m \times n$ matrix and $\mathbf{y} \in \mathbb{R}^m$. The vectors $\mathbf{w}$ that minimize $\|A\mathbf{w} - \mathbf{y}\|$ are exactly the vectors $\mathbf{w} = A^+\mathbf{y} + \mathbf{v}$ with $\mathbf{v} \in \mathrm{null}(A)$, and among them, $A^+\mathbf{y}$ is the only one with the smallest norm.*

Proof. Let $A = U\Sigma V^\top$, and write $\mathbf{z} = V^\top\mathbf{w}$ and $\mathbf{c} = U^\top\mathbf{y}$. Since $U^\top$ preserves lengths (Theorem 2.4.7) and $U^\top A = \Sigma V^\top$,

$$\|A\mathbf{w} - \mathbf{y}\|^2 = \|U^\top(A\mathbf{w} - \mathbf{y})\|^2 = \|\Sigma\mathbf{z} - \mathbf{c}\|^2 = \sum_{i=1}^{r}(\sigma_i z_i - c_i)^2 + \sum_{i=r+1}^{m} c_i^2.$$

This is smallest if and only if $z_i = c_i/\sigma_i$ for $i \le r$, with $z_{r+1}, \dots, z_n$ arbitrary. For $z_{r+1} = \cdots = z_n = 0$, this is $\mathbf{z} = \Sigma^+\mathbf{c}$, i.e., $\mathbf{w} = V\Sigma^+U^\top\mathbf{y} = A^+\mathbf{y}$. In general, $\mathbf{w} = V\mathbf{z} = A^+\mathbf{y} + \sum_{i>r} z_i\mathbf{v}_i$, and the vectors $\sum_{i>r} z_i\mathbf{v}_i$ are exactly the elements of $\mathrm{null}(A)$ (Proposition 2.11.4). Finally, $\|\mathbf{w}\|^2 = \|\mathbf{z}\|^2 = \sum_{i\le r}(c_i/\sigma_i)^2 + \sum_{i>r} z_i^2$ is smallest exactly when $z_{r+1} = \cdots = z_n = 0$. □

NumPy's `np.linalg.pinv` and `np.linalg.lstsq` use the SVD in this way.

2.11.5 The condition number

Data are never exact, and computers round every number to about 16 significant digits. How much can such small errors change the solution of a linear system? The singular values give the answer.

Definition 2.11.13 (Condition number)**.** Let A be an $m \times n$ matrix with linearly independent columns, and singular values $\sigma_1 \ge \cdots \ge \sigma_n > 0$. Its condition number is $\kappa(A) = \frac{\sigma_1}{\sigma_n} \ge 1$.

For an invertible $n \times n$ matrix, $A^{-1} = V\Sigma^{-1}U^\top$ has the singular values $\frac{1}{\sigma_i}$, so $\kappa(A) = \|A\|_2\|A^{-1}\|_2$ by Proposition 2.11.6. For a symmetric matrix $A = Q\Lambda Q^\top$, the singular values are the numbers $|\lambda_i|$ (as $A^\top A = Q\Lambda^2 Q^\top$), so $\kappa(A) = \frac{\max_i |\lambda_i|}{\min_i |\lambda_i|}$. Orthogonal matrices have $\kappa(Q) = 1$, the best possible value.

Theorem 2.11.14 (Sensitivity of linear systems)**.** *Let A be invertible, $A\mathbf{x} = \mathbf{b} \neq \mathbf{0}$, and $A(\mathbf{x} + \boldsymbol{\delta}\mathbf{x}) = \mathbf{b} + \boldsymbol{\delta}\mathbf{b}$. Then*

$$\frac{\|\boldsymbol{\delta}\mathbf{x}\|}{\|\mathbf{x}\|} \leq \kappa(A)\,\frac{\|\boldsymbol{\delta}\mathbf{b}\|}{\|\mathbf{b}\|},$$

with equality for $\mathbf{b} = \mathbf{u}_1$ and $\boldsymbol{\delta}\mathbf{b} = t\mathbf{u}_n$.

Proof. $\boldsymbol{\delta}\mathbf{x} = A^{-1}\boldsymbol{\delta}\mathbf{b}$, so $\|\boldsymbol{\delta}\mathbf{x}\| \leq \|A^{-1}\|_2\|\boldsymbol{\delta}\mathbf{b}\| = \frac{1}{\sigma_n}\|\boldsymbol{\delta}\mathbf{b}\|$, and $\|\mathbf{b}\| = \|A\mathbf{x}\| \leq \sigma_1\|\mathbf{x}\|$. Dividing the first inequality by $\|\mathbf{x}\| \geq \frac{1}{\sigma_1}\|\mathbf{b}\|$ gives the claim. For $\mathbf{b} = \mathbf{u}_1$, $\mathbf{x} = \frac{1}{\sigma_1}\mathbf{v}_1$, and for $\boldsymbol{\delta}\mathbf{b} = t\mathbf{u}_n$, $\boldsymbol{\delta}\mathbf{x} = \frac{t}{\sigma_n}\mathbf{v}_n$, so both inequalities are equalities. □

A relative error in $\mathbf{b}$ can thus be amplified by the factor $\kappa(A)$ in the solution. As a rule of thumb, solving a system with a condition number of 10^k in floating-point arithmetic loses about k of the 16 significant digits.

Example 2.11.15. Let $A = \begin{pmatrix} 1 & 1 \\ 1 & 1.001 \end{pmatrix}$, whose columns are almost parallel. Then $\kappa(A) \approx 4002$. The system $A\mathbf{x} = (2, 2.001)^\top$ has the solution $\mathbf{x} = (1, 1)^\top$, but changing the right side by 0.001 to $(2, 2.002)^\top$ gives $\mathbf{x} = (0, 2)^\top$: a relative change of 0.035% in $\mathbf{b}$ changes $\mathbf{x}$ by 100%.

The condition number explains how least squares problems should be solved. The singular values of $A^\top A$ are σ_i^2, so

$$\kappa(A^\top A) = \kappa(A)^2.$$

Solving the normal equations $A^\top A\mathbf{w} = A^\top\mathbf{y}$ squares the condition number, and may lose twice as many digits as necessary. With a QR decomposition $A = QR$ (Proposition 2.5.9), the normal equations become $R^\top R\mathbf{w} = R^\top Q^\top\mathbf{y}$, i.e., $R\mathbf{w} = Q^\top\mathbf{y}$, and $\kappa(R) = \kappa(A)$, since multiplying by Q does not change lengths. This is why `np.linalg.lstsq` uses the SVD and never forms $A^\top A$. Regularization also helps: ridge regression solves $(A^\top A + \lambda I)\mathbf{w} = A^\top\mathbf{y}$, whose matrix has the condition number $\frac{\sigma_1^2+\lambda}{\sigma_n^2+\lambda} \leq \kappa(A)^2$. In Lecture 3, the condition number of the Hessian determines how fast gradient descent converges.

2.11.6 Principal component analysis

Principal component analysis (PCA) is the most widely used method to reduce the dimension of data. Let $\mathbf{x}_1, \ldots, \mathbf{x}_N \in \mathbb{R}^d$ be data points, which we assume to be centered, i.e., $\sum_i \mathbf{x}_i = \mathbf{0}$ (otherwise, subtract the mean from every data point first). Let X be the $N \times d$ data matrix whose rows are $\mathbf{x}_1^\top, \ldots, \mathbf{x}_N^\top$, and let

$$S = \frac{1}{N}X^\top X = \frac{1}{N}\sum_{i=1}^{N}\mathbf{x}_i\mathbf{x}_i^\top$$

be the (sample) covariance matrix, which is symmetric positive semidefinite.

We look for a k-dimensional subspace U such that the projected data points $\pi_U(\mathbf{x}_i)$ are as close as possible to the original ones, i.e., such that the reconstruction error $\sum_i\|\mathbf{x}_i - \pi_U(\mathbf{x}_i)\|^2$ is as small as possible. By Pythagoras, $\|\mathbf{x}_i\|^2 = \|\pi_U(\mathbf{x}_i)\|^2 + \|\mathbf{x}_i - \pi_U(\mathbf{x}_i)\|^2$, so equivalently, we look for the subspace for which the projected data has the largest spread $\sum_i\|\pi_U(\mathbf{x}_i)\|^2$.

Theorem 2.11.16 (PCA)**.** *Let $\mathbf{q}_1, \dots, \mathbf{q}_d$ be orthonormal eigenvectors of S with eigenvalues $\lambda_1 \geq \cdots \geq \lambda_d \geq 0$. Among all k-dimensional subspaces U of $\mathbb{R}^d$, the reconstruction error is smallest for $U = \operatorname{span}\{\mathbf{q}_1, \dots, \mathbf{q}_k\}$, and then*

$$\frac{1}{N}\sum_{i=1}^{N} \|\mathbf{x}_i - \pi_U(\mathbf{x}_i)\|^2 = \lambda_{k+1} + \cdots + \lambda_d.$$

Proof. Let $\mathbf{b}_1, \dots, \mathbf{b}_k$ be an orthonormal basis of U. Then $\pi_U(\mathbf{x}_i) = \sum_j (\mathbf{b}_j^\top \mathbf{x}_i)\mathbf{b}_j$, so $\|\pi_U(\mathbf{x}_i)\|^2 = \sum_j (\mathbf{b}_j^\top \mathbf{x}_i)^2$ by Pythagoras, and

$$\frac{1}{N}\sum_{i=1}^{N} \|\pi_U(\mathbf{x}_i)\|^2 = \sum_{j=1}^{k} \frac{1}{N}\sum_{i=1}^{N} (\mathbf{b}_j^\top \mathbf{x}_i)^2 = \sum_{j=1}^{k} \frac{1}{N}\|X\mathbf{b}_j\|^2 = \sum_{j=1}^{k} \mathbf{b}_j^\top S \mathbf{b}_j,$$

which by Proposition 2.10.8 is at most $\lambda_1 + \cdots + \lambda_k$, with equality for $\mathbf{b}_j = \mathbf{q}_j$. For $k = 1$, this is just Proposition 2.10.7. The error formula follows from Pythagoras, since $\frac{1}{N}\sum_i \|\mathbf{x}_i\|^2 = \frac{1}{N}\operatorname{tr}(X^\top X) = \operatorname{tr}(S) = \lambda_1 + \cdots + \lambda_d$. □

The vectors $\mathbf{q}_1, \mathbf{q}_2, \dots$ are called the principal components, or principal directions, of the data, and $\lambda_i = \mathbf{q}_i^\top S \mathbf{q}_i$ is the variance of the data in the direction $\mathbf{q}_i$. The fraction $\frac{\lambda_1 + \cdots + \lambda_k}{\lambda_1 + \cdots + \lambda_d}$ is the proportion of the variance explained by the first k principal components. In practice, PCA is computed with the SVD of the data matrix: if $X = \tilde{U}\tilde{\Sigma}\tilde{V}^\top$, then

$$S = \frac{1}{N}X^\top X = \tilde{V}\,\frac{\tilde{\Sigma}^\top \tilde{\Sigma}}{N}\,\tilde{V}^\top,$$

so the principal components are the right singular vectors of X, and $\lambda_i = \sigma_i^2/N$. The coordinates of the data in the principal directions are the entries of $X\tilde{V}$.

Example 2.11.17. Consider the four centered data points $(2,2)$, $(-2,-2)$, $(1,-1)$ and $(-1,1)$ in $\mathbb{R}^2$. Then

$$X^\top X = \begin{pmatrix} 10 & 6 \\ 6 & 10 \end{pmatrix}, \qquad S = \frac{1}{4}X^\top X = \begin{pmatrix} 2.5 & 1.5 \\ 1.5 & 2.5 \end{pmatrix},$$

with eigenvalues 4 and 1 and eigenvectors $\mathbf{q}_1 = \frac{1}{\sqrt{2}}(1,1)^\top$ and $\mathbf{q}_2 = \frac{1}{\sqrt{2}}(1,-1)^\top$. The first principal component explains $\frac{4}{4+1} = 80\%$ of the variance. Projecting onto $\operatorname{span}\{\mathbf{q}_1\}$ keeps the first two points, and maps the last two to $\mathbf{0}$; the average reconstruction error is $\frac{1}{4}(0+0+2+2) = 1 = \lambda_2$, as predicted by Theorem 2.11.16.

2.12 Key Takeaways

- A norm measures length: it is positive definite, absolutely homogeneous, and satisfies the triangle inequality. Important examples on $\mathbb{R}^n$ are the ℓ_1, ℓ_2 and ℓ_∞ norms, whose unit balls are a diamond, a ball and a cube.
- An inner product is a symmetric, positive definite, bilinear function. On an n-dimensional space with a fixed basis, inner products correspond to symmetric positive definite matrices: $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top A \mathbf{y}$.
- Every inner product induces a norm $\|x\| = \sqrt{\langle x, x \rangle}$ and a distance $d(x,y) = \|x - y\|$. The Cauchy–Schwarz inequality $|\langle x, y \rangle| \leq \|x\|\|y\|$ gives the triangle inequality and allows us to define angles by $\cos\omega = \frac{\langle x,y \rangle}{\|x\|\|y\|}$.
- Vectors are orthogonal if their inner product is 0. Orthogonal matrices ($Q^\top Q = I$) are exactly the matrices that preserve dot products, lengths and angles.

- In an orthonormal basis, coordinates are inner products: $v = \sum_i \langle v, b_i \rangle b_i$. The Gram–Schmidt process turns any basis into an orthonormal basis, and gives the QR decomposition.
- Every subspace U of a finite-dimensional inner product space has an orthogonal complement with $V = U \oplus U^\perp$. For a matrix, $\operatorname{null}(A) = \operatorname{row}(A)^\perp$ and $\operatorname{null}(A^\top) = \operatorname{col}(A)^\perp$.
- A hyperplane $\{\mathbf{x} : \mathbf{w}^\top \mathbf{x} = c\}$ has normal vector $\mathbf{w}$; linear classifiers use hyperplanes as decision boundaries, and the distance of $\mathbf{x}$ to the hyperplane is $\frac{|\mathbf{w}^\top \mathbf{x} - c|}{\|\mathbf{w}\|}$.
- The orthogonal projection onto $U = \operatorname{col}(B)$ is $\pi_U(\mathbf{x}) = B(B^\top B)^{-1} B^\top \mathbf{x}$; it is the point of U closest to $\mathbf{x}$. Least squares regression projects the target vector onto the column space of the design matrix, which leads to the normal equations $A^\top A \mathbf{w} = A^\top \mathbf{y}$.
- Rotations are orthogonal matrices with determinant 1. They preserve lengths, angles and orientation; rotations of the plane commute, but rotations of $\mathbb{R}^3$ do not.
- Spectral theorem: a symmetric matrix has real eigenvalues and an orthonormal basis of eigenvectors, $A = Q \Lambda Q^\top = \sum_i \lambda_i \mathbf{q}_i \mathbf{q}_i^\top$. It is positive definite if and only if all eigenvalues are positive, and $\max_{\|\mathbf{x}\|=1} \mathbf{x}^\top A \mathbf{x} = \lambda_{\max}$; more generally, λ_k is the max-min of the Rayleigh quotient over k-dimensional subspaces (Courant–Fischer), so eigenvalues of symmetric matrices move by at most ε under perturbations with eigenvalues in $[-\varepsilon, \varepsilon]$ (Weyl).
- A symmetric matrix is positive definite if and only if $A = LL^\top$ with a lower triangular L with positive diagonal (Cholesky), the standard way to solve systems with such matrices and to compute $\det(A)$. Positive semidefinite matrices have a unique positive semidefinite square root $A^{1/2} = Q \Lambda^{1/2} Q^\top$.
- The Laplacian $L = D - W$ of a weighted graph satisfies $\mathbf{x}^\top L \mathbf{x} = \sum_{i<j} w_{ij}(x_i - x_j)^2$, so it is positive semidefinite. The multiplicity of its eigenvalue 0 is the number of connected components, and its second smallest eigenvalue λ_2 is a lower bound for the ratio cut of every split of the vertices. Spectral clustering splits the vertices by the signs of the Fiedler vector $\mathbf{q}_2$.
- Every matrix has a singular value decomposition $A = U \Sigma V^\top = \sum_{i=1}^{r} \sigma_i \mathbf{u}_i \mathbf{v}_i^\top$: rotate, scale the axes by the singular values, rotate. The σ_i^2 are the eigenvalues of $A^\top A$, $r = \operatorname{rank}(A)$, $\|A\|_2 = \sigma_1$ and $\|A\|_F^2 = \sum_i \sigma_i^2$.
- The condition number $\kappa(A) = \sigma_1 / \sigma_n$ bounds the amplification of relative errors in linear systems, $\frac{\|\boldsymbol{\delta}\mathbf{x}\|}{\|\mathbf{x}\|} \le \kappa(A) \frac{\|\boldsymbol{\delta}\mathbf{b}\|}{\|\mathbf{b}\|}$. Since $\kappa(A^\top A) = \kappa(A)^2$, least squares problems are solved with QR or the SVD rather than the normal equations.
- Keeping the k largest singular values gives the best rank-k approximation (Eckart–Young), with applications to compression and recommender systems. PCA projects centered data onto the eigenvectors of the covariance matrix with the largest eigenvalues, i.e., onto the top right singular vectors of the data matrix.

2.13 Exercises

1. Let $\mathbf{x} = (1, -2, 2)^\top$ and $\mathbf{y} = (3, 0, 1)^\top$. Compute $\|\mathbf{x}\|_1$, $\|\mathbf{x}\|_2$ and $\|\mathbf{x}\|_\infty$, and the distance between $\mathbf{x}$ and $\mathbf{y}$ in each of the three norms.
2. Show that $\|\mathbf{x}\|_2 \le \sqrt{n}\, \|\mathbf{x}\|_\infty$ for all $\mathbf{x} \in \mathbb{R}^n$. For which vectors $\mathbf{x}$ does equality hold in $\|\mathbf{x}\|_1 \le \sqrt{n}\, \|\mathbf{x}\|_2$?
3. Show that the ℓ_∞ norm on $\mathbb{R}^2$ does not satisfy the parallelogram law, and conclude that it does not come from an inner product.

4. Which of the following matrices are symmetric positive definite? For each one that is, write down the corresponding inner product $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top A \mathbf{y}$ on $\mathbb{R}^2$ explicitly.

$$\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \qquad \begin{pmatrix} 4 & -2 \\ -2 & 1 \end{pmatrix}, \qquad \begin{pmatrix} 3 & 1 \\ 1 & 1 \end{pmatrix}.$$

5. Find the angle between the vectors: (a) $(1,1,0)^\top$ and $(1,0,1)^\top$; (b) $(1,2,3)^\top$ and $(-3,0,1)^\top$; (c) $(1,0)^\top$ and $(-1,\sqrt{3})^\top$.

6. (a) Find all vectors in $\mathbb{R}^3$ that are orthogonal to both $(1,1,1)^\top$ and $(1,-1,1)^\top$. (Hint: this is the null space of a matrix.)

   (b) The planes $x + y + z = 3$ and $x - y + z = 5$ intersect in a line ℓ. Find an equation of the plane through the origin that is perpendicular to ℓ.

7. Apply the Gram–Schmidt process to $\mathbf{v}_1 = (1,2,2)^\top$ and $\mathbf{v}_2 = (1,0,1)^\top$. Then find a third vector that extends the result to an orthonormal basis of $\mathbb{R}^3$, and write $(1,1,1)^\top$ in this basis. Check Parseval's identity.

8. Find a basis of the orthogonal complement of $U = \operatorname{span}\{(1,0,1,0)^\top, (0,1,0,1)^\top\}$ in $\mathbb{R}^4$.

9. Let $\mathbf{x} = (1,0,2)^\top$. Compute the orthogonal projection of $\mathbf{x}$ onto (a) the line spanned by $(1,1,0)^\top$, and (b) the plane $x_1 + x_2 + x_3 = 0$. Write down the projection matrices of both projections.

10. Find the point of the plane $2x + y + 3z = 3$ that is closest to the origin, and the distance from the origin to this plane.

11. Let P be a symmetric $n \times n$ matrix with $P^2 = P$. Show that P is the matrix of the orthogonal projection onto $\operatorname{col}(P)$. (Hint: show that $\mathbf{x} - P\mathbf{x} \in \operatorname{col}(P)^\perp = \operatorname{null}(P^\top)$.)

12. Let $\mathbf{u} \in \mathbb{R}^n$ be a unit vector, and let $H = I_n - 2\mathbf{u}\mathbf{u}^\top$.

    (a) Show that H is symmetric and orthogonal, and that $H^2 = I_n$.

    (b) Show that $H\mathbf{u} = -\mathbf{u}$, and that $H\mathbf{x} = \mathbf{x}$ for every $\mathbf{x} \perp \mathbf{u}$. Describe H geometrically. (Such matrices are called Householder reflections; `np.linalg.qr` uses them.)

13. Find the polynomial p of degree at most 1 that minimizes $\int_{-1}^{1} \left(x^3 - p(x)\right)^2 dx$, by projecting x^3 onto $\operatorname{span}\{1, x\}$ in $C[-1,1]$. Compare with Example 2.7.3.

14. Find the least squares line for the data points $(0,1)$, $(1,1)$ and $(2,3)$, and check that the residual vector is orthogonal to the columns of the design matrix.

15. Show that the product of two orthogonal 2×2 matrices with determinant -1 (for example, two reflections) is a rotation. Compute $R_1(\theta)^\top R_1(\theta)$ and $\det R_1(\theta)$.

16. Use the Cauchy–Schwarz inequality to show that $(x_1 + \cdots + x_n)^2 \le n(x_1^2 + \cdots + x_n^2)$. Conclude that if $a, b, c \ge 0$ and $a + b + c = 1$, then $a^2 + b^2 + c^2 \ge \frac{1}{3}$. When does equality hold?

17. Find the spectral decomposition $A = Q\Lambda Q^\top$ of $A = \begin{pmatrix} 5 & 2 \\ 2 & 2 \end{pmatrix}$, and write A as $\lambda_1 \mathbf{q}_1 \mathbf{q}_1^\top + \lambda_2 \mathbf{q}_2 \mathbf{q}_2^\top$.

    Is A positive definite? Conversely, show that every matrix of the form $Q\Lambda Q^\top$, with Q orthogonal and Λ diagonal, is symmetric.

18. Find the largest and the smallest value of $\mathbf{x}^\top A \mathbf{x}$ over all unit vectors $\mathbf{x} \in \mathbb{R}^2$, for $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, and the unit vectors where they are attained.

19. (Cholesky decomposition and square roots.)

    (a) Find the Cholesky decomposition of $\begin{pmatrix} 4 & 2 \\ 2 & 10 \end{pmatrix}$. What goes wrong when you apply the algorithm to $\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$, and why?

    (b) Compute the square root of $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$.

    (c) Let A be symmetric positive definite with the Cholesky decomposition $A = LL^\top$. Show that $Q = L^{-1}A^{1/2}$ is orthogonal, so that $A^{1/2} = LQ$. More generally, show that if $A = BB^\top = CC^\top$ with invertible matrices B and C, then $B^{-1}C$ is orthogonal.

20. (Courant–Fischer.) Let A be a symmetric $n \times n$ matrix with eigenvalues $\lambda_1 \geq \cdots \geq \lambda_n$, and let A' be its upper left $(n-1) \times (n-1)$ submatrix, with eigenvalues $\mu_1 \geq \cdots \geq \mu_{n-1}$.

    (a) Show that $\lambda_k \geq \mu_k$ for $k \leq n-1$. (Hint: the subspaces of $\mathbb{R}^{n-1}$ are subspaces of $\mathbb{R}^n$ after appending a zero coordinate.)

    (b) Apply (a) to $-A$ to show that $\mu_k \geq \lambda_{k+1}$. So the eigenvalues of A' interlace with those of A (Cauchy's interlacing theorem).

    (c) Check the interlacing for the matrix $J + 2I_3$ of Example 2.10.2.

21. Let L be the Laplacian of a graph with weights w_{ij}.

    (a) Show that $L = \sum_{i<j} w_{ij}(\mathbf{e}_i - \mathbf{e}_j)(\mathbf{e}_i - \mathbf{e}_j)^\top$, and use this to give another proof of Proposition 2.10.13.

    (b) Find the Laplacian and its eigenvalues for the star graph with the center 1 and the leaves $2, \ldots, n$, where $n \geq 3$ and all weights are 1. (Hint: consider the vectors with $x_1 = 0$ whose entries add up to 0, and the vector $(n-1, -1, \ldots, -1)^\top$.) Why is the Fiedler vector not unique, and what does this say about clusters in a star?

22. Compute an SVD of (a) $\begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix}$ and (b) $\begin{pmatrix} 2 & 2 \\ 1 & 1 \end{pmatrix}$. For (b), write A as $\sigma_1 \mathbf{u}_1 \mathbf{v}_1^\top$, and compute the pseudo-inverse A^+.

23. Show that:

    (a) all singular values of an orthogonal matrix are equal to 1;

    (b) the singular values of a symmetric matrix are the absolute values of its nonzero eigenvalues;

    (c) $A^\top A$ and $AA^\top$ have the same nonzero eigenvalues;

    (d) $\|A\|_2 = \|A^\top\|_2$.

24. (Condition numbers.)

    (a) Show that $\kappa(cA) = \kappa(A)$ for $c \neq 0$, that $\kappa(QA) = \kappa(A)$ for an orthogonal matrix Q, and that $\kappa(A^{-1}) = \kappa(A)$ for an invertible matrix A.

    (b) Compute the condition numbers of $\begin{pmatrix} 1 & 0 \\ 0 & \varepsilon \end{pmatrix}$ for $0 < \varepsilon < 1$, and of the shear $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

    (c) Show that $\kappa(A^\top A + \lambda I) = \frac{\sigma_1^2 + \lambda}{\sigma_n^2 + \lambda}$ for $\lambda > 0$, and that this is a decreasing function of λ if $\sigma_1 > \sigma_n$.

25. Carry out PCA by hand for the four data points $(3,3)$, $(-3,-3)$, $(1,-1)$, $(-1,1)$: compute the covariance matrix, the principal components, and the proportion of the variance explained by the first principal component. Project the data onto the first principal component, and compute the average reconstruction error.

2.14 Programming Exercises

As in Lecture 1, implement the algorithms yourself with NumPy, and use NumPy's built-in routines only to check your results. You need Matplotlib for P1, P5, P7, P10 and P11. Start each file with

```
import numpy as np
rng = np.random.default_rng(0)  # random number generator for test data
```

P1. Norms and unit balls. Write a function `p_norm(x, p)` for $p \geq 1$ and $p = \infty$ (use `np.inf`), and compare it with `np.linalg.norm(x, ord=p)`.

- For 10 000 random vectors in $\mathbb{R}^5$, check the inequalities $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1 \leq \sqrt{5}\,\|\mathbf{x}\|_2$.
- Plot the unit circles $\{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\|_p = 1\}$ for $p = 1, 1.5, 2, 4, \infty$ in one picture, as in Figure 2.1. (Hint: for each angle t, rescale the vector $(\cos t, \sin t)$ to have p-norm 1.)

P2. Inner products and angles. Write a function `inner(x, y, A)` that computes $\mathbf{x}^\top A\mathbf{y}$, and functions `norm_A(x, A)` and `angle_A(x, y, A)` for the induced norm and angle.

- Check Example 2.4.4: the angle between $(1,1)$ and $(-1,1)$ is $90°$ for $A = I_2$, but about $109.5°$ for $A = \operatorname{diag}(2,1)$.
- Write a function `is_spd(A)` that tests whether a symmetric 2×2 matrix is positive definite, using Proposition 2.2.7. For larger matrices, NumPy's `np.linalg.cholesky(A)` succeeds exactly for symmetric positive definite matrices (Theorem 2.2.10) and raises `np.linalg.LinAlgError` otherwise; use it to test the matrices of Exercise 4.
- For random vectors, check the Cauchy–Schwarz inequality, and the parallelogram law for the ℓ_2 norm (but not for the ℓ_1 norm).

P3. Gram–Schmidt, and how it can fail. Implement the classical Gram–Schmidt process `gram_schmidt(V)`, which returns a matrix Q whose columns are the orthonormal vectors obtained from the columns of `V`, exactly as in Theorem 2.5.6.

- Check your function on Example 2.5.7, and compute $R = Q^\top V$; it should be upper triangular, with $V = QR$.
- The modified Gram–Schmidt process subtracts the components one at a time, from the updated vector: for each $i < j$, replace $\mathbf{u}_j$ by $\mathbf{u}_j - \langle \mathbf{u}_j, \mathbf{e}_i \rangle \mathbf{e}_i$. In exact arithmetic, both versions give the same result. Implement it as well.
- For the Hilbert matrices H_n with entries $\frac{1}{i+j-1}$ (whose columns are nearly linearly dependent) and $n = 6, 8, 10, 12$, compute the loss of orthogonality `np.linalg.norm(Q.T @ Q - np.eye(n))` for the classical version, the modified version, and `np.linalg.qr`. What do you observe?

P4. Projections. Write a function `project(x, B)` that returns the orthogonal projection of `x` onto the column space of `B`, by solving the normal equations, and a function `projection_matrix(B)`.

- Check your functions on Example 2.8.6.

- For a random 6×3 matrix B and a random $\mathbf{x} \in \mathbb{R}^6$, check that $P^2 = P$, $P^\top = P$, and that $\mathbf{x} - P\mathbf{x}$ is orthogonal to every column of B.
- Check Theorem 2.8.8 numerically: compare $\|\mathbf{x} - P\mathbf{x}\|$ with $\|\mathbf{x} - B\boldsymbol{\lambda}\|$ for 1000 random vectors $\boldsymbol{\lambda} \in \mathbb{R}^3$.
- Orthonormalize the columns of B with your function from P3, and check that $QQ^\top$ is the same projection matrix.

P5. Polynomial regression. Generate 20 noisy data points

```
x = np.linspace(-1, 1, 20)
y = np.sin(3 * x) + rng.normal(0, 0.1, size=20)
```

- For $d = 1, 3, 5, 15$, fit a polynomial of degree d by least squares: build the design matrix with columns $1, x, \ldots, x^d$ and solve the normal equations. Compare your coefficients with `np.linalg.lstsq`.
- Plot the data and the four fitted polynomials. Which degree fits the training data best? Which one would you trust for new data? (This is the phenomenon of overfitting.)
- Check that for each d, the residual vector is orthogonal to all columns of the design matrix.

P6. Nearest neighbors with different metrics. Generate two classes of points in $\mathbb{R}^2$:

```
X0 = rng.normal([0, 0], [1.0, 0.3], size=(100, 2))   # class 0
X1 = rng.normal([1.5, 1.0], [1.0, 0.3], size=(100, 2))  # class 1
```

and split the 200 points randomly into 150 training points and 50 test points.

- Implement the 1-nearest neighbor classifier with the ℓ_1, ℓ_2 and ℓ_∞ distances, and with the cosine similarity (largest similarity instead of smallest distance). Compare their accuracies on the test points.
- Multiply the second coordinate of all points by 1000 (as if it were measured in different units) and repeat. What happens to the accuracies of the distance-based and the cosine-based classifiers, and why? What happens if you standardize each feature (subtract its mean, divide by its standard deviation) first?
- Rotate all points by a random rotation R_θ and repeat. Which classifiers give exactly the same predictions, and why?

P7. Rotations. Write functions `R1(t)`, `R2(t)` and `R3(t)` that return the rotation matrices of Section 2.9, and a function `givens(n, i, j, t)`.

- Check that all of them are orthogonal with determinant 1, and that they preserve the lengths of random vectors.
- Check that `R1(t) @ R3(s)` and `R3(s) @ R1(t)` differ for most angles, while `R3(t) @ R3(s)` equals `R3(t + s)`.
- Let `Q, R = np.linalg.qr(rng.normal(size=(3, 3)))`. Then `Q` is orthogonal, but its determinant may be -1. How can you turn it into a rotation by changing the sign of one column? Generate many random rotations this way, apply them to $\mathbf{e}_3$, and plot the resulting points on the sphere (Matplotlib can make 3D scatter plots).

P8. The perceptron and its margin. The perceptron algorithm learns a linear classifier $f(\mathbf{x}) = \operatorname{sign}(\mathbf{w}^\top \mathbf{x} + b)$ from labeled data $(\mathbf{x}_i, y_i)$ with $y_i \in \{+1, -1\}$: start with $\mathbf{w} = \mathbf{0}$ and $b = 0$, and repeatedly go through the data; whenever a point is misclassified, i.e., $y_i(\mathbf{w}^\top \mathbf{x}_i + b) \le 0$, update $\mathbf{w} \leftarrow \mathbf{w} + y_i \mathbf{x}_i$ and $b \leftarrow b + y_i$. Stop when all points are classified correctly.

- Generate linearly separable data, e.g., 50 points on each side of the line $\mathbf{x}_1 + \mathbf{x}_2 = 2$ (discard points that are closer than 0.2 to the line), and run the perceptron.
- Using Theorem 2.8.13, compute the margin of the learned classifier, i.e., the smallest distance of a training point to the decision boundary. Compare it with the margin of the classifier $\mathbf{w} = (1, 1)$, $b = -2$.
- Count the number of updates the perceptron needs. How does it change when you discard points closer than 0.5 (instead of 0.2) to the line?

P9. The spectral theorem in practice. For symmetric matrices, use `np.linalg.eigh`, which returns real eigenvalues in increasing order and orthonormal eigenvectors.

- For a random symmetric matrix `A = B + B.T` (with `B = rng.normal(size=(5, 5))`), check that `Q.T @ Q` is close to I_5 and that `Q @ np.diag(lam) @ Q.T` is close to A. Check the spectral decomposition $A = \sum_i \lambda_i \mathbf{q}_i \mathbf{q}_i^\top$ as well.
- Write a function `is_spd(A)` based on Corollary 2.10.3, and compare it with your test from P2.
- Check Proposition 2.10.7 numerically: evaluate the Rayleigh quotient $\frac{\mathbf{x}^\top A \mathbf{x}}{\mathbf{x}^\top \mathbf{x}}$ at 10 000 random vectors, and compare the largest and smallest values you find with the largest and smallest eigenvalues.

P10. Image compression with the SVD. Load a grayscale image as a 2D array `A`, for example with `A = plt.imread("photo.png").mean(axis=2)` for a color image (with `import matplotlib.pyplot as plt`), or create one, e.g., from the function $\sin(x)\cos(y) + \frac{x}{10}$ on a 200×300 grid plus some noise.

- Compute `U, s, Vt = np.linalg.svd(A, full_matrices=False)` and plot the singular values on a logarithmic scale. How fast do they decay?
- For $k = 1, 5, 20, 50$, compute the rank-k approximation A_k (Theorem 2.11.7) and display it as an image. Compare `np.linalg.norm(A - A_k)` with $\sqrt{\sigma_{k+1}^2 + \cdots + \sigma_r^2}$, and compute the storage ratio $\frac{k(m+n+1)}{mn}$.
- Check numerically that $\|A\|_2 = \sigma_1$ (`np.linalg.norm(A, 2)`) and $\|A\|_F^2 = \sum_i \sigma_i^2$.

P11. Principal component analysis. Generate 500 data points from a correlated Gaussian distribution:

```
C = np.array([[3.0, 1.5], [1.5, 1.0]])  # covariance matrix
X = rng.multivariate_normal([2, -1], C, size=500)
```

- Center the data, and compute the principal components in two ways: as eigenvectors of the covariance matrix $S = \frac{1}{N} X^\top X$ (with `np.linalg.eigh`), and as right singular vectors of X (with `np.linalg.svd`). Check that they agree (up to sign), and that $\lambda_i = \sigma_i^2 / N$. How do they compare with the eigenvectors of C?
- Plot the data together with the two principal directions, scaled by $\sqrt{\lambda_i}$. What proportion of the variance does the first principal component explain?
- Project the data onto the first principal component, plot the projected points, and check that the average reconstruction error is λ_2 (Theorem 2.11.16).
- Generate 500 points in $\mathbb{R}^{10}$ that lie close to a 2-dimensional subspace, for example

```
X = rng.normal(size=(500, 2)) @ rng.normal(size=(2, 10))
X = X + 0.05 * rng.normal(size=(500, 10))  # small noise
```

Plot the 10 eigenvalues of the covariance matrix. How can you see from this plot that the data is essentially 2-dimensional?

P12. Spectral clustering. Generate 100 points on each of two concentric circles with the radii 1 and 3:

```
t = rng.uniform(0, 2 * np.pi, 200)
r = np.repeat([1.0, 3.0], 100)
X = np.c_[r * np.cos(t), r * np.sin(t)] + 0.1 * rng.normal(size=(200, 2))
```

- Build the weights $w_{ij} = e^{-\|\mathbf{x}_i - \mathbf{x}_j\|^2/(2s^2)}$ with $s = 0.5$ (and $w_{ii} = 0$), the Laplacian L (Definition 2.10.12), and its eigenvalues and eigenvectors with `np.linalg.eigh`. Check that the smallest eigenvalue is 0, with the eigenvector $\frac{1}{\sqrt{n}}\mathbf{1}$ (up to sign).
- Cluster the points by the signs of the Fiedler vector, and plot them in two colors. Does spectral clustering find the two circles? Why would a method that only splits the plane by a line fail here?
- Repeat with $s = 2$ and $s = 0.2$. How do the second and third smallest eigenvalues change, and how does the clustering change? Relate your observations to Theorem 2.10.14.

P13. Conditioning and least squares.

- Write a function `cholesky(A)` following the proof of Theorem 2.2.10, and compare it with `np.linalg.cholesky` on a random matrix $B^\top B + I$. Use it to solve $A\mathbf{x} = \mathbf{b}$ by forward and back substitution.
- The Hilbert matrix H_n has the entries $\frac{1}{i+j-1}$. For $n = 4, 6, \ldots, 14$, compute $\kappa(H_n)$ with `np.linalg.cond`, solve $H_n\mathbf{x} = H_n\mathbf{1}$, and plot the error $\|\mathbf{x} - \mathbf{1}\|$ against $\kappa(H_n)$ on logarithmic axes. How many correct digits does the rule of thumb after Theorem 2.11.14 predict?
- Fit a polynomial of degree 12 to 50 points $(t_i, \sin 4t_i)$ with t_i equally spaced in $[0, 1]$, using the columns $1, t, \ldots, t^{12}$ of a design matrix A. Compare $\kappa(A)$ with $\kappa(A^\top A)$, and compare the residuals of three solutions: the normal equations with `np.linalg.solve`, the QR decomposition, and `np.linalg.lstsq`.

Lecture 3

Vector Calculus and Optimization

Introduction

Almost every machine learning model is trained by solving an optimization problem. A model has parameters, such as the weights of a linear model or the millions of weights of a neural network, which we collect in a vector $\boldsymbol{\theta}$. A loss function $L(\boldsymbol{\theta})$ measures how badly the model with these parameters fits the training data, and training means finding parameters with a small loss. There are far too many parameters to try out all possible values. Instead, we start with some parameters and improve them step by step, and for this we need to know how the loss changes when we change the parameters slightly. This information is contained in the derivative of L. Vector calculus is the study of derivatives and integrals of functions of several variables. In this lecture:

- we extend the derivative to functions $f : \mathbb{R}^n \to \mathbb{R}^m$: partial derivatives, gradients and Jacobian matrices;
- we prove the chain rule, which says that the derivative of a composition of functions is the product of their Jacobian matrices; it is the basis of backpropagation, the algorithm that computes the gradients of neural networks;
- we compute gradients with respect to vectors and matrices, such as the weight matrices of a neural network;
- we use second derivatives (the Hessian matrix) and Taylor's theorem to describe the shape of a function near a point;
- we apply all of this to optimization: critical points, gradient descent with its convergence rates and line search, Newton's method, Gauss–Newton for nonlinear least squares, Hessian-vector products, convexity and strong convexity, subgradients and the proximal gradient method for nonsmooth losses such as the lasso, and optimization with constraints (Lagrange multipliers, the KKT conditions, duality, convex conjugates, and linear and quadratic programs);
- we study vector fields with their divergence and curl, and integrals over regions of $\mathbb{R}^n$: Fubini's theorem, the change of variables formula, line integrals, and the divergence theorem, which leads to integration by parts in $\mathbb{R}^n$; these are the tools for probability densities in Lecture 4 and for generative models;
- we solve ordinary differential equations (ODEs) $\mathbf{x}' = F(\mathbf{x})$, in particular linear systems, with the eigenvalues and eigenvectors of Lecture 1, and see that gradient descent is Euler's method for the gradient flow $\mathbf{x}' = -\nabla f(\mathbf{x})$, which explains its step size limit;

- we study how the flow of a vector field changes volumes and probability densities, using the divergence; this is the basis of continuous normalizing flows, and, with noise added, of the stochastic differential equations of Lecture 4.

We use the linear algebra of Lectures 1 and 2 throughout: derivatives are linear maps, given by matrices; the direction of steepest ascent comes from the Cauchy–Schwarz inequality; and the shape of a function near a minimum is described by the eigenvalues of a symmetric matrix. Some results from analysis, such as the existence of maxima on closed and bounded sets, the implicit function theorem and the existence of solutions of differential equations, are only stated here; Appendix A on topology, differential topology and functional analysis proves them and puts them in a wider context.

Notation. As in Lectures 1 and 2, vectors of $\mathbb{R}^n$ are written in bold, $\mathbf{x} = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$, to distinguish them from numbers, and they are columns; $\mathbf{e}_1, \dots, \mathbf{e}_n$ is the standard basis of $\mathbb{R}^n$, $\|\cdot\|$ is the Euclidean norm, and log is the natural logarithm.

3.1 Multivariable Functions

Definition 3.1.1. A function $f : \mathbb{R}^n \to \mathbb{R}^m$ is a mapping that takes an input vector $\mathbf{x} \in \mathbb{R}^n$ and produces an output vector $\mathbf{y} = f(\mathbf{x}) \in \mathbb{R}^m$. The components of the output vector are given by $y_i = f_i(\mathbf{x})$ for $i = 1, 2, \dots, m$, where each $f_i : \mathbb{R}^n \to \mathbb{R}$ is a scalar-valued function.

Example 3.1.2. $f(x, y) = x^2 + y^2$ is a function from $\mathbb{R}^2$ to $\mathbb{R}$, which takes a point (x, y) in the plane and maps it to the sum of the squares of its coordinates.

Example 3.1.3 (Functions in machine learning)**.**

1. A linear model with weight vector $\mathbf{w} \in \mathbb{R}^n$ and bias $b \in \mathbb{R}$ is the function $f : \mathbb{R}^n \to \mathbb{R}$, $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x} + b$.

2. A layer of a neural network is a function $f : \mathbb{R}^n \to \mathbb{R}^m$, $f(\mathbf{x}) = \sigma(W\mathbf{x} + \mathbf{b})$, where $W \in \mathbb{R}^{m \times n}$ is a weight matrix, $\mathbf{b} \in \mathbb{R}^m$ is a bias vector, and an activation function $\sigma : \mathbb{R} \to \mathbb{R}$ is applied to each entry of $W\mathbf{x} + \mathbf{b}$. Common activation functions are the sigmoid function $\sigma(t) = \frac{1}{1+e^{-t}}$, the hyperbolic tangent tanh, and the rectified linear unit $\mathrm{ReLU}(t) = \max(0, t)$.

3. The softmax function $\mathrm{softmax} : \mathbb{R}^K \to \mathbb{R}^K$,
$$\mathrm{softmax}(\mathbf{z})_i = \frac{e^{z_i}}{e^{z_1} + \cdots + e^{z_K}}, \qquad i = 1, \dots, K,$$
turns a vector of scores $\mathbf{z}$ into a vector of probabilities: its entries are positive and sum to 1. Classifiers with K classes use it to predict the probability of each class.

4. Given training data $(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_N, y_N)$ with $\mathbf{x}_i \in \mathbb{R}^n$ and $y_i \in \mathbb{R}$, the mean squared error of a linear model without bias,
$$L(\mathbf{w}) = \frac{1}{N} \sum_{i=1}^{N} (\mathbf{w}^\top \mathbf{x}_i - y_i)^2,$$
is a function $L : \mathbb{R}^n \to \mathbb{R}$ of the weights.

The last example shows an important change of perspective. When we use a model, its parameters are fixed and the input $\mathbf{x}$ is the variable. When we train the model, the data are fixed and the parameters are the variables. The derivatives that we need for training are derivatives with respect to the parameters.

Functions are not always defined on all of $\mathbb{R}^n$; for example, $f(\mathbf{x}) = \log(x_1) + x_2$ is only defined when $x_1 > 0$. We then write $f : D \to \mathbb{R}^m$, where $D \subseteq \mathbb{R}^n$ is the domain of f. To take derivatives at a point $\mathbf{a} \in D$, we need to be able to move away from $\mathbf{a}$ a little in every direction without leaving D.

Definition 3.1.4 (Open sets)**.** The open ball with center $\mathbf{a} \in \mathbb{R}^n$ and radius $r > 0$ is $B(\mathbf{a}, r) = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{a}\| < r\}$. A set $D \subseteq \mathbb{R}^n$ is open if for every $\mathbf{a} \in D$, there is an $r > 0$ with $B(\mathbf{a}, r) \subseteq D$.

For example, $\mathbb{R}^n$, every open ball, and the half-plane $\{\mathbf{x} \in \mathbb{R}^2 : x_1 > 0\}$ are open. The closed half-plane $\{\mathbf{x} \in \mathbb{R}^2 : x_1 \geq 0\}$ is not open, since every ball around a point with $x_1 = 0$ contains points with $x_1 < 0$.

3.1.1 Graphs and level sets

Definition 3.1.5 (Graph and level sets)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$. The graph of f is the set

$$\{(\mathbf{x}, f(\mathbf{x})) : \mathbf{x} \in D\} \subseteq \mathbb{R}^{n+1},$$

and for $c \in \mathbb{R}$, the level set of f at level c is the set $\{\mathbf{x} \in D : f(\mathbf{x}) = c\}$. For $n = 2$, the level sets are usually curves, called level curves or contour lines, and a picture of several level curves is a contour plot.

For $n = 2$, the graph is a surface in $\mathbb{R}^3$, a landscape whose height above the point (x, y) is $f(x, y)$. A contour plot is like a topographic map of this landscape. For $n \geq 3$, we cannot draw the graph, but level sets and contour plots of two-dimensional slices are still useful; for example, contour plots of loss functions are used to visualize the training of neural networks.

Example 3.1.6. For $f(x, y) = x^2 + y^2$ (Example 3.1.2), the graph is a bowl-shaped surface, a paraboloid. The level set at level c is the circle of radius $\sqrt{c}$ around the origin if $c > 0$, the single point $(0, 0)$ if $c = 0$, and empty if $c < 0$. The function $g(x, y) = x^2 - y^2$ has a saddle-shaped graph: along the x-axis, it increases away from the origin, and along the y-axis, it decreases. Its level sets are hyperbolas for $c \neq 0$, and the pair of lines $y = \pm x$ for $c = 0$. See Figure 3.1.




Figure 3.1: Left: the graph of $f(x, y) = x^2 + y^2$, a paraboloid. Middle: its level curves for the levels $c = \frac{1}{2}, 1, \frac{3}{2}, \ldots, 4$, which are circles of radius $\sqrt{c}$. They get closer together as we move away from the origin, because f gets steeper. Right: level curves of $g(x, y) = x^2 - y^2$ for $c = \pm\frac{1}{2}, \pm\frac{3}{2}, \pm\frac{5}{2}, \pm\frac{7}{2}$ (red: $c > 0$, blue: $c < 0$), and the level set $c = 0$ (dashed).

3.1.2 Limits and continuity

Definition 3.1.7 (Limits and continuity)**.** Let $f : D \to \mathbb{R}^m$ with $D \subseteq \mathbb{R}^n$, and let $\mathbf{a} \in \mathbb{R}^n$ be a point such that all points $\mathbf{x} \neq \mathbf{a}$ of some ball $B(\mathbf{a}, r)$ lie in D. We say that $f(\mathbf{x})$ converges to $\mathbf{L} \in \mathbb{R}^m$ as $\mathbf{x}$ tends to $\mathbf{a}$, and write $\lim_{\mathbf{x} \to \mathbf{a}} f(\mathbf{x}) = \mathbf{L}$, if for every $\varepsilon > 0$ there is a $\delta > 0$ such that

$$\|f(\mathbf{x}) - \mathbf{L}\| < \varepsilon \qquad \text{for all } \mathbf{x} \in D \text{ with } 0 < \|\mathbf{x} - \mathbf{a}\| < \delta.$$

If $\mathbf{a} \in D$, then f is continuous at $\mathbf{a}$ if $\lim_{\mathbf{x} \to \mathbf{a}} f(\mathbf{x}) = f(\mathbf{a})$. The function f is continuous if it is continuous at every point of D.

By the inequalities $\|\mathbf{x}\|_\infty \le \|\mathbf{x}\|_2 \le \|\mathbf{x}\|_1 \le n\|\mathbf{x}\|_\infty$ of Lecture 2, we could use the ℓ_1 or the ℓ_∞ norm instead of the Euclidean norm in this definition, without changing its meaning. In particular, since

$$|f_i(\mathbf{x}) - L_i| \le \|f(\mathbf{x}) - \mathbf{L}\| \le |f_1(\mathbf{x}) - L_1| + \cdots + |f_m(\mathbf{x}) - L_m|,$$

$f(\mathbf{x})$ converges to $\mathbf{L}$ if and only if every component $f_i(\mathbf{x})$ converges to L_i, and f is continuous if and only if all its components are continuous.

As for functions of one variable, sums, products, quotients (where the denominator is not zero) and compositions of continuous functions are continuous, and the coordinate functions $\mathbf{x} \mapsto x_i$ are continuous, since $|x_i - a_i| \le \|\mathbf{x} - \mathbf{a}\|$. Hence polynomials in $x_1, \ldots, x_n$, linear maps $\mathbf{x} \mapsto A\mathbf{x}$, and functions such as $e^{x_1}\sin(x_2)$ or the functions of Example 3.1.3 are continuous.

Example 3.1.8. Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$, and $f(0, 0) = 0$. On the coordinate axes, f is 0. On the line $y = x$, we have $f(t, t) = \frac{t^2}{2t^2} = \frac{1}{2}$ for $t \neq 0$. So arbitrarily close to the origin, there are points where f is 0 and points where f is $\frac{1}{2}$, and $f(\mathbf{x})$ has no limit as $\mathbf{x} \to \mathbf{0}$. In particular, f is not continuous at the origin, although the functions $t \mapsto f(t, 0)$ and $t \mapsto f(0, t)$ are continuous (they are constant). More generally, f is constant on every line through the origin: $f(t, mt) = \frac{m}{1+m^2}$ for $t \neq 0$.

The lesson of this example is that to understand a function near a point, it is not enough to look at it along the coordinate axes. We will meet this function again when we discuss derivatives.

3.2 Derivatives of Functions of One Variable

We first recall derivatives of functions $f : \mathbb{R} \to \mathbb{R}$, and rewrite the definition in a form that generalizes to several variables.

Definition 3.2.1 (Derivative)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}$ open, and let $x \in D$. The function f is differentiable at x if the limit

$$f'(x) = \lim_{h \to 0} \frac{f(x + h) - f(x)}{h}$$

exists. Then $f'(x)$, also written $\frac{df}{dx}(x)$, is the derivative of f at x. The function f is differentiable if it is differentiable at every point of D.

The quotient $\frac{f(x+h)-f(x)}{h}$ is the slope of the line through the points $(x, f(x))$ and $(x+h, f(x+h))$ of the graph, and its limit $f'(x)$ is the slope of the tangent line to the graph at x.

Proposition 3.2.2 (Linear approximation)**.** *Let $c \in \mathbb{R}$. Then f is differentiable at x with $f'(x) = c$ if and only if*

$$f(x + h) = f(x) + ch + r(h), \qquad \textit{where } \lim_{h \to 0} \frac{r(h)}{h} = 0.$$

Proof. Let $r(h) = f(x + h) - f(x) - ch$. Then $\frac{r(h)}{h} = \frac{f(x+h)-f(x)}{h} - c$, which converges to 0 if and only if the difference quotient converges to c. □

We write $r(h) = o(|h|)$, read "little o of $|h|$", if $\frac{r(h)}{|h|} \to 0$ as $h \to 0$: the error $r(h)$ is small even compared with the small number $|h|$. So f is differentiable at x if and only if near x, it is approximated by the affine function $h \mapsto f(x)+f'(x)h$, with an error that is $o(|h|)$. This description of the derivative, as the slope of the best linear approximation, is the one that generalizes to functions of several variables.

Proposition 3.2.3 (Rules of differentiation)**.** *Let f and g be differentiable at x, and let $\lambda \in \mathbb{R}$. Then:*

1. $(f+g)'(x) = f'(x) + g'(x)$ *and* $(\lambda f)'(x) = \lambda f'(x)$;
2. *product rule:* $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$;
3. *quotient rule: if* $g(x) \neq 0$, *then* $\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$;
4. *chain rule: if* φ *is differentiable at* $g(x)$, *then* $(\varphi \circ g)'(x) = \varphi'(g(x))\, g'(x)$.

We assume these rules from calculus, together with the derivatives of the standard functions: $(x^k)' = kx^{k-1}$, $(e^x)' = e^x$, $(\log x)' = \frac{1}{x}$ for $x > 0$, $\sin' = \cos$ and $\cos' = -\sin$. Part (1) says that differentiation is a linear map, as in Lecture 1.

Example 3.2.4 (Activation functions)**.** The sigmoid function $\sigma(x) = \frac{1}{1+e^{-x}}$ has the derivative

$$\sigma'(x) = \frac{e^{-x}}{(1+e^{-x})^2} = \sigma(x)(1-\sigma(x)),$$

since $1 - \sigma(x) = \frac{e^{-x}}{1+e^{-x}}$. So the derivative can be computed from the value $\sigma(x)$, which saves work when training neural networks. Since $t(1-t) \leq \frac{1}{4}$ for all t, we have $0 < \sigma'(x) \leq \frac{1}{4}$, with the maximum at $x = 0$. Similarly, $\tanh'(x) = 1 - \tanh(x)^2$. The ReLU function has derivative 0 for $x < 0$ and 1 for $x > 0$, but it is not differentiable at 0: the difference quotient $\frac{\mathrm{ReLU}(h) - \mathrm{ReLU}(0)}{h}$ is 1 for $h > 0$ and 0 for $h < 0$. In practice, software simply uses the derivative 0 at $x = 0$; since an input is very rarely exactly 0, this causes no problems.

If f' is differentiable, its derivative is the second derivative f'', and similarly we define the k-th derivative $f^{(k)}$. A function is k times continuously differentiable if $f^{(k)}$ exists and is continuous.

3.2.1 Taylor polynomials

The linear approximation of f near x_0 only uses $f(x_0)$ and $f'(x_0)$. Using higher derivatives, we get better approximations by polynomials.

Definition 3.2.5 (Taylor polynomial)**.** Let f be k times differentiable at x_0. The Taylor polynomial of degree k of f at x_0 is

$$\begin{aligned} T_k(x) &= \sum_{j=0}^{k} \frac{f^{(j)}(x_0)}{j!}(x - x_0)^j \\ &= f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \cdots + \frac{f^{(k)}(x_0)}{k!}(x - x_0)^k. \end{aligned}$$

The coefficients are chosen such that T_k and f have the same derivatives of orders $0, 1, \ldots, k$ at x_0: the j-th derivative of $\frac{(x-x_0)^i}{i!}$ at x_0 is 1 if $i = j$, and 0 otherwise. The Taylor polynomial $T_1(x) = f(x_0) + f'(x_0)(x - x_0)$ is the tangent line.

Theorem 3.2.6 (Taylor's theorem)**.** *Let* f *be* $k+1$ *times differentiable on an open interval* I *that contains* x_0 *and* x. *Then there is a number* ξ *between* x_0 *and* x *such that*

$$f(x) = T_k(x) + \frac{f^{(k+1)}(\xi)}{(k+1)!}(x - x_0)^{k+1}.$$

Proof. We use Rolle's theorem: if g is differentiable on an open interval that contains $[a, b]$, and $g(a) = g(b)$, then $g'(\eta) = 0$ for some η strictly between a and b. Indeed, g is continuous, so it attains a maximum and a minimum on $[a, b]$ (Theorem A.3.5). If both are attained at the endpoints, then they are equal, since $g(a) = g(b)$, so g is constant, and every η works. Otherwise, say, the maximum

is attained at some $\eta \in (a, b)$. Then the difference quotients $\frac{g(\eta+h)-g(\eta)}{h}$ are ≤ 0 for $h > 0$ and ≥ 0 for $h < 0$, so their limit $g'(\eta)$ is 0; the same argument works for a minimum.

For $x = x_0$, there is nothing to prove. For $x \neq x_0$, choose the number K with $f(x) = T_k(x) + \frac{K}{(k+1)!}(x - x_0)^{k+1}$, and let

$$g(t) = f(t) - T_k(t) - \frac{K}{(k+1)!}(t - x_0)^{k+1}.$$

Since T_k has the same derivatives of orders $0, \ldots, k$ at x_0 as f, and these derivatives of $(t - x_0)^{k+1}$ vanish at x_0, we have $g(x_0) = g'(x_0) = \cdots = g^{(k)}(x_0) = 0$, and $g(x) = 0$ by the choice of K. By Rolle's theorem, $g'(\xi_1) = 0$ for some ξ_1 strictly between x_0 and x. Since $g'(x_0) = 0 = g'(\xi_1)$, Rolle's theorem for g' gives a zero ξ_2 of g'' between x_0 and ξ_1, and so on, until we find a zero $\xi = \xi_{k+1}$ of $g^{(k+1)}$ between x_0 and x. Since T_k is a polynomial of degree at most k, $g^{(k+1)}(t) = f^{(k+1)}(t) - K$, so $K = f^{(k+1)}(\xi)$. □

For $k = 0$, this is the mean value theorem $f(x) - f(x_0) = f'(\xi)(x - x_0)$. Taylor's theorem says that if $|f^{(k+1)}| \leq M$ on I, then the error of the approximation $f(x) \approx T_k(x)$ is at most $\frac{M}{(k+1)!}|x - x_0|^{k+1}$, which is tiny when x is close to x_0.

Example 3.2.7. All derivatives of e^x are e^x, so its Taylor polynomials at $x_0 = 0$ are $T_k(x) = 1 + x + \frac{x^2}{2!} + \cdots + \frac{x^k}{k!}$. For example, $e^{0.1} \approx T_2(0.1) = 1 + 0.1 + 0.005 = 1.105$. By Taylor's theorem, the error is $\frac{e^\xi}{3!}(0.1)^3$ for some $0 < \xi < 0.1$, which is at most $\frac{e^{0.1}}{6000} < 0.0002$. Indeed, $e^{0.1} = 1.10517\ldots$.

Example 3.2.8. The derivatives of cos at 0 are $1, 0, -1, 0, 1, 0, -1, 0, \ldots$, so the Taylor polynomials of cos at 0 are

$$T_{2k}(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots + (-1)^k \frac{x^{2k}}{(2k)!}.$$

Figure 3.2 shows that they approximate cos on larger and larger intervals. Since all derivatives of cos are bounded by 1, Taylor's theorem gives $|\cos x - T_k(x)| \leq \frac{|x|^{k+1}}{(k+1)!}$, which tends to 0 as $k \to \infty$ for every fixed x, since factorials grow faster than powers. Hence $\cos x = \sum_{j=0}^{\infty}(-1)^j \frac{x^{2j}}{(2j)!}$ for all $x \in \mathbb{R}$: the Taylor series of cos converges to cos everywhere. Not all functions behave this nicely. For example, the Taylor series $x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots$ of $\log(1 + x)$ at 0 only converges for $-1 < x \leq 1$.




Figure 3.2: The function $\cos x$ (black) and its Taylor polynomials T_2, T_4 and T_6 at 0. Near 0, all of them are good approximations, and higher degrees give good approximations on larger intervals.

Taylor's theorem also tells us how to compute derivatives numerically, and how accurate the result is.

Proposition 3.2.9 (Finite differences)**.** *Let $h > 0$, and let f be three times differentiable on an open interval containing $[x-h, x+h]$, with $|f''| \le M_2$ and $|f'''| \le M_3$ there. Then*

$$\left|\frac{f(x+h)-f(x)}{h} - f'(x)\right| \le \frac{M_2}{2}h \qquad \text{and} \qquad \left|\frac{f(x+h)-f(x-h)}{2h} - f'(x)\right| \le \frac{M_3}{6}h^2.$$

Proof. By Taylor's theorem with $k=1$, $f(x+h) = f(x) + f'(x)h + \frac{f''(\xi)}{2}h^2$, so the error of the forward difference is $\frac{f''(\xi)}{2}h$. With $k=2$,

$$f(x \pm h) = f(x) \pm f'(x)h + \frac{f''(x)}{2}h^2 \pm \frac{f'''(\xi_\pm)}{6}h^3$$

for some ξ_+ and ξ_-. Subtracting the two equations gives $f(x+h) - f(x-h) = 2f'(x)h + \frac{f'''(\xi_+)+f'''(\xi_-)}{6}h^3$. We divide by $2h$; the error is $\frac{f'''(\xi_+)+f'''(\xi_-)}{12}h^2$, which is at most $\frac{M_3}{6}h^2$. □

The central difference $\frac{f(x+h)-f(x-h)}{2h}$ is much more accurate than the forward difference: halving h divides its error by 4 instead of 2. In machine learning, it is used for gradient checking, i.e., to test a program that computes derivatives. The step h cannot be chosen arbitrarily small, though: computers store numbers with about 16 significant digits, and in the difference $f(x+h) - f(x-h)$ of two almost equal numbers, most of these digits cancel (see Programming Exercise P1).

3.3 Partial Derivatives and Gradients

For a function of several variables, the simplest way to take a derivative is to vary only one variable and to keep all others fixed.

Definition 3.3.1 (Partial derivatives and gradient)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$ open, and let $\mathbf{x} \in D$. The partial derivative of f with respect to x_i at $\mathbf{x}$ is

$$\frac{\partial f}{\partial x_i}(\mathbf{x}) = \lim_{h \to 0} \frac{f(\mathbf{x} + h\mathbf{e}_i) - f(\mathbf{x})}{h},$$

if this limit exists. If all partial derivatives of f at $\mathbf{x}$ exist, the gradient of f at $\mathbf{x}$ is the column vector

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f}{\partial x_1}(\mathbf{x}) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\mathbf{x}) \end{pmatrix} \in \mathbb{R}^n.$$

The partial derivative $\frac{\partial f}{\partial x_i}(\mathbf{x})$ is the ordinary derivative at $t = x_i$ of the function of one variable

$$t \mapsto f(x_1, \ldots, x_{i-1}, t, x_{i+1}, \ldots, x_n).$$

So we compute it with the rules of Section 3.2, treating all variables other than x_i as constants.

Example 3.3.2. Let $f(x_1, x_2) = x_1^2 x_2 + x_1 x_2^3$. Treating x_2 as a constant, and then x_1 as a constant, we get

$$\frac{\partial f}{\partial x_1} = 2x_1x_2 + x_2^3, \qquad \frac{\partial f}{\partial x_2} = x_1^2 + 3x_1x_2^2.$$

At $\mathbf{x} = (1,2)^\top$, the gradient is $\nabla f(\mathbf{x}) = (4+8,\ 1+12)^\top = (12, 13)^\top$.

Proposition 3.3.3. *Let $\mathbf{a} \in \mathbb{R}^n$ and $A \in \mathbb{R}^{n \times n}$. Then for all $\mathbf{x} \in \mathbb{R}^n$:*

1. $\nabla(\mathbf{a}^\top \mathbf{x}) = \mathbf{a}$;

2. $\nabla(\mathbf{x}^\top A\mathbf{x}) = (A + A^\top)\mathbf{x}$; *in particular,* $\nabla(\mathbf{x}^\top A\mathbf{x}) = 2A\mathbf{x}$ *if* A *is symmetric, and* $\nabla\|\mathbf{x}\|^2 = 2\mathbf{x}$;

3. $\nabla\|\mathbf{x}\| = \frac{\mathbf{x}}{\|\mathbf{x}\|}$ *for* $\mathbf{x} \neq \mathbf{0}$.

Proof. (1) Since $\mathbf{a}^\top\mathbf{x} = \sum_j a_j x_j$, the partial derivative with respect to x_k is a_k.

(2) We have $\mathbf{x}^\top A\mathbf{x} = \sum_{i,j} A_{ij}x_i x_j$. The terms that contain x_k are those with $i = k$ or $j = k$, and differentiating them gives

$$\frac{\partial}{\partial x_k}\mathbf{x}^\top A\mathbf{x} = \sum_j A_{kj}x_j + \sum_i A_{ik}x_i = (A\mathbf{x})_k + (A^\top\mathbf{x})_k.$$

(The term $A_{kk}x_k^2$ has the derivative $2A_{kk}x_k$, which is counted once in each sum.) For $A = I_n$, we get $\nabla\|\mathbf{x}\|^2 = \nabla(\mathbf{x}^\top\mathbf{x}) = 2\mathbf{x}$.

(3) By the chain rule, $\frac{\partial}{\partial x_k}(x_1^2 + \cdots + x_n^2)^{1/2} = \frac{1}{2}(x_1^2 + \cdots + x_n^2)^{-1/2} \cdot 2x_k = \frac{x_k}{\|\mathbf{x}\|}$. □

These formulas look like their one-variable versions $(ax)' = a$ and $(ax^2)' = 2ax$, and they will be used constantly. Note that the gradient of the Euclidean norm is a unit vector, which points away from the origin.

3.3.1 Differentiability

Partial derivatives only describe f along the coordinate axes, and by Example 3.1.8, this is not enough. The function $f(x, y) = \frac{xy}{x^2+y^2}$ of that example is 0 on both axes, so both partial derivatives at the origin exist and are 0. But f is not even continuous at the origin, so it would be strange to call it differentiable there. The right definition generalizes Proposition 3.2.2: f should be approximated by a linear function near $\mathbf{x}$, in all directions at once. As before, we write $r(\mathbf{h}) = o(\|\mathbf{h}\|)$ if $\frac{r(\mathbf{h})}{\|\mathbf{h}\|} \to 0$ as $\mathbf{h} \to \mathbf{0}$.

Definition 3.3.4 (Differentiable function)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$ open, and let $\mathbf{x} \in D$. The function f is differentiable at $\mathbf{x}$ if there is a row vector $\mathbf{c} \in \mathbb{R}^{1\times n}$ such that

$$f(\mathbf{x} + \mathbf{h}) = f(\mathbf{x}) + \mathbf{c}\mathbf{h} + r(\mathbf{h}), \qquad \text{where } r(\mathbf{h}) = o(\|\mathbf{h}\|).$$

The row vector $\mathbf{c}$, or the linear map $\mathbf{h} \mapsto \mathbf{c}\mathbf{h}$, is then called the derivative of f at $\mathbf{x}$, and is denoted by $Df(\mathbf{x})$. The function f is differentiable if it is differentiable at every point of D.

Proposition 3.3.5. *Let* f *be differentiable at* $\mathbf{x}$*, with derivative* $Df(\mathbf{x}) = \mathbf{c}$*. Then:*

1. *for every vector* $\mathbf{v} \in \mathbb{R}^n$*, the directional derivative of* f *at* $\mathbf{x}$ *in the direction* $\mathbf{v}$*,*

$$D_\mathbf{v} f(\mathbf{x}) = \lim_{t\to 0} \frac{f(\mathbf{x} + t\mathbf{v}) - f(\mathbf{x})}{t},$$

   exists and is equal to $\mathbf{c}\mathbf{v}$;

2. *all partial derivatives of* f *at* $\mathbf{x}$ *exist, and* $\mathbf{c} = \left(\frac{\partial f}{\partial x_1}(\mathbf{x}), \ldots, \frac{\partial f}{\partial x_n}(\mathbf{x})\right)$*. In other words,* $Df(\mathbf{x}) = \nabla f(\mathbf{x})^\top$*, and* $D_\mathbf{v} f(\mathbf{x}) = \nabla f(\mathbf{x})^\top\mathbf{v}$;

3. f *is continuous at* $\mathbf{x}$.

Proof. (1) For $\mathbf{v} = \mathbf{0}$, both sides are 0. For $\mathbf{v} \neq \mathbf{0}$ and $t \neq 0$, we apply the definition with $\mathbf{h} = t\mathbf{v}$:

$$\left|\frac{f(\mathbf{x} + t\mathbf{v}) - f(\mathbf{x})}{t} - \mathbf{c}\mathbf{v}\right| = \frac{|r(t\mathbf{v})|}{|t|} = \|\mathbf{v}\|\frac{|r(t\mathbf{v})|}{\|t\mathbf{v}\|},$$

which tends to 0 as $t \to 0$.

(2) The partial derivative $\frac{\partial f}{\partial x_i}(\mathbf{x})$ is the directional derivative in the direction $\mathbf{e}_i$, which by (1) is $\mathbf{c}\mathbf{e}_i = c_i$.

(3) We have $|f(\mathbf{x} + \mathbf{h}) - f(\mathbf{x})| \le |\mathbf{c}\mathbf{h}| + |r(\mathbf{h})| \le \|\mathbf{c}\|\|\mathbf{h}\| + |r(\mathbf{h})|$ by the Cauchy–Schwarz inequality, and both terms tend to 0 as $\mathbf{h} \to \mathbf{0}$. □

By (3), the function of Example 3.1.8 is not differentiable at the origin, although its partial derivatives exist there. So differentiability is a stronger condition than the existence of partial derivatives. Fortunately, the following theorem shows that almost all functions that we meet in practice are differentiable. Its proof, which uses the mean value theorem in each coordinate, is given in Proposition A.5.2.

Theorem 3.3.6. *Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$ open. If all partial derivatives of f exist in a ball around $\mathbf{x}$ and are continuous at $\mathbf{x}$, then f is differentiable at $\mathbf{x}$.*

A function whose partial derivatives exist and are continuous on D is called continuously differentiable. By Theorem 3.3.6, polynomials, and functions built from exp, log, sin, the sigmoid function, etc., by sums, products, quotients and compositions, are differentiable wherever they are defined. Some important functions in machine learning are differentiable only at most points, such as ReLU, the ℓ_1 norm, and the hinge loss $\max(0, 1 - t)$; at the few points where they are not differentiable, software uses a reasonable substitute for the derivative.

Remark 3.3.7 (Notation). In this lecture, as in the remark on gradients in the section on dual spaces in Lecture 1, the derivative $Df(\mathbf{x})$ of a function $f : \mathbb{R}^n \to \mathbb{R}$ is a row vector, and the gradient $\nabla f(\mathbf{x}) = Df(\mathbf{x})^\top$ is a column vector. Some books call the row vector the gradient. In machine learning software such as PyTorch, the gradient with respect to a parameter always has the same shape as the parameter, so it is a column vector for a vector of parameters, and a matrix for a matrix of parameters (Section 3.6).

Remark 3.3.8 (The gradient depends on the inner product). The derivative $Df(\mathbf{x})$ is a linear functional $\mathbb{R}^n \to \mathbb{R}$, i.e., an element of the dual space $(\mathbb{R}^n)^*$ of Lecture 1. The gradient is the vector that represents this linear functional with respect to the dot product:

$$Df(\mathbf{x})\mathbf{h} = \langle \nabla f(\mathbf{x}), \mathbf{h} \rangle \qquad \text{for all } \mathbf{h} \in \mathbb{R}^n.$$

In every finite-dimensional inner product space, every linear functional is represented in this way by a unique vector; this is the Riesz representation theorem. If we use a different inner product on $\mathbb{R}^n$, such as $\langle \mathbf{x}, \mathbf{y} \rangle_A = \mathbf{x}^\top A \mathbf{y}$ for a symmetric positive definite matrix A (Lecture 2), then the vector that represents $Df(\mathbf{x})$ changes: since

$$\langle A^{-1} \nabla f(\mathbf{x}), \mathbf{h} \rangle_A = \nabla f(\mathbf{x})^\top A^{-1} A \mathbf{h} = Df(\mathbf{x})\mathbf{h},$$

the gradient with respect to $\langle \cdot, \cdot \rangle_A$ is $A^{-1}\nabla f(\mathbf{x})$. So the derivative is determined by f alone, but the gradient also depends on how we measure lengths and angles. Changing the inner product is the idea behind preconditioning and the natural gradient method, and Newton's method in Section 3.10.3 is an example.

3.3.2 Directional derivatives and steepest ascent

By Proposition 3.3.5, the rate of change of a differentiable function in the direction $\mathbf{v}$ is $D_{\mathbf{v}} f(\mathbf{x}) = \nabla f(\mathbf{x})^\top \mathbf{v}$. In which direction does f increase fastest?

Theorem 3.3.9 (Steepest ascent)**.** *Let f be differentiable at $\mathbf{x}$, with $\nabla f(\mathbf{x}) \neq \mathbf{0}$. Then for every unit vector $\mathbf{v}$,*

$$-\|\nabla f(\mathbf{x})\| \leq D_{\mathbf{v}} f(\mathbf{x}) \leq \|\nabla f(\mathbf{x})\|.$$

The maximum is attained only for $\mathbf{v} = \frac{\nabla f(\mathbf{x})}{\|\nabla f(\mathbf{x})\|}$, and the minimum only for $\mathbf{v} = -\frac{\nabla f(\mathbf{x})}{\|\nabla f(\mathbf{x})\|}$.

Proof. By the Cauchy–Schwarz inequality (Lecture 2), $|D_{\mathbf{v}} f(\mathbf{x})| = |\nabla f(\mathbf{x})^\top \mathbf{v}| \leq \|\nabla f(\mathbf{x})\| \|\mathbf{v}\| = \|\nabla f(\mathbf{x})\|$, with equality if and only if $\mathbf{v}$ is a multiple of $\nabla f(\mathbf{x})$. The only unit vectors that are multiples of $\nabla f(\mathbf{x})$ are $\pm \frac{\nabla f(\mathbf{x})}{\|\nabla f(\mathbf{x})\|}$, and they give $D_{\mathbf{v}} f(\mathbf{x}) = \pm \|\nabla f(\mathbf{x})\|$. □

So the gradient points in the direction in which f increases fastest, and its length is the rate of increase in this direction. The negative gradient $-\nabla f(\mathbf{x})$ is the direction of steepest descent, which is the direction that gradient descent uses to decrease a loss function (Section 3.10). In directions orthogonal to $\nabla f(\mathbf{x})$, the rate of change is 0; we will see in Section 3.5 that these are the directions tangent to the level set through $\mathbf{x}$.

Example 3.3.10. For the function $f(x_1, x_2) = x_1^2 x_2 + x_1 x_2^3$ of Example 3.3.2, at $\mathbf{x} = (1, 2)^\top$, we found $\nabla f(\mathbf{x}) = (12, 13)^\top$. In the direction of the unit vector $\mathbf{v} = \frac{1}{5}(3, 4)^\top$, the rate of change is $D_\mathbf{v} f(\mathbf{x}) = \frac{1}{5}(36+52) = 17.6$. The largest possible rate is $\|\nabla f(\mathbf{x})\| = \sqrt{313} \approx 17.69$, in the direction $\frac{1}{\sqrt{313}}(12, 13)^\top$, and in the orthogonal direction $\frac{1}{\sqrt{313}}(13, -12)^\top$ the rate of change is 0.

3.4 Jacobians: Derivatives of Vector-Valued Functions

Many functions in machine learning have vector outputs, such as the layers of a neural network. Their derivatives are matrices.

Definition 3.4.1 (Jacobian)**.** Let $f : D \to \mathbb{R}^m$ with $D \subseteq \mathbb{R}^n$ open, and let $\mathbf{x} \in D$. The function f is differentiable at $\mathbf{x}$ if there is an $m \times n$ matrix J such that

$$f(\mathbf{x} + \mathbf{h}) = f(\mathbf{x}) + J\mathbf{h} + \mathbf{r}(\mathbf{h}), \qquad \text{where } \lim_{\mathbf{h} \to \mathbf{0}} \frac{\|\mathbf{r}(\mathbf{h})\|}{\|\mathbf{h}\|} = 0.$$

The matrix J is called the Jacobian matrix, or the derivative, of f at $\mathbf{x}$, and is denoted by $Df(\mathbf{x})$ or $J_f(\mathbf{x})$.

For $m = 1$, this is Definition 3.3.4. The next proposition shows that the Jacobian consists of the derivatives of the components of f.

Proposition 3.4.2. *A function $f = (f_1, \ldots, f_m)^\top$ is differentiable at $\mathbf{x}$ if and only if all its components f_i are differentiable at $\mathbf{x}$. In this case, the i-th row of $Df(\mathbf{x})$ is $Df_i(\mathbf{x}) = \nabla f_i(\mathbf{x})^\top$, i.e.,*

$$Df(\mathbf{x}) = \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(\mathbf{x}) & \cdots & \frac{\partial f_1}{\partial x_n}(\mathbf{x}) \\ \vdots & & \vdots \\ \frac{\partial f_m}{\partial x_1}(\mathbf{x}) & \cdots & \frac{\partial f_m}{\partial x_n}(\mathbf{x}) \end{pmatrix}, \qquad \textit{that is,} \quad \big(Df(\mathbf{x})\big)_{ij} = \frac{\partial f_i}{\partial x_j}(\mathbf{x}).$$

Proof. The i-th component of the equation in Definition 3.4.1 is $f_i(\mathbf{x} + \mathbf{h}) = f_i(\mathbf{x}) + J_i\mathbf{h} + r_i(\mathbf{h})$, where J_i is the i-th row of J. Since $|r_i(\mathbf{h})| \le \|\mathbf{r}(\mathbf{h})\| \le |r_1(\mathbf{h})| + \cdots + |r_m(\mathbf{h})|$, we have $\frac{\|\mathbf{r}(\mathbf{h})\|}{\|\mathbf{h}\|} \to 0$ if and only if $\frac{r_i(\mathbf{h})}{\|\mathbf{h}\|} \to 0$ for all i. So f is differentiable with Jacobian J if and only if every f_i is differentiable with derivative J_i, and by Proposition 3.3.5, $J_i = \nabla f_i(\mathbf{x})^\top$. □

So the rows of the Jacobian correspond to the outputs of f, and the columns to the inputs: the j-th column $\frac{\partial f}{\partial x_j}(\mathbf{x})$ describes how the output vector changes when we change x_j. In the language of Lecture 1, the derivative of f at $\mathbf{x}$ is the linear transformation $\mathbf{h} \mapsto Df(\mathbf{x})\mathbf{h}$ from $\mathbb{R}^n$ to $\mathbb{R}^m$ that best approximates the change $f(\mathbf{x} + \mathbf{h}) - f(\mathbf{x})$, and the Jacobian is its matrix with respect to the standard bases. Near $\mathbf{x}$,

$$f(\mathbf{x} + \mathbf{h}) \approx f(\mathbf{x}) + Df(\mathbf{x})\mathbf{h} :$$

a differentiable function looks like an affine function if we zoom in far enough.

Example 3.4.3. Let $f(x_1, x_2) = (x_1^2 x_2,\ 3x_1 + x_2^2)^\top$. Then

$$Df(\mathbf{x}) = \begin{pmatrix} 2x_1x_2 & x_1^2 \\ 3 & 2x_2 \end{pmatrix}, \qquad \text{and at } \mathbf{a} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} : \quad f(\mathbf{a}) = \begin{pmatrix} 2 \\ 7 \end{pmatrix}, \quad Df(\mathbf{a}) = \begin{pmatrix} 4 & 1 \\ 3 & 4 \end{pmatrix}.$$

The linear approximation gives $f(1.1, 1.9) \approx f(\mathbf{a}) + Df(\mathbf{a}) \begin{pmatrix} 0.1 \\ -0.1 \end{pmatrix} = \begin{pmatrix} 2.3 \\ 6.9 \end{pmatrix}$, while the exact value is $(2.299,\, 6.91)^\top$.

Example 3.4.4 (Affine maps)**.** Let $f(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ with $A \in \mathbb{R}^{m\times n}$ and $\mathbf{b} \in \mathbb{R}^m$. Then $f(\mathbf{x}+\mathbf{h}) = f(\mathbf{x}) + A\mathbf{h}$ exactly, so $Df(\mathbf{x}) = A$ for every $\mathbf{x}$: the best linear approximation of an affine map is its linear part. For $m = 1$ and $f(\mathbf{x}) = \mathbf{a}^\top\mathbf{x}$, this is Proposition 3.3.3(1) again.

Example 3.4.5 (Functions applied entrywise)**.** Let $\varphi : \mathbb{R} \to \mathbb{R}$ be differentiable, such as an activation function, and let $f : \mathbb{R}^n \to \mathbb{R}^n$ apply φ to each entry, $f(\mathbf{x}) = (\varphi(x_1), \dots, \varphi(x_n))^\top$. Since f_i only depends on x_i, we have $\frac{\partial f_i}{\partial x_j} = 0$ for $i \neq j$, so

$$Df(\mathbf{x}) = \operatorname{diag}\left(\varphi'(x_1), \dots, \varphi'(x_n)\right).$$

Multiplying a vector by a diagonal matrix multiplies its entries: $Df(\mathbf{x})\mathbf{v} = \varphi'(\mathbf{x}) \odot \mathbf{v}$, where $\varphi'(\mathbf{x}) = (\varphi'(x_1), \dots, \varphi'(x_n))^\top$, and $\odot$ is the entrywise product (or Hadamard product) $(\mathbf{a} \odot \mathbf{b})_i = a_i b_i$. Computing the product in this way costs n multiplications instead of n^2.

Example 3.4.6 (Softmax)**.** Let $\mathbf{s} = \operatorname{softmax}(\mathbf{z})$ with $\mathbf{z} \in \mathbb{R}^K$, i.e., $s_i = \frac{e^{z_i}}{S}$ with $S = e^{z_1} + \cdots + e^{z_K}$. Since $\frac{\partial S}{\partial z_j} = e^{z_j}$, the quotient rule gives

$$\frac{\partial s_i}{\partial z_j} = \frac{\delta_{ij} e^{z_i} S - e^{z_i} e^{z_j}}{S^2} = s_i\delta_{ij} - s_i s_j, \qquad \text{where } \delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

So $D\operatorname{softmax}(\mathbf{z}) = \operatorname{diag}(\mathbf{s}) - \mathbf{s}\mathbf{s}^\top$. For example, for $\mathbf{z} = (2, 1, 0)^\top$, we have $\mathbf{s} \approx (0.665, 0.245, 0.090)^\top$ and

$$D\operatorname{softmax}(\mathbf{z}) \approx \begin{pmatrix} 0.223 & -0.163 & -0.060 \\ -0.163 & 0.185 & -0.022 \\ -0.060 & -0.022 & 0.082 \end{pmatrix}.$$

The Jacobian is symmetric, and its rows sum to 0: with $\mathbf{1} = (1, \dots, 1)^\top$, we have $D\operatorname{softmax}(\mathbf{z})\mathbf{1} = \mathbf{s} - \mathbf{s}(\mathbf{s}^\top\mathbf{1}) = \mathbf{0}$, since the entries of $\mathbf{s}$ sum to 1. This reflects a fact from the section on quotient spaces in Lecture 1: $\operatorname{softmax}(\mathbf{z} + c\mathbf{1}) = \operatorname{softmax}(\mathbf{z})$ for all $c \in \mathbb{R}$, so moving $\mathbf{z}$ in the direction $\mathbf{1}$ does not change the output, and $\mathbf{1}$ lies in the null space of the Jacobian.

3.4.1 The Jacobian determinant

If $m = n$, the Jacobian is a square matrix, and its determinant describes how f changes areas and volumes near a point. By Lecture 1, the linear map $\mathbf{h} \mapsto A\mathbf{h}$ multiplies areas (for $n = 2$) and volumes (for $n \geq 3$) by $|\det A|$. Near $\mathbf{x}$, f is approximately the affine map $\mathbf{h} \mapsto f(\mathbf{x}) + Df(\mathbf{x})\mathbf{h}$. So a small square with side length ε at $\mathbf{x}$ is mapped approximately to the parallelogram spanned by $\varepsilon Df(\mathbf{x})\mathbf{e}_1$ and $\varepsilon Df(\mathbf{x})\mathbf{e}_2$, whose area is $|\det Df(\mathbf{x})|\,\varepsilon^2$: the number $|\det Df(\mathbf{x})|$ is the factor by which f scales small areas near $\mathbf{x}$.

Example 3.4.7 (Polar coordinates)**.** The map $f(r, \varphi) = (r\cos\varphi,\, r\sin\varphi)^\top$ turns polar coordinates into Cartesian coordinates. Its Jacobian and Jacobian determinant are

$$Df(r, \varphi) = \begin{pmatrix} \cos\varphi & -r\sin\varphi \\ \sin\varphi & r\cos\varphi \end{pmatrix}, \qquad \det Df(r, \varphi) = r\cos^2\varphi + r\sin^2\varphi = r.$$

A small rectangle $[r, r+\Delta r] \times [\varphi, \varphi+\Delta\varphi]$ is mapped to a piece of an annulus, which is approximately a rectangle with sides Δr and $r\Delta\varphi$ (an arc of a circle of radius r), with area approximately $r\,\Delta r\,\Delta\varphi$; see Figure 3.3. This explains the factor r in the substitution rule for integrals in polar coordinates,

$$\iint_{\mathbb{R}^2} g(x, y)\,dx\,dy = \int_0^{2\pi}\int_0^{\infty} g(r\cos\varphi, r\sin\varphi)\,r\,dr\,d\varphi.$$

A famous application is the Gaussian integral. Since $e^{-(x^2+y^2)} = e^{-x^2}e^{-y^2}$,

$$\left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)^2 = \iint_{\mathbb{R}^2} e^{-(x^2+y^2)}\,dx\,dy = \int_0^{2\pi}\int_0^{\infty} e^{-r^2} r\,dr\,d\varphi = 2\pi\left[-\tfrac{1}{2}e^{-r^2}\right]_0^{\infty} = \pi,$$

so $\int_{-\infty}^{\infty} e^{-x^2}dx = \sqrt{\pi}$. This is where the factor $\frac{1}{\sqrt{2\pi}}$ in the density of the normal distribution comes from.




Figure 3.3: Polar coordinates $f(r,\varphi) = (r\cos\varphi, r\sin\varphi)$. The grid of rectangles on the left is mapped to the grid of annulus pieces on the right. The two shaded rectangles have the same area, but their images do not: since $\det Df(r,\varphi) = r$, areas are stretched more far from the origin. (The image of $[r_1,r_2]\times[\varphi_1,\varphi_2]$ has the area $\frac{r_1+r_2}{2}(r_2-r_1)(\varphi_2-\varphi_1)$, so the orange image is $\frac{1.625}{0.625} = 2.6$ times as large as the blue one.)

Remark 3.4.8 (Change of variables). More generally, if f is a differentiable bijection between open subsets U and V of $\mathbb{R}^n$ with a differentiable inverse, then $\int_V g(\mathbf{y})\,d\mathbf{y} = \int_U g(f(\mathbf{x}))\,|\det Df(\mathbf{x})|\,d\mathbf{x}$ (Theorem 3.14.4). In probability, it says that if a random vector X has the density p_X, then $Y = f(X)$ has the density p_Y with $p_Y(f(\mathbf{x})) = \frac{p_X(\mathbf{x})}{|\det Df(\mathbf{x})|}$. Normalizing flows are generative models that turn simple random vectors (e.g., Gaussian ones) into complicated ones in this way. They use neural networks f whose Jacobian determinant is cheap to compute, for example networks with triangular Jacobians, whose determinant is the product of the diagonal entries (Lecture 1).

3.5 The Chain Rule

Most functions in machine learning are compositions of simpler functions: a neural network is a composition of layers, and its loss is a composition of the network with a loss function. The chain rule tells us how to differentiate compositions.

Theorem 3.5.1 (Chain rule)**.** *Let $g: D \to \mathbb{R}^m$ with $D \subseteq \mathbb{R}^n$ open, and $f: E \to \mathbb{R}^k$ with $E \subseteq \mathbb{R}^m$ open and $g(D) \subseteq E$. If g is differentiable at $\mathbf{x}$ and f is differentiable at $g(\mathbf{x})$, then $f\circ g$ is differentiable at $\mathbf{x}$, and*

$$D(f\circ g)(\mathbf{x}) = Df(g(\mathbf{x}))\,Dg(\mathbf{x}).$$

The sizes of the matrices are $(k\times n) = (k\times m)(m\times n)$. In terms of the entries,

$$\frac{\partial (f\circ g)_i}{\partial x_j}(\mathbf{x}) = \sum_{l=1}^{m} \frac{\partial f_i}{\partial y_l}(g(\mathbf{x}))\,\frac{\partial g_l}{\partial x_j}(\mathbf{x}),$$

where $y_1,\dots,y_m$ denote the variables of f.

Proof. Let $\mathbf{y} = g(\mathbf{x})$, $A = Dg(\mathbf{x})$ and $B = Df(\mathbf{y})$. By assumption,

$$g(\mathbf{x}+\mathbf{h}) = \mathbf{y} + \mathbf{k}(\mathbf{h}) \quad \text{with } \mathbf{k}(\mathbf{h}) = A\mathbf{h} + \mathbf{r}(\mathbf{h}), \qquad f(\mathbf{y}+\mathbf{k}) = f(\mathbf{y}) + B\mathbf{k} + \mathbf{s}(\mathbf{k}),$$

where $\frac{\|\mathbf{r}(\mathbf{h})\|}{\|\mathbf{h}\|} \to 0$ as $\mathbf{h} \to \mathbf{0}$ and $\frac{\|\mathbf{s}(\mathbf{k})\|}{\|\mathbf{k}\|} \to 0$ as $\mathbf{k} \to \mathbf{0}$, with $\mathbf{s}(\mathbf{0}) = \mathbf{0}$. Hence

$$f(g(\mathbf{x}+\mathbf{h})) = f(\mathbf{y}) + BA\mathbf{h} + \big(B\mathbf{r}(\mathbf{h}) + \mathbf{s}(\mathbf{k}(\mathbf{h}))\big),$$

and we have to show that the error term in parentheses is $o(\|\mathbf{h}\|)$. We use the spectral norm of Lecture 2, which satisfies $\|M\mathbf{v}\| \le \|M\|_2\|\mathbf{v}\|$. First, $\|B\mathbf{r}(\mathbf{h})\| \le \|B\|_2\|\mathbf{r}(\mathbf{h})\|$, which is $o(\|\mathbf{h}\|)$. Second, for small $\mathbf{h}$, we have $\|\mathbf{r}(\mathbf{h})\| \le \|\mathbf{h}\|$ and therefore $\|\mathbf{k}(\mathbf{h})\| \le (\|A\|_2 + 1)\|\mathbf{h}\|$. Let $\varepsilon > 0$. For small $\mathbf{k}$, we have $\|\mathbf{s}(\mathbf{k})\| \le \varepsilon\|\mathbf{k}\|$, so for small $\mathbf{h}$,

$$\|\mathbf{s}(\mathbf{k}(\mathbf{h}))\| \le \varepsilon\|\mathbf{k}(\mathbf{h})\| \le \varepsilon(\|A\|_2 + 1)\|\mathbf{h}\|.$$

Since ε was arbitrary, $\mathbf{s}(\mathbf{k}(\mathbf{h})) = o(\|\mathbf{h}\|)$ as well. □

The chain rule says that the linear approximation of $f \circ g$ is the composition of the linear approximations of f and g. Since the matrix of a composition of linear maps is the product of their matrices (Lecture 1), the Jacobian of $f \circ g$ is the product of the Jacobians. For $n = m = k = 1$, it is the familiar rule $(f \circ g)'(x) = f'(g(x))g'(x)$. Two special cases are particularly important.

Corollary 3.5.2. *1. (Functions along curves.) Let $\mathbf{x} : I \to \mathbb{R}^n$, $t \mapsto \mathbf{x}(t) = (x_1(t), \dots, x_n(t))^\top$, be differentiable on an open interval I, and let $f : \mathbb{R}^n \to \mathbb{R}$ be differentiable. Then*

$$\frac{d}{dt} f(\mathbf{x}(t)) = \nabla f(\mathbf{x}(t))^\top \mathbf{x}'(t) = \sum_{i=1}^{n} \frac{\partial f}{\partial x_i}(\mathbf{x}(t))\, x_i'(t), \qquad \text{where } \mathbf{x}'(t) = (x_1'(t), \dots, x_n'(t))^\top.$$

2. (Gradients of compositions.) Let $g : \mathbb{R}^n \to \mathbb{R}^m$ and $f : \mathbb{R}^m \to \mathbb{R}$ be differentiable. Then

$$\nabla(f \circ g)(\mathbf{x}) = Dg(\mathbf{x})^\top \nabla f(g(\mathbf{x})).$$

Proof. (1) The Jacobian of $t \mapsto \mathbf{x}(t)$ is the $n \times 1$ matrix $\mathbf{x}'(t)$, so this is the chain rule with $Df(\mathbf{x}(t)) = \nabla f(\mathbf{x}(t))^\top$. (2) By the chain rule, $D(f \circ g)(\mathbf{x}) = Df(g(\mathbf{x}))Dg(\mathbf{x}) = \nabla f(g(\mathbf{x}))^\top Dg(\mathbf{x})$; now take the transpose. □

Part (2) says that gradients are pulled back through a function by multiplying with the transposed Jacobian. This is the key step of backpropagation (Section 3.7).

Example 3.5.3. Let $f(x_1, x_2) = x_1^2 + 2x_2$, and let $\mathbf{x}(t) = (\sin t, \cos t)^\top$ move on the unit circle. Then $\nabla f(\mathbf{x}) = (2x_1, 2)^\top$ and $\mathbf{x}'(t) = (\cos t, -\sin t)^\top$, so

$$\frac{d}{dt} f(\mathbf{x}(t)) = 2\sin t \cos t - 2\sin t.$$

We can check this directly: $f(\mathbf{x}(t)) = \sin^2 t + 2\cos t$ has the derivative $2\sin t\cos t - 2\sin t$.

Corollary 3.5.4 (The gradient is orthogonal to level sets)**.** *Let $f : \mathbb{R}^n \to \mathbb{R}$ be differentiable, and let $\mathbf{x} : I \to \mathbb{R}^n$ be a differentiable curve in a level set of f, i.e., $f(\mathbf{x}(t)) = c$ for all $t \in I$. Then $\nabla f(\mathbf{x}(t))^\top \mathbf{x}'(t) = 0$ for all $t \in I$.*

Proof. By Corollary 3.5.2(1), $\nabla f(\mathbf{x}(t))^\top \mathbf{x}'(t)$ is the derivative of the constant function $t \mapsto c$, which is 0. □

The vector $\mathbf{x}'(t)$ is tangent to the curve, so the gradient is orthogonal to every curve in the level set, i.e., orthogonal to the level set itself; see Figure 3.4. Together with Theorem 3.3.9, this gives a good picture: the steepest way up a hill is perpendicular to the contour lines of the map, and the gradient is longer where the contour lines are closer together.




Figure 3.4: Level curves $x_1^2 + 4x_2^2 = c$ of $f(\mathbf{x}) = x_1^2 + 4x_2^2$ for $c = 1, 4, 9, 16$, and the gradients $\nabla f(\mathbf{x}) = (2x_1, 8x_2)^\top$ (scaled by 0.1) at points of the level curve $c = 9$. The gradients are orthogonal to the level curve (dashed: the tangent line at one point), and they are longest where the level curves are closest together, on the x_2-axis.

Example 3.5.5 (The gradient of least squares)**.** Let $\Phi \in \mathbb{R}^{N\times d}$ be a design matrix and $\mathbf{y} \in \mathbb{R}^N$ a vector of targets, as in the section on least squares in Lecture 2, and let

$$L(\boldsymbol{\theta}) = \|\mathbf{y} - \Phi\boldsymbol{\theta}\|^2, \qquad \boldsymbol{\theta} \in \mathbb{R}^d.$$

We write $L = f \circ \mathbf{e}$ with the residual $\mathbf{e}(\boldsymbol{\theta}) = \mathbf{y} - \Phi\boldsymbol{\theta}$ and $f(\mathbf{e}) = \|\mathbf{e}\|^2$. By Example 3.4.4 and Proposition 3.3.3, $D\mathbf{e}(\boldsymbol{\theta}) = -\Phi$ and $Df(\mathbf{e}) = 2\mathbf{e}^\top$. By the chain rule,

$$DL(\boldsymbol{\theta}) = \underbrace{2(\mathbf{y} - \Phi\boldsymbol{\theta})^\top}_{1\times N}\,\underbrace{(-\Phi)}_{N\times d}, \qquad \text{so} \qquad \nabla L(\boldsymbol{\theta}) = -2\Phi^\top(\mathbf{y} - \Phi\boldsymbol{\theta}) = 2\Phi^\top(\Phi\boldsymbol{\theta} - \mathbf{y}).$$

Setting the gradient to zero gives $\Phi^\top\Phi\boldsymbol{\theta} = \Phi^\top\mathbf{y}$, the normal equations, which we found in Lecture 2 with orthogonal projections. Note how writing the sizes of the matrices under the factors checks that the product makes sense. This simple test catches many mistakes in gradient computations.

Example 3.5.6 (Logistic regression)**.** In binary classification, each data point $\mathbf{x} \in \mathbb{R}^n$ has a label $y \in \{0, 1\}$. Logistic regression predicts the probability $p = \sigma(\mathbf{w}^\top\mathbf{x})$ that $y = 1$, where σ is the sigmoid function, and it is trained by minimizing the cross-entropy loss, which for one data point is

$$\ell(\mathbf{w}) = -y \log p - (1 - y)\log(1 - p), \qquad p = \sigma(\mathbf{w}^\top\mathbf{x}).$$

It is small if the predicted probability of the correct label is close to 1. We compute the gradient with the chain rule, following the computation $\mathbf{w} \mapsto z = \mathbf{w}^\top\mathbf{x} \mapsto p = \sigma(z) \mapsto \ell$. By Example 3.2.4, $\frac{dp}{dz} = p(1 - p)$, so

$$\frac{d\ell}{dz} = \left(-\frac{y}{p} + \frac{1 - y}{1 - p}\right) p(1 - p) = -y(1 - p) + (1 - y)p = p - y,$$

and since $Dz(\mathbf{w}) = \mathbf{x}^\top$,

$$\nabla\ell(\mathbf{w}) = (\sigma(\mathbf{w}^\top\mathbf{x}) - y)\,\mathbf{x}:$$

the prediction error times the input. For example, for $\mathbf{x} = (1, 2)^\top$, $y = 1$ and $\mathbf{w} = (0.5, -0.25)^\top$, we have $z = 0$, $p = \frac{1}{2}$, $\ell = \log 2$, and $\nabla\ell(\mathbf{w}) = -\frac{1}{2}(1, 2)^\top = (-0.5, -1)^\top$. A step in the direction $-\nabla\ell(\mathbf{w})$ increases $\mathbf{w}^\top\mathbf{x}$, and hence the predicted probability of the correct label $y = 1$. For a data set $(\mathbf{x}_1, y_1), \ldots, (\mathbf{x}_N, y_N)$ with the average loss $L(\mathbf{w}) = \frac{1}{N}\sum_i \ell_i(\mathbf{w})$, we get

$$\nabla L(\mathbf{w}) = \frac{1}{N}\sum_{i=1}^{N}(\sigma(\mathbf{w}^\top\mathbf{x}_i) - y_i)\mathbf{x}_i = \frac{1}{N}X^\top(\sigma(X\mathbf{w}) - \mathbf{y}),$$

where $X \in \mathbb{R}^{N\times n}$ has the rows $\mathbf{x}_i^\top$, and σ is applied entrywise.

3.6 Gradients with Respect to Matrices

The parameters of a layer of a neural network include a weight matrix W, so we need derivatives of functions $f : \mathbb{R}^{m\times n} \to \mathbb{R}$. By Lecture 1, $\mathbb{R}^{m\times n}$ is a vector space isomorphic to $\mathbb{R}^{mn}$, so f is a function of the mn entries of X, and we can take the partial derivatives $\frac{\partial f}{\partial X_{ij}}$.

Definition 3.6.1 (Gradient with respect to a matrix)**.** Let $f : \mathbb{R}^{m\times n} \to \mathbb{R}$ be differentiable. The gradient of f at X is the $m \times n$ matrix $\nabla f(X)$ with the entries

$$(\nabla f(X))_{ij} = \frac{\partial f}{\partial X_{ij}}(X).$$

The gradient has the same shape as X. Recall from Lecture 2 the Frobenius inner product $\langle A, B\rangle = \operatorname{tr}(A^\top B) = \sum_{i,j} A_{ij}B_{ij}$ on $\mathbb{R}^{m\times n}$, which is the dot product of the lists of entries, and the Frobenius norm $\|A\|_F = \sqrt{\langle A, A\rangle}$. Since f is a differentiable function of mn variables, Proposition 3.3.5 gives

$$f(X + H) = f(X) + \langle \nabla f(X), H\rangle + o(\|H\|_F),$$

and conversely, if $f(X + H) = f(X) + \langle G, H\rangle + o(\|H\|_F)$ for some matrix G, then $G = \nabla f(X)$. This gives a convenient method to compute gradients: expand $f(X + H)$, collect the terms that are linear in H, and write them in the form $\langle G, H\rangle = \operatorname{tr}(G^\top H)$. The following rules for the trace (Lecture 1) are useful: $\operatorname{tr}(A) = \operatorname{tr}(A^\top)$ and $\operatorname{tr}(AB) = \operatorname{tr}(BA)$, and hence

$$\operatorname{tr}(MH) = \operatorname{tr}(H^\top M^\top) = \langle M^\top, H\rangle \qquad \text{for all } M \in \mathbb{R}^{n\times m},\ H \in \mathbb{R}^{m\times n}.$$

Proposition 3.6.2. *Let $X \in \mathbb{R}^{m\times n}$. Then:*

1. $\nabla_X \operatorname{tr}(AX) = A^\top$ *for* $A \in \mathbb{R}^{n\times m}$, *and* $\nabla_X(\mathbf{a}^\top X\mathbf{b}) = \mathbf{a}\mathbf{b}^\top$ *for* $\mathbf{a} \in \mathbb{R}^m$, $\mathbf{b} \in \mathbb{R}^n$;
2. $\nabla_X\|AX - B\|_F^2 = 2A^\top(AX - B)$ *for* $A \in \mathbb{R}^{p\times m}$, $B \in \mathbb{R}^{p\times n}$; *in particular,* $\nabla_X\|X\|_F^2 = 2X$;
3. $\nabla_X \operatorname{tr}(X^\top AX) = (A + A^\top)X$ *for* $A \in \mathbb{R}^{m\times m}$.

Here the subscript in ∇_X indicates the variable.

Proof. (1) $\operatorname{tr}(A(X + H)) = \operatorname{tr}(AX) + \operatorname{tr}(AH)$, and $\operatorname{tr}(AH) = \langle A^\top, H\rangle$. For the second formula, $\mathbf{a}^\top X\mathbf{b}$ is a 1×1 matrix, so it equals its trace, and $\operatorname{tr}(\mathbf{a}^\top X\mathbf{b}) = \operatorname{tr}(\mathbf{b}\mathbf{a}^\top X)$; now use the first formula with $A = \mathbf{b}\mathbf{a}^\top$.

(2) Let $R = AX - B$. Then

$$\|A(X + H) - B\|_F^2 = \|R + AH\|_F^2 = \|R\|_F^2 + 2\langle R, AH\rangle + \|AH\|_F^2.$$

The last term is quadratic in H, so it is $o(\|H\|_F)$, and $\langle R, AH\rangle = \operatorname{tr}(R^\top AH) = \langle A^\top R, H\rangle$. For $A = I$ and $B = 0$, we get $\nabla\|X\|_F^2 = 2X$.

(3) $\operatorname{tr}((X + H)^\top A(X + H)) = \operatorname{tr}(X^\top AX) + \operatorname{tr}(H^\top AX) + \operatorname{tr}(X^\top AH) + \operatorname{tr}(H^\top AH)$. The last term is quadratic in H, $\operatorname{tr}(H^\top AX) = \langle H, AX\rangle = \langle AX, H\rangle$, and $\operatorname{tr}(X^\top AH) = \langle A^\top X, H\rangle$. □

For $n = 1$, part (3) is Proposition 3.3.3(2). The gradient of the determinant is a little harder to compute.

Lemma 3.6.3. *For every $E \in \mathbb{R}^{n\times n}$, $\det(I + tE) = 1 + t\operatorname{tr}(E) + t^2q(t)$ for some polynomial q. In particular, $\frac{d}{dt}\det(I + tE)\big|_{t=0} = \operatorname{tr}(E)$.*

Proof. Let $\mu_1, \dots, \mu_n$ be the complex eigenvalues of E, so that $\det(E - \lambda I) = \prod_i (\mu_i - \lambda)$, and $\mu_1 + \cdots + \mu_n = \operatorname{tr}(E)$ by Lecture 1. For $t \neq 0$, we take $\lambda = -\frac{1}{t}$ and get

$$\det(I + tE) = t^n \det\Big(E + \frac{1}{t} I\Big) = t^n \prod_{i=1}^{n} \Big(\mu_i + \frac{1}{t}\Big) = \prod_{i=1}^{n} (1 + t\mu_i),$$

and this also holds for $t = 0$. Multiplying out the product gives

$$\det(I + tE) = 1 + t(\mu_1 + \cdots + \mu_n) + t^2 q(t) = 1 + t \operatorname{tr}(E) + t^2 q(t),$$

where q collects the terms with at least two factors $t\mu_i$. (Its coefficients are real, since $t^2 q(t)$ is the difference of the real polynomials $\det(I + tE)$ and $1 + t \operatorname{tr}(E)$.) □

Proposition 3.6.4 (Jacobi's formula)**.** *For every invertible matrix $X \in \mathbb{R}^{n \times n}$,*

$$\nabla_X \det(X) = \det(X)\, X^{-\top} \qquad \textit{and} \qquad \nabla_X \log|\det(X)| = X^{-\top},$$

where $X^{-\top} = (X^{-1})^\top$.

Proof. The determinant is a polynomial in the entries of X, so it is differentiable by Theorem 3.3.6. For every $H \in \mathbb{R}^{n \times n}$, by Lemma 3.6.3 with $E = X^{-1}H$,

$$\det(X + tH) = \det(X) \det(I + tX^{-1}H) = \det(X)\big(1 + t \operatorname{tr}(X^{-1}H) + t^2 q(t)\big).$$

So the directional derivative of det at X in the direction H is $\det(X) \operatorname{tr}(X^{-1}H) = \langle \det(X) X^{-\top}, H \rangle$. By Proposition 3.3.5, it is also equal to $\langle \nabla \det(X), H \rangle$. Taking for H the matrices with a single entry 1 and all other entries 0, we see that the two gradients have the same entries. For the second formula, $\frac{d}{du} \log|u| = \frac{1}{u}$ for $u \neq 0$, so by the chain rule $\nabla \log|\det(X)| = \frac{1}{\det(X)} \nabla \det(X) = X^{-\top}$. □

For $n = 1$, this says that $\frac{d}{dx} \log|x| = \frac{1}{x}$. There is also a direct way to see the first formula: by the cofactor expansion along the i-th row, $\frac{\partial \det(X)}{\partial X_{ij}}$ is the cofactor C_{ij}, and by Lecture 1, $X^{-1} = \frac{1}{\det(X)} C^\top$, i.e., $\det(X) X^{-\top} = C$. The gradient of $\log|\det|$ appears in the training of Gaussian models and normalizing flows (Remark 3.4.8).

Example 3.6.5 (A linear layer)**.** Let $\mathbf{y} = W\mathbf{x} + \mathbf{b}$ with $W \in \mathbb{R}^{m \times n}$, $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{b} \in \mathbb{R}^m$, and let L be a loss that depends on $\mathbf{y}$ (and maybe on other things). Let $\boldsymbol{\delta} = \nabla_{\mathbf{y}} L \in \mathbb{R}^m$ be the gradient of the loss with respect to the output of the layer. Then

$$\nabla_W L = \boldsymbol{\delta}\, \mathbf{x}^\top, \qquad \nabla_{\mathbf{b}} L = \boldsymbol{\delta}, \qquad \nabla_{\mathbf{x}} L = W^\top \boldsymbol{\delta}.$$

Indeed, $y_i = \sum_j W_{ij} x_j + b_i$, and the entry W_{ij} only influences L through y_i. So by the chain rule, $\frac{\partial L}{\partial W_{ij}} = \frac{\partial L}{\partial y_i} x_j = \delta_i x_j$, which is the (i, j) entry of the outer product $\boldsymbol{\delta}\mathbf{x}^\top$. Similarly, $\frac{\partial L}{\partial b_i} = \delta_i$, and the last formula is Corollary 3.5.2(2) with $D\mathbf{y}(\mathbf{x}) = W$. The sizes fit: $\boldsymbol{\delta}\mathbf{x}^\top$ is $m \times n$, like W.

In practice, a layer processes a batch of N inputs at once. If the inputs are the rows of $X \in \mathbb{R}^{N \times n}$, the outputs are the rows of $Y = XW^\top + \mathbf{1}\mathbf{b}^\top \in \mathbb{R}^{N \times m}$, and the loss is the sum of the losses of the data points, then the gradient is the sum of the gradients $\boldsymbol{\delta}_k \mathbf{x}_k^\top$ of the data points:

$$\nabla_W L = \sum_{k=1}^{N} \boldsymbol{\delta}_k \mathbf{x}_k^\top = \Delta^\top X, \qquad \nabla_{\mathbf{b}} L = \Delta^\top \mathbf{1},$$

where $\Delta \in \mathbb{R}^{N \times m}$ has the rows $\boldsymbol{\delta}_k^\top$.

Example 3.6.6 (Softmax with cross-entropy)**.** A classifier with K classes computes a vector of scores $\mathbf{z} \in \mathbb{R}^K$ and predicts the probabilities $\mathbf{p} = \mathrm{softmax}(\mathbf{z})$. If the correct class is c, the cross-entropy loss is

$$\ell(\mathbf{z}) = -\log p_c = -z_c + \log\left(e^{z_1} + \cdots + e^{z_K}\right).$$

Since $\frac{\partial}{\partial z_k} \log\left(\sum_j e^{z_j}\right) = \frac{e^{z_k}}{\sum_j e^{z_j}} = p_k$, the gradient is

$$\nabla \ell(\mathbf{z}) = \mathbf{p} - \mathbf{e}_c,$$

the predicted probabilities minus the "one-hot" vector of the correct class. This has the same form as for logistic regression. For example, for $\mathbf{z} = (2, 1, 0)^\top$ and $c = 1$, we get $\mathbf{p} \approx (0.665, 0.245, 0.090)^\top$, $\ell \approx 0.408$, and $\nabla \ell(\mathbf{z}) \approx (-0.335, 0.245, 0.090)^\top$. Gradient descent increases the score of the correct class and decreases the other scores. If the scores come from a linear layer $\mathbf{z} = W\mathbf{x} + \mathbf{b}$, then by Example 3.6.5, $\nabla_W \ell = (\mathbf{p} - \mathbf{e}_c)\mathbf{x}^\top$.

We also found that the gradient of the log-sum-exp function $\log(e^{z_1} + \cdots + e^{z_K})$ is $\mathrm{softmax}(\mathbf{z})$. Table 3.1 collects the gradients of this and the previous sections.

function	gradient	assumptions
$\mathbf{a}^\top \mathbf{x}$	$\mathbf{a}$	
$\mathbf{x}^\top A \mathbf{x}$	$(A + A^\top)\mathbf{x}$	$(= 2A\mathbf{x}$ if A is symmetric)
$\lVert\mathbf{x}\rVert^2$	$2\mathbf{x}$	
$\lVert\mathbf{x}\rVert$	$\mathbf{x}/\lVert\mathbf{x}\rVert$	$\mathbf{x} \neq \mathbf{0}$
$\lVert A\mathbf{x} - \mathbf{b}\rVert^2$	$2A^\top(A\mathbf{x} - \mathbf{b})$	
$f(g(\mathbf{x}))$	$Dg(\mathbf{x})^\top \nabla f(g(\mathbf{x}))$	f scalar-valued
$\log(e^{z_1} + \cdots + e^{z_K})$	$\mathrm{softmax}(\mathbf{z})$	
$-\log \mathrm{softmax}(\mathbf{z})_c$	$\mathrm{softmax}(\mathbf{z}) - \mathbf{e}_c$	
$\mathrm{tr}(AX)$	$A^\top$	
$\mathbf{a}^\top X \mathbf{b}$	$\mathbf{a}\mathbf{b}^\top$	
$\lVert AX - B\rVert_F^2$	$2A^\top(AX - B)$	
$\mathrm{tr}(X^\top A X)$	$(A + A^\top)X$	
$\det(X)$	$\det(X)X^{-\top}$	X invertible
$\log\lvert\det(X)\rvert$	$X^{-\top}$	X invertible

Table 3.1: Some gradients with respect to a vector $\mathbf{x}$ or $\mathbf{z}$, or a matrix X.

Remark 3.6.7 (Derivatives of matrix-valued functions). The derivative of a function $F : \mathbb{R}^{m\times n} \to \mathbb{R}^{p\times q}$ at X is a linear map from $\mathbb{R}^{m\times n}$ to $\mathbb{R}^{p\times q}$. In the language of the section on tensors in Lecture 1, it is an element of $\mathbb{R}^{p\times q} \otimes (\mathbb{R}^{m\times n})^*$, a tensor of order 4 with the entries $\frac{\partial F_{ij}}{\partial X_{kl}}$; arranged as a matrix, it is the $pq \times mn$ Jacobian of the entries. For example, $F(X) = AX$ has $\frac{\partial F_{ij}}{\partial X_{kl}} = A_{ik}\delta_{jl}$. For a scalar loss $L(F(X))$ with $G = \nabla_F L$, the chain rule contracts this tensor with G: in Einstein notation, $(\nabla_X L)_{kl} = G_{ij}\frac{\partial F_{ij}}{\partial X_{kl}}$, which is `np.einsum("ij,ijkl->kl", G, DF)`. For $F(X) = AX$, this is $\sum_i G_{il}A_{ik} = (A^\top G)_{kl}$ (compare Example 3.6.5). In machine learning, we rarely form these large tensors: we only need the gradient of a scalar loss, and the chain rule lets us pass gradients through each function without ever writing down its full Jacobian. This is how backpropagation works, which is the topic of the next section.

3.7 Backpropagation and Automatic Differentiation

To train a neural network, we need the gradient of the loss with respect to millions of parameters, at every step of training. There are three ways to let a computer compute derivatives.

- Symbolic differentiation, as in computer algebra systems, applies the rules of differentiation to formulas. The resulting formulas can become huge, and many programs, with loops and if-statements, are not given by a single formula.
- Numerical differentiation uses finite differences (Proposition 3.2.9). For a function of n variables, the gradient needs at least $n+1$ evaluations of the function, which is far too slow for $n = 10^6$, and the result is only approximate.
- Automatic differentiation (AD) uses the fact that every program computes a function by a sequence of elementary operations, such as $+$, $\times$, exp and log, whose derivatives are known. It applies the chain rule to this sequence, with numbers instead of formulas. The result is exact up to rounding errors, and in reverse mode, the cost of the gradient is only a small multiple of the cost of evaluating the function.

Backpropagation is reverse-mode automatic differentiation, applied to neural networks.

3.7.1 Forward mode and reverse mode

Let $f = f_L \circ \cdots \circ f_2 \circ f_1$ be a composition of functions $f_l : \mathbb{R}^{n_{l-1}} \to \mathbb{R}^{n_l}$, with $n_0 = n$ and $n_L = m$, such as the layers of a neural network. We evaluate f step by step: $\mathbf{x}_0 = \mathbf{x}$ and $\mathbf{x}_l = f_l(\mathbf{x}_{l-1})$ for $l = 1, \dots, L$. By the chain rule,

$$Df(\mathbf{x}) = J_L J_{L-1} \cdots J_1, \qquad \text{where } J_l = Df_l(\mathbf{x}_{l-1}) \in \mathbb{R}^{n_l \times n_{l-1}}.$$

Matrix multiplication is associative, so we can compute this product in any order, but the cost depends on the order: multiplying a $p \times q$ matrix by a $q \times r$ matrix costs pqr multiplications.

- Forward mode computes $J_L(\cdots(J_2(J_1\mathbf{v})))$ for a vector $\mathbf{v} \in \mathbb{R}^n$, i.e., the Jacobian-vector product $Df(\mathbf{x})\mathbf{v}$, from right to left. Each multiplication can be done together with the evaluation of f_l, in the forward direction. For $\mathbf{v} = \mathbf{e}_j$, we get the j-th column of $Df(\mathbf{x})$, so the full Jacobian needs n passes.
- Reverse mode computes $((\mathbf{u}^\top J_L)J_{L-1})\cdots J_1$ for a row vector $\mathbf{u}^\top \in \mathbb{R}^{1\times m}$, i.e., the vector-Jacobian product $\mathbf{u}^\top Df(\mathbf{x})$, from left to right. This goes backwards, from the output to the input, so we first evaluate f and store the intermediate values $\mathbf{x}_0, \dots, \mathbf{x}_{L-1}$, which we need to compute the Jacobians. For $\mathbf{u} = \mathbf{e}_i$, we get the i-th row of $Df(\mathbf{x})$, so the full Jacobian needs m passes.

For a loss function, $m = 1$, and a single backward pass with $\mathbf{u} = 1$ gives the whole gradient. For example, if $n_0 = \cdots = n_{L-1} = n$ and $n_L = 1$, then computing the gradient in forward mode needs n passes, each with about Ln^2 multiplications, so about Ln^3 in total. Reverse mode needs one pass with L vector-matrix products, about Ln^2 multiplications, which is comparable to the cost of evaluating f if the f_l are linear layers. For $n = 1000$ and $L = 10$, this is 10^{10} versus 10^7 multiplications. In general, the cost of reverse mode for a scalar function is a small multiple (typically between 2 and 5) of the cost of evaluating the function, no matter how many variables it has. The price is memory, since all intermediate values have to be stored. In the language of Lecture 1, reverse mode applies the transposes, or dual maps, $J_1^\top \cdots J_L^\top \mathbf{u}$, which is the rule $(S \circ T)^* = T^* \circ S^*$.

3.7.2 Computation graphs

Most programs are not a simple chain of functions. In general, every intermediate variable v_j is computed from some earlier variables by an elementary operation, which we draw as a computation graph, with an arrow $v_i \to v_j$ if v_j is computed from v_i. The two modes of automatic differentiation work as follows.

- Forward mode: choose a direction $\dot{\mathbf{x}} = \mathbf{v}$, and compute the derivatives $\dot{v}_j$ of all intermediate variables in this direction, in the order of evaluation:

$$\dot{v}_j = \sum_{i : v_i \to v_j} \frac{\partial v_j}{\partial v_i} \dot{v}_i.$$

- Reverse mode: first evaluate all v_j, and then compute the adjoints $\bar{v}_i = \frac{\partial f}{\partial v_i}$, which describe how the output depends on v_i, in the reverse order, starting with $\bar{f} = 1$:

$$\bar{v}_i = \sum_{j : v_i \to v_j} \bar{v}_j \frac{\partial v_j}{\partial v_i}.$$

  If v_i is used by several later variables, their contributions are added; this is the sum in the chain rule (Theorem 3.5.1).

Example 3.7.1. Let $f(x_1, x_2) = \log x_1 + x_1 x_2 - \sin x_2$. Figure 3.5 shows how a computer evaluates f, and Table 3.2 shows both modes of automatic differentiation at $(x_1, x_2) = (2, 5)$. Forward mode with the direction $\dot{\mathbf{x}} = \mathbf{e}_1$ gives $\frac{\partial f}{\partial x_1} = 5.5$; for $\frac{\partial f}{\partial x_2}$, we need a second pass with $\dot{\mathbf{x}} = \mathbf{e}_2$. Reverse mode gives both partial derivatives in one pass. Note that $\bar{x}_1$ collects contributions from v_1 and v_2, since x_1 is used to compute both. We check the result: $\frac{\partial f}{\partial x_1} = \frac{1}{x_1} + x_2 = 5.5$ and $\frac{\partial f}{\partial x_2} = x_1 - \cos x_2 \approx 2 - 0.284 = 1.716$.




Figure 3.5: The computation graph of $f(x_1, x_2) = \log x_1 + x_1 x_2 - \sin x_2$. Each arrow $v_i \to v_j$ is labeled with the local partial derivative $\frac{\partial v_j}{\partial v_i}$.

3.7.3 Backpropagation in a neural network

Consider a neural network with one hidden layer, which maps an input $\mathbf{x} \in \mathbb{R}^n$ to an output $\hat{\mathbf{y}} \in \mathbb{R}^m$, and its squared loss for a target $\mathbf{y} \in \mathbb{R}^m$:

$$\mathbf{z} = W_1 \mathbf{x} + \mathbf{b}_1, \qquad \mathbf{a} = \sigma(\mathbf{z}), \qquad \hat{\mathbf{y}} = W_2 \mathbf{a} + \mathbf{b}_2, \qquad L = \tfrac{1}{2} \|\hat{\mathbf{y}} - \mathbf{y}\|^2,$$

evaluation		forward mode, $\dot{\mathbf{x}} = \mathbf{e}_1$	reverse mode
$x_1 = 2$		$\dot{x}_1 = 1$	$\bar{v}_5 = 1$
$x_2 = 5$		$\dot{x}_2 = 0$	$\bar{v}_4 = \bar{v}_5 = 1$
$v_1 = \log x_1$	$= 0.693$	$\dot{v}_1 = \dot{x}_1/x_1 = 0.5$	$\bar{v}_3 = -\bar{v}_5 = -1$
$v_2 = x_1 x_2$	$= 10$	$\dot{v}_2 = \dot{x}_1 x_2 + x_1 \dot{x}_2 = 5$	$\bar{v}_2 = \bar{v}_4 = 1$
$v_3 = \sin x_2$	$= -0.959$	$\dot{v}_3 = \cos(x_2)\, \dot{x}_2 = 0$	$\bar{v}_1 = \bar{v}_4 = 1$
$v_4 = v_1 + v_2$	$= 10.693$	$\dot{v}_4 = \dot{v}_1 + \dot{v}_2 = 5.5$	$\bar{x}_2 = \bar{v}_2 x_1 + \bar{v}_3 \cos x_2 = 1.716$
$v_5 = v_4 - v_3$	$= 11.652$	$\dot{v}_5 = \dot{v}_4 - \dot{v}_3 = 5.5$	$\bar{x}_1 = \bar{v}_1/x_1 + \bar{v}_2 x_2 = 5.5$

Table 3.2: Automatic differentiation of $f(x_1, x_2) = \log x_1 + x_1 x_2 - \sin x_2$ at $(2, 5)$. Left: evaluation and forward mode, from top to bottom. Right: reverse mode, from top to bottom, i.e., backwards through the computation graph.

where $W_1 \in \mathbb{R}^{h \times n}$, $\mathbf{b}_1 \in \mathbb{R}^h$, $W_2 \in \mathbb{R}^{m \times h}$, $\mathbf{b}_2 \in \mathbb{R}^m$, and σ is applied entrywise. The forward pass computes and stores $\mathbf{z}$, $\mathbf{a}$, $\hat{\mathbf{y}}$ and L. The backward pass computes, using Examples 3.6.5 and 3.4.5:

$$\begin{aligned}
\boldsymbol{\delta}_2 &= \nabla_{\hat{\mathbf{y}}} L = \hat{\mathbf{y}} - \mathbf{y}, \\
\nabla_{W_2} L &= \boldsymbol{\delta}_2 \mathbf{a}^\top, \qquad \nabla_{\mathbf{b}_2} L = \boldsymbol{\delta}_2, \\
\nabla_{\mathbf{a}} L &= W_2^\top \boldsymbol{\delta}_2, \\
\boldsymbol{\delta}_1 &= \nabla_{\mathbf{z}} L = D\sigma(\mathbf{z})^\top \nabla_{\mathbf{a}} L = \sigma'(\mathbf{z}) \odot (W_2^\top \boldsymbol{\delta}_2), \\
\nabla_{W_1} L &= \boldsymbol{\delta}_1 \mathbf{x}^\top, \qquad \nabla_{\mathbf{b}_1} L = \boldsymbol{\delta}_1.
\end{aligned}$$

Each line only uses values from the forward pass and from the previous lines. For a deeper network with layers $\mathbf{z}_l = W_l \mathbf{a}_{l-1} + \mathbf{b}_l$, $\mathbf{a}_l = \sigma(\mathbf{z}_l)$, the same pattern repeats: the "error signals" $\boldsymbol{\delta}_l = \nabla_{\mathbf{z}_l} L$ satisfy

$$\boldsymbol{\delta}_l = \sigma'(\mathbf{z}_l) \odot (W_{l+1}^\top \boldsymbol{\delta}_{l+1}), \qquad \nabla_{W_l} L = \boldsymbol{\delta}_l \mathbf{a}_{l-1}^\top, \qquad \nabla_{\mathbf{b}_l} L = \boldsymbol{\delta}_l.$$

The error signals are propagated backwards through the network, which explains the name backpropagation. The backward pass of a layer costs about twice as much as its forward pass: it computes $W_{l+1}^\top \boldsymbol{\delta}_{l+1}$ and the outer product $\boldsymbol{\delta}_l \mathbf{a}_{l-1}^\top$, and each of them costs about as much as the product $W_l \mathbf{a}_{l-1}$ in the forward pass. Machine learning libraries such as PyTorch and JAX implement reverse-mode automatic differentiation for general programs: they record the computation graph during the forward pass, and each elementary operation knows how to compute its vector-Jacobian product.

Remark 3.7.2 (Vanishing and exploding gradients). By the recursion above, $\boldsymbol{\delta}_1$ is obtained from $\boldsymbol{\delta}_L$ by multiplying with the matrices $D\sigma(\mathbf{z}_l)^\top W_{l+1}^\top = \operatorname{diag}(\sigma'(\mathbf{z}_l)) W_{l+1}^\top$, once for every layer. For the sigmoid function, $\sigma' \le \frac{1}{4}$ (Example 3.2.4), so unless the weights are large, the gradients of the first layers become tiny in deep networks (vanishing gradients), and these layers hardly learn. If the weight matrices have large norms, the gradients can instead grow exponentially (exploding gradients). Remedies include the ReLU activation function (whose derivative is 1 for active units), careful initialization of the weights, and residual connections $\mathbf{a}_l = \mathbf{a}_{l-1} + f_l(\mathbf{a}_{l-1})$, whose Jacobian $I + Df_l$ is close to the identity matrix.

Remark 3.7.3 (Gradient checking). Hand-written backward passes are easy to get wrong. To check them, compare the computed gradient $\mathbf{g}$ with central differences $\tilde{g}_i = \frac{f(\mathbf{x}+h\mathbf{e}_i) - f(\mathbf{x}-h\mathbf{e}_i)}{2h}$ for a few random inputs, and compute the relative error $\frac{\|\mathbf{g}-\tilde{\mathbf{g}}\|}{\|\mathbf{g}\|+\|\tilde{\mathbf{g}}\|}$. With $h \approx 10^{-5}$, a relative error around 10^{-8} or smaller indicates a correct gradient, while a mistake typically gives a relative error of order 1.

3.8 Higher Derivatives and Taylor's Theorem

The gradient describes how a function changes to first order. To see whether a critical point is a minimum, and how fast optimization algorithms converge, we need the second derivatives, which describe the curvature of the function.

Definition 3.8.1 (Hessian matrix)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$ open. If the partial derivative $\frac{\partial f}{\partial x_i}$ has a partial derivative with respect to x_j, we write

$$\frac{\partial^2 f}{\partial x_j \partial x_i} = \frac{\partial}{\partial x_j}\left(\frac{\partial f}{\partial x_i}\right), \qquad \frac{\partial^2 f}{\partial x_i^2} = \frac{\partial}{\partial x_i}\left(\frac{\partial f}{\partial x_i}\right).$$

The function f is twice continuously differentiable if all these second partial derivatives exist and are continuous on D. The Hessian matrix of f at $\mathbf{x}$ is the $n \times n$ matrix $\nabla^2 f(\mathbf{x})$ with the entries

$$\left(\nabla^2 f(\mathbf{x})\right)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x}).$$

Theorem 3.8.2 (Schwarz)**.** *If f is twice continuously differentiable on D, then $\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$ for all i, j. In particular, the Hessian matrix is symmetric.*

Proof. Let $i \neq j$ and $\mathbf{x} \in D$, and let $h \neq 0$ be so small that $\mathbf{x} + s\mathbf{e}_i + t\mathbf{e}_j \in D$ whenever $|s|, |t| \leq |h|$. Let

$$\Delta(h) = f(\mathbf{x} + h\mathbf{e}_i + h\mathbf{e}_j) - f(\mathbf{x} + h\mathbf{e}_i) - f(\mathbf{x} + h\mathbf{e}_j) + f(\mathbf{x}).$$

With $\varphi(s) = f(\mathbf{x} + s\mathbf{e}_i + h\mathbf{e}_j) - f(\mathbf{x} + s\mathbf{e}_i)$, we have $\Delta(h) = \varphi(h) - \varphi(0)$. By the mean value theorem, $\Delta(h) = h\varphi'(s_1)$ for some s_1 between 0 and h, i.e.,

$$\Delta(h) = h\Big(\frac{\partial f}{\partial x_i}(\mathbf{x} + s_1\mathbf{e}_i + h\mathbf{e}_j) - \frac{\partial f}{\partial x_i}(\mathbf{x} + s_1\mathbf{e}_i)\Big) = h^2\,\frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{x} + s_1\mathbf{e}_i + t_1\mathbf{e}_j),$$

where we applied the mean value theorem again, to $t \mapsto \frac{\partial f}{\partial x_i}(\mathbf{x} + s_1\mathbf{e}_i + t\mathbf{e}_j)$, with some t_1 between 0 and h. Since $\Delta(h)$ is symmetric in i and j, the same argument with the roles of i and j exchanged gives $\Delta(h) = h^2\,\frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x} + s_2\mathbf{e}_i + t_2\mathbf{e}_j)$ with s_2 and t_2 between 0 and h. Divide both expressions by h^2 and let $h \to 0$. Both points tend to $\mathbf{x}$, so by the continuity of the second partial derivatives, $\frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{x}) = \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x})$. □

The proof shows that $\frac{\Delta(h)}{h^2}$, a second difference quotient, approximates both mixed partial derivatives. Without the continuity of the second partial derivatives, the mixed partial derivatives can differ: for $f(x, y) = \frac{xy(x^2 - y^2)}{x^2 + y^2}$ and $f(0, 0) = 0$, one finds $\frac{\partial^2 f}{\partial y \partial x}(0, 0) = -1$ and $\frac{\partial^2 f}{\partial x \partial y}(0, 0) = 1$. The Hessian is the Jacobian of the gradient, $\nabla^2 f(\mathbf{x}) = D(\nabla f)(\mathbf{x})$: the i-th row of $D(\nabla f)(\mathbf{x})$ is the derivative of $\frac{\partial f}{\partial x_i}$, with the entries $\frac{\partial^2 f}{\partial x_j \partial x_i}$, which are equal to $\frac{\partial^2 f}{\partial x_i \partial x_j}$ by Schwarz's theorem.

Example 3.8.3. For $f(x_1, x_2) = x_1^2 x_2 + x_1 x_2^3$ (Example 3.3.2), we have $\frac{\partial f}{\partial x_1} = 2x_1 x_2 + x_2^3$ and $\frac{\partial f}{\partial x_2} = x_1^2 + 3x_1 x_2^2$, so

$$\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 2x_2 & 2x_1 + 3x_2^2 \\ 2x_1 + 3x_2^2 & 6x_1 x_2 \end{pmatrix}, \qquad \nabla^2 f(1, 2) = \begin{pmatrix} 4 & 14 \\ 14 & 12 \end{pmatrix}.$$

As predicted by Schwarz's theorem, the two mixed partial derivatives are equal.

Example 3.8.4 (Hessians in machine learning)**.**

1. For a quadratic function $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x} + c$ with a symmetric matrix A, Proposition 3.3.3 gives $\nabla f(\mathbf{x}) = A\mathbf{x} - \mathbf{b}$, and by Example 3.4.4, $\nabla^2 f(\mathbf{x}) = A$ for all $\mathbf{x}$. For example, the least squares loss $L(\boldsymbol{\theta}) = \|\mathbf{y} - \Phi\boldsymbol{\theta}\|^2$ has the gradient $2\Phi^\top\Phi\boldsymbol{\theta} - 2\Phi^\top\mathbf{y}$ and the Hessian $2\Phi^\top\Phi$.

2. For logistic regression (Example 3.5.6), $\nabla \ell(\mathbf{w}) = (\sigma(\mathbf{w}^\top \mathbf{x}) - y)\mathbf{x}$. Its Jacobian is $\mathbf{x} \cdot \sigma'(\mathbf{w}^\top \mathbf{x})\mathbf{x}^\top$, so $\nabla^2 \ell(\mathbf{w}) = p(1-p)\mathbf{x}\mathbf{x}^\top$ with $p = \sigma(\mathbf{w}^\top \mathbf{x})$. For a data set, $\nabla^2 L(\mathbf{w}) = \frac{1}{N}\sum_i p_i(1-p_i)\mathbf{x}_i\mathbf{x}_i^\top = \frac{1}{N} X^\top S X$, where $S = \operatorname{diag}(p_1(1-p_1), \dots, p_N(1-p_N))$.
3. The log-sum-exp function $\operatorname{lse}(\mathbf{z}) = \log(e^{z_1} + \cdots + e^{z_K})$ has the gradient $\operatorname{softmax}(\mathbf{z})$ (Example 3.6.6), so its Hessian is the Jacobian of the softmax function, $\nabla^2 \operatorname{lse}(\mathbf{z}) = \operatorname{diag}(\mathbf{p}) - \mathbf{p}\mathbf{p}^\top$ with $\mathbf{p} = \operatorname{softmax}(\mathbf{z})$ (Example 3.4.6).

3.8.1 Taylor's theorem in several variables

Theorem 3.8.5 (Second-order Taylor expansion)**.** *Let f be twice continuously differentiable on an open set D that contains the line segment from $\mathbf{x}$ to $\mathbf{x} + \mathbf{h}$. Then there is a $\tau \in (0,1)$ such that*

$$f(\mathbf{x}+\mathbf{h}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top \mathbf{h} + \frac{1}{2}\mathbf{h}^\top \nabla^2 f(\mathbf{x}+\tau\mathbf{h})\,\mathbf{h}.$$

Consequently,

$$f(\mathbf{x}+\mathbf{h}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top \mathbf{h} + \frac{1}{2}\mathbf{h}^\top \nabla^2 f(\mathbf{x})\,\mathbf{h} + o(\|\mathbf{h}\|^2).$$

Proof. We reduce to one variable. Let $g(t) = f(\mathbf{x} + t\mathbf{h})$ for t in an open interval containing $[0,1]$. By the chain rule (Corollary 3.5.2(1)),

$$g'(t) = \nabla f(\mathbf{x}+t\mathbf{h})^\top \mathbf{h} = \sum_{i=1}^{n} \frac{\partial f}{\partial x_i}(\mathbf{x}+t\mathbf{h})\, h_i,$$

and applying the chain rule again to each $\frac{\partial f}{\partial x_i}$,

$$g''(t) = \sum_{i=1}^{n}\sum_{j=1}^{n} \frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{x}+t\mathbf{h})\, h_j h_i = \mathbf{h}^\top \nabla^2 f(\mathbf{x}+t\mathbf{h})\,\mathbf{h}.$$

By Taylor's theorem in one variable (Theorem 3.2.6 with $k=1$), $g(1) = g(0) + g'(0) + \frac{1}{2}g''(\tau)$ for some $\tau \in (0,1)$, which is the first formula. For the second formula, the difference between the two quadratic terms is

$$\tfrac{1}{2}|\mathbf{h}^\top(\nabla^2 f(\mathbf{x}+\tau\mathbf{h}) - \nabla^2 f(\mathbf{x}))\mathbf{h}| \le \tfrac{1}{2}\|\nabla^2 f(\mathbf{x}+\tau\mathbf{h}) - \nabla^2 f(\mathbf{x})\|_2 \|\mathbf{h}\|^2$$

by the Cauchy–Schwarz inequality, and the spectral norm tends to 0 as $\mathbf{h} \to \mathbf{0}$, since the second partial derivatives are continuous (and the spectral norm is at most the Frobenius norm). □

So near $\mathbf{x}$, f is approximated by the quadratic function

$$f(\mathbf{x}+\mathbf{h}) \approx f(\mathbf{x}) + \nabla f(\mathbf{x})^\top \mathbf{h} + \tfrac{1}{2}\mathbf{h}^\top \nabla^2 f(\mathbf{x})\mathbf{h}.$$

The gradient gives the slope, and the Hessian gives the curvature: for a unit vector $\mathbf{v}$, the number $\mathbf{v}^\top \nabla^2 f(\mathbf{x})\mathbf{v}$ is the second derivative of $t \mapsto f(\mathbf{x}+t\mathbf{v})$ at $t=0$, i.e., the curvature of f in the direction $\mathbf{v}$. The eigenvalues and eigenvectors of the Hessian describe the curvature in all directions at once; we work this out in Section 3.9.1, where it is the key to classifying critical points.

Example 3.8.6. Let $f(x,y) = e^x \cos y$. Then

$$\nabla f(x,y) = \begin{pmatrix} e^x \cos y \\ -e^x \sin y \end{pmatrix}, \qquad \nabla^2 f(x,y) = \begin{pmatrix} e^x \cos y & -e^x \sin y \\ -e^x \sin y & -e^x \cos y \end{pmatrix},$$

so at the origin, $f = 1$, $\nabla f = (1,0)^\top$ and $\nabla^2 f = \operatorname{diag}(1,-1)$. The second-order approximation is

$$e^x \cos y \approx 1 + x + \tfrac{1}{2}(x^2 - y^2).$$

At $(0.1, 0.2)$, it gives 1.085, while the linear approximation $1 + x$ gives 1.1, and the exact value is $1.0831\ldots$.

There are also Taylor expansions of higher order, which use all partial derivatives of order k. They are rarely needed in machine learning, and we do not discuss them.

3.9 Unconstrained Optimization

We now use derivatives to minimize functions, which is how models are trained. Maximizing f is the same as minimizing $-f$, so we only discuss minimization.

Definition 3.9.1 (Minima and critical points)**.** Let $f : D \to \mathbb{R}$ with $D \subseteq \mathbb{R}^n$ open.

1. A point $\mathbf{x}^* \in D$ is a global minimum (or global minimizer) of f if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in D$, and a local minimum if there is an $r > 0$ such that $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in D$ with $\|\mathbf{x}-\mathbf{x}^*\| < r$. It is a strict local minimum if moreover $f(\mathbf{x}^*) < f(\mathbf{x})$ for these $\mathbf{x} \neq \mathbf{x}^*$. Maxima are defined in the same way.

2. If f is differentiable, a point $\mathbf{x}^*$ with $\nabla f(\mathbf{x}^*) = \mathbf{0}$ is called a critical point (or stationary point) of f.

We write $\arg\min_{\mathbf{x}} f(\mathbf{x})$ for the set of global minima of f, or for the global minimum if it is unique.

Theorem 3.9.2 (First-order condition)**.** *If f is differentiable and $\mathbf{x}^*$ is a local minimum of f, then $\nabla f(\mathbf{x}^*) = \mathbf{0}$.*

Proof. Suppose that $\mathbf{g} = \nabla f(\mathbf{x}^*) \neq \mathbf{0}$. By Proposition 3.3.5, the directional derivative of f at $\mathbf{x}^*$ in the direction $-\mathbf{g}$ is $\nabla f(\mathbf{x}^*)^\top(-\mathbf{g}) = -\|\mathbf{g}\|^2 < 0$. So $\frac{f(\mathbf{x}^*-t\mathbf{g})-f(\mathbf{x}^*)}{t} < 0$ for all small $t > 0$, i.e., $f(\mathbf{x}^* - t\mathbf{g}) < f(\mathbf{x}^*)$, and $\mathbf{x}^*$ is not a local minimum. □

The same holds for local maxima. The converse is false: a critical point can be a local minimum, a local maximum, or neither. For example, the origin is a critical point of $g(x, y) = x^2 - y^2$ (Figure 3.1), but g increases along the x-axis and decreases along the y-axis. A critical point that is neither a local minimum nor a local maximum is called a saddle point. To tell these cases apart, we look at the second-order term of the Taylor expansion, and in particular at the eigenvalues of the Hessian.

3.9.1 The second-order test

Let $\nabla f(\mathbf{x}^*) = \mathbf{0}$, and let $H = \nabla^2 f(\mathbf{x}^*)$. Since the gradient vanishes, Theorem 3.8.5 gives

$$f(\mathbf{x}^* + \mathbf{h}) = f(\mathbf{x}^*) + \tfrac{1}{2}\mathbf{h}^\top H\mathbf{h} + o(\|\mathbf{h}\|^2),$$

so near $\mathbf{x}^*$, the shape of f is described by the quadratic form $\mathbf{h}^\top H\mathbf{h}$. The Hessian is symmetric, so by the spectral theorem (Lecture 2) it has an orthonormal basis of eigenvectors $\mathbf{q}_1, \ldots, \mathbf{q}_n$, with real eigenvalues $\lambda_1 \geq \cdots \geq \lambda_n$, and $H = Q\Lambda Q^\top$. Let us measure the displacement $\mathbf{h}$ in this basis: $\mathbf{h} = y_1\mathbf{q}_1 + \cdots + y_n\mathbf{q}_n$, i.e., $\mathbf{y} = Q^\top\mathbf{h}$. Then $H\mathbf{h} = \sum_i \lambda_i y_i \mathbf{q}_i$, and since the $\mathbf{q}_i$ are orthonormal,

$$\mathbf{h}^\top H\mathbf{h} = \lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2 \qquad \text{and} \qquad \|\mathbf{h}\|^2 = y_1^2 + \cdots + y_n^2.$$

Therefore, near the critical point,

$$f(\mathbf{x}^* + y_1\mathbf{q}_1 + \cdots + y_n\mathbf{q}_n) \approx f(\mathbf{x}^*) + \tfrac{1}{2}(\lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2).$$

In the coordinates given by the eigenvectors, the mixed terms $y_i y_j$ disappear: near a critical point, f looks like a sum of n independent parabolas, one for each eigenvector. Along the line through $\mathbf{x}^*$ in the direction $\mathbf{q}_i$,

$$f(\mathbf{x}^* + t\mathbf{q}_i) \approx f(\mathbf{x}^*) + \tfrac{1}{2}\lambda_i t^2,$$

a parabola that opens upward if $\lambda_i > 0$ and downward if $\lambda_i < 0$. So the eigenvalue λ_i is the curvature of f in the direction $\mathbf{q}_i$, i.e., the second derivative of $t \mapsto f(\mathbf{x}^* + t\mathbf{q}_i)$ at $t = 0$. The eigenvectors are called the principal directions (or principal axes) of f at $\mathbf{x}^*$. In any other unit

direction $\mathbf{v}$, with $\mathbf{y} = Q^\top \mathbf{v}$, the curvature is $\mathbf{v}^\top H \mathbf{v} = \sum_i \lambda_i y_i^2$ with $\sum_i y_i^2 = 1$, a weighted average of the eigenvalues; so it lies between λ_n and λ_1, as for the Rayleigh quotient in Lecture 2.

Hence the signs of the eigenvalues determine the shape of f near a critical point (Figure 3.6 and Table 3.3). If all eigenvalues are positive, f curves upward in every direction, like a bowl, and $\mathbf{x}^*$ is a minimum. If all are negative, f curves downward in every direction, like an upside-down bowl, and $\mathbf{x}^*$ is a maximum. If there are eigenvalues of both signs, f curves upward in some directions and downward in others, like a saddle or a mountain pass. And if some eigenvalues are 0, then f is flat to second order in these directions, like a trough, and the quadratic approximation cannot decide what happens: this depends on the terms of higher order.




Figure 3.6: The quadratic approximation $f(\mathbf{x}^*) + \frac{1}{2}(\lambda_1 y_1^2 + \lambda_2 y_2^2)$ of a function near a critical point, in the coordinates y_1, y_2 along the eigenvectors of the Hessian. The curves are the parabolas $\frac{1}{2}\lambda_i t^2$ along the eigenvectors: orange for $\lambda_i > 0$, violet for $\lambda_i < 0$, and gray for $\lambda_i = 0$. Left: $\lambda_1, \lambda_2 > 0$, a bowl (local minimum). Middle: $\lambda_1 > 0 > \lambda_2$, a saddle (saddle point). Right: $\lambda_1 > 0$ and $\lambda_2 = 0$, a trough (the second-order test is inconclusive).

Theorem 3.9.3 (Second-order test)**.** *Let f be twice continuously differentiable on D, let $\nabla f(\mathbf{x}^*) = \mathbf{0}$, and let $\lambda_1 \geq \cdots \geq \lambda_n$ be the eigenvalues of the Hessian $\nabla^2 f(\mathbf{x}^*)$.*

1. *If all eigenvalues are positive ($\lambda_n > 0$), then $\mathbf{x}^*$ is a strict local minimum.*
2. *If all eigenvalues are negative ($\lambda_1 < 0$), then $\mathbf{x}^*$ is a strict local maximum.*
3. *If there are positive and negative eigenvalues ($\lambda_1 > 0 > \lambda_n$), then $\mathbf{x}^*$ is a saddle point: f increases along every eigenvector with a positive eigenvalue, and decreases along every eigenvector with a negative eigenvalue.*
4. *Conversely, if $\mathbf{x}^*$ is a local minimum, then all eigenvalues are ≥ 0, and if $\mathbf{x}^*$ is a local maximum, then all eigenvalues are ≤ 0.*

Proof. Let $H = \nabla^2 f(\mathbf{x}^*)$, and write $f(\mathbf{x}^* + \mathbf{h}) = f(\mathbf{x}^*) + \frac{1}{2}\mathbf{h}^\top H \mathbf{h} + r(\mathbf{h})$ with $r(\mathbf{h}) = o(\|\mathbf{h}\|^2)$, as above.

(1) In the coordinates $\mathbf{y} = Q^\top \mathbf{h}$, we have $\mathbf{h}^\top H \mathbf{h} = \sum_i \lambda_i y_i^2 \geq \lambda_n \sum_i y_i^2 = \lambda_n \|\mathbf{h}\|^2$: in every direction, f curves upward at least as much as along $\mathbf{q}_n$. For all sufficiently small $\mathbf{h} \neq \mathbf{0}$, we have $|r(\mathbf{h})| \leq \frac{1}{4}\lambda_n \|\mathbf{h}\|^2$, and hence $f(\mathbf{x}^* + \mathbf{h}) \geq f(\mathbf{x}^*) + \frac{1}{4}\lambda_n \|\mathbf{h}\|^2 > f(\mathbf{x}^*)$.

(2) Apply (1) to $-f$, whose Hessian has the eigenvalues $-\lambda_i$.

(3) Let $\mathbf{q}$ be a unit eigenvector with eigenvalue $\lambda \neq 0$. Then $f(\mathbf{x}^* + t\mathbf{q}) = f(\mathbf{x}^*) + \frac{1}{2}\lambda t^2 + r(t\mathbf{q})$, and $|r(t\mathbf{q})| < \frac{1}{2}|\lambda| t^2$ for all small $t \neq 0$. So for all small $t \neq 0$, $f(\mathbf{x}^* + t\mathbf{q}) > f(\mathbf{x}^*)$ if $\lambda > 0$, and $f(\mathbf{x}^* + t\mathbf{q}) < f(\mathbf{x}^*)$ if $\lambda < 0$. If both signs occur, there are points with larger and points with smaller values arbitrarily close to $\mathbf{x}^*$, so $\mathbf{x}^*$ is neither a local minimum nor a local maximum.

(4) If some eigenvalue were negative, then by the argument of (3), there would be points with smaller values arbitrarily close to $\mathbf{x}^*$, so $\mathbf{x}^*$ would not be a local minimum. The case of a local maximum is the same. □

In the language of Lecture 2, the Hessian in (1) is positive definite, in (2) negative definite, and in (3) indefinite, and (4) says that the Hessian at a local minimum is positive semidefinite. If some eigenvalues are 0 and all others have the same sign, the test is inconclusive: x^4, $-x^4$ and x^3 all have the critical point 0 with second derivative 0, but the first has a minimum there, the second a maximum, and the third neither (see also Exercise 16).

For $n = 2$, we do not even need to compute the eigenvalues, since $\lambda_1\lambda_2 = \det(H)$ and $\lambda_1 + \lambda_2 = \operatorname{tr}(H)$ (Lecture 1). If $\det(H) < 0$, the eigenvalues have opposite signs, and the critical point is a saddle point. If $\det(H) > 0$, they have the same sign, which is the sign of $\operatorname{tr}(H)$: the critical point is a strict local minimum if $\operatorname{tr}(H) > 0$, and a strict local maximum if $\operatorname{tr}(H) < 0$.

eigenvalues of $\nabla^2 f(\mathbf{x}^*)$	graph near $\mathbf{x}^*$	level curves	$\mathbf{x}^*$ is a
all positive	bowl	ellipses	strict local minimum
all negative	upside-down bowl	ellipses	strict local maximum
positive and negative	saddle	hyperbolas	saddle point
some zero, the rest of one sign	trough	parallel lines	(test inconclusive)

Table 3.3: The second-order test at a critical point $\mathbf{x}^*$. The shape of the graph and the level curves (for $n = 2$) are those of the quadratic approximation $f(\mathbf{x}^*) + \frac{1}{2}(\lambda_1 y_1^2 + \cdots + \lambda_n y_n^2)$.

The eigenvalues also describe the level sets near a strict local minimum. For $c > 0$, the level set $\frac{1}{2}(\lambda_1 y_1^2 + \cdots + \lambda_n y_n^2) = c$ of the quadratic approximation is an ellipse (for $n = 2$) or an ellipsoid, whose axes point in the directions of the eigenvectors $\mathbf{q}_i$ and have the half-lengths $\sqrt{2c/\lambda_i}$, as in the section on the spectral theorem in Lecture 2. So the level curves around a minimum are approximately ellipses that are long in directions of small curvature and short in directions of large curvature. The ratio of the longest and the shortest axis is $\sqrt{\lambda_1/\lambda_n}$, the square root of the condition number of the Hessian, and we will see in Section 3.10 that this elongation determines how fast gradient descent converges. Around a saddle point in the plane, the level curves of the quadratic approximation are hyperbolas, as for $x^2 - y^2$ in Figure 3.1.

Example 3.9.4. Let $f(x, y) = x^3 + y^3 - 3xy$. Its gradient is $\nabla f(x, y) = (3x^2 - 3y,\ 3y^2 - 3x)^\top$, so at a critical point, $y = x^2$ and $x = y^2 = x^4$. Hence $x = 0$ or $x^3 = 1$, i.e., $x = 1$, and the critical points are $(0, 0)$ and $(1, 1)$. The Hessian is

$$\nabla^2 f(x, y) = \begin{pmatrix} 6x & -3 \\ -3 & 6y \end{pmatrix}.$$

- At $(1, 1)$, the Hessian $\begin{pmatrix} 6 & -3 \\ -3 & 6 \end{pmatrix}$ has the eigenvector $(1, 1)^\top$ with eigenvalue 3, and the eigenvector $(1, -1)^\top$ with eigenvalue 9. Both eigenvalues are positive, so $(1, 1)$ is a strict local minimum, with $f(1, 1) = -1$. The curvature across the diagonal is three times as large as the curvature along it, so the level curves around $(1, 1)$ are approximately ellipses that are elongated in the direction of the diagonal, with axes in the ratio $\sqrt{9/3} = \sqrt{3}$ (Figure 3.7). The minimum is not global, since $f(x, 0) = x^3 \to -\infty$ as $x \to -\infty$.
- At $(0, 0)$, the Hessian $\begin{pmatrix} 0 & -3 \\ -3 & 0 \end{pmatrix}$ has the eigenvector $(1, 1)^\top$ with eigenvalue -3, and the eigenvector $(1, -1)^\top$ with eigenvalue 3. So the origin is a saddle point: f decreases along the diagonal $y = x$ and increases along the diagonal $y = -x$. Indeed, $f(t, t) = 2t^3 - 3t^2 \approx -3t^2$ and $f(t, -t) = 3t^2$ for small t, which are the parabolas $\frac{1}{2}\lambda t^2 \|\mathbf{v}\|^2$ for $\mathbf{v} = (1, \pm 1)^\top$. Note that both diagonal entries of the Hessian are 0: along the coordinate axes, $f(t, 0) = f(0, t) = t^3$ has

no curvature at $t = 0$, and we would not see the saddle at all. As in Example 3.1.8, looking along the coordinate axes is not enough; the eigenvectors show the directions in which f really curves.

The level set $f = 0$ is the folium of Descartes $x^3 + y^3 = 3xy$, which crosses itself at the saddle point. Near the origin, it looks like the zero set of the quadratic approximation $\frac{1}{2}\mathbf{h}^\top H\mathbf{h} = -3xy$, which consists of the two coordinate axes.




Figure 3.7: Level curves of $f(x, y) = x^3 + y^3 - 3xy$ for the levels $-0.9, -0.6, -0.3$ (blue), 0 (black: the folium of Descartes) and $0.6, 1.5, 3$ (red), and the eigenvectors of the Hessian at the two critical points (orange: positive eigenvalue, violet: negative eigenvalue). Around the minimum $(1, 1)$, the level curves are approximately ellipses with axes along the eigenvectors, longer in the direction of the smaller eigenvalue. At the saddle point $(0, 0)$, f decreases in the violet direction and increases in the orange one.

Example 3.9.5 (Quadratic functions)**.** Let $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$, where A is symmetric positive definite. Then $\nabla f(\mathbf{x}) = A\mathbf{x} - \mathbf{b}$, so the only critical point is $\mathbf{x}^* = A^{-1}\mathbf{b}$, and the Hessian A is positive definite. In fact, $\mathbf{x}^*$ is the unique global minimum: using $\mathbf{b} = A\mathbf{x}^*$, one checks that

$$f(\mathbf{x}) - f(\mathbf{x}^*) = \tfrac{1}{2}(\mathbf{x} - \mathbf{x}^*)^\top A(\mathbf{x} - \mathbf{x}^*),$$

which is positive for $\mathbf{x} \neq \mathbf{x}^*$. The level sets of f are ellipsoids centered at $\mathbf{x}^*$, whose axes point in the directions of the eigenvectors of A (Lecture 2). So solving the linear system $A\mathbf{x} = \mathbf{b}$ with a symmetric positive definite matrix is the same as minimizing a quadratic function; this is the starting point of iterative methods for large linear systems, such as the conjugate gradient method. For example, the least squares loss $\|\mathbf{y} - \Phi\boldsymbol{\theta}\|^2 = \boldsymbol{\theta}^\top\Phi^\top\Phi\boldsymbol{\theta} - 2\mathbf{y}^\top\Phi\boldsymbol{\theta} + \mathbf{y}^\top\mathbf{y}$ is of this form with $A = 2\Phi^\top\Phi$ and $\mathbf{b} = 2\Phi^\top\mathbf{y}$, plus a constant. If Φ has linearly independent columns, then $\Phi^\top\Phi$ is positive definite (since $\boldsymbol{\theta}^\top\Phi^\top\Phi\boldsymbol{\theta} = \|\Phi\boldsymbol{\theta}\|^2 > 0$ for $\boldsymbol{\theta} \neq \mathbf{0}$), and the unique minimum is $\boldsymbol{\theta}^* = (\Phi^\top\Phi)^{-1}\Phi^\top\mathbf{y}$, as in Lecture 2.

3.10 Gradient Descent and Newton's Method

For most models, such as logistic regression and neural networks, the equation $\nabla f(\mathbf{x}) = \mathbf{0}$ cannot be solved by a formula. Instead, we start with a guess $\mathbf{x}_0$ and improve it step by step.

Definition 3.10.1 (Gradient descent)**.** Let $f : \mathbb{R}^n \to \mathbb{R}$ be differentiable, let $\mathbf{x}_0 \in \mathbb{R}^n$, and let $\eta > 0$. Gradient descent with step size (or learning rate) η computes

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \eta \nabla f(\mathbf{x}_k), \qquad k = 0, 1, 2, \ldots$$

By Theorem 3.3.9, $-\nabla f(\mathbf{x}_k)$ is the direction of steepest descent, and by the linear approximation,

$$f(\mathbf{x}_k - \eta \nabla f(\mathbf{x}_k)) = f(\mathbf{x}_k) - \eta \|\nabla f(\mathbf{x}_k)\|^2 + o(\eta),$$

so a small enough step decreases f, unless $\nabla f(\mathbf{x}_k) = \mathbf{0}$. But how small is small enough? If η is too small, we make little progress; if it is too large, the linear approximation is no longer valid, and f can increase. The answer depends on how fast the gradient changes.

Example 3.10.2 (Gradient descent for linear regression)**.** For linear regression with the mean squared error $L(\mathbf{w}) = \frac{1}{N}\|X\mathbf{w} - \mathbf{y}\|^2 = \frac{1}{N}\sum_i (\mathbf{w}^\top \mathbf{x}_i - y_i)^2$, Example 3.5.5 gives $\nabla L(\mathbf{w}) = \frac{2}{N} X^\top (X\mathbf{w} - \mathbf{y})$. So gradient descent is

$$\mathbf{w}_{k+1} = \mathbf{w}_k - \eta \frac{2}{N} \sum_{i=1}^{N} (\mathbf{w}_k^\top \mathbf{x}_i - y_i)\, \mathbf{x}_i.$$

Each data point pulls the weights in the direction of its input $\mathbf{x}_i$, in proportion to its residual: badly predicted points have the most influence, and perfectly predicted points none. Unlike the normal equations, each step only needs two matrix-vector products, which cost $O(Nd)$ for N data points with d features, so the method works for very large data sets, and with mini-batches (Remark 3.10.10) even for data that do not fit into memory.

For example, for the model $y = wx$ with a single weight and the data $(1, 2)$, $(2, 3)$, $(3, 5)$, we have $L'(w) = \frac{2}{3}\sum_i (wx_i - y_i)x_i = \frac{2}{3}(14w - 23)$, so the minimum is $w^* = \frac{23}{14} \approx 1.643$. With $\eta = 0.1$ and $w_0 = 0$, the iterates are $w_1 = 1.533$, $w_2 = 1.636$, $w_3 = 1.642$: the error is multiplied by $1 - 0.1 \cdot \frac{28}{3} = \frac{1}{15}$ in every step. Since $L'' = \frac{28}{3}$, gradient descent converges if and only if $\eta < \frac{2}{28/3} = \frac{3}{14} \approx 0.21$ (Theorem 3.10.7 below); for $\eta = 0.3$, the error is multiplied by $1 - 2.8 = -1.8$ in every step, and the iterates diverge.

3.10.1 Convergence for smooth functions

Definition 3.10.3 (L-smooth function)**.** A differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ is L-smooth, for a number $L > 0$, if its gradient is Lipschitz continuous with constant L:

$$\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\| \le L\|\mathbf{x} - \mathbf{y}\| \qquad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

For example, the quadratic function $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top \mathbf{x}$ with a symmetric matrix A has $\nabla f(\mathbf{x}) - \nabla f(\mathbf{y}) = A(\mathbf{x} - \mathbf{y})$, so it is L-smooth with $L = \|A\|_2$, which is the largest absolute value of an eigenvalue of A. More generally, a twice continuously differentiable function is L-smooth if and only if $\|\nabla^2 f(\mathbf{x})\|_2 \le L$ for all $\mathbf{x}$: the curvature is bounded by L. Indeed, if the Hessians are bounded, then $\nabla f(\mathbf{x}) - \nabla f(\mathbf{y}) = \int_0^1 \nabla^2 f(\mathbf{y} + t(\mathbf{x} - \mathbf{y}))(\mathbf{x} - \mathbf{y})\, dt$ by the fundamental theorem of calculus, applied to each entry, and the norm of this integral is at most $L\|\mathbf{x} - \mathbf{y}\|$ (the norm of an integral of a vector-valued function is at most the integral of its norm, as we show in the proof of Theorem 3.10.11). Conversely, $\nabla^2 f(\mathbf{x})\mathbf{v} = \lim_{t\to 0} \frac{1}{t}(\nabla f(\mathbf{x} + t\mathbf{v}) - \nabla f(\mathbf{x}))$ has norm at most $L\|\mathbf{v}\|$ if f is L-smooth.

Lemma 3.10.4 (Descent lemma)**.** *If f is L-smooth, then for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$,*

$$f(\mathbf{y}) \le f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{L}{2}\|\mathbf{y} - \mathbf{x}\|^2.$$

Proof. Let $\mathbf{d} = \mathbf{y} - \mathbf{x}$ and $g(t) = f(\mathbf{x} + t\mathbf{d})$. By the chain rule, $g'(t) = \nabla f(\mathbf{x} + t\mathbf{d})^\top \mathbf{d}$, which is continuous in t, since ∇f is Lipschitz continuous. By the fundamental theorem of calculus,

$$f(\mathbf{y}) - f(\mathbf{x}) - \nabla f(\mathbf{x})^\top \mathbf{d} = \int_0^1 \big(g'(t) - g'(0)\big)\,dt = \int_0^1 \big(\nabla f(\mathbf{x} + t\mathbf{d}) - \nabla f(\mathbf{x})\big)^\top \mathbf{d}\,dt.$$

By the Cauchy–Schwarz inequality and L-smoothness, the integrand is at most $\|\nabla f(\mathbf{x} + t\mathbf{d}) - \nabla f(\mathbf{x})\|\|\mathbf{d}\| \le Lt\|\mathbf{d}\|^2$, and $\int_0^1 Lt\|\mathbf{d}\|^2\,dt = \frac{L}{2}\|\mathbf{d}\|^2$. □

So an L-smooth function lies below the quadratic function on the right, which touches it at $\mathbf{x}$. Gradient descent with step size $\frac{1}{L}$ minimizes this quadratic upper bound: the right-hand side is a quadratic function of $\mathbf{y}$ with gradient $\nabla f(\mathbf{x}) + L(\mathbf{y} - \mathbf{x})$, which is zero for $\mathbf{y} = \mathbf{x} - \frac{1}{L}\nabla f(\mathbf{x})$.

Theorem 3.10.5. *Let f be L-smooth and bounded below, $f(\mathbf{x}) \ge f^*$ for all $\mathbf{x}$, and let $0 < \eta \le \frac{1}{L}$. Then gradient descent satisfies*

$$f(\mathbf{x}_{k+1}) \le f(\mathbf{x}_k) - \frac{\eta}{2}\|\nabla f(\mathbf{x}_k)\|^2 \qquad \text{and} \qquad \min_{0 \le k < K} \|\nabla f(\mathbf{x}_k)\|^2 \le \frac{2\big(f(\mathbf{x}_0) - f^*\big)}{\eta K}.$$

Proof. By the descent lemma with $\mathbf{x} = \mathbf{x}_k$ and $\mathbf{y} = \mathbf{x}_{k+1} = \mathbf{x}_k - \eta\nabla f(\mathbf{x}_k)$,

$$f(\mathbf{x}_{k+1}) \le f(\mathbf{x}_k) - \eta\|\nabla f(\mathbf{x}_k)\|^2 + \frac{L\eta^2}{2}\|\nabla f(\mathbf{x}_k)\|^2 = f(\mathbf{x}_k) - \eta\Big(1 - \frac{L\eta}{2}\Big)\|\nabla f(\mathbf{x}_k)\|^2,$$

and $1 - \frac{L\eta}{2} \ge \frac{1}{2}$ since $L\eta \le 1$. Adding these inequalities for $k = 0, \dots, K-1$ gives

$$\frac{\eta}{2}\sum_{k=0}^{K-1} \|\nabla f(\mathbf{x}_k)\|^2 \le f(\mathbf{x}_0) - f(\mathbf{x}_K) \le f(\mathbf{x}_0) - f^*,$$

and the minimum of the K numbers $\|\nabla f(\mathbf{x}_k)\|^2$ is at most their average. □

So with a small enough step size, gradient descent decreases f in every step, and it finds points where the gradient is small. It does not guarantee that we find a global minimum, or even a local minimum: it can approach a saddle point, or a local minimum that is much worse than the global one. For the non-convex loss functions of neural networks, this is essentially the best general guarantee. For convex functions (Section 3.11), the situation is much better.

In practice, L is unknown, and a step size that works in one region may be too large in another. Backtracking line search chooses the step size in every iteration: starting from a trial step $\eta = \bar{\eta}$, it halves η until the sufficient decrease condition (Armijo condition)

$$f\big(\mathbf{x}_k - \eta\nabla f(\mathbf{x}_k)\big) \le f(\mathbf{x}_k) - \frac{\eta}{2}\|\nabla f(\mathbf{x}_k)\|^2$$

holds, and then takes the step $\mathbf{x}_{k+1} = \mathbf{x}_k - \eta\nabla f(\mathbf{x}_k)$.

Proposition 3.10.6 (Backtracking line search)**.** *Let f be L-smooth and bounded below by f^*. Then in every iteration, backtracking stops after finitely many halvings, with a step size $\eta_k \ge \eta_{\min} = \min\big(\bar{\eta}, \frac{1}{2L}\big)$, and*

$$\min_{0 \le k < K} \|\nabla f(\mathbf{x}_k)\|^2 \le \frac{2\big(f(\mathbf{x}_0) - f^*\big)}{\eta_{\min} K}.$$

Proof. By the proof of Theorem 3.10.5, the sufficient decrease condition holds for every $\eta \le \frac{1}{L}$. So the halving stops at the latest at the first trial step $\bar{\eta}2^{-j}$ that is at most $\frac{1}{L}$. This step is either $\bar{\eta}$ itself, or it is larger than $\frac{1}{2L}$, since twice this step was larger than $\frac{1}{L}$. Hence $\eta_k \ge \eta_{\min}$, and $f(\mathbf{x}_{k+1}) \le f(\mathbf{x}_k) - \frac{\eta_{\min}}{2}\|\nabla f(\mathbf{x}_k)\|^2$. Adding these inequalities for $k = 0, \dots, K-1$ gives the bound, as in the proof of Theorem 3.10.5. □

So backtracking does almost as well as the best constant step size $\frac{1}{L}$ without knowing L, and it takes larger steps where f is flat. The price is a few extra evaluations of f per iteration. In practice, the factor $\frac{1}{2}$ in the condition is often replaced by a smaller constant, such as 10^{-4}, and η is multiplied by some $\beta \in (0, 1)$ instead of $\frac{1}{2}$ (Exercise 19).

3.10.2 Gradient descent on quadratic functions

Quadratic functions show precisely how the speed of gradient descent depends on the shape of the function. They are also important in general, since every twice continuously differentiable function is approximately quadratic near a minimum, with the eigenvalues of the Hessian as the curvatures along its principal directions (Section 3.9.1).

Let $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$, where A is symmetric positive definite with eigenvalues $\lambda_{\max} = \lambda_1 \geq \cdots \geq \lambda_n = \lambda_{\min} > 0$ and orthonormal eigenvectors $\mathbf{q}_1, \ldots, \mathbf{q}_n$. By Example 3.9.5, the minimum is $\mathbf{x}^* = A^{-1}\mathbf{b}$, and $\nabla f(\mathbf{x}) = A\mathbf{x} - \mathbf{b} = A(\mathbf{x} - \mathbf{x}^*)$. So the error $\mathbf{e}_k = \mathbf{x}_k - \mathbf{x}^*$ satisfies

$$\mathbf{e}_{k+1} = \mathbf{e}_k - \eta A\mathbf{e}_k = (I - \eta A)\mathbf{e}_k, \qquad \text{hence} \qquad \mathbf{e}_k = (I - \eta A)^k \mathbf{e}_0.$$

The matrix $I - \eta A$ has the same eigenvectors as A, with the eigenvalues $1 - \eta\lambda_i$. So if we write $\mathbf{e}_0 = \sum_i c_i \mathbf{q}_i$, then

$$\mathbf{e}_k = \sum_{i=1}^{n} (1 - \eta\lambda_i)^k c_i \mathbf{q}_i :$$

in each eigenvector direction, the error is multiplied by $1 - \eta\lambda_i$ in every step.

Theorem 3.10.7. *Let f be as above, and let $\kappa = \frac{\lambda_{\max}}{\lambda_{\min}}$ be the condition number of A.*

1. *$\|\mathbf{e}_k\| \leq \rho^k \|\mathbf{e}_0\|$, where $\rho = \max_i |1 - \eta\lambda_i| = \max\left(|1 - \eta\lambda_{\max}|, |1 - \eta\lambda_{\min}|\right)$.*
2. *Gradient descent converges to $\mathbf{x}^*$ for every $\mathbf{x}_0$ if and only if $0 < \eta < \frac{2}{\lambda_{\max}}$.*
3. *The rate ρ is smallest for $\eta = \frac{2}{\lambda_{\max} + \lambda_{\min}}$, and then $\rho = \frac{\kappa - 1}{\kappa + 1}$.*

Proof. (1) Since the $\mathbf{q}_i$ are orthonormal, $\|\mathbf{e}_k\|^2 = \sum_i (1 - \eta\lambda_i)^{2k} c_i^2 \leq \rho^{2k} \sum_i c_i^2 = \rho^{2k} \|\mathbf{e}_0\|^2$. The maximum of $|1 - \eta\lambda|$ over $\lambda \in [\lambda_{\min}, \lambda_{\max}]$ is attained at one of the endpoints.

(2) If $0 < \eta < \frac{2}{\lambda_{\max}}$, then $-1 < 1 - \eta\lambda_i < 1$ for all i, so $\rho < 1$ and $\mathbf{e}_k \to \mathbf{0}$ by (1). Conversely, if $\eta \geq \frac{2}{\lambda_{\max}}$, then $1 - \eta\lambda_1 \leq -1$, and for $\mathbf{x}_0 = \mathbf{x}^* + \mathbf{q}_1$, the error $\mathbf{e}_k = (1 - \eta\lambda_1)^k \mathbf{q}_1$ does not tend to $\mathbf{0}$.

(3) Let $\eta^* = \frac{2}{\lambda_{\max} + \lambda_{\min}}$. Then $1 - \eta^*\lambda_{\min} = \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} = \frac{\kappa - 1}{\kappa + 1}$ and $1 - \eta^*\lambda_{\max} = -\frac{\kappa - 1}{\kappa + 1}$, so $\rho(\eta^*) = \frac{\kappa - 1}{\kappa + 1}$. For $\eta < \eta^*$, we have $|1 - \eta\lambda_{\min}| \geq 1 - \eta\lambda_{\min} > 1 - \eta^*\lambda_{\min}$, and for $\eta > \eta^*$, we have $|1 - \eta\lambda_{\max}| \geq \eta\lambda_{\max} - 1 > \eta^*\lambda_{\max} - 1$. In both cases, $\rho(\eta) > \rho(\eta^*)$. □

Example 3.10.8. Let $f(\mathbf{x}) = \frac{1}{2}(x_1^2 + 10x_2^2)$, i.e., $A = \operatorname{diag}(1, 10)$, $\mathbf{b} = \mathbf{0}$ and $\mathbf{x}^* = \mathbf{0}$. Gradient descent multiplies x_1 by $1 - \eta$ and x_2 by $1 - 10\eta$ in every step. It converges if and only if $\eta < 0.2$; for $\eta = 0.21$, for example, $|x_2|$ grows by the factor 1.1 in every step. The best step size is $\eta = \frac{2}{11}$, with $\rho = \frac{9}{11} \approx 0.82$: then x_1 is multiplied by $\frac{9}{11}$, and x_2 by $-\frac{9}{11}$, so the iterates jump back and forth across the valley, while they slowly move along it (Figure 3.8). To reduce the error by a factor 10^{-6}, we need k steps with $(\frac{9}{11})^k \leq 10^{-6}$, i.e., $k \geq 69$. For $A = I$, the step size $\eta = 1$ reaches the minimum in a single step.

For large κ, the rate $\frac{\kappa - 1}{\kappa + 1} \approx 1 - \frac{2}{\kappa}$ is close to 1, and the number of steps needed for a given accuracy grows proportionally to κ. Ill-conditioned problems, whose level sets are long and thin ellipsoids, are slow for gradient descent. This is the zig-zagging that was mentioned at the end of the section on the spectral theorem in Lecture 2. Section 3.15.2 explains the step size limit $\frac{2}{\lambda_{\max}}$ from another angle: gradient descent is Euler's method for the gradient flow $\mathbf{x}' = -\nabla f(\mathbf{x})$, and $\frac{2}{\lambda_{\max}}$ is the limit at which Euler's method becomes unstable.

Remark 3.10.9 (Making gradient descent faster). 1. Feature scaling. For least squares, the Hessian is $2\Phi^\top\Phi$. If the features (the columns of Φ) are measured on very different scales, then κ is large. Standardizing each feature, i.e., subtracting its mean and dividing by its standard deviation, often reduces κ dramatically.




Figure 3.8: Gradient descent on $f(\mathbf{x}) = \frac{1}{2}(x_1^2 + 10x_2^2)$ with the optimal step size $\eta = \frac{2}{11}$, starting at $\mathbf{x}_0 = (8, 2)^\top$ (red: the first 14 steps). The ellipses are level curves of f. Since the condition number is $\kappa = 10$, the level curves are elongated, the gradient points mostly across the valley instead of towards the minimum, and the iterates zig-zag. Newton's method (dashed, Section 3.10.3) reaches the minimum in one step.

2. Momentum. The heavy ball method $\mathbf{x}_{k+1} = \mathbf{x}_k - \eta\nabla f(\mathbf{x}_k) + \beta(\mathbf{x}_k - \mathbf{x}_{k-1})$ keeps some of the previous step. This damps the zig-zagging and accelerates the motion along the valley. For quadratic functions, with well-chosen η and β, the error decreases roughly like $(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1})^k$, so the number of steps grows only like $\sqrt{\kappa}$.

3. Preconditioning. Gradient descent with respect to a different inner product $\langle \mathbf{x}, \mathbf{y}\rangle_P = \mathbf{x}^\top P\mathbf{y}$ uses the step $-\eta P^{-1}\nabla f(\mathbf{x})$ (Remark 3.3.8), which is the same as gradient descent after a linear change of variables. A good choice of P makes the level sets rounder. Optimizers such as Adam use a diagonal matrix P that is adapted during training. The extreme case $P = \nabla^2 f(\mathbf{x})$ is Newton's method.

Remark 3.10.10 (Stochastic gradient descent). In machine learning, the loss is an average over the training data, $L(\boldsymbol{\theta}) = \frac{1}{N}\sum_{i=1}^N \ell_i(\boldsymbol{\theta})$, where ℓ_i is the loss on the i-th data point. Computing the full gradient needs a pass over all N data points, which is expensive for large data sets. Stochastic gradient descent (SGD) instead chooses a small random subset $B \subseteq \{1, \dots, N\}$ (a mini-batch) in every step and uses the mini-batch gradient

$$\mathbf{g} = \frac{1}{|B|}\sum_{i\in B} \nabla \ell_i(\boldsymbol{\theta}),$$

which is cheap to compute. If B is chosen uniformly at random, then the expected value of $\mathbf{g}$ is the full gradient $\nabla L(\boldsymbol{\theta})$, so on average SGD moves in the right direction. The noise in $\mathbf{g}$ means that SGD does not settle exactly at a minimum unless the step size is decreased during training. Theorem 4.7.16 proves that SGD converges for strongly convex functions (Section 3.11.1) with the step sizes $\eta_k = \frac{1}{\mu(k+1)}$, at the rate $\frac{1}{k}$, and that with a constant step size it approaches a neighborhood of the minimum whose size is proportional to the step size. In practice, SGD and its variants with momentum and preconditioning are used to train almost all neural networks.

3.10.3 Newton's method

Gradient descent only uses the first-order approximation of f. Newton's method uses the second-order approximation: at $\mathbf{x}_k$, it minimizes the quadratic model

$$m_k(\mathbf{h}) = f(\mathbf{x}_k) + \nabla f(\mathbf{x}_k)^\top \mathbf{h} + \tfrac{1}{2}\mathbf{h}^\top \nabla^2 f(\mathbf{x}_k)\mathbf{h}$$

of Theorem 3.8.5. If the Hessian $H_k = \nabla^2 f(\mathbf{x}_k)$ is positive definite, then by Example 3.9.5 (with $A = H_k$ and $\mathbf{b} = -\nabla f(\mathbf{x}_k)$), the minimum of m_k is $\mathbf{h} = -H_k^{-1}\nabla f(\mathbf{x}_k)$. This gives Newton's method:

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \nabla^2 f(\mathbf{x}_k)^{-1}\nabla f(\mathbf{x}_k).$$

In practice, we do not invert the Hessian, but solve the linear system $H_k\mathbf{h} = -\nabla f(\mathbf{x}_k)$ (Lecture 1). For $n = 1$, Newton's method is $x_{k+1} = x_k - \frac{f'(x_k)}{f''(x_k)}$, i.e., Newton's method for finding a zero of f'.

- If f is a quadratic function with a positive definite matrix A, then $m_0 = f$, and Newton's method finds the minimum in a single step, no matter how large the condition number is (Figure 3.8).
- Near a strict local minimum $\mathbf{x}^*$ with a positive definite Hessian, Newton's method converges quadratically: roughly, the error is squared in every step, $\|\mathbf{x}_{k+1} - \mathbf{x}^*\| \le C\|\mathbf{x}_k - \mathbf{x}^*\|^2$, so the number of correct digits doubles in every step (Theorem 3.10.11).

Theorem 3.10.11 (Quadratic convergence of Newton's method)**.** *Let f be twice continuously differentiable on an open set that contains the closed ball $B = \{\mathbf{x} : \|\mathbf{x} - \mathbf{x}^*\| \le r\}$, and let $\nabla f(\mathbf{x}^*) = \mathbf{0}$. Suppose that $\nabla^2 f(\mathbf{x}^*)$ is positive definite, with smallest eigenvalue $\mu > 0$, and that the Hessian is Lipschitz continuous on B, i.e., $\|\nabla^2 f(\mathbf{x}) - \nabla^2 f(\mathbf{y})\|_2 \le L\|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in B$, where $\|M\|_2 = \max_{\mathbf{u}\neq\mathbf{0}} \frac{\|M\mathbf{u}\|}{\|\mathbf{u}\|}$ is the spectral norm of Lecture 2. If $\|\mathbf{x}_0 - \mathbf{x}^*\| \le \rho = \min\left(r, \frac{\mu}{2L}\right)$, then all Newton iterates $\mathbf{x}_k$ are defined and lie in B, and*

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\| \le \frac{L}{\mu}\|\mathbf{x}_k - \mathbf{x}^*\|^2 \le \frac{1}{2}\|\mathbf{x}_k - \mathbf{x}^*\|.$$

Proof. Let $\mathbf{e}_k = \mathbf{x}_k - \mathbf{x}^*$, and suppose that $\|\mathbf{e}_k\| \le \rho$. First, the Hessian $H_k = \nabla^2 f(\mathbf{x}_k)$ is close to $\nabla^2 f(\mathbf{x}^*)$. For every $\mathbf{u}$, the Rayleigh quotient (Lecture 2) and the Cauchy–Schwarz inequality give

$$\mathbf{u}^\top H_k\mathbf{u} = \mathbf{u}^\top\nabla^2 f(\mathbf{x}^*)\mathbf{u} + \mathbf{u}^\top(H_k - \nabla^2 f(\mathbf{x}^*))\mathbf{u} \ge \mu\|\mathbf{u}\|^2 - L\|\mathbf{e}_k\|\|\mathbf{u}\|^2 \ge \frac{\mu}{2}\|\mathbf{u}\|^2.$$

So H_k is positive definite, hence invertible, and $\frac{\mu}{2}\|\mathbf{u}\|^2 \le \mathbf{u}^\top H_k\mathbf{u} \le \|\mathbf{u}\|\|H_k\mathbf{u}\|$ shows that $\|H_k\mathbf{u}\| \ge \frac{\mu}{2}\|\mathbf{u}\|$, i.e., $\|H_k^{-1}\mathbf{w}\| \le \frac{2}{\mu}\|\mathbf{w}\|$ for all $\mathbf{w}$.

Second, the derivative of $t \mapsto \nabla f(\mathbf{x}^* + t\mathbf{e}_k)$ is $\nabla^2 f(\mathbf{x}^* + t\mathbf{e}_k)\mathbf{e}_k$, by the chain rule, so the fundamental theorem of calculus, applied to each entry, gives

$$\nabla f(\mathbf{x}_k) = \nabla f(\mathbf{x}_k) - \nabla f(\mathbf{x}^*) = \int_0^1 \nabla^2 f(\mathbf{x}^* + t\mathbf{e}_k)\mathbf{e}_k\, dt.$$

Hence

$$\mathbf{e}_{k+1} = \mathbf{e}_k - H_k^{-1}\nabla f(\mathbf{x}_k) = H_k^{-1}(H_k\mathbf{e}_k - \nabla f(\mathbf{x}_k)) = H_k^{-1}\int_0^1 (\nabla^2 f(\mathbf{x}_k) - \nabla^2 f(\mathbf{x}^* + t\mathbf{e}_k))\mathbf{e}_k\, dt.$$

The points $\mathbf{x}_k$ and $\mathbf{x}^* + t\mathbf{e}_k$ lie in B and have the distance $(1-t)\|\mathbf{e}_k\|$, so the integrand has norm at most $L(1-t)\|\mathbf{e}_k\|^2$. The norm of the integral of a vector-valued function is at most the integral of its norm (for $\mathbf{w} = \int_0^1 \mathbf{v}(t)\, dt$, the Cauchy–Schwarz inequality gives $\|\mathbf{w}\|^2 = \int_0^1 \mathbf{w}^\top\mathbf{v}(t)\, dt \le \|\mathbf{w}\|\int_0^1\|\mathbf{v}(t)\|\, dt$). Therefore

$$\|\mathbf{e}_{k+1}\| \le \frac{2}{\mu}\int_0^1 L(1-t)\|\mathbf{e}_k\|^2\, dt = \frac{L}{\mu}\|\mathbf{e}_k\|^2 \le \frac{L\rho}{\mu}\|\mathbf{e}_k\| \le \frac{1}{2}\|\mathbf{e}_k\|.$$

In particular, $\|\mathbf{e}_{k+1}\| \le \rho$, and the claim follows by induction on k. □

So once the iterates are close to $\mathbf{x}^*$, the error at least halves in every step, and soon it is squared in every step: if $\frac{L}{\mu}\|\mathbf{e}_k\| \le 10^{-2}$, then $\frac{L}{\mu}\|\mathbf{e}_{k+1}\| \le 10^{-4}$ and $\frac{L}{\mu}\|\mathbf{e}_{k+2}\| \le 10^{-8}$. For a quadratic function, the Hessian is constant, $L = 0$, and the theorem says again that Newton's method converges in one step. Far from $\mathbf{x}^*$, Newton's method may diverge, or converge to a saddle point or a maximum, where the gradient is also $\mathbf{0}$; in practice, one damps the step, $\mathbf{x}_{k+1} = \mathbf{x}_k - \eta_k H_k^{-1}\nabla f(\mathbf{x}_k)$, or modifies the Hessian to make it positive definite.

Example 3.10.12. Let $f(x) = e^x - 2x$. Then $f'(x) = e^x - 2$ and $f''(x) = e^x > 0$, so the minimum is at $x^* = \log 2 = 0.693147\ldots$, and Newton's method is $x_{k+1} = x_k - \frac{e^{x_k}-2}{e^{x_k}} = x_k - 1 + 2e^{-x_k}$. Starting at $x_0 = 0$:

k	0	1	2	3	4	5
x_k	0	1	0.735759	0.694042	0.6931476	0.693147180560
$\lvert x_k - x^*\rvert$	0.69	0.31	$4.3 \cdot 10^{-2}$	$9.0 \cdot 10^{-4}$	$4.0 \cdot 10^{-7}$	$8.0 \cdot 10^{-14}$

Once the iterates are close to x^*, the exponent of the error roughly doubles in every step.

Newton's method has two drawbacks. First, for n parameters, the Hessian has n^2 entries, and solving the linear system costs about n^3 operations, which is impossible for neural networks with millions of parameters. Second, far from a minimum, the Hessian need not be positive definite, and Newton's method can move to a saddle point or a maximum, since it only looks for a critical point. Quasi-Newton methods such as BFGS and L-BFGS build approximations of the Hessian (or of its inverse) from the gradients of previous steps, and are popular for problems with up to a few thousand parameters. Note also that the Newton step $-H_k^{-1}\nabla f(\mathbf{x}_k)$ is the gradient step with respect to the inner product $\langle \mathbf{x}, \mathbf{y}\rangle_{H_k} = \mathbf{x}^\top H_k \mathbf{y}$ defined by the Hessian (Remark 3.3.8): Newton's method measures distances in a way that makes the level sets of the quadratic model round.

Nonlinear least squares. Many fitting problems minimize $f(\boldsymbol{\theta}) = \frac{1}{2}\|\mathbf{r}(\boldsymbol{\theta})\|^2 = \frac{1}{2}\sum_{i=1}^m r_i(\boldsymbol{\theta})^2$, where the residuals $r_i(\boldsymbol{\theta}) = g(\mathbf{x}_i;\boldsymbol{\theta}) - y_i$ measure how far a model g with the parameters $\boldsymbol{\theta} \in \mathbb{R}^n$ is from the data. With the $m \times n$ Jacobian $J = D\mathbf{r}(\boldsymbol{\theta})$, the chain rule gives

$$\nabla f(\boldsymbol{\theta}) = J^\top \mathbf{r}(\boldsymbol{\theta}), \qquad \nabla^2 f(\boldsymbol{\theta}) = J^\top J + \sum_{i=1}^m r_i(\boldsymbol{\theta})\nabla^2 r_i(\boldsymbol{\theta}).$$

The Gauss–Newton method drops the second term, which is small when the residuals are small, and zero for a linear model. Its step solves $J^\top J\mathbf{h} = -J^\top \mathbf{r}$, i.e., $\mathbf{h}$ is the least squares solution of $J\mathbf{h} \approx -\mathbf{r}$ (Lecture 2): Gauss–Newton replaces $\mathbf{r}$ by its linear approximation $\mathbf{r}(\boldsymbol{\theta}+\mathbf{h}) \approx \mathbf{r}(\boldsymbol{\theta}) + J\mathbf{h}$ and minimizes $\|\mathbf{r} + J\mathbf{h}\|^2$. It only needs first derivatives, and if J has linearly independent columns and $\nabla f(\boldsymbol{\theta}) \neq \mathbf{0}$, the step goes downhill: $\nabla f(\boldsymbol{\theta})^\top \mathbf{h} = \mathbf{r}^\top J\mathbf{h} = -\mathbf{h}^\top J^\top J\mathbf{h} < 0$. For a linear model $\mathbf{r}(\boldsymbol{\theta}) = A\boldsymbol{\theta} - \mathbf{y}$, it finds the least squares solution in one step. The Levenberg–Marquardt method solves $(J^\top J + \lambda I)\mathbf{h} = -J^\top \mathbf{r}$ instead: for small λ, this is a Gauss–Newton step, and for large λ, $\mathbf{h} \approx -\frac{1}{\lambda}\nabla f(\boldsymbol{\theta})$ is a short gradient step. The damping λ is decreased after every step that decreases f and increased otherwise; it is the ridge penalty of Lecture 2 for the linearized problem.

Example 3.10.13. Fit the model $y = e^{\theta x}$ to the data $(0,1)$, $(1,2)$, $(2,4)$, which it fits exactly for $\theta^* = \log 2$. Here $r_i = e^{\theta x_i} - y_i$ and $J_{i1} = x_i e^{\theta x_i}$, and the Gauss–Newton step for one parameter is $h = -\frac{\sum_i J_{i1} r_i}{\sum_i J_{i1}^2}$. Starting at $\theta_0 = 1$, the iterates are 0.7695, 0.69856, 0.693175, 0.6931471813, with errors 0.31, 0.076, $5.4 \cdot 10^{-3}$, $2.8 \cdot 10^{-5}$ and $7.8 \cdot 10^{-10}$: the convergence is quadratic, as for Newton's method, because the residuals vanish at θ^*, and so does the dropped term of the Hessian.

Hessian-vector products. Newton-type methods for large n never form the Hessian. The conjugate gradient method (mentioned after the Cayley–Hamilton theorem in Lecture 1) solves $H\mathbf{h} = -\nabla f$ using only products $H\mathbf{v}$, and automatic differentiation computes such products at the cost of a few gradients, without computing H. The Hessian is the Jacobian of the map $\mathbf{x} \mapsto \nabla f(\mathbf{x})$, so

$$\nabla^2 f(\mathbf{x})\mathbf{v} = \frac{d}{dt}\nabla f(\mathbf{x} + t\mathbf{v})\Big|_{t=0}$$

is a Jacobian-vector product of the gradient. It is computed by applying forward mode to the reverse-mode program that computes ∇f ("forward over reverse"), or, since the Hessian is symmetric, as the gradient of the scalar function $\mathbf{x} \mapsto \nabla f(\mathbf{x})^\top \mathbf{v}$. Hessian-vector products are used in Hessian-free optimization, to estimate the largest eigenvalue of the Hessian of a neural network loss by power iteration (Lecture 1), which limits the stable step size of gradient descent, and, with Hutchinson's estimator (Exercise 32), to estimate the trace of the Hessian.

3.11 Convexity

Gradient descent and Newton's method only find critical points, and a critical point need not be a global minimum. For an important class of functions, the convex functions, this problem disappears.

Definition 3.11.1 (Convex sets and functions)**.** A set $C \subseteq \mathbb{R}^n$ is convex if for all $\mathbf{x}, \mathbf{y} \in C$ and all $\theta \in [0,1]$, also $\theta\mathbf{x} + (1-\theta)\mathbf{y} \in C$: with any two points, C contains the line segment between them. A function $f : C \to \mathbb{R}$ on a convex set C is convex if

$$f(\theta\mathbf{x} + (1-\theta)\mathbf{y}) \le \theta f(\mathbf{x}) + (1-\theta) f(\mathbf{y}) \qquad \text{for all } \mathbf{x}, \mathbf{y} \in C,\ \theta \in [0,1],$$

and strictly convex if the inequality is strict whenever $\mathbf{x} \neq \mathbf{y}$ and $0 < \theta < 1$. A function f is concave if $-f$ is convex.

Geometrically, f is convex if the line segment between any two points of its graph lies above the graph (Figure 3.9). Examples of convex sets are $\mathbb{R}^n$, subspaces, affine subspaces, half-spaces $\{\mathbf{x} : \mathbf{w}^\top \mathbf{x} \le c\}$, balls $\{\mathbf{x} : \|\mathbf{x} - \mathbf{a}\| \le r\}$ for any norm, and intersections of convex sets. A circle is not convex.

Example 3.11.2.

1. Affine functions $f(\mathbf{x}) = \mathbf{a}^\top \mathbf{x} + b$ are convex (and concave), with equality in the definition.
2. Every norm is convex: by the triangle inequality and homogeneity, $\|\theta\mathbf{x} + (1-\theta)\mathbf{y}\| \le \theta\|\mathbf{x}\| + (1-\theta)\|\mathbf{y}\|$ for $\theta \in [0,1]$.
3. The functions x^2, $|x|$, e^x on $\mathbb{R}$, and $-\log x$ on $(0,\infty)$ are convex. The function x^3 is not convex on $\mathbb{R}$.

Theorem 3.11.3 (First-order characterization)**.** *Let $C \subseteq \mathbb{R}^n$ be open and convex, and let $f : C \to \mathbb{R}$ be differentiable. Then f is convex if and only if*

$$f(\mathbf{y}) \ge f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) \qquad \textit{for all } \mathbf{x}, \mathbf{y} \in C.$$

Proof. Suppose that f is convex, and let $\mathbf{x}, \mathbf{y} \in C$. For $0 < t \le 1$, convexity with $\theta = 1 - t$ gives $f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) \le f(\mathbf{x}) + t(f(\mathbf{y}) - f(\mathbf{x}))$, so

$$\frac{f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) - f(\mathbf{x})}{t} \le f(\mathbf{y}) - f(\mathbf{x}).$$




Figure 3.9: Left: a convex function lies below every chord, i.e., $f(\theta x+(1-\theta)y) \leq \theta f(x)+(1-\theta)f(y)$. Right: a differentiable convex function lies above every tangent line (Theorem 3.11.3).

As $t \to 0$, the left side tends to the directional derivative $\nabla f(\mathbf{x})^\top(\mathbf{y}-\mathbf{x})$ (Proposition 3.3.5).

Conversely, suppose that the inequality holds, let $\theta \in [0,1]$, and let $\mathbf{z} = \theta\mathbf{x} + (1-\theta)\mathbf{y}$. Then $f(\mathbf{x}) \geq f(\mathbf{z}) + \nabla f(\mathbf{z})^\top(\mathbf{x}-\mathbf{z})$ and $f(\mathbf{y}) \geq f(\mathbf{z}) + \nabla f(\mathbf{z})^\top(\mathbf{y}-\mathbf{z})$. Multiplying the first inequality by θ and the second by $1-\theta$ and adding, we get

$$\theta f(\mathbf{x}) + (1-\theta)f(\mathbf{y}) \geq f(\mathbf{z}) + \nabla f(\mathbf{z})^\top(\theta\mathbf{x} + (1-\theta)\mathbf{y} - \mathbf{z}) = f(\mathbf{z}). \qquad \square$$

So the linear approximation of a convex function is a global lower bound. This has a remarkable consequence.

Corollary 3.11.4. *Let f be convex and differentiable on an open convex set C. If $\nabla f(\mathbf{x}^*) = \mathbf{0}$, then $\mathbf{x}^*$ is a global minimum of f.*

Proof. By Theorem 3.11.3, $f(\mathbf{y}) \geq f(\mathbf{x}^*) + \mathbf{0}^\top(\mathbf{y}-\mathbf{x}^*) = f(\mathbf{x}^*)$ for all $\mathbf{y} \in C$. $\square$

Proposition 3.11.5. *Let $f : C \to \mathbb{R}$ be convex. Then every local minimum of f is a global minimum, and the set of global minima is convex. If f is strictly convex, it has at most one global minimum.*

Proof. Let $\mathbf{x}^*$ be a local minimum, and suppose that $f(\mathbf{y}) < f(\mathbf{x}^*)$ for some $\mathbf{y} \in C$. For $0 < t \leq 1$, the point $\mathbf{x}_t = (1-t)\mathbf{x}^* + t\mathbf{y}$ satisfies $f(\mathbf{x}_t) \leq (1-t)f(\mathbf{x}^*) + tf(\mathbf{y}) < f(\mathbf{x}^*)$, and $\mathbf{x}_t$ is arbitrarily close to $\mathbf{x}^*$ for small t. This contradicts the local minimality of $\mathbf{x}^*$. If $\mathbf{x}$ and $\mathbf{y}$ are global minima with $f(\mathbf{x}) = f(\mathbf{y}) = m$, then $f(\theta\mathbf{x}+(1-\theta)\mathbf{y}) \leq m$, so all points of the segment are global minima. For strictly convex f and $\mathbf{x} \neq \mathbf{y}$, we would get $f(\frac{1}{2}\mathbf{x} + \frac{1}{2}\mathbf{y}) < m$, which is impossible. $\square$

So for convex functions, there are no bad local minima and no saddle points: every critical point is a global minimum, and gradient descent with a suitable step size converges to the minimum value. (Theorem 3.11.12 below shows that for a convex L-smooth function with a minimum $\mathbf{x}^*$, gradient descent with $\eta = \frac{1}{L}$ satisfies $f(\mathbf{x}_k) - f(\mathbf{x}^*) \leq \frac{L\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{2k}$.) For twice differentiable functions, convexity can be checked with the Hessian.

Theorem 3.11.6 (Second-order characterization)**.** *Let $C \subseteq \mathbb{R}^n$ be open and convex, and let $f : C \to \mathbb{R}$ be twice continuously differentiable. Then f is convex if and only if $\nabla^2 f(\mathbf{x})$ is positive semidefinite for all $\mathbf{x} \in C$. If $\nabla^2 f(\mathbf{x})$ is positive definite for all $\mathbf{x} \in C$, then f is strictly convex.*

Proof. If all Hessians are positive semidefinite, then by Theorem 3.8.5, for all $\mathbf{x}, \mathbf{y} \in C$,

$$f(\mathbf{y}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top(\mathbf{y}-\mathbf{x}) + \tfrac{1}{2}(\mathbf{y}-\mathbf{x})^\top \nabla^2 f(\mathbf{z})(\mathbf{y}-\mathbf{x}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top(\mathbf{y}-\mathbf{x}),$$

where $\mathbf{z}$ is a point on the segment between $\mathbf{x}$ and $\mathbf{y}$, so f is convex by Theorem 3.11.3. (If all Hessians are positive definite, the inequality is strict for $\mathbf{y} \neq \mathbf{x}$, and the proof of Theorem 3.11.3

then gives strict convexity.) Conversely, suppose that $\mathbf{v}^\top \nabla^2 f(\mathbf{x})\mathbf{v} < 0$ for some $\mathbf{x} \in C$ and $\mathbf{v}$. By the second form of Theorem 3.8.5, for small $t \neq 0$,

$$f(\mathbf{x}+t\mathbf{v}) = f(\mathbf{x}) + t\nabla f(\mathbf{x})^\top \mathbf{v} + \tfrac{1}{2}t^2 \mathbf{v}^\top \nabla^2 f(\mathbf{x})\mathbf{v} + o(t^2) < f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (t\mathbf{v}),$$

which contradicts the inequality of Theorem 3.11.3 for $\mathbf{y} = \mathbf{x} + t\mathbf{v}$. □

Example 3.11.7 (Convex loss functions)**.**

1. The least squares loss $\|\mathbf{y} - \Phi\boldsymbol{\theta}\|^2$ has the Hessian $2\Phi^\top\Phi$, which is positive semidefinite, since $\mathbf{v}^\top\Phi^\top\Phi\mathbf{v} = \|\Phi\mathbf{v}\|^2 \geq 0$. So it is convex, and strictly convex if Φ has linearly independent columns. Adding a ridge penalty $\lambda\|\boldsymbol{\theta}\|^2$ with $\lambda > 0$ adds $2\lambda I$ to the Hessian, which makes it positive definite, so ridge regression always has a unique minimum.

2. The logistic regression loss has the Hessian $\frac{1}{N}\sum_i p_i(1-p_i)\mathbf{x}_i\mathbf{x}_i^\top$ (Example 3.8.4). It is positive semidefinite, since $\mathbf{v}^\top\mathbf{x}_i\mathbf{x}_i^\top\mathbf{v} = (\mathbf{x}_i^\top\mathbf{v})^2 \geq 0$ and $p_i(1-p_i) > 0$. So the loss is convex, and every critical point is a global minimum.

3. The log-sum-exp function has the Hessian $\operatorname{diag}(\mathbf{p}) - \mathbf{p}\mathbf{p}^\top$, where $\mathbf{p}$ has positive entries with sum 1. For every $\mathbf{v}$, by the Cauchy–Schwarz inequality applied to the vectors with entries $\sqrt{p_i}$ and $\sqrt{p_i}v_i$,
$$\Big(\sum_i p_i v_i\Big)^2 \leq \Big(\sum_i p_i\Big)\Big(\sum_i p_i v_i^2\Big) = \sum_i p_i v_i^2,$$
so $\mathbf{v}^\top\big(\operatorname{diag}(\mathbf{p}) - \mathbf{p}\mathbf{p}^\top\big)\mathbf{v} = \sum_i p_i v_i^2 - \big(\sum_i p_i v_i\big)^2 \geq 0$. So log-sum-exp is convex, and hence so is the softmax cross-entropy loss of Example 3.6.6.

Proposition 3.11.8 (Operations that preserve convexity)**.**

1. *If f and g are convex and $\alpha, \beta \geq 0$, then $\alpha f + \beta g$ is convex.*

2. *If f is convex, then $\mathbf{x} \mapsto f(A\mathbf{x} + \mathbf{b})$ is convex.*

3. *If f and g are convex, then $\mathbf{x} \mapsto \max(f(\mathbf{x}), g(\mathbf{x}))$ is convex.*

Proof. (1) Add the inequalities for f and g, multiplied by α and β. (2) The map $\mathbf{x} \mapsto A\mathbf{x}+\mathbf{b}$ maps $\theta\mathbf{x} + (1-\theta)\mathbf{y}$ to $\theta(A\mathbf{x}+\mathbf{b}) + (1-\theta)(A\mathbf{y}+\mathbf{b})$; now use the convexity of f. (3) $f(\theta\mathbf{x} + (1-\theta)\mathbf{y}) \leq \theta f(\mathbf{x}) + (1-\theta)f(\mathbf{y}) \leq \theta \max(f,g)(\mathbf{x}) + (1-\theta)\max(f,g)(\mathbf{y})$, and similarly for g. □

For example, the hinge loss $\max(0, 1 - y\,\mathbf{w}^\top\mathbf{x})$ of support vector machines is convex in $\mathbf{w}$, as the maximum of two affine functions, and so are the ℓ_1 and ℓ_2 regularization terms $\lambda\|\mathbf{w}\|_1$ and $\lambda\|\mathbf{w}\|^2$.

Remark 3.11.9 (Neural networks are not convex). The loss of a neural network with a hidden layer is not convex as a function of the weights. One reason is symmetry: if we swap two hidden units, i.e., swap two rows of W_1, the corresponding entries of $\mathbf{b}_1$, and the corresponding columns of W_2, the network computes the same function, so the loss does not change. If $\boldsymbol{\theta}^*$ is a global minimum, then so is the swapped parameter vector $\boldsymbol{\theta}'$. If the loss were convex, then by Proposition 3.11.5 the midpoint $\frac{1}{2}(\boldsymbol{\theta}^* + \boldsymbol{\theta}')$ would be a global minimum too. But in the midpoint, the two hidden units have identical weights, so the network effectively has one hidden unit less, and its loss is usually larger. Nevertheless, gradient-based methods work remarkably well for neural networks in practice, which is an active area of research.

3.11.1 Strong convexity and convergence rates

Convexity guarantees that gradient descent heads for a global minimum. How fast it gets there depends on how curved the function is, from below and from above.

Definition 3.11.10 (Strong convexity)**.** Let $\mu > 0$. A differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ is μ-strongly convex if

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top(\mathbf{y} - \mathbf{x}) + \frac{\mu}{2}\|\mathbf{y} - \mathbf{x}\|^2 \qquad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

So a strongly convex function lies above a quadratic function at every point, just as an L-smooth function lies below one (Lemma 3.10.4). Since $\|\mathbf{y}\|^2 - \|\mathbf{x}\|^2 - 2\mathbf{x}^\top(\mathbf{y} - \mathbf{x}) = \|\mathbf{y} - \mathbf{x}\|^2$, the inequality says exactly that $g(\mathbf{x}) = f(\mathbf{x}) - \frac{\mu}{2}\|\mathbf{x}\|^2$ satisfies the inequality of Theorem 3.11.3. Hence f is μ-strongly convex if and only if $f - \frac{\mu}{2}\|\cdot\|^2$ is convex, and for a twice continuously differentiable f, by Theorem 3.11.6, if and only if $\nabla^2 f(\mathbf{x}) - \mu I$ is positive semidefinite, i.e., all eigenvalues of every Hessian are at least μ. For example, $\frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$ with a symmetric positive definite A is $\lambda_{\min}$-strongly convex and $\lambda_{\max}$-smooth, and the ridge regression loss $\|X\mathbf{w} - \mathbf{y}\|^2 + \lambda\|\mathbf{w}\|^2$ is 2λ-strongly convex, whatever the data. A strongly convex function has a unique global minimum: $f(\mathbf{y}) \geq f(\mathbf{0}) + \nabla f(\mathbf{0})^\top\mathbf{y} + \frac{\mu}{2}\|\mathbf{y}\|^2 \to \infty$ as $\|\mathbf{y}\| \to \infty$, so a minimum exists (by Proposition A.3.6), and it is unique, since f is strictly convex.

Lemma 3.11.11. *Let f be μ-strongly convex with the minimum value $f^* = f(\mathbf{x}^*)$. Then for all $\mathbf{x}$,*

$$\nabla f(\mathbf{x})^\top(\mathbf{x} - \mathbf{x}^*) \geq \mu\|\mathbf{x} - \mathbf{x}^*\|^2 \qquad \text{and} \qquad f(\mathbf{x}) - f^* \leq \frac{1}{2\mu}\|\nabla f(\mathbf{x})\|^2.$$

Proof. Adding the inequalities of Definition 3.11.10 for the pairs $(\mathbf{x}, \mathbf{x}^*)$ and $(\mathbf{x}^*, \mathbf{x})$, and using $\nabla f(\mathbf{x}^*) = \mathbf{0}$, gives $0 \geq \nabla f(\mathbf{x})^\top(\mathbf{x}^* - \mathbf{x}) + \mu\|\mathbf{x} - \mathbf{x}^*\|^2$, which is the first inequality. For the second, the right side of the definition is a quadratic function of $\mathbf{y}$, which is smallest for $\mathbf{y} - \mathbf{x} = -\frac{1}{\mu}\nabla f(\mathbf{x})$, so $f(\mathbf{y}) \geq f(\mathbf{x}) - \frac{1}{2\mu}\|\nabla f(\mathbf{x})\|^2$ for all $\mathbf{y}$, in particular for $\mathbf{y} = \mathbf{x}^*$. □

The second inequality says that a small gradient guarantees a small error in the function value, which is false for general convex functions (think of $f(x) = e^x$). It is called the Polyak–Łojasiewicz inequality.

Theorem 3.11.12 (Convergence rates of gradient descent)**.** *Let f be convex and L-smooth, with a minimum $\mathbf{x}^*$ and $f^* = f(\mathbf{x}^*)$, and let $\mathbf{x}_k$ be the iterates of gradient descent with step size $\eta = \frac{1}{L}$.*

1. *For all $k \geq 1$, $f(\mathbf{x}_k) - f^* \leq \dfrac{L\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{2k}$.*
2. *If f is also μ-strongly convex, then $f(\mathbf{x}_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(\mathbf{x}_0) - f^*)$.*

Proof. Let $\mathbf{g}_k = \nabla f(\mathbf{x}_k)$. By Theorem 3.10.5, $f(\mathbf{x}_{k+1}) \leq f(\mathbf{x}_k) - \frac{1}{2L}\|\mathbf{g}_k\|^2$.

(2) By Lemma 3.11.11, $\|\mathbf{g}_k\|^2 \geq 2\mu(f(\mathbf{x}_k) - f^*)$, so $f(\mathbf{x}_{k+1}) - f^* \leq \left(1 - \frac{\mu}{L}\right)(f(\mathbf{x}_k) - f^*)$; repeat this k times.

(1) By Theorem 3.11.3, $f^* \geq f(\mathbf{x}_k) + \mathbf{g}_k^\top(\mathbf{x}^* - \mathbf{x}_k)$. Together with the descent inequality and $\mathbf{x}_{k+1} = \mathbf{x}_k - \frac{1}{L}\mathbf{g}_k$,

$$f(\mathbf{x}_{k+1}) - f^* \leq \mathbf{g}_k^\top(\mathbf{x}_k - \mathbf{x}^*) - \frac{1}{2L}\|\mathbf{g}_k\|^2 = \frac{L}{2}\Big(\|\mathbf{x}_k - \mathbf{x}^*\|^2 - \|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2\Big),$$

as one checks by expanding $\|\mathbf{x}_k - \mathbf{x}^* - \frac{1}{L}\mathbf{g}_k\|^2$. Adding these inequalities for $k = 0, \ldots, K-1$, the right side telescopes, and $\sum_{k=1}^{K}\left(f(\mathbf{x}_k) - f^*\right) \leq \frac{L}{2}\|\mathbf{x}_0 - \mathbf{x}^*\|^2$. Since the values $f(\mathbf{x}_k)$ decrease, the left side is at least $K\big(f(\mathbf{x}_K) - f^*\big)$. □

With $\kappa = \frac{L}{\mu}$, the condition number, the second rate is $(1 - \frac{1}{\kappa})^k \leq e^{-k/\kappa}$: the error decreases geometrically ("linear convergence"), and about $\kappa \log \frac{1}{\varepsilon}$ steps reduce it by the factor ε, as for quadratic functions (Theorem 3.10.7). Without strong convexity, the guarantee is only $O(\frac{1}{k})$: to halve the error, we have to double the number of steps. Adding a small ridge penalty $\lambda\|\boldsymbol{\theta}\|^2$ to a convex loss makes it 2λ-strongly convex, which gives the linear rate of part 2 instead of the rate of part 1, at the price of a slightly different minimum.

3.11.2 Nonsmooth convex functions and the proximal gradient method

The ℓ_1 norm, the hinge loss and ReLU are convex, but not differentiable everywhere. For convex functions, the gradient has a substitute.

Definition 3.11.13 (Subgradient)**.** Let $f : \mathbb{R}^n \to \mathbb{R}$ be convex. A vector $\mathbf{g}$ is a subgradient of f at $\mathbf{x}$ if

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \mathbf{g}^\top(\mathbf{y} - \mathbf{x}) \qquad \text{for all } \mathbf{y} \in \mathbb{R}^n.$$

The set $\partial f(\mathbf{x})$ of all subgradients at $\mathbf{x}$ is the subdifferential of f at $\mathbf{x}$.

A subgradient defines an affine function that lies below f and touches it at $\mathbf{x}$. By the definition, $\mathbf{x}^*$ is a global minimum of f if and only if $\mathbf{0} \in \partial f(\mathbf{x}^*)$.

Proposition 3.11.14. *1. If f is convex and differentiable, then $\partial f(\mathbf{x}) = \{\nabla f(\mathbf{x})\}$ for every $\mathbf{x}$.*

2. For $f(x) = |x|$ on $\mathbb{R}$: $\partial f(x) = \{1\}$ for $x > 0$, $\{-1\}$ for $x < 0$, and $\partial f(0) = [-1, 1]$.

3. If $f(\mathbf{x}) = f_1(x_1) + \cdots + f_n(x_n)$ with convex functions f_i of one variable, then $\mathbf{g} \in \partial f(\mathbf{x})$ if and only if $g_i \in \partial f_i(x_i)$ for all i. In particular, $\partial\|\cdot\|_1(\mathbf{x})$ consists of the vectors $\mathbf{g}$ with $g_i = \operatorname{sign}(x_i)$ if $x_i \neq 0$ and $g_i \in [-1, 1]$ if $x_i = 0$.

Proof. (1) By Theorem 3.11.3, $\nabla f(\mathbf{x}) \in \partial f(\mathbf{x})$. If $\mathbf{g} \in \partial f(\mathbf{x})$, then for every $\mathbf{v}$ and $t > 0$, $\frac{f(\mathbf{x}+t\mathbf{v})-f(\mathbf{x})}{t} \geq \mathbf{g}^\top\mathbf{v}$, and as $t \to 0$, $\nabla f(\mathbf{x})^\top\mathbf{v} \geq \mathbf{g}^\top\mathbf{v}$. With $-\mathbf{v}$ instead of $\mathbf{v}$, we get equality, so $\mathbf{g} = \nabla f(\mathbf{x})$.

(2) At 0, $|y| \geq gy$ for all y if and only if $|g| \leq 1$. For $x > 0$, $g = 1$ is a subgradient, since $|y| \geq y = x+(y-x)$. Conversely, if g is a subgradient at $x > 0$, then for $|t| < x$, $x+t = |x+t| \geq x+gt$, i.e., $t \geq gt$ for positive and negative t, so $g = 1$. The case $x < 0$ is symmetric.

(3) If $g_i \in \partial f_i(x_i)$ for all i, add the inequalities $f_i(y_i) \geq f_i(x_i) + g_i(y_i - x_i)$. Conversely, if $\mathbf{g} \in \partial f(\mathbf{x})$, use the definition with a vector $\mathbf{y}$ that differs from $\mathbf{x}$ only in its i-th entry. The last statement follows with (2). □

Machine learning losses are often a sum of a smooth part and a simple nonsmooth part, such as the lasso

$$F(\mathbf{w}) = \frac{1}{2}\|X\mathbf{w} - \mathbf{y}\|^2 + \lambda\|\mathbf{w}\|_1,$$

whose ℓ_1 penalty produces sparse weights, i.e., many weights that are exactly 0. For such sums, the optimality condition $\mathbf{0} \in \partial F(\mathbf{x})$ splits into the two parts.

Theorem 3.11.15. *Let $f : \mathbb{R}^n \to \mathbb{R}$ be convex and differentiable, and let $h : \mathbb{R}^n \to \mathbb{R}$ be convex. Then $\mathbf{x}^*$ minimizes $f + h$ if and only if $-\nabla f(\mathbf{x}^*) \in \partial h(\mathbf{x}^*)$.*

Proof. If $-\nabla f(\mathbf{x}^*) \in \partial h(\mathbf{x}^*)$, add the inequalities $f(\mathbf{y}) \geq f(\mathbf{x}^*)+\nabla f(\mathbf{x}^*)^\top(\mathbf{y}-\mathbf{x}^*)$ (Theorem 3.11.3) and $h(\mathbf{y}) \geq h(\mathbf{x}^*) - \nabla f(\mathbf{x}^*)^\top(\mathbf{y} - \mathbf{x}^*)$. Conversely, let $\mathbf{x}^*$ be a minimum, let $\mathbf{y} \in \mathbb{R}^n$, and let $\mathbf{x}_t = \mathbf{x}^* + t(\mathbf{y} - \mathbf{x}^*)$ for $0 < t \leq 1$. By convexity, $h(\mathbf{x}_t) \leq h(\mathbf{x}^*) + t\big(h(\mathbf{y}) - h(\mathbf{x}^*)\big)$, so

$$0 \leq \frac{(f + h)(\mathbf{x}_t) - (f + h)(\mathbf{x}^*)}{t} \leq \frac{f(\mathbf{x}_t) - f(\mathbf{x}^*)}{t} + h(\mathbf{y}) - h(\mathbf{x}^*).$$

As $t \to 0$, we get $0 \leq \nabla f(\mathbf{x}^*)^\top(\mathbf{y} - \mathbf{x}^*) + h(\mathbf{y}) - h(\mathbf{x}^*)$, i.e., $-\nabla f(\mathbf{x}^*) \in \partial h(\mathbf{x}^*)$. □

Definition 3.11.16 (Proximal operator)**.** Let h be convex and $\eta > 0$. The proximal operator of ηh maps $\mathbf{v}$ to the minimum

$$\operatorname{prox}_{\eta h}(\mathbf{v}) = \arg\min_{\mathbf{u}} \Big(h(\mathbf{u}) + \frac{1}{2\eta}\|\mathbf{u} - \mathbf{v}\|^2\Big),$$

whenever it exists; it is unique, since the function in brackets is strictly convex.

By Theorem 3.11.15 with $f(\mathbf{u}) = \frac{1}{2\eta}\|\mathbf{u}-\mathbf{v}\|^2$, whose gradient is $\frac{1}{\eta}(\mathbf{u}-\mathbf{v})$, a point $\mathbf{p}$ is $\operatorname{prox}_{\eta h}(\mathbf{v})$ if and only if $\frac{1}{\eta}(\mathbf{v}-\mathbf{p}) \in \partial h(\mathbf{p})$. For $h = 0$, the proximal operator is the identity; it moves $\mathbf{v}$ towards smaller values of h, but not too far from $\mathbf{v}$.

Proposition 3.11.17 (Soft thresholding)**.** *The proximal operator of $\eta\lambda\|\cdot\|_1$ acts on each entry by soft thresholding:*

$$\operatorname{prox}_{\eta\lambda\|\cdot\|_1}(\mathbf{v})_i = S_{\eta\lambda}(v_i), \qquad S_\tau(v) = \begin{cases} v-\tau & \text{if } v > \tau, \\ 0 & \text{if } |v| \le \tau, \\ v+\tau & \text{if } v < -\tau. \end{cases}$$

Proof. Let $p_i = S_{\eta\lambda}(v_i)$. By the criterion above and Proposition 3.11.14, we must check that $\frac{1}{\eta}(v_i - p_i) \in \lambda\,\partial|\cdot|(p_i)$ for every i. If $v_i > \eta\lambda$, then $p_i > 0$ and $\frac{1}{\eta}(v_i - p_i) = \lambda = \lambda\operatorname{sign}(p_i)$; the case $v_i < -\eta\lambda$ is symmetric. If $|v_i| \le \eta\lambda$, then $p_i = 0$ and $\frac{1}{\eta}v_i \in [-\lambda, \lambda]$. □

The proximal gradient method takes a gradient step for the smooth part and a proximal step for the nonsmooth part:

$$\mathbf{x}_{k+1} = \operatorname{prox}_{\eta h}\left(\mathbf{x}_k - \eta\nabla f(\mathbf{x}_k)\right).$$

For the lasso, this is the iterative soft-thresholding algorithm (ISTA), $\mathbf{w}_{k+1} = S_{\eta\lambda}(\mathbf{w}_k - \eta X^\top(X\mathbf{w}_k - \mathbf{y}))$, applied entrywise. For $h = 0$, it is gradient descent.

Theorem 3.11.18 (Proximal gradient method)**.** *Let f be convex and L-smooth, let h be convex, let $F = f + h$, let $0 < \eta \le \frac{1}{L}$, and suppose that the proximal steps exist.*

1. *$\mathbf{x}^*$ minimizes F if and only if $\mathbf{x}^* = \operatorname{prox}_{\eta h}\left(\mathbf{x}^* - \eta\nabla f(\mathbf{x}^*)\right)$.*
2. *The method decreases F: $F(\mathbf{x}_{k+1}) \le F(\mathbf{x}_k) - \frac{1}{2\eta}\|\mathbf{x}_{k+1} - \mathbf{x}_k\|^2$.*

Proof. (1) By the criterion for the proximal operator, $\mathbf{x}^* = \operatorname{prox}_{\eta h}(\mathbf{x}^* - \eta\nabla f(\mathbf{x}^*))$ if and only if $\frac{1}{\eta}(\mathbf{x}^* - \eta\nabla f(\mathbf{x}^*) - \mathbf{x}^*) = -\nabla f(\mathbf{x}^*) \in \partial h(\mathbf{x}^*)$, which is the condition of Theorem 3.11.15.

(2) Let $\mathbf{p} = \mathbf{x}_{k+1}$ and $\mathbf{d} = \mathbf{p} - \mathbf{x}_k$. By the criterion, $\mathbf{g} = \frac{1}{\eta}(\mathbf{x}_k - \eta\nabla f(\mathbf{x}_k) - \mathbf{p}) = -\frac{1}{\eta}\mathbf{d} - \nabla f(\mathbf{x}_k)$ is a subgradient of h at $\mathbf{p}$, so $h(\mathbf{x}_k) \ge h(\mathbf{p}) - \mathbf{g}^\top\mathbf{d}$. By the descent lemma, $f(\mathbf{p}) \le f(\mathbf{x}_k) + \nabla f(\mathbf{x}_k)^\top\mathbf{d} + \frac{L}{2}\|\mathbf{d}\|^2$. Adding,

$$F(\mathbf{p}) \le F(\mathbf{x}_k) + \left(\mathbf{g} + \nabla f(\mathbf{x}_k)\right)^\top\mathbf{d} + \frac{L}{2}\|\mathbf{d}\|^2 = F(\mathbf{x}_k) - \Big(\frac{1}{\eta} - \frac{L}{2}\Big)\|\mathbf{d}\|^2,$$

and $\frac{1}{\eta} - \frac{L}{2} \ge \frac{1}{2\eta}$ since $\eta L \le 1$. □

Example 3.11.19 (Why the lasso gives sparse weights)**.** Suppose that the columns of X are orthonormal, $X^\top X = I$. Then $\frac{1}{2}\|X\mathbf{w} - \mathbf{y}\|^2 = \frac{1}{2}\|\mathbf{w} - X^\top\mathbf{y}\|^2 + c$, where $c = \frac{1}{2}(\|\mathbf{y}\|^2 - \|X^\top\mathbf{y}\|^2)$ does not depend on $\mathbf{w}$ (expand both sides). So the lasso solution is $\mathbf{w}^* = \operatorname{prox}_{\lambda\|\cdot\|_1}(X^\top\mathbf{y})$, i.e., $w_i^* = S_\lambda((X^\top\mathbf{y})_i)$: the least squares weights $(X^\top\mathbf{y})_i$ are shrunk towards 0 by λ, and those with $|(X^\top\mathbf{y})_i| \le \lambda$ become exactly 0. For example, least squares weights $(3, -0.5, 1.2, 0.1)$ with $\lambda = 1$ give the lasso weights $(2, 0, 0.2, 0)$. Ridge regression instead gives $\frac{1}{1+2\lambda}(X^\top\mathbf{y})$ for the penalty $\lambda\|\mathbf{w}\|^2$, which shrinks all weights by the same factor, but sets none of them to 0.

Other proximal operators are just as simple. For the indicator function of a closed convex set C (which is 0 on C and $+\infty$ outside, so we have to allow the value $+\infty$), the proximal operator is the orthogonal projection onto C, and the proximal gradient method becomes projected gradient descent (Example 3.12.7). Automatic differentiation libraries use subgradients as well: for ReLU at 0, they return a fixed element of $\partial\,\mathrm{ReLU}(0) = [0, 1]$, usually 0.

3.12 Constrained Optimization

Many optimization problems in machine learning have constraints. In principal component analysis, we maximize the variance over unit vectors; a vector of probabilities must have nonnegative entries that sum to 1; and support vector machines maximize a margin subject to classifying all training points correctly. We first consider equality constraints:

$$\text{minimize } f(\mathbf{x}) \quad \text{subject to} \quad g_1(\mathbf{x}) = 0,\ \ldots,\ g_m(\mathbf{x}) = 0.$$

3.12.1 Lagrange multipliers

Consider first one constraint $g(\mathbf{x}) = 0$ in $\mathbb{R}^2$, which is usually a curve. If we walk along the curve, then at a local minimum of f on the curve, the derivative of f in the direction of the curve must be zero, i.e., $\nabla f(\mathbf{x}^*)$ is orthogonal to the curve. By Corollary 3.5.4, $\nabla g(\mathbf{x}^*)$ is also orthogonal to the curve, which is a level set of g. In the plane, this means that $\nabla f(\mathbf{x}^*)$ and $\nabla g(\mathbf{x}^*)$ are parallel: the level curve of f through $\mathbf{x}^*$ touches the constraint curve (Figure 3.10). At a point where ∇f is not orthogonal to the curve, we can decrease f by moving along the curve.

Theorem 3.12.1 (Lagrange multipliers)**.** *Let $f, g_1, \ldots, g_m : D \to \mathbb{R}$ be continuously differentiable on an open set $D \subseteq \mathbb{R}^n$, and let $\mathbf{x}^*$ be a local minimum of f on the set $S = \{\mathbf{x} \in D : g_1(\mathbf{x}) = \cdots = g_m(\mathbf{x}) = 0\}$. If the gradients $\nabla g_1(\mathbf{x}^*), \ldots, \nabla g_m(\mathbf{x}^*)$ are linearly independent, then there are unique numbers $\lambda_1, \ldots, \lambda_m \in \mathbb{R}$, called Lagrange multipliers, such that*

$$\nabla f(\mathbf{x}^*) + \lambda_1 \nabla g_1(\mathbf{x}^*) + \cdots + \lambda_m \nabla g_m(\mathbf{x}^*) = \mathbf{0}.$$

Sketch of proof. Let $U = \operatorname{span}\{\nabla g_1(\mathbf{x}^*), \ldots, \nabla g_m(\mathbf{x}^*)\}$. If $\mathbf{x}(t)$ is a differentiable curve in S with $\mathbf{x}(0) = \mathbf{x}^*$, then $g_i(\mathbf{x}(t)) = 0$ for all t, so $\nabla g_i(\mathbf{x}^*)^\top \mathbf{x}'(0) = 0$ by the chain rule, i.e., the tangent vector $\mathbf{x}'(0)$ lies in $U^\perp$. Conversely, the implicit function theorem from analysis, which we do not prove, shows that, since the $\nabla g_i(\mathbf{x}^*)$ are linearly independent, every vector $\mathbf{v} \in U^\perp$ is the tangent vector $\mathbf{x}'(0)$ of such a curve. Since $t \mapsto f(\mathbf{x}(t))$ has a local minimum at $t = 0$, its derivative $\nabla f(\mathbf{x}^*)^\top \mathbf{x}'(0)$ is 0. Hence $\nabla f(\mathbf{x}^*)$ is orthogonal to every vector in $U^\perp$, i.e., $\nabla f(\mathbf{x}^*) \in (U^\perp)^\perp = U$ (Lecture 2). So $\nabla f(\mathbf{x}^*)$ is a linear combination of the $\nabla g_i(\mathbf{x}^*)$, and the coefficients are unique, since the $\nabla g_i(\mathbf{x}^*)$ are linearly independent. □

The same condition holds at local maxima (apply the theorem to $-f$; the multipliers can have any sign). It is convenient to write it with the Lagrangian

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = f(\mathbf{x}) + \lambda_1 g_1(\mathbf{x}) + \cdots + \lambda_m g_m(\mathbf{x}).$$

The condition of the theorem says that $\nabla_{\mathbf{x}} \mathcal{L}(\mathbf{x}^*, \boldsymbol{\lambda}) = \mathbf{0}$, and the constraints say that $\nabla_{\boldsymbol{\lambda}} \mathcal{L}(\mathbf{x}^*, \boldsymbol{\lambda}) = (g_1(\mathbf{x}^*), \ldots, g_m(\mathbf{x}^*))^\top = \mathbf{0}$. So we look for critical points of $\mathcal{L}$, which gives $n + m$ equations for the $n + m$ unknowns $\mathbf{x}$ and $\boldsymbol{\lambda}$. If S is closed and bounded (such as a sphere), then f attains its minimum and maximum on S (Theorem A.3.5), and we find them by comparing the values of f at all solutions of these equations.

Example 3.12.2. We find the maximum and minimum of $f(x, y) = x + y$ on the unit circle $g(x, y) = x^2 + y^2 - 1 = 0$. The condition $\nabla f + \lambda \nabla g = \mathbf{0}$ is $(1, 1) + \lambda(2x, 2y) = (0, 0)$, so $x = y = -\frac{1}{2\lambda}$, and with $x^2 + y^2 = 1$, we get $x = y = \pm\frac{1}{\sqrt{2}}$. The maximum is $f(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}) = \sqrt{2}$, and the minimum is $-\sqrt{2}$. This agrees with the Cauchy–Schwarz inequality: $x + y = (1, 1)(x, y)^\top \le \sqrt{2}\sqrt{x^2 + y^2} = \sqrt{2}$. See Figure 3.10.




Figure 3.10: Maximizing $f(x, y) = x + y$ on the unit circle $g(x, y) = x^2 + y^2 - 1 = 0$. The dashed lines are level lines of f. At the maximum $\mathbf{x}^*$, the level line $x + y = \sqrt{2}$ touches the circle, and ∇f and ∇g are parallel. At the other marked point, they are not parallel: ∇f has a component along the circle, so moving along the circle in that direction increases f.

Example 3.12.3 (Eigenvalues and PCA)**.** Let A be a symmetric $n \times n$ matrix. We maximize $f(\mathbf{x}) = \mathbf{x}^\top A\mathbf{x}$ subject to $g(\mathbf{x}) = 1 - \mathbf{x}^\top\mathbf{x} = 0$. Here $\nabla g(\mathbf{x}) = -2\mathbf{x} \neq \mathbf{0}$ on the unit sphere, and the Lagrange condition $2A\mathbf{x} - 2\lambda\mathbf{x} = \mathbf{0}$ says that

$$A\mathbf{x} = \lambda\mathbf{x}\,:$$

the candidates are exactly the unit eigenvectors of A, and the Lagrange multiplier is the eigenvalue. At a unit eigenvector, $f(\mathbf{x}) = \mathbf{x}^\top(\lambda\mathbf{x}) = \lambda$. Since the sphere is closed and bounded, the maximum exists, and it is the largest eigenvalue $\lambda_{\max}$, attained at a corresponding eigenvector. This is the Rayleigh quotient result of Lecture 2, found with calculus instead of the spectral theorem; in PCA, it gives the first principal component.

Example 3.12.4 (The minimum norm solution)**.** Let $A \in \mathbb{R}^{m\times n}$ have linearly independent rows $\mathbf{a}_1^\top, \ldots, \mathbf{a}_m^\top$ (so $m \leq n$), and let $\mathbf{b} \in \mathbb{R}^m$. The system $A\mathbf{x} = \mathbf{b}$ has infinitely many solutions if $m < n$, and we look for the one with the smallest norm: minimize $\|\mathbf{x}\|^2$ subject to $g_i(\mathbf{x}) = \mathbf{a}_i^\top\mathbf{x} - b_i = 0$ for $i = 1, \ldots, m$. The gradients $\nabla g_i = \mathbf{a}_i$ are linearly independent, and the Lagrange condition is

$$2\mathbf{x} + \sum_{i=1}^{m} \lambda_i \mathbf{a}_i = 2\mathbf{x} + A^\top\boldsymbol{\lambda} = \mathbf{0}, \qquad \text{so} \qquad \mathbf{x} = A^\top\boldsymbol{\mu} \text{ with } \boldsymbol{\mu} = -\tfrac{1}{2}\boldsymbol{\lambda}.$$

So $\mathbf{x}$ lies in the row space of A. Substituting into $A\mathbf{x} = \mathbf{b}$ gives $AA^\top\boldsymbol{\mu} = \mathbf{b}$. The $m \times m$ matrix $AA^\top$ is invertible (it is positive definite, since $\mathbf{v}^\top AA^\top\mathbf{v} = \|A^\top\mathbf{v}\|^2 > 0$ for $\mathbf{v} \neq \mathbf{0}$), so

$$\mathbf{x}^* = A^\top(AA^\top)^{-1}\mathbf{b}.$$

This is indeed the global minimum: every solution of $A\mathbf{x} = \mathbf{b}$ is $\mathbf{x} = \mathbf{x}^* + \mathbf{v}$ with $\mathbf{v} \in \operatorname{null}(A)$, and since $\mathbf{x}^* \in \operatorname{row}(A) = \operatorname{null}(A)^\perp$, Pythagoras gives $\|\mathbf{x}\|^2 = \|\mathbf{x}^*\|^2 + \|\mathbf{v}\|^2 \geq \|\mathbf{x}^*\|^2$. By the proposition on the pseudo-inverse in Lecture 2, applied to $A^\top$ (which has linearly independent columns), $(A^\top)^+ = (AA^\top)^{-1}A$. By the definition of the pseudo-inverse with the SVD, $(A^\top)^+ = (A^+)^\top$, so taking transposes gives $A^+ = A^\top(AA^\top)^{-1}$. So $\mathbf{x}^* = A^+\mathbf{b}$, which gives a second proof, for matrices with linearly independent rows, of the result of Lecture 2 that $A^+\mathbf{b}$ is the solution with the smallest norm.

3.12.2 Inequality constraints and the KKT conditions

Now consider inequality constraints:

$$\text{minimize } f(\mathbf{x}) \quad \text{subject to} \quad g_1(\mathbf{x}) \le 0,\ \dots,\ g_m(\mathbf{x}) \le 0.$$

At a feasible point $\mathbf{x}^*$, the constraint g_i is active if $g_i(\mathbf{x}^*) = 0$, and inactive if $g_i(\mathbf{x}^*) < 0$. Inactive constraints do not matter near $\mathbf{x}^*$, since they remain satisfied when we move $\mathbf{x}^*$ a little. Active constraints behave like equality constraints, with one difference: moving into the feasible region, where $g_i < 0$, is allowed. At a minimum, this must not decrease f, and this forces the multipliers to be nonnegative.

Theorem 3.12.5 (Karush–Kuhn–Tucker conditions)**.** *Let $f, g_1, \dots, g_m$ be continuously differentiable, and let $\mathbf{x}^*$ be a local minimum of f subject to $g_1(\mathbf{x}) \le 0, \dots, g_m(\mathbf{x}) \le 0$, such that the gradients of the active constraints at $\mathbf{x}^*$ are linearly independent. Then there are numbers $\mu_1, \dots, \mu_m$ such that:*

1. *(stationarity) $\nabla f(\mathbf{x}^*) + \mu_1 \nabla g_1(\mathbf{x}^*) + \cdots + \mu_m \nabla g_m(\mathbf{x}^*) = \mathbf{0}$;*
2. *(primal feasibility) $g_i(\mathbf{x}^*) \le 0$ for all i;*
3. *(dual feasibility) $\mu_i \ge 0$ for all i;*
4. *(complementary slackness) $\mu_i g_i(\mathbf{x}^*) = 0$ for all i, i.e., $\mu_i = 0$ for every inactive constraint.*

Conversely, if f and all g_i are convex and differentiable, then every $\mathbf{x}^*$ that satisfies (1)–(4) for some $\mu_1, \dots, \mu_m$ is a global minimum.

Proof of the converse. The first part is Theorem A.8.2, which is proved with the implicit function theorem. For the converse, let $\mathbf{x}$ be feasible. The function $\mathcal{L}(\mathbf{z}) = f(\mathbf{z}) + \sum_i \mu_i g_i(\mathbf{z})$ is convex by Proposition 3.11.8, since $\mu_i \ge 0$, and $\nabla \mathcal{L}(\mathbf{x}^*) = \mathbf{0}$ by (1). By Corollary 3.11.4, $\mathcal{L}(\mathbf{x}) \ge \mathcal{L}(\mathbf{x}^*)$. Since $\mu_i g_i(\mathbf{x}) \le 0$ for feasible $\mathbf{x}$ and $\mu_i g_i(\mathbf{x}^*) = 0$ by (4),

$$f(\mathbf{x}) \ge f(\mathbf{x}) + \sum_i \mu_i g_i(\mathbf{x}) = \mathcal{L}(\mathbf{x}) \ge \mathcal{L}(\mathbf{x}^*) = f(\mathbf{x}^*). \qquad \square$$

Equality constraints can be added to the KKT conditions with multipliers of any sign, as in Theorem 3.12.1.

Example 3.12.6. Minimize $f(x) = (x-2)^2$ subject to $x \le 1$, i.e., $g(x) = x - 1 \le 0$. The KKT conditions are $2(x-2) + \mu = 0$, $x \le 1$, $\mu \ge 0$ and $\mu(x-1) = 0$. If $\mu = 0$, then $x = 2$, which is not feasible. So the constraint is active, $x = 1$, and $\mu = 2 \ge 0$. Since f and g are convex, $x^* = 1$ is the global minimum. The multiplier measures how much the constraint costs: if we relax it to $x \le 1 + \varepsilon$, the minimum value $(1 + \varepsilon - 2)^2 = 1 - 2\varepsilon + \varepsilon^2$ decreases at the rate $\mu = 2$. For the constraint $x \le 3$ instead, the minimum $x^* = 2$ satisfies the constraint, which is inactive, and $\mu = 0$.

Example 3.12.7 (Projection onto a ball)**.** Let $\mathbf{p} \in \mathbb{R}^n$ with $\|\mathbf{p}\| > 1$. We find the point of the unit ball closest to $\mathbf{p}$: minimize $\|\mathbf{x} - \mathbf{p}\|^2$ subject to $\|\mathbf{x}\|^2 - 1 \le 0$. Stationarity is $2(\mathbf{x} - \mathbf{p}) + 2\mu\mathbf{x} = \mathbf{0}$, i.e., $\mathbf{x} = \frac{\mathbf{p}}{1+\mu}$. If $\mu = 0$, then $\mathbf{x} = \mathbf{p}$ is not feasible, so the constraint is active, $\|\mathbf{x}\| = 1$, which gives $1 + \mu = \|\mathbf{p}\|$, $\mu = \|\mathbf{p}\| - 1 > 0$, and

$$\mathbf{x}^* = \frac{\mathbf{p}}{\|\mathbf{p}\|}.$$

For example, for $\mathbf{p} = (3, 4)^\top$, we get $\mathbf{x}^* = (0.6, 0.8)^\top$ and $\mu = 4$. This projection is used in projected gradient descent, which alternates gradient steps with projections onto the feasible set.

Remark 3.12.8 (Support vector machines). For linearly separable data $(\mathbf{x}_i, y_i)$ with labels $y_i = \pm 1$, the support vector machine finds the separating hyperplane $\mathbf{w}^\top \mathbf{x} + b = 0$ with the largest margin. By the formula for the distance to a hyperplane in Lecture 2, this leads to the convex problem

$$\text{minimize } \tfrac{1}{2}\|\mathbf{w}\|^2 \quad \text{subject to} \quad 1 - y_i(\mathbf{w}^\top \mathbf{x}_i + b) \le 0 \quad \text{for } i = 1, \dots, N,$$

whose margin is $\frac{1}{\|\mathbf{w}\|}$. The stationarity condition gives $\mathbf{w} = \sum_i \mu_i y_i \mathbf{x}_i$, and by complementary slackness, $\mu_i > 0$ only for the points with $y_i(\mathbf{w}^\top \mathbf{x}_i + b) = 1$, which lie exactly on the margin. These points are the support vectors: the classifier only depends on them.

3.12.3 Lagrangian duality

The multipliers of the KKT conditions are more than auxiliary variables: they solve an optimization problem of their own. Consider again the problem

$$\text{minimize } f(\mathbf{x}) \quad \text{subject to} \quad g_1(\mathbf{x}) \le 0, \ \dots, \ g_m(\mathbf{x}) \le 0,$$

with the optimal value $p^* = \inf\{f(\mathbf{x}) : \mathbf{x} \text{ feasible}\}$, the primal problem.

Definition 3.12.9 (Dual problem)**.** The Lagrangian of the problem is $\mathcal{L}(\mathbf{x}, \boldsymbol{\mu}) = f(\mathbf{x}) + \sum_{i=1}^m \mu_i g_i(\mathbf{x})$, the dual function is

$$d(\boldsymbol{\mu}) = \inf_{\mathbf{x} \in \mathbb{R}^n} \mathcal{L}(\mathbf{x}, \boldsymbol{\mu}) \qquad \text{for } \boldsymbol{\mu} \ge \mathbf{0}$$

(which may be $-\infty$), and the dual problem is to maximize $d(\boldsymbol{\mu})$ subject to $\boldsymbol{\mu} \ge \mathbf{0}$, i.e., $\mu_i \ge 0$ for all i, with the optimal value $d^* = \sup_{\boldsymbol{\mu} \ge \mathbf{0}} d(\boldsymbol{\mu})$.

For each fixed $\mathbf{x}$, $\boldsymbol{\mu} \mapsto \mathcal{L}(\mathbf{x}, \boldsymbol{\mu})$ is affine, and an infimum of affine functions is concave: if $\boldsymbol{\mu} = \theta \boldsymbol{\mu}_1 + (1-\theta)\boldsymbol{\mu}_2$, then $\mathcal{L}(\mathbf{x}, \boldsymbol{\mu}) = \theta \mathcal{L}(\mathbf{x}, \boldsymbol{\mu}_1) + (1-\theta)\mathcal{L}(\mathbf{x}, \boldsymbol{\mu}_2) \ge \theta d(\boldsymbol{\mu}_1) + (1-\theta) d(\boldsymbol{\mu}_2)$ for every $\mathbf{x}$. So the dual problem is always a concave maximization problem, even if the primal problem is not convex.

Theorem 3.12.10 (Weak and strong duality)**.**

1. *(Weak duality.)* $d(\boldsymbol{\mu}) \le f(\mathbf{x})$ *for every feasible* $\mathbf{x}$ *and every* $\boldsymbol{\mu} \ge \mathbf{0}$*. Hence* $d^* \le p^*$*.*
2. *(Strong duality.) Let* $f, g_1, \dots, g_m$ *be convex and differentiable, and let* $\mathbf{x}^*$ *and* $\boldsymbol{\mu}^*$ *satisfy the KKT conditions of Theorem 3.12.5. Then* $\mathbf{x}^*$ *solves the primal problem,* $\boldsymbol{\mu}^*$ *solves the dual problem, and* $d(\boldsymbol{\mu}^*) = f(\mathbf{x}^*)$*, so* $d^* = p^*$*.*

Proof. (1) For feasible $\mathbf{x}$, $\mu_i g_i(\mathbf{x}) \le 0$, so $d(\boldsymbol{\mu}) \le \mathcal{L}(\mathbf{x}, \boldsymbol{\mu}) \le f(\mathbf{x})$.

(2) The function $\mathcal{L}(\cdot, \boldsymbol{\mu}^*)$ is convex (Proposition 3.11.8, since $\mu_i^* \ge 0$), and its gradient at $\mathbf{x}^*$ is $\mathbf{0}$ by stationarity, so $\mathbf{x}^*$ minimizes it (Corollary 3.11.4). By complementary slackness, $d(\boldsymbol{\mu}^*) = \mathcal{L}(\mathbf{x}^*, \boldsymbol{\mu}^*) = f(\mathbf{x}^*)$. By weak duality, $d(\boldsymbol{\mu}) \le f(\mathbf{x}^*) = d(\boldsymbol{\mu}^*)$ for all $\boldsymbol{\mu} \ge \mathbf{0}$, and $f(\mathbf{x}) \ge d(\boldsymbol{\mu}^*) = f(\mathbf{x}^*)$ for all feasible $\mathbf{x}$. □

Together with Theorem 3.12.5, strong duality holds for every convex problem whose solution has linearly independent gradients of the active constraints. Duality is useful in several ways. Every $\boldsymbol{\mu} \ge \mathbf{0}$ gives a lower bound $d(\boldsymbol{\mu})$ for the optimal value, so the duality gap $f(\mathbf{x}) - d(\boldsymbol{\mu}) \ge f(\mathbf{x}) - p^*$ certifies how far a feasible point $\mathbf{x}$ is from being optimal; solvers stop when it is small. The dual problem may also be easier, for example when it has fewer variables, or simpler constraints.

Example 3.12.11. For the problem of Example 3.12.6, minimize $(x-2)^2$ subject to $x - 1 \le 0$, the Lagrangian $(x-2)^2 + \mu(x-1)$ is smallest at $x = 2 - \frac{\mu}{2}$, so $d(\mu) = \frac{\mu^2}{4} + \mu(1 - \frac{\mu}{2}) = \mu - \frac{\mu^2}{4}$. The concave function d is largest at $\mu = 2$, with $d^* = 1 = p^*$, and the maximizer is the KKT multiplier $\mu = 2$ found there.

Example 3.12.12 (The dual of the support vector machine)**.** For the support vector machine of the remark above, with the variables $\mathbf{w}$ and b, the Lagrangian is

$$\mathcal{L}(\mathbf{w}, b, \boldsymbol{\mu}) = \frac{1}{2}\|\mathbf{w}\|^2 + \sum_{i=1}^{N} \mu_i(1 - y_i(\mathbf{w}^\top \mathbf{x}_i + b)) = \frac{1}{2}\|\mathbf{w}\|^2 - \mathbf{w}^\top \sum_i \mu_i y_i \mathbf{x}_i - b\sum_i \mu_i y_i + \sum_i \mu_i.$$

As a function of b, it is affine, so its infimum over b is $-\infty$ unless $\sum_i \mu_i y_i = 0$. As a function of $\mathbf{w}$, it is a convex quadratic function, which is smallest where its gradient $\mathbf{w} - \sum_i \mu_i y_i \mathbf{x}_i$ vanishes. Substituting $\mathbf{w} = \sum_i \mu_i y_i \mathbf{x}_i$ gives the dual problem

$$\text{maximize } \sum_{i=1}^{N} \mu_i - \frac{1}{2}\sum_{i,j=1}^{N} \mu_i \mu_j y_i y_j \, \mathbf{x}_i^\top \mathbf{x}_j \quad \text{subject to} \quad \boldsymbol{\mu} \geq \mathbf{0}, \quad \sum_{i=1}^{N} \mu_i y_i = 0.$$

Whenever the KKT conditions hold at the solution, Theorem 3.12.10 shows that solving the dual problem solves the support vector machine, with $\mathbf{w} = \sum_i \mu_i y_i \mathbf{x}_i$. The dual has one variable per data point instead of per feature, and it only depends on the data through the inner products $\mathbf{x}_i^\top \mathbf{x}_j$. Replacing them by a kernel $k(\mathbf{x}_i, \mathbf{x}_j)$ (the remark on kernels in Lecture 2) gives the kernel support vector machine, which separates the data by a nonlinear boundary.

For example, for the two points $\mathbf{x}_1 = (1, 0)^\top$ with $y_1 = 1$ and $\mathbf{x}_2 = (-1, 0)^\top$ with $y_2 = -1$, the constraint $\mu_1 - \mu_2 = 0$ gives $\mu_1 = \mu_2 = m$, and $\sum_i \mu_i y_i \mathbf{x}_i = (2m, 0)^\top$, so the dual function is $2m - 2m^2$. It is largest at $m = \frac{1}{2}$, with the value $\frac{1}{2}$. Then $\mathbf{w} = (1, 0)^\top$, and $b = 0$ from the active constraint $y_1(\mathbf{w}^\top \mathbf{x}_1 + b) = 1$. The separating line is $x_1 = 0$, with margin $\frac{1}{\|\mathbf{w}\|} = 1$, and the primal value $\frac{1}{2}\|\mathbf{w}\|^2 = \frac{1}{2}$ equals the dual value.

For problems with linear constraints, the dual function can be written with the following transform of f.

Definition 3.12.13 (Convex conjugate)**.** The convex conjugate of $f : \mathbb{R}^n \to \mathbb{R}$ is the function

$$f^*(\mathbf{y}) = \sup_{\mathbf{x} \in \mathbb{R}^n} \left(\mathbf{y}^\top \mathbf{x} - f(\mathbf{x})\right) \qquad (\mathbf{y} \in \mathbb{R}^n),$$

which may take the value $+\infty$.

So $f^*(\mathbf{y})$ is the largest amount by which the linear function $\mathbf{y}^\top \mathbf{x}$ exceeds f. For $n = 1$: the line with slope y that touches the graph of f from below crosses the vertical axis at $-f^*(y)$.

Proposition 3.12.14. *1. The set where f^* is finite is convex, and f^* is convex on it.*

2. (Fenchel–Young inequality.) $f(\mathbf{x}) + f^*(\mathbf{y}) \geq \mathbf{x}^\top \mathbf{y}$ for all $\mathbf{x}$ and $\mathbf{y}$.

3. If f is convex and differentiable, then $f^*(\nabla f(\mathbf{x})) = \mathbf{x}^\top \nabla f(\mathbf{x}) - f(\mathbf{x})$ for every $\mathbf{x}$, and $f^{**} = f$, i.e., $f(\mathbf{x}) = \sup_{\mathbf{y}} \left(\mathbf{x}^\top \mathbf{y} - f^*(\mathbf{y})\right)$.

Proof. (1) If $f^*(\mathbf{y}_1)$ and $f^*(\mathbf{y}_2)$ are finite and $\theta \in [0, 1]$, then for every $\mathbf{x}$,

$$(\theta\mathbf{y}_1 + (1-\theta)\mathbf{y}_2)^\top \mathbf{x} - f(\mathbf{x}) = \theta(\mathbf{y}_1^\top \mathbf{x} - f(\mathbf{x})) + (1-\theta)(\mathbf{y}_2^\top \mathbf{x} - f(\mathbf{x})) \leq \theta f^*(\mathbf{y}_1) + (1-\theta) f^*(\mathbf{y}_2),$$

and the supremum over $\mathbf{x}$ gives $f^*(\theta\mathbf{y}_1 + (1-\theta)\mathbf{y}_2) \leq \theta f^*(\mathbf{y}_1) + (1-\theta) f^*(\mathbf{y}_2) < \infty$.

(2) $\mathbf{y}^\top \mathbf{x} - f(\mathbf{x}) \leq f^*(\mathbf{y})$ by the definition of the supremum.

(3) Let $\mathbf{y} = \nabla f(\mathbf{x})$. By Theorem 3.11.3, $f(\mathbf{z}) \geq f(\mathbf{x}) + \mathbf{y}^\top(\mathbf{z} - \mathbf{x})$, i.e., $\mathbf{y}^\top \mathbf{z} - f(\mathbf{z}) \leq \mathbf{y}^\top \mathbf{x} - f(\mathbf{x})$ for all $\mathbf{z}$, with equality for $\mathbf{z} = \mathbf{x}$; so $f^*(\mathbf{y}) = \mathbf{y}^\top \mathbf{x} - f(\mathbf{x})$. Hence $\sup_{\mathbf{y}}(\mathbf{x}^\top \mathbf{y} - f^*(\mathbf{y})) \geq \mathbf{x}^\top \nabla f(\mathbf{x}) - f^*(\nabla f(\mathbf{x})) = f(\mathbf{x})$, and the reverse inequality is (2). □

Part 3 says that a convex differentiable function is the supremum of its tangent functions $\mathbf{x} \mapsto f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0)^\top(\mathbf{x} - \mathbf{x}_0)$, and that f^* records the tangents by their slopes: f and f^* contain the same information. (This is the Legendre transform of physics.)

Example 3.12.15. 1. For $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top Q\mathbf{x}$ with Q symmetric positive definite, the supremum of $\mathbf{y}^\top\mathbf{x} - \frac{1}{2}\mathbf{x}^\top Q\mathbf{x}$ is attained where $\mathbf{y} = Q\mathbf{x}$, so $f^*(\mathbf{y}) = \frac{1}{2}\mathbf{y}^\top Q^{-1}\mathbf{y}$. In particular, $\frac{1}{2}\|\mathbf{x}\|^2$ is its own conjugate.

2. For $f(x) = e^x$, $xy - e^x$ is largest at $x = \log y$ if $y > 0$, so $f^*(y) = y\log y - y$; $f^*(0) = 0$; and $f^*(y) = \infty$ for $y < 0$, since $xy - e^x \to \infty$ as $x \to -\infty$.

3. For $f(\mathbf{x}) = \|\mathbf{x}\|_1$: if $|y_i| \le 1$ for all i, then $\mathbf{y}^\top\mathbf{x} - \|\mathbf{x}\|_1 = \sum_i (y_i x_i - |x_i|) \le 0$, with equality for $\mathbf{x} = \mathbf{0}$, so $f^*(\mathbf{y}) = 0$; if $|y_j| > 1$, then $\mathbf{x} = t\,\operatorname{sign}(y_j)\mathbf{e}_j$ gives $t(|y_j| - 1) \to \infty$. So f^* is 0 on the unit ball of the ℓ_∞ norm and ∞ outside it.

4. For the log-sum-exp function $\operatorname{lse}(\mathbf{x}) = \log(e^{x_1} + \cdots + e^{x_n})$, which is convex with the gradient $\operatorname{softmax}(\mathbf{x})$: if $\mathbf{p}$ has positive entries with sum 1, then $\operatorname{softmax}(\mathbf{x}) = \mathbf{p}$ for $x_i = \log p_i$, and part 3 gives $\operatorname{lse}^*(\mathbf{p}) = \sum_i p_i \log p_i - \log \sum_i p_i = \sum_i p_i \log p_i$, the negative entropy of the distribution $\mathbf{p}$ (Lecture 4).

Now consider the problem of minimizing $f(\mathbf{x})$ subject to $A\mathbf{x} \le \mathbf{b}$ (i.e., $g_i(\mathbf{x}) = \mathbf{a}_i^\top\mathbf{x} - b_i \le 0$ for the rows $\mathbf{a}_i^\top$ of A). Its dual function is

$$d(\boldsymbol{\mu}) = \inf_{\mathbf{x}} \left(f(\mathbf{x}) + \boldsymbol{\mu}^\top(A\mathbf{x} - \mathbf{b})\right) = -\mathbf{b}^\top\boldsymbol{\mu} - \sup_{\mathbf{x}} \left((-A^\top\boldsymbol{\mu})^\top\mathbf{x} - f(\mathbf{x})\right) = -\mathbf{b}^\top\boldsymbol{\mu} - f^*(-A^\top\boldsymbol{\mu}).$$

Example 3.12.16 (Linear and quadratic programs)**.** 1. A linear program minimizes $f(\mathbf{x}) = \mathbf{c}^\top\mathbf{x}$ subject to $A\mathbf{x} \le \mathbf{b}$. Here $f^*(\mathbf{y}) = \sup_{\mathbf{x}}(\mathbf{y} - \mathbf{c})^\top\mathbf{x}$ is 0 for $\mathbf{y} = \mathbf{c}$ and ∞ otherwise. So the dual problem is again a linear program: maximize $-\mathbf{b}^\top\boldsymbol{\mu}$ subject to $A^\top\boldsymbol{\mu} = -\mathbf{c}$ and $\boldsymbol{\mu} \ge \mathbf{0}$. For example, minimize $-x_1 - x_2$ subject to $0 \le x_1 \le 1$, $0 \le x_2 \le 1$ and $x_1 + x_2 \le 1.5$. The optimal value is -1.5, attained at $(1, 0.5)$, for example. The dual point with the multiplier 1 for $x_1 + x_2 \le 1.5$ and 0 for the other constraints is feasible, with the dual value -1.5: it is a certificate of optimality, since adding the constraint $x_1 + x_2 \le 1.5$ with the weight 1 proves $-x_1 - x_2 \ge -1.5$ for every feasible $\mathbf{x}$.

2. A quadratic program minimizes $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top Q\mathbf{x} + \mathbf{c}^\top\mathbf{x}$ with Q symmetric positive definite, subject to $A\mathbf{x} \le \mathbf{b}$. As in Example 3.12.15, $f^*(\mathbf{y}) = \frac{1}{2}(\mathbf{y}-\mathbf{c})^\top Q^{-1}(\mathbf{y}-\mathbf{c})$, so the dual problem is to maximize $-\mathbf{b}^\top\boldsymbol{\mu} - \frac{1}{2}(A^\top\boldsymbol{\mu} + \mathbf{c})^\top Q^{-1}(A^\top\boldsymbol{\mu} + \mathbf{c})$ subject to $\boldsymbol{\mu} \ge \mathbf{0}$. It has only the simple constraints $\mu_i \ge 0$, which projected gradient ascent handles by setting negative entries to 0. The dual of the support vector machine is of this kind.

Remark 3.12.17 (Constraint qualifications). Theorem 3.12.10 needs a point that satisfies the KKT conditions, which Theorem 3.12.5 guarantees when the gradients of the active constraints are linearly independent. Strong duality holds more generally: for linear programs whenever p^* is finite, and for convex problems that have a strictly feasible point, with $g_i(\bar{\mathbf{x}}) < 0$ for all i (Slater's condition). Both facts are proved with separating hyperplanes, and we do not prove them here.

3.13 Vector Fields: Divergence and Curl

Definition 3.13.1 (Vector field)**.** A vector field on an open set $D \subseteq \mathbb{R}^n$ is a function $F : D \to \mathbb{R}^n$. We picture it by attaching to every point $\mathbf{x} \in D$ the arrow $F(\mathbf{x})$, starting at $\mathbf{x}$.

The most important examples for us are gradient fields: the gradient ∇f of a differentiable function $f : D \to \mathbb{R}$ is a vector field, whose arrows point in the direction of steepest ascent of f and are orthogonal to its level sets (Section 3.3). Another example is the velocity field of a fluid, which tells us how fast and in which direction a particle at $\mathbf{x}$ moves. This motion is described by a differential equation, the subject of Section 3.15.

The derivative of a vector field is a square matrix, the Jacobian $DF(\mathbf{x}) \in \mathbb{R}^{n\times n}$, and in this section we extract two geometric quantities from it: the divergence, which measures how much of F flows out of small regions, and the curl, which measures how F rotates.

Example 3.13.2. 1. The gradient field of $f(\mathbf{x}) = \frac{1}{2}\|\mathbf{x}\|^2$ is $F(\mathbf{x}) = \mathbf{x}$, which points away from the origin. For a symmetric matrix A, the field $A\mathbf{x}$ is the gradient field of $\frac{1}{2}\mathbf{x}^\top A\mathbf{x}$ (Proposition 3.3.3).

2. The rotation field $F(x, y) = (-y, x)^\top$ is orthogonal to the position vector $(x, y)^\top$ everywhere: its arrows go around the origin counterclockwise (Figure 3.11).

3. Let p be a positive probability density on $\mathbb{R}^n$. Its score is the vector field $\mathbf{s}(\mathbf{x}) = \nabla \log p(\mathbf{x}) = \frac{\nabla p(\mathbf{x})}{p(\mathbf{x})}$, which points in the direction in which the density increases fastest. For the Gaussian density $p(\mathbf{x}) \propto \exp\big(-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})\big)$ of Lecture 4, $\mathbf{s}(\mathbf{x}) = -\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})$ points towards the mean. Score-based generative models, such as diffusion models, learn the score of the distribution of the training data with a neural network $\mathbf{s}_{\boldsymbol{\theta}} : \mathbb{R}^n \to \mathbb{R}^n$ (Lecture 4).

Near a point $\mathbf{x}$, $F(\mathbf{x}+\mathbf{h}) \approx F(\mathbf{x}) + DF(\mathbf{x})\mathbf{h}$, and every square matrix is the sum of a symmetric and a skew-symmetric matrix:

$$DF(\mathbf{x}) = S + W, \qquad S = \tfrac{1}{2}\big(DF(\mathbf{x}) + DF(\mathbf{x})^\top\big), \qquad W = \tfrac{1}{2}\big(DF(\mathbf{x}) - DF(\mathbf{x})^\top\big).$$

By the spectral theorem (Lecture 2), S has an orthonormal basis of eigenvectors: the symmetric part stretches or compresses a small neighborhood of $\mathbf{x}$ along these orthogonal directions, at rates given by the eigenvalues. The skew-symmetric part satisfies $\mathbf{h}^\top W\mathbf{h} = 0$, so $W\mathbf{h}$ is orthogonal to $\mathbf{h}$: it is an infinitesimal rotation. The divergence summarizes the symmetric part, and the curl is the skew-symmetric part.

Definition 3.13.3 (Divergence)**.** The divergence of a differentiable vector field F is the function

$$\operatorname{div} F(\mathbf{x}) = \frac{\partial F_1}{\partial x_1}(\mathbf{x}) + \cdots + \frac{\partial F_n}{\partial x_n}(\mathbf{x}) = \operatorname{tr} DF(\mathbf{x}).$$

It is also written $\nabla \cdot F$, as a formal dot product of $\nabla = (\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_n})^\top$ with F.

Since $\operatorname{tr} W = 0$, the divergence only depends on the symmetric part: $\operatorname{div} F(\mathbf{x}) = \operatorname{tr} S$ is the sum of the stretching rates. To see what it measures, think of F as the velocity field of a fluid, and consider a small box Q around $\mathbf{x}$ with the side lengths $h_1, \ldots, h_n$. The two faces of Q perpendicular to $\mathbf{e}_1$ have the area $\frac{\operatorname{vol}(Q)}{h_1}$ and the outward normals $\pm\mathbf{e}_1$, so the fluid flows out through them at the rates $F_1 \cdot \frac{\operatorname{vol}(Q)}{h_1}$ and $-F_1 \cdot \frac{\operatorname{vol}(Q)}{h_1}$, evaluated at the centers of the faces. The net outflow through the two faces is approximately

$$\Big(F_1\big(\mathbf{x} + \tfrac{h_1}{2}\mathbf{e}_1\big) - F_1\big(\mathbf{x} - \tfrac{h_1}{2}\mathbf{e}_1\big)\Big)\frac{\operatorname{vol}(Q)}{h_1} \approx \frac{\partial F_1}{\partial x_1}(\mathbf{x})\operatorname{vol}(Q).$$

Adding the contributions of all n pairs of faces, we find that the net outflow from Q is approximately $\operatorname{div} F(\mathbf{x})$ times the volume of Q. In other words,

$$\operatorname{div} F(\mathbf{x}) \textit{ is the net outflow of } F \textit{ per unit volume near } \mathbf{x}.$$

The divergence theorem (Theorem 3.14.12) makes this precise. A vector field with positive divergence at $\mathbf{x}$ acts like a source there, one with negative divergence like a sink, and a field with $\operatorname{div} F = 0$ describes an incompressible fluid, in which as much flows into every region as flows out. In Section 3.16, we will see that the divergence is also the rate at which the flow of the differential equation $\mathbf{x}' = F(\mathbf{x})$ expands volumes.

Example 3.13.4. 1. $F(\mathbf{x}) = \mathbf{x}$ has $\operatorname{div} F = n$: the field points away from the origin and grows in length, so more flows out of every small region than flows in.

2. The rotation field $(-y, x)^\top$ has $\operatorname{div} F = 0 + 0 = 0$: the fluid circulates without being compressed.

3. For $F(\mathbf{x}) = A\mathbf{x}$, we have $DF = A$ and $\operatorname{div} F = \operatorname{tr} A$. For example, the field of $A = \left(\begin{smallmatrix} -2 & 1 \\ 1 & -2 \end{smallmatrix}\right)$ has $\operatorname{div} F = -4$ everywhere: it is a sink.

4. The field $-\nabla f$, which points downhill, has $\operatorname{div}(-\nabla f) = -\operatorname{tr} \nabla^2 f$. Near a strict local minimum, where the Hessian is positive definite, it is a sink.

Definition 3.13.5 (Laplacian)**.** The Laplacian of a twice differentiable function $f : D \to \mathbb{R}$ is

$$\Delta f = \operatorname{div} \nabla f = \frac{\partial^2 f}{\partial x_1^2} + \cdots + \frac{\partial^2 f}{\partial x_n^2} = \operatorname{tr} \nabla^2 f.$$

Functions with $\Delta f = 0$ are called harmonic.

The trace of the Hessian is the sum of its eigenvalues, so $\Delta f(\mathbf{x})$ is the sum of the curvatures of f in the principal directions (Section 3.9.1), and $\Delta f \geq 0$ at a local minimum. For example, $\Delta \|\mathbf{x}\|^2 = 2n$, while $x^2 - y^2$ is harmonic: the curvatures 2 and -2 of the saddle cancel. The Laplacian also describes diffusion, such as the spreading of heat, or of the probability density of a randomly moving particle (Lecture 4).

Definition 3.13.6 (Curl)**.** The curl of a differentiable vector field F on $D \subseteq \mathbb{R}^2$ is the function $\operatorname{curl} F = \frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2}$. For $D \subseteq \mathbb{R}^3$, it is the vector field

$$\operatorname{curl} F = \nabla \times F = \Big(\frac{\partial F_3}{\partial x_2} - \frac{\partial F_2}{\partial x_3},\ \frac{\partial F_1}{\partial x_3} - \frac{\partial F_3}{\partial x_1},\ \frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2}\Big)^\top.$$

Up to the factor 2, the entries of the curl are the entries $W_{ij} = \frac{1}{2}(\frac{\partial F_i}{\partial x_j} - \frac{\partial F_j}{\partial x_i})$ of the skew-symmetric part. For $n = 2$,

$$W = \frac{\operatorname{curl} F(\mathbf{x})}{2} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

so on top of the stretching by S, the field rotates a small neighborhood of $\mathbf{x}$ counterclockwise with the angular velocity $\frac{1}{2} \operatorname{curl} F(\mathbf{x})$: a small paddle wheel in a fluid with the velocity field F spins at this rate. For $n = 3$, one checks that $W\mathbf{h} = \frac{1}{2} \operatorname{curl} F(\mathbf{x}) \times \mathbf{h}$, a rotation about the axis $\operatorname{curl} F(\mathbf{x})$ with the angular velocity $\frac{1}{2}\|\operatorname{curl} F(\mathbf{x})\|$. In n dimensions, the skew-symmetric matrix W has $\frac{n(n-1)}{2}$ independent entries, which is 1 for $n = 2$ and 3 for $n = 3$; only in three dimensions can they be arranged into a vector field.

Example 3.13.7.

1. The rotation field $(-y, x)^\top$ has $\operatorname{curl} F = 1 - (-1) = 2$, so it rotates with the angular velocity 1. The field $\mathbf{x}$ has $\operatorname{curl} F = 0$.


2. The shear field $F(x, y) = (y, 0)^\top$ (Figure 3.11) has $\operatorname{div} F = 0$ and $\operatorname{curl} F = -1$. Its Jacobian splits as

$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix} + \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix}:$$

the shear of Lecture 1 combines a stretching along $(1, 1)$ and a compression along $(1, -1)$ (the eigenvalues $\pm\frac{1}{2}$ of S) with a clockwise rotation with the angular velocity $\frac{1}{2}$.

Proposition 3.13.8. *Let f and F be twice continuously differentiable.*

1. *The Jacobian of ∇f is the symmetric matrix $\nabla^2 f$. In particular,* $\operatorname{curl} \nabla f = 0$ *for $n = 2$ and $n = 3$.*



source $F = (x, y)$
$\operatorname{div} F = 2,\ \operatorname{curl} F = 0$

rotation $F = (-y, x)$
$\operatorname{div} F = 0,\ \operatorname{curl} F = 2$

shear $F = (y, 0)$
$\operatorname{div} F = 0,\ \operatorname{curl} F = -1$

Figure 3.11: Three linear vector fields. The source expands areas and does not rotate, the rotation turns without changing areas, and the shear is a combination of a stretching (along the diagonal) and a clockwise rotation, which preserves areas.

2. For $n = 3$, $\operatorname{div}\operatorname{curl} F = 0$.

Proof. (1) This is Schwarz's theorem (Theorem 3.8.2): the entries of $\operatorname{curl}\nabla f$ are differences $\frac{\partial^2 f}{\partial x_i \partial x_j} - \frac{\partial^2 f}{\partial x_j \partial x_i} = 0$. (2) By definition,

$$\operatorname{div}\operatorname{curl} F = \frac{\partial^2 F_3}{\partial x_1 \partial x_2} - \frac{\partial^2 F_2}{\partial x_1 \partial x_3} + \frac{\partial^2 F_1}{\partial x_2 \partial x_3} - \frac{\partial^2 F_3}{\partial x_2 \partial x_1} + \frac{\partial^2 F_2}{\partial x_3 \partial x_1} - \frac{\partial^2 F_1}{\partial x_3 \partial x_2},$$

and the terms cancel in pairs by Schwarz's theorem. □

So a gradient field has no rotational part. Conversely, a vector field on $\mathbb{R}^n$ whose Jacobian is symmetric everywhere is a gradient field (Theorem 3.14.10 below), while a field with a nonzero curl is not the gradient of any function. This matters whenever we follow a vector field and hope to minimize something.

Remark 3.13.9 (Vector fields in machine learning). 1. (Games.) Generative adversarial networks (GANs) train two networks against each other: a generator with the parameters $\mathbf{x}$ minimizes a function $f(\mathbf{x}, \mathbf{y})$, which a discriminator with the parameters $\mathbf{y}$ maximizes. Simultaneous gradient descent–ascent follows the vector field $F = (-\nabla_{\mathbf{x}} f, \nabla_{\mathbf{y}} f)$, which is in general not a gradient field. For $f(x, y) = xy$, $F(x, y) = (-y, x)$ is the rotation field, which is not a gradient field, since its curl is 2. Gradient descent–ascent with the step size η moves (x, y) to $(x - \eta y,\ y + \eta x)$, which multiplies the distance from the equilibrium $(0, 0)$ by $\sqrt{1 + \eta^2}$: the iterates spiral outwards instead of converging (Example 3.16.2 studies the continuous-time version). The oscillations of GAN training come from such rotational parts of the dynamics, and methods such as extragradient are designed to damp them.

2. (Scores.) A neural network $\mathbf{s}_{\boldsymbol{\theta}} : \mathbb{R}^n \to \mathbb{R}^n$ that approximates a score $\nabla \log p$ (Example 3.13.2) need not have a symmetric Jacobian, so in general it is not exactly a gradient field. Energy-based models instead learn a scalar function $E_{\boldsymbol{\theta}}$ and use the gradient field $-\nabla E_{\boldsymbol{\theta}}$.

3. (Computing divergences.) For a vector field $F : \mathbb{R}^n \to \mathbb{R}^n$ given by a neural network, computing $\operatorname{div} F = \operatorname{tr} DF$ exactly needs n passes of automatic differentiation, one for each diagonal entry. Hutchinson's estimator uses that $\mathbb{E}[\boldsymbol{\varepsilon}^\top A \boldsymbol{\varepsilon}] = \operatorname{tr}(A\,\mathbb{E}[\boldsymbol{\varepsilon}\boldsymbol{\varepsilon}^\top]) = \operatorname{tr} A$ for every random vector $\boldsymbol{\varepsilon}$ with $\mathbb{E}[\boldsymbol{\varepsilon}\boldsymbol{\varepsilon}^\top] = I$, for example with independent random signs as entries (expected values are treated in Lecture 4; see also Exercise 32). So $\boldsymbol{\varepsilon}^\top DF(\mathbf{x})\boldsymbol{\varepsilon}$ is on average equal to $\operatorname{div} F(\mathbf{x})$, and $\boldsymbol{\varepsilon}^\top DF(\mathbf{x})$ is a single vector-Jacobian product, computed by one pass of reverse mode (Section 3.7). Averages over a few random vectors are used in score matching (Section 3.14) and in continuous normalizing flows (Section 3.16).

3.14 Multiple Integrals and the Divergence Theorem

Probabilities and expectations of random vectors are integrals of densities over regions of $\mathbb{R}^n$ (Lecture 4), and we have already used integrals over the plane for the change of variables in Section 3.4. In this section, we define integrals of functions of several variables, compute them as iterated one-dimensional integrals, and prove the analogue of the fundamental theorem of calculus for vector fields, the divergence theorem. Its main consequence for us is a rule for integration by parts in $\mathbb{R}^n$, which is behind score matching and, in Section 3.16, behind the evolution of probability densities under a flow.

3.14.1 Integrals over boxes and Fubini's theorem

A box is a product of intervals $Q = [a_1, b_1] \times \cdots \times [a_n, b_n]$, with the volume $\operatorname{vol}(Q) = (b_1 - a_1) \cdots (b_n - a_n)$. Dividing each interval into k intervals of equal length divides Q into k^n small boxes $Q_1, \ldots, Q_{k^n}$, each with the volume $\frac{\operatorname{vol}(Q)}{k^n}$.

Definition 3.14.1 (Integral over a box)**.** A bounded function $f : Q \to \mathbb{R}$ is integrable if the Riemann sums $\sum_{j=1}^{k^n} f(\boldsymbol{\xi}_j) \operatorname{vol}(Q_j)$ converge as $k \to \infty$, to the same limit for all choices of points $\boldsymbol{\xi}_j \in Q_j$. The limit is the integral $\int_Q f(\mathbf{x})\, d\mathbf{x}$, also written $\iint_Q f(x, y)\, dx\, dy$ for $n = 2$.

As in one variable, continuous functions on boxes are integrable, the integral is linear, and $f \le g$ implies $\int_Q f \le \int_Q g$. For $f \ge 0$, the integral is the volume of the region below the graph of f. To integrate over a bounded set D, we choose a box $Q \supseteq D$ and integrate the function that agrees with f on D and is 0 outside, $\int_D f(\mathbf{x})\, d\mathbf{x} = \int_Q f(\mathbf{x}) \mathbf{1}_D(\mathbf{x})\, d\mathbf{x}$, where $\mathbf{1}_D$ is the indicator function of D. This works, for example, for continuous functions f and regions D bounded by graphs of continuous functions. In particular, $\operatorname{vol}(D) = \int_D 1\, d\mathbf{x}$. Integrals over unbounded sets, such as $\mathbb{R}^n$, are defined as limits of integrals over growing boxes, like improper integrals in one variable.

Integrals of several variables are computed one variable at a time.

Theorem 3.14.2 (Fubini)**.** *Let f be continuous on the rectangle $Q = [a, b] \times [c, d]$. Then*

$$\iint_Q f(x, y)\, dx\, dy = \int_a^b \Big(\int_c^d f(x, y)\, dy \Big) dx = \int_c^d \Big(\int_a^b f(x, y)\, dx \Big) dy.$$

The same holds for boxes in $\mathbb{R}^n$, integrating over one variable after the other in any order.

We do not give a complete proof, but the idea is simple. With the grid points x_i and y_j, a Riemann sum is a double sum $\sum_i \sum_j f(x_i, y_j)\, \Delta x\, \Delta y$. Summing over j first gives

$$\sum_i \Big(\sum_j f(x_i, y_j)\, \Delta y \Big) \Delta x \approx \sum_i \Big(\int_c^d f(x_i, y)\, dy \Big) \Delta x,$$

a Riemann sum of the iterated integral with y first; summing over i first gives the other order. Geometrically, we compute a volume by cutting it into slices and integrating the areas of the slices. For a region $D = \{(x, y) : a \le x \le b,\ g_1(x) \le y \le g_2(x)\}$ between the graphs of continuous functions $g_1 \le g_2$, Fubini's theorem gives

$$\iint_D f(x, y)\, dx\, dy = \int_a^b \Big(\int_{g_1(x)}^{g_2(x)} f(x, y)\, dy \Big) dx.$$

Example 3.14.3. 1. For the triangle $T = \{(x, y) : 0 \le y \le x \le 1\}$,

$$\iint_T xy\, dx\, dy = \int_0^1 \int_0^x xy\, dy\, dx = \int_0^1 \frac{x^3}{2}\, dx = \frac{1}{8}.$$

Describing T by $0 \le y \le 1$ and $y \le x \le 1$ gives the same result, $\int_0^1 \int_y^1 xy\, dx\, dy = \int_0^1 \frac{y(1-y^2)}{2}\, dy = \frac{1}{8}$.

2. (A probability.) Let X and Y be independent waiting times with the exponential densities $\lambda e^{-\lambda x}$ and $\mu e^{-\mu y}$ on $[0,\infty)$, so that (X,Y) has the joint density $\lambda\mu e^{-\lambda x-\mu y}$ (Lecture 4). The probability that Y ends first is the integral of the joint density over $\{0 \le y < x\}$:
$$\begin{aligned} P(Y<X) &= \int_0^\infty \lambda e^{-\lambda x}\Big(\int_0^x \mu e^{-\mu y}\,dy\Big)dx = \int_0^\infty \lambda e^{-\lambda x}(1-e^{-\mu x})\,dx \\ &= 1-\frac{\lambda}{\lambda+\mu} = \frac{\mu}{\lambda+\mu}. \end{aligned}$$

3. (The order can matter.) Let $f(x,y) = \frac{x^2-y^2}{(x^2+y^2)^2}$ on $(0,1]^2$. Since $\frac{\partial}{\partial y}\frac{y}{x^2+y^2} = f(x,y)$, we get $\int_0^1 f(x,y)\,dy = \frac{1}{1+x^2}$ and $\int_0^1\int_0^1 f(x,y)\,dy\,dx = \frac{\pi}{4}$. Since $f(y,x) = -f(x,y)$, the other order gives $-\frac{\pi}{4}$. Fubini's theorem does not apply, because f is unbounded near the origin; in fact, $\iint |f| = \infty$.

For unbounded functions or regions, Fubini's theorem remains true if $f \ge 0$ or if $\int |f| < \infty$ (the Fubini–Tonelli theorem, Theorem B.6.1), which covers the integrals of probability theory.

3.14.2 Change of variables

Theorem 3.14.4 (Change of variables)**.** *Let $\Phi : U \to V$ be a bijection between open sets $U, V \subseteq \mathbb{R}^n$, such that Φ and Φ^{-1} are continuously differentiable. Then for every integrable function g on V,*
$$\int_V g(\mathbf{y})\,d\mathbf{y} = \int_U g(\Phi(\mathbf{x}))\,|\det D\Phi(\mathbf{x})|\,d\mathbf{x}.$$

This is the formula of Remark 3.4.8, and it comes from the interpretation of $|\det D\Phi(\mathbf{x})|$ as the factor by which Φ scales small volumes (Section 3.4): the images $\Phi(Q_j)$ of small boxes Q_j in U cover V and have the volumes $|\det D\Phi(\boldsymbol{\xi}_j)|\,\mathrm{vol}(Q_j)$ approximately, so Riemann sums for the two sides are approximately equal. We do not make this precise.

Example 3.14.5 (The Gaussian integral in n dimensions)**.** Let Σ be symmetric positive definite. We show that
$$\int_{\mathbb{R}^n} \exp\big(-\tfrac{1}{2}\mathbf{x}^\top\Sigma^{-1}\mathbf{x}\big)\,d\mathbf{x} = (2\pi)^{n/2}\sqrt{\det\Sigma},$$
which gives the normalizing constant of the multivariate Gaussian distribution in Lecture 4. For $\Sigma = I$, the integrand is the product $e^{-x_1^2/2}\cdots e^{-x_n^2/2}$, so by Fubini's theorem the integral is the product of n one-dimensional integrals $\int_{-\infty}^{\infty} e^{-t^2/2}\,dt = \sqrt{2\pi}$ (Example 3.4.7, with the substitution $t = \sqrt{2}x$). In general, write $\Sigma = AA^\top$ with an invertible matrix A, for example $A = Q\Lambda^{1/2}$ from the spectral decomposition $\Sigma = Q\Lambda Q^\top$ (Lecture 2), where $\Lambda^{1/2}$ is the diagonal matrix of the square roots of the positive eigenvalues. Substitute $\mathbf{x} = A\mathbf{z}$. Then $\mathbf{x}^\top\Sigma^{-1}\mathbf{x} = \mathbf{z}^\top A^\top A^{-\top}A^{-1}A\mathbf{z} = \|\mathbf{z}\|^2$ and $|\det A| = \sqrt{\det\Sigma}$, so the integral is $\sqrt{\det\Sigma}\int_{\mathbb{R}^n} e^{-\|\mathbf{z}\|^2/2}\,d\mathbf{z} = (2\pi)^{n/2}\sqrt{\det\Sigma}$.

3.14.3 Differentiating under the integral sign

Proposition 3.14.6 (Leibniz rule)**.** *Let $f(x,\theta)$ and $\frac{\partial f}{\partial\theta}(x,\theta)$ be continuous for $x \in [a,b]$ and θ in an open interval I. Then*
$$\frac{d}{d\theta}\int_a^b f(x,\theta)\,dx = \int_a^b \frac{\partial f}{\partial\theta}(x,\theta)\,dx \qquad \text{for all } \theta \in I.$$

Proof. The function $g(\theta) = \int_a^b \frac{\partial f}{\partial\theta}(x,\theta)\,dx$ is continuous, since $\frac{\partial f}{\partial\theta}$ is uniformly continuous on closed and bounded sets (Corollary A.3.9). Fix $\theta_0 \in I$. By Fubini's theorem and the fundamental theorem of calculus,
$$\int_{\theta_0}^{\theta} g(s)\,ds = \int_a^b\int_{\theta_0}^{\theta}\frac{\partial f}{\partial\theta}(x,s)\,ds\,dx = \int_a^b f(x,\theta)\,dx - \int_a^b f(x,\theta_0)\,dx.$$

Differentiating with respect to θ gives the claim, again by the fundamental theorem of calculus. □

The same holds for integrals over boxes in $\mathbb{R}^n$ and for several parameters, and, under an additional condition (for example, if $|\frac{\partial f}{\partial \theta}|$ is bounded by an integrable function that does not depend on θ), for integrals over unbounded regions such as $\mathbb{R}^n$ (Example B.4.3 proves this with the dominated convergence theorem). In machine learning, we often need the gradient of an expectation $\int g(\mathbf{x}) p_{\boldsymbol{\theta}}(\mathbf{x})\, d\mathbf{x}$ with respect to the parameters $\boldsymbol{\theta}$ of a distribution, for example of a policy in reinforcement learning. Differentiating under the integral sign and using $\nabla_{\boldsymbol{\theta}} p_{\boldsymbol{\theta}} = p_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \log p_{\boldsymbol{\theta}}$ gives

$$\nabla_{\boldsymbol{\theta}} \int g(\mathbf{x}) p_{\boldsymbol{\theta}}(\mathbf{x})\, d\mathbf{x} = \int g(\mathbf{x})\, \nabla_{\boldsymbol{\theta}} \log p_{\boldsymbol{\theta}}(\mathbf{x})\, p_{\boldsymbol{\theta}}(\mathbf{x})\, d\mathbf{x},$$

again an expectation with respect to $p_{\boldsymbol{\theta}}$, which can be estimated by averaging over samples. This is the score function (or REINFORCE) estimator of policy gradient methods.

3.14.4 Line integrals and gradient fields

Definition 3.14.7 (Line integral)**.** Let $\gamma : [a, b] \to D$ be a continuously differentiable curve, and let F be a continuous vector field on D. The line integral of F along γ is

$$\int_\gamma F \cdot d\mathbf{x} = \int_a^b F(\gamma(t))^\top \gamma'(t)\, dt.$$

If F is a force, the line integral is the work done by it along the path. By the substitution rule, the line integral does not change if we reparametrize the curve in the same direction, and it changes its sign if we reverse the direction. For gradient fields, it only depends on the endpoints.

Theorem 3.14.8 (Fundamental theorem for line integrals)**.** *If f is continuously differentiable on D, then for every continuously differentiable curve $\gamma : [a, b] \to D$,*

$$\int_\gamma \nabla f \cdot d\mathbf{x} = f(\gamma(b)) - f(\gamma(a)).$$

In particular, the line integral of a gradient field along a closed curve ($\gamma(a) = \gamma(b)$) is 0*.*

Proof. By the chain rule (Corollary 3.5.2), $\frac{d}{dt} f(\gamma(t)) = \nabla f(\gamma(t))^\top \gamma'(t)$; now integrate from a to b. □

Example 3.14.9 (Integrated gradients)**.** To explain the prediction $f(\mathbf{x})$ of a model, the method of integrated gradients compares $\mathbf{x}$ with a baseline input $\mathbf{x}'$, such as a black image, and assigns to the i-th input feature the attribution

$$A_i = (x_i - x_i') \int_0^1 \frac{\partial f}{\partial x_i}(\mathbf{x}' + t(\mathbf{x} - \mathbf{x}'))\, dt.$$

The sum of the attributions is the line integral of ∇f along the segment $\gamma(t) = \mathbf{x}' + t(\mathbf{x} - \mathbf{x}')$, so $A_1 + \cdots + A_n = f(\mathbf{x}) - f(\mathbf{x}')$ by Theorem 3.14.8: the attributions account exactly for the change of the output. For example, for $f(\mathbf{x}) = x_1 x_2$, $\mathbf{x} = (2, 3)^\top$ and $\mathbf{x}' = \mathbf{0}$, we get $A_1 = 2\int_0^1 3t\, dt = 3$ and $A_2 = 3\int_0^1 2t\, dt = 3$, which add up to $f(\mathbf{x}) = 6$. The simpler attributions $x_i \frac{\partial f}{\partial x_i}(\mathbf{x})$ ("gradient times input") are 6 and 6, which add up to 12.

Conversely, if the line integrals of F only depend on the endpoints, then F is a gradient field: fix a point $\mathbf{x}_0$, and let $f(\mathbf{x})$ be the line integral of F along any curve from $\mathbf{x}_0$ to $\mathbf{x}$. On $\mathbb{R}^n$, a symmetric Jacobian is enough.

Theorem 3.14.10 (Poincaré lemma)**.** *Let $F : \mathbb{R}^n \to \mathbb{R}^n$ be continuously differentiable, with a symmetric Jacobian $DF(\mathbf{x})$ for every $\mathbf{x}$. Then $F = \nabla f$ for the function*

$$f(\mathbf{x}) = \int_0^1 F(t\mathbf{x})^\top \mathbf{x}\, dt,$$

the line integral of F along the segment from $\mathbf{0}$ to $\mathbf{x}$.

Proof. By the Leibniz rule and the product rule, and since $\frac{\partial F_j}{\partial x_i} = \frac{\partial F_i}{\partial x_j}$,

$$\frac{\partial f}{\partial x_i}(\mathbf{x}) = \int_0^1 \Big(F_i(t\mathbf{x}) + \sum_{j=1}^n t\, \frac{\partial F_j}{\partial x_i}(t\mathbf{x})\, x_j \Big)\, dt = \int_0^1 \Big(F_i(t\mathbf{x}) + t \sum_{j=1}^n \frac{\partial F_i}{\partial x_j}(t\mathbf{x})\, x_j \Big)\, dt.$$

By the chain rule, the integrand is $\frac{d}{dt}\big(tF_i(t\mathbf{x})\big)$, so $\frac{\partial f}{\partial x_i}(\mathbf{x}) = 1 \cdot F_i(\mathbf{x}) - 0 = F_i(\mathbf{x})$. □

Example 3.14.11 (A field with zero curl that is not a gradient field)**.** On $D = \mathbb{R}^2 \setminus \{\mathbf{0}\}$, let $F(x,y) = \frac{1}{x^2+y^2}(-y, x)^\top$. A short computation gives $\frac{\partial F_2}{\partial x} = \frac{y^2-x^2}{(x^2+y^2)^2} = \frac{\partial F_1}{\partial y}$, so $\operatorname{curl} F = 0$. But along the unit circle $\gamma(t) = (\cos t, \sin t)^\top$, $0 \le t \le 2\pi$, we have $F(\gamma(t)) = (-\sin t, \cos t)^\top = \gamma'(t)$, so

$$\int_\gamma F \cdot d\mathbf{x} = \int_0^{2\pi} (\sin^2 t + \cos^2 t)\, dt = 2\pi \neq 0,$$

and F is not a gradient field on D. Near each point, F is the gradient of the polar angle, but the angle cannot be defined continuously along a closed curve around the origin. The Poincaré lemma does not apply because D has a hole: whether vector fields with zero curl are gradient fields depends on the topology of the domain.

3.14.5 The divergence theorem

Let $V \subseteq \mathbb{R}^n$ be a bounded region whose boundary ∂V is smooth, such as a ball, or a box (whose boundary is smooth except at the edges), and let $\mathbf{n}(\mathbf{x})$ be the outward unit normal vector at $\mathbf{x} \in \partial V$. We also need integrals $\int_{\partial V} g\, dS$ over the boundary, with respect to its $(n-1)$-dimensional volume (length for $n = 2$, area for $n = 3$). We do not define them in general. For $n = 2$ and a curve $\gamma : [a,b] \to \mathbb{R}^2$ that traverses ∂V once, $\int_{\partial V} g\, dS = \int_a^b g(\gamma(t))\|\gamma'(t)\|\, dt$; and the faces of a box are boxes of dimension $n-1$.

Theorem 3.14.12 (Divergence theorem)**.** *For every continuously differentiable vector field F on an open set that contains V and ∂V,*

$$\int_V \operatorname{div} F(\mathbf{x})\, d\mathbf{x} = \int_{\partial V} F^\top \mathbf{n}\, dS.$$

The right side is the flux of F through the boundary: if F is the velocity field of a fluid, it is the net amount of fluid that flows out of V per unit time. So the theorem says that the total strength of the sources inside V equals the net outflow through its boundary, which shows again that the divergence is the outflow per unit volume. For $n = 1$ and $V = [a,b]$, with the outward normals $+1$ at b and -1 at a, it is the fundamental theorem of calculus $\int_a^b F'(x)\, dx = F(b) - F(a)$.

Proof for a box. Let $V = [a_1, b_1] \times \cdots \times [a_n, b_n]$, and consider the term $\frac{\partial F_1}{\partial x_1}$ of $\operatorname{div} F$. Write $\mathbf{x} = (x_1, \tilde{\mathbf{x}})$ with $\tilde{\mathbf{x}}$ in the box $\tilde{Q} = [a_2, b_2] \times \cdots \times [a_n, b_n]$. By Fubini's theorem and the fundamental theorem of calculus,

$$\int_V \frac{\partial F_1}{\partial x_1}\, d\mathbf{x} = \int_{\tilde{Q}} \Big(\int_{a_1}^{b_1} \frac{\partial F_1}{\partial x_1}(x_1, \tilde{\mathbf{x}})\, dx_1 \Big) d\tilde{\mathbf{x}} = \int_{\tilde{Q}} F_1(b_1, \tilde{\mathbf{x}})\, d\tilde{\mathbf{x}} - \int_{\tilde{Q}} F_1(a_1, \tilde{\mathbf{x}})\, d\tilde{\mathbf{x}}.$$

This is the flux of the vector field $F_1\mathbf{e}_1$ through the boundary of V: through the face $x_1 = b_1$ with the outward normal $\mathbf{e}_1$, through the face $x_1 = a_1$ with the outward normal $-\mathbf{e}_1$, and nothing through the other faces, whose normals $\pm\mathbf{e}_j$ ($j \neq 1$) are orthogonal to $\mathbf{e}_1$. Adding the same identities for $F_2\mathbf{e}_2, \ldots, F_n\mathbf{e}_n$ proves the theorem for V. □

For a general region, one approximates V by many small boxes. When we add the identities for all boxes, the fluxes through the faces shared by two neighboring boxes cancel, since the outward normal of one box is the inward normal of the other, and only the fluxes through the outer boundary remain. Making this precise requires more care, and we omit it.

Example 3.14.13. For $F(\mathbf{x}) = \mathbf{x}$, $\operatorname{div} F = n$, and on the sphere of radius R, $\mathbf{n} = \frac{\mathbf{x}}{R}$ and $F^\top \mathbf{n} = R$. So the divergence theorem for the ball B_R of radius R says that $n \operatorname{vol}(B_R) = R \operatorname{area}(\partial B_R)$: for $n = 2$, $2\pi R^2 = R \cdot 2\pi R$, and for $n = 3$, $3 \cdot \frac{4}{3}\pi R^3 = R \cdot 4\pi R^2$.

In the plane, the divergence theorem has a second form. If γ traverses ∂V counterclockwise (with V on its left), then the unit tangent vector is $\boldsymbol{\tau} = \frac{\gamma'}{\|\gamma'\|}$, and the outward normal is $\boldsymbol{\tau}$ rotated clockwise by 90°, $\mathbf{n} = (\tau_2, -\tau_1)^\top$. For the vector field $G = (F_2, -F_1)^\top$, we have $\operatorname{div} G = \operatorname{curl} F$ and $G^\top \mathbf{n} = F^\top \boldsymbol{\tau}$, so the divergence theorem for G gives the following.

Theorem 3.14.14 (Green)**.** *Let $V \subseteq \mathbb{R}^2$ be a bounded region whose boundary is a closed curve γ, traversed counterclockwise. Then for every continuously differentiable vector field F on an open set that contains V and ∂V,*

$$\int_\gamma F \cdot d\mathbf{x} = \iint_V \operatorname{curl} F \, dx \, dy.$$

The left side is the circulation of F around V, so the curl is the circulation per unit area. For example, $F = \frac{1}{2}(-y, x)^\top$ has $\operatorname{curl} F = 1$, so the area of V is the line integral of F along its boundary. (The vortex field of Example 3.14.11 is not differentiable at the origin, so Green's theorem does not apply to it on the unit disk.)

Remark 3.14.15 (Stokes' theorem). In three dimensions, Stokes' theorem says that $\int_{\partial S} F \cdot d\mathbf{x} = \int_S (\operatorname{curl} F)^\top \mathbf{n} \, dS$ for a surface S with the boundary curve ∂S; for a flat surface, it is Green's theorem. The fundamental theorem of calculus, the fundamental theorem for line integrals, Green's theorem, the divergence theorem and Stokes' theorem are all special cases of one theorem about differential forms on manifolds, the general Stokes theorem: the integral of a derivative over a region equals an integral over its boundary. Differential forms and this general Stokes theorem are treated in Appendix A on topology, differential topology and functional analysis. For machine learning, the most important of these theorems is the divergence theorem, through its consequence, integration by parts.

3.14.6 Integration by parts and its applications

Corollary 3.14.16 (Integration by parts)**.** *Let $g : \mathbb{R}^n \to \mathbb{R}$ and $F : \mathbb{R}^n \to \mathbb{R}^n$ be continuously differentiable. Then*

$$\int_V g \operatorname{div} F \, d\mathbf{x} = \int_{\partial V} g F^\top \mathbf{n} \, dS - \int_V \nabla g^\top F \, d\mathbf{x}.$$

If gF vanishes outside a bounded set, or decays so fast that its flux through spheres of radius R tends to 0 as $R \to \infty$, then $\int_{\mathbb{R}^n} g \operatorname{div} F \, d\mathbf{x} = -\int_{\mathbb{R}^n} \nabla g^\top F \, d\mathbf{x}$.

Proof. By the product rule, $\frac{\partial}{\partial x_i}(gF_i) = \frac{\partial g}{\partial x_i} F_i + g \frac{\partial F_i}{\partial x_i}$, and summing over i gives $\operatorname{div}(gF) = \nabla g^\top F + g \operatorname{div} F$. Now apply the divergence theorem to gF. For the second statement, take for V balls with growing radii. □

For $n = 1$, this is the familiar rule $\int_a^b gF' \, dx = [gF]_a^b - \int_a^b g'F \, dx$. We give an application to machine learning; in Section 3.16, the same rule describes how probability densities move under the flow of a differential equation.

Example 3.14.17 (Score matching)**.** We want to fit a model $\mathbf{s}_{\boldsymbol{\theta}}$ of the score $\nabla \log p$ of an unknown data density p, given samples from p (Example 3.13.2). The natural loss

$$J(\boldsymbol{\theta}) = \frac{1}{2} \int p(\mathbf{x}) \|\mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}) - \nabla \log p(\mathbf{x})\|^2 \, d\mathbf{x}$$

seems useless, since it contains the unknown score. Expanding the square gives $\frac{1}{2}\int p\|\mathbf{s}_{\boldsymbol{\theta}}\|^2 \, d\mathbf{x} - \int p\, \mathbf{s}_{\boldsymbol{\theta}}^\top \nabla \log p \, d\mathbf{x} + C$, where C does not depend on $\boldsymbol{\theta}$. Since $p\nabla \log p = \nabla p$, integration by parts (with $g = p$ and $F = \mathbf{s}_{\boldsymbol{\theta}}$, assuming that $p\,\mathbf{s}_{\boldsymbol{\theta}}$ decays fast enough) turns the middle term into

$$-\int \mathbf{s}_{\boldsymbol{\theta}}^\top \nabla p \, d\mathbf{x} = \int p \operatorname{div} \mathbf{s}_{\boldsymbol{\theta}} \, d\mathbf{x}.$$

So $J(\boldsymbol{\theta}) = \int p(\mathbf{x}) \big(\frac{1}{2}\|\mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x})\|^2 + \operatorname{div} \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x})\big) \, d\mathbf{x} + C$: an average over the data distribution, in which the unknown score no longer appears. It is estimated by the average of $\frac{1}{2}\|\mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}_i)\|^2 + \operatorname{div} \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}_i)$ over the training data $\mathbf{x}_1, \dots, \mathbf{x}_N$, with Hutchinson's estimator for the divergence (Remark 3.13.9). This is Hyvärinen's score matching; diffusion models use a variant of it, denoising score matching.

3.15 Ordinary Differential Equations

A differential equation describes how a state changes continuously in time: at every moment, the rate of change of the state is a given function of the state. This function is a vector field, and the state moves along its arrows, with the velocity given by the arrow at its current position. Differential equations appear in machine learning in the analysis of training dynamics, in neural networks with continuous depth, and, with randomness added, in diffusion models and sampling algorithms.

Definition 3.15.1 (Ordinary differential equation)**.** Let $F : \mathbb{R}^n \to \mathbb{R}^n$ be a vector field (Definition 3.13.1). A solution of the ordinary differential equation (ODE) $\mathbf{x}' = F(\mathbf{x})$ with the initial value $\mathbf{x}(0) = \mathbf{x}_0$ is a differentiable curve $\mathbf{x}(t)$ with $\mathbf{x}(0) = \mathbf{x}_0$ and $\mathbf{x}'(t) = F(\mathbf{x}(t))$ for all t.

Geometrically, a solution is a curve that is tangent to the vector field at each of its points: at the point $\mathbf{x}$, it moves with the velocity $F(\mathbf{x})$.

Theorem 3.15.2 (Existence and uniqueness)**.** *If F is Lipschitz continuous, i.e., $\|F(\mathbf{x}) - F(\mathbf{y})\| \le L\|\mathbf{x}-\mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y}$, then for every $\mathbf{x}_0$, the ODE $\mathbf{x}' = F(\mathbf{x})$ has exactly one solution with $\mathbf{x}(0) = \mathbf{x}_0$, defined for all $t \in \mathbb{R}$.*

This is the Picard–Lindelöf theorem. The idea of the proof is that $\mathbf{x}$ solves the problem if and only if $\mathbf{x}(t) = \mathbf{x}_0 + \int_0^t F(\mathbf{x}(s))\, ds$ for all t, so a solution is a fixed point of the map that sends a curve $\mathbf{x}$ to the right-hand side; with respect to a suitable norm, this map is a contraction, and the Banach fixed-point theorem applies. Theorem A.12.4 carries out this proof.

Example 3.15.3.

1. The solution of $x' = ax$ with $x(0) = x_0$ is $x(t) = e^{at}x_0$, since $\frac{d}{dt}e^{at}x_0 = ae^{at}x_0$. It grows exponentially if $a > 0$ and decays to 0 if $a < 0$.
2. The logistic equation $x' = x(1-x)$ models growth with a limited capacity. By Example 3.2.4, the sigmoid function satisfies $\sigma' = \sigma(1-\sigma)$, so $x(t) = \sigma(t + c)$ is a solution, with c chosen such that $\sigma(c) = x_0$. For $0 < x_0 < 1$, the solution increases and tends to 1.

3.15.1 Linear systems

For a matrix $A \in \mathbb{R}^{n\times n}$, the linear ODE $\mathbf{x}' = A\mathbf{x}$ couples the coordinates, and eigenvectors decouple them (Lecture 1).

Proposition 3.15.4. *Let $A = V\Lambda V^{-1}$ be diagonalizable, with eigenvalues $\lambda_1, \dots, \lambda_n$ and eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ (the columns of V), and write $\mathbf{x}_0 = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$. Then the solution of $\mathbf{x}' = A\mathbf{x}$ with $\mathbf{x}(0) = \mathbf{x}_0$ is*

$$\mathbf{x}(t) = c_1 e^{\lambda_1 t}\mathbf{v}_1 + \cdots + c_n e^{\lambda_n t}\mathbf{v}_n.$$

Proof. Each term satisfies $\frac{d}{dt} c_i e^{\lambda_i t}\mathbf{v}_i = \lambda_i c_i e^{\lambda_i t}\mathbf{v}_i = A(c_i e^{\lambda_i t}\mathbf{v}_i)$, so the sum is a solution, and it has the right initial value. It is the only one by Theorem 3.15.2, since $\|A\mathbf{x} - A\mathbf{y}\| \le \|A\|_2\|\mathbf{x} - \mathbf{y}\|$. (Equivalently, the coordinates $\mathbf{y} = V^{-1}\mathbf{x}$ satisfy the decoupled equations $y_i' = \lambda_i y_i$.) □

More generally, the solution is $\mathbf{x}(t) = e^{At}\mathbf{x}_0$ with the matrix exponential $e^{At} = \sum_{k=0}^{\infty} \frac{(At)^k}{k!}$, which converges for every matrix and satisfies $\frac{d}{dt}e^{At} = Ae^{At}$; for diagonalizable A, $e^{At} = Ve^{\Lambda t}V^{-1}$ with $e^{\Lambda t} = \operatorname{diag}(e^{\lambda_1 t}, \dots, e^{\lambda_n t})$. So the eigenvalues determine the long-term behavior: if all (complex) eigenvalues have negative real parts, every solution tends to $\mathbf{0}$, and if some eigenvalue has a positive real part, solutions starting in the direction of its eigenvector grow exponentially. This is the continuous-time analogue of the powers P^t of a matrix, which govern gradient descent (Theorem 3.10.7) and Markov chains (Lecture 4).

Example 3.15.5.

1. Let $A = \begin{pmatrix} -2 & 1 \\ 1 & -2 \end{pmatrix}$, with the eigenvalues -1 and -3 and the eigenvectors $\mathbf{v}_1 = (1,1)^\top$ and $\mathbf{v}_2 = (1,-1)^\top$. For $\mathbf{x}_0 = (2,0)^\top = \mathbf{v}_1 + \mathbf{v}_2$, the solution is $\mathbf{x}(t) = e^{-t}(1,1)^\top + e^{-3t}(1,-1)^\top$; for example, $\mathbf{x}(1) \approx (0.418, 0.318)^\top$. The component along $\mathbf{v}_2$ decays faster, so all solutions approach the origin along the slow direction $\mathbf{v}_1$ (Figure 3.12).

2. Let $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, with the complex eigenvalues $\pm i$. Then $e^{At} = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}$ is the rotation by the angle t (Lecture 2): one checks that this matrix is I at $t = 0$ and that its derivative is A times itself. The solutions move on circles. Indeed, for every skew-symmetric A, $\frac{d}{dt}\|\mathbf{x}\|^2 = 2\mathbf{x}^\top A\mathbf{x} = 0$.




Figure 3.12: The vector field $A\mathbf{x}$ (gray) and solutions of $\mathbf{x}' = A\mathbf{x}$ (blue) for $A = \left(\begin{smallmatrix} -2 & 1 \\ 1 & -2 \end{smallmatrix}\right)$, starting on the circle of radius 2. The component along the eigenvector $\mathbf{v}_2$ (eigenvalue -3) dies out quickly, and the solutions approach the origin along $\mathbf{v}_1$ (eigenvalue -1).

3.15.2 Gradient flow and Euler's method

Definition 3.15.6 (Gradient flow)**.** The gradient flow of a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ is the ODE $\mathbf{x}' = -\nabla f(\mathbf{x})$.

Proposition 3.15.7. *Along every solution of the gradient flow, $\frac{d}{dt}f(\mathbf{x}(t)) = -\|\nabla f(\mathbf{x}(t))\|^2 \le 0$: the value of f decreases, and strictly unless $\mathbf{x}(t)$ is a critical point.*

Proof. By the chain rule (Corollary 3.5.2), $\frac{d}{dt}f(\mathbf{x}(t)) = \nabla f(\mathbf{x}(t))^\top \mathbf{x}'(t) = -\|\nabla f(\mathbf{x}(t))\|^2$. □

For the quadratic function $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$ with symmetric positive definite A, the error $\mathbf{e} = \mathbf{x} - \mathbf{x}^*$ satisfies $\mathbf{e}' = -A\mathbf{e}$, so by Proposition 3.15.4, $\mathbf{e}(t) = \sum_i c_i e^{-\lambda_i t}\mathbf{q}_i$ in an orthonormal eigenbasis of A. The gradient flow converges to the minimum for every positive definite A, and the slowest direction decays like $e^{-\lambda_{\min} t}$.

To solve an ODE $\mathbf{x}' = F(\mathbf{x})$ numerically, Euler's method follows the tangent for a short time h:

$$\mathbf{x}_{k+1} = \mathbf{x}_k + hF(\mathbf{x}_k), \qquad \text{so that } \mathbf{x}_k \approx \mathbf{x}(kh).$$

For a smooth F, its error on a fixed time interval is proportional to h. For example, for $x' = -x$ with $x(0) = 1$, Euler's method gives $x_k = (1-h)^k$, so it approximates $x(1) = e^{-1} \approx 0.368$ by 0.25 for $h = 0.5$, and by $0.9^{10} \approx 0.349$ for $h = 0.1$.

For the gradient flow, Euler's method is $\mathbf{x}_{k+1} = \mathbf{x}_k - h\nabla f(\mathbf{x}_k)$: gradient descent is Euler's method for the gradient flow, with the step size $\eta = h$. This explains the results of Section 3.10. For the quadratic function, the flow converges for every positive definite A, but Euler's method multiplies the error in the direction $\mathbf{q}_i$ by $1 - h\lambda_i$, which is stable only if $h < \frac{2}{\lambda_{\max}}$. The directions of large curvature force small steps, while the directions of small curvature need the time $\frac{1}{\lambda_{\min}}$, so about $\kappa = \frac{\lambda_{\max}}{\lambda_{\min}}$ steps are needed. Such ODEs are called stiff, and the zig-zagging of Figure 3.8 is an instability of the discretization, not of the flow.

Remark 3.15.8 (ODEs in machine learning).

1. The layers of a residual network compute $\mathbf{x}_{l+1} = \mathbf{x}_l + f_l(\mathbf{x}_l)$, which is an Euler step of an ODE $\mathbf{x}' = f(\mathbf{x}, t)$ with $h = 1$. Neural ODEs replace the layers by an ODE solver, and compute gradients by solving an adjoint ODE backwards in time, which is reverse-mode differentiation (Section 3.7) in continuous time.

2. Gradient descent with momentum is a discretization of the second-order ODE $\mathbf{x}'' + \gamma\mathbf{x}' + \nabla f(\mathbf{x}) = \mathbf{0}$ of a heavy ball with friction rolling in the landscape f; with $\mathbf{v} = \mathbf{x}'$, it is a first-order system for $(\mathbf{x}, \mathbf{v})$.

3. Adding random noise to an ODE gives a stochastic differential equation. Langevin dynamics and diffusion models, which sample from probability distributions in this way, are discussed in Lecture 4.

3.16 Flows, Volumes and Densities

Instead of following a single solution, we can move all points at once. Let $\varphi_t(\mathbf{x}_0)$ be the solution of $\mathbf{x}' = F(\mathbf{x})$ with $\mathbf{x}(0) = \mathbf{x}_0$ at the time t. The map φ_t, the flow of F, moves every point along the vector field for the time t; for example, the flow of $F(\mathbf{x}) = A\mathbf{x}$ is $\varphi_t(\mathbf{x}) = e^{At}\mathbf{x}$. In Section 3.13, we saw that the divergence $\operatorname{div} F = \operatorname{tr} DF$ is the net outflow of F per unit volume. For the flow, this means that the divergence is the rate at which the flow expands volumes. For linear vector fields, the effect of the flow on volumes can be computed exactly.

Proposition 3.16.1 (Liouville's formula)**.** *For every diagonalizable $A \in \mathbb{R}^{n\times n}$, $\det e^{At} = e^{t\operatorname{tr} A}$. So the flow $\varphi_t(\mathbf{x}) = e^{At}\mathbf{x}$ of $F(\mathbf{x}) = A\mathbf{x}$ multiplies all volumes by $e^{t\operatorname{div} F}$.*

Proof. With $A = V\Lambda V^{-1}$, $\det e^{At} = \det(V)\det(e^{\Lambda t})\det(V)^{-1} = e^{\lambda_1 t}\cdots e^{\lambda_n t} = e^{t(\lambda_1+\cdots+\lambda_n)} = e^{t\operatorname{tr} A}$, and the linear map e^{At} multiplies volumes by $|\det e^{At}|$ (Lecture 1). □

(The formula holds for every square matrix; Exercise 38 checks it for a matrix that is not diagonalizable.) For a general vector field and a small t, the flow $\varphi_t(\mathbf{y}) \approx \mathbf{y} + tF(\mathbf{y})$ has approximately the Jacobian $I + tDF(\mathbf{y})$, whose determinant is $1 + t \operatorname{div} F(\mathbf{y})$ up to terms of order t^2 (Lemma 3.6.3). So a small region near $\mathbf{x}$ with the volume v has, after the short time t, approximately the volume $(1 + t \operatorname{div} F(\mathbf{x}))v$:

$$\operatorname{div} F(\mathbf{x}) \text{ is the rate at which the flow of } F \text{ expands volumes near } \mathbf{x}.$$

In particular, the flow of a vector field with $\operatorname{div} F = 0$ preserves volumes.

Example 3.16.2.

1. $F(\mathbf{x}) = \mathbf{x}$ has $\operatorname{div} F = n$. Its flow $\varphi_t(\mathbf{x}) = e^t\mathbf{x}$ multiplies lengths by e^t and volumes by e^{nt}.
2. The rotation field $(-y, x)^\top$ has $\operatorname{div} F = 0$, and its flow, the rotation by the angle t (Example 3.15.5), preserves areas. It is the continuous-time version of gradient descent–ascent for the game $\min_x \max_y xy$ (Remark 3.13.9): the flow circles around the equilibrium forever, and Euler's method, i.e., gradient descent–ascent with a constant step size, spirals outwards (Exercise 37).
3. For the field $A\mathbf{x}$ of Figure 3.12, $\operatorname{div} F = \operatorname{tr} A = -4$, so the flow shrinks areas by the factor e^{-4t}, in agreement with the eigenvalues -1 and -3.
4. The gradient flow $F = -\nabla f$ has $\operatorname{div} F = -\Delta f = -\operatorname{tr} \nabla^2 f$. Near a strict local minimum, where the Hessian is positive definite, the gradient flow compresses volumes, and nearby solutions are drawn together.

3.16.1 Transport of densities

If the initial value of an ODE is random, then so is the solution at every time t, and the divergence also tells us how its probability density changes.

Proposition 3.16.3 (Continuity equation)**.** *Let the initial value of the ODE $\mathbf{x}' = F(\mathbf{x})$ be random, with the density p_0, and let p_t be the density of $\mathbf{x}(t)$. Then, if all functions are smooth enough,*

$$\frac{\partial p_t}{\partial t} = -\operatorname{div}(p_t F), \qquad \text{and along every solution,} \qquad \frac{d}{dt} \log p_t(\mathbf{x}(t)) = -\operatorname{div} F(\mathbf{x}(t)).$$

Sketch of proof. Probability is neither created nor destroyed; it moves with the velocity F. So for a fixed region V, the probability $\int_V p_t \, d\mathbf{x}$ of V changes by the flux of $p_t F$ out of V, and by the divergence theorem (Theorem 3.14.12),

$$\frac{d}{dt} \int_V p_t \, d\mathbf{x} = -\int_{\partial V} p_t F^\top \mathbf{n} \, dS = -\int_V \operatorname{div}(p_t F) \, d\mathbf{x}.$$

Differentiating under the integral sign on the left, the integrals of $\frac{\partial p_t}{\partial t}$ and of $-\operatorname{div}(p_t F)$ agree on every region V, so the continuous functions agree. For the second formula, the chain rule and $\operatorname{div}(p_t F) = \nabla p_t^\top F + p_t \operatorname{div} F$ give

$$\frac{d}{dt} p_t(\mathbf{x}(t)) = \frac{\partial p_t}{\partial t}(\mathbf{x}(t)) + \nabla p_t(\mathbf{x}(t))^\top F(\mathbf{x}(t)) = -p_t(\mathbf{x}(t)) \operatorname{div} F(\mathbf{x}(t)),$$

and we divide by $p_t(\mathbf{x}(t))$. □

The second formula is the continuous-time version of the change of variables formula of Remark 3.4.8: where the flow expands volumes ($\operatorname{div} F > 0$), the density along the solution decreases, and conversely. Continuous normalizing flows, which are generative models, transform a Gaussian

distribution by the flow of an ODE $\mathbf{x}' = F(\mathbf{x})$ whose vector field is a neural network, and compute densities by

$$\log p_T(\mathbf{x}(T)) = \log p_0(\mathbf{x}(0)) - \int_0^T \operatorname{div} F(\mathbf{x}(t))\, dt,$$

which replaces the determinant of a Jacobian by the integral of a trace, estimated with Hutchinson's estimator (Remark 3.13.9). If noise is added to the ODE, the continuity equation becomes the Fokker–Planck equation (Lecture 4), which explains why Langevin dynamics samples from a given distribution.

3.17 Key Takeaways

- A function $f : D \to \mathbb{R}^m$ on an open set $D \subseteq \mathbb{R}^n$ is differentiable at $\mathbf{x}$ if it is approximated by a linear map near $\mathbf{x}$: $f(\mathbf{x}+\mathbf{h}) = f(\mathbf{x}) + Df(\mathbf{x})\mathbf{h} + o(\|\mathbf{h}\|)$. The Jacobian $Df(\mathbf{x})$ is the $m \times n$ matrix of partial derivatives $\frac{\partial f_i}{\partial x_j}$; its rows belong to the outputs, its columns to the inputs. Existence of the partial derivatives alone is not enough, but continuous partial derivatives imply differentiability.

- For $f : \mathbb{R}^n \to \mathbb{R}$, the derivative is a row vector (a linear functional), and the gradient $\nabla f(\mathbf{x}) = Df(\mathbf{x})^\top$ is the vector that represents it with the dot product. The directional derivative is $\nabla f(\mathbf{x})^\top \mathbf{v}$; by Cauchy–Schwarz, the gradient points in the direction of steepest ascent, and it is orthogonal to the level sets.

- Chain rule: $D(f \circ g)(\mathbf{x}) = Df(g(\mathbf{x}))\, Dg(\mathbf{x})$, the product of the Jacobians. For a scalar function, gradients are pulled back with transposed Jacobians: $\nabla(f \circ g)(\mathbf{x}) = Dg(\mathbf{x})^\top \nabla f(g(\mathbf{x}))$. Writing the sizes under each factor is a good check.

- Important gradients: $\nabla(\mathbf{a}^\top \mathbf{x}) = \mathbf{a}$, $\nabla(\mathbf{x}^\top A \mathbf{x}) = (A + A^\top)\mathbf{x}$, $\nabla\|A\mathbf{x} - \mathbf{b}\|^2 = 2A^\top(A\mathbf{x} - \mathbf{b})$; for logistic regression and softmax with cross-entropy, the gradient with respect to the scores is "prediction minus target". For a linear layer $\mathbf{y} = W\mathbf{x} + \mathbf{b}$, $\nabla_W L = \boldsymbol{\delta}\mathbf{x}^\top$ and $\nabla_{\mathbf{x}} L = W^\top \boldsymbol{\delta}$ with $\boldsymbol{\delta} = \nabla_{\mathbf{y}} L$.

- Gradients with respect to matrices have the shape of the matrix, and satisfy $f(X+H) \approx f(X) + \langle \nabla f(X), H \rangle$ with the Frobenius inner product. Examples: $\nabla \operatorname{tr}(AX) = A^\top$, $\nabla\|AX - B\|_F^2 = 2A^\top(AX - B)$, $\nabla \log|\det X| = X^{-\top}$.

- The Jacobian determinant $|\det Df(\mathbf{x})|$ is the local factor by which f scales areas and volumes; it appears in the change of variables formula, e.g., the factor r in polar coordinates.

- Automatic differentiation applies the chain rule to the elementary operations of a program. Forward mode computes Jacobian-vector products; reverse mode computes vector-Jacobian products and gives the whole gradient of a scalar function at a small multiple of the cost of evaluating it. Backpropagation is reverse mode for neural networks: $\boldsymbol{\delta}_l = \sigma'(\mathbf{z}_l) \odot (W_{l+1}^\top \boldsymbol{\delta}_{l+1})$.

- The Hessian $\nabla^2 f(\mathbf{x})$ is the symmetric matrix of second partial derivatives. Second-order Taylor: $f(\mathbf{x}+\mathbf{h}) \approx f(\mathbf{x}) + \nabla f(\mathbf{x})^\top \mathbf{h} + \frac{1}{2}\mathbf{h}^\top \nabla^2 f(\mathbf{x}) \mathbf{h}$.

- At a local minimum, $\nabla f = \mathbf{0}$. At a critical point, the eigenvectors of the Hessian are the principal directions, and its eigenvalues are the curvatures along them: in these coordinates, $f \approx f(\mathbf{x}^*) + \frac{1}{2}\sum_i \lambda_i y_i^2$. All eigenvalues positive: strict local minimum (a bowl); all negative: strict local maximum; both signs: saddle point; some zero: the test is inconclusive.

- Gradient descent $\mathbf{x}_{k+1} = \mathbf{x}_k - \eta \nabla f(\mathbf{x}_k)$ decreases an L-smooth function for $\eta \le \frac{1}{L}$ and finds approximate critical points. On quadratic functions, it converges for $\eta < \frac{2}{\lambda_{\max}}$, at best with the rate

$\frac{\kappa-1}{\kappa+1}$, where κ is the condition number: ill-conditioned problems zig-zag. Backtracking line search finds step sizes of order $\frac{1}{L}$ without knowing L. Newton's method $\mathbf{x}_{k+1} = \mathbf{x}_k - \nabla^2 f(\mathbf{x}_k)^{-1} \nabla f(\mathbf{x}_k)$ converges quadratically near a minimum, but is too expensive for large models; Gauss–Newton approximates the Hessian of $\frac{1}{2}\|\mathbf{r}\|^2$ by $J^\top J$, and Levenberg–Marquardt damps it to $J^\top J + \lambda I$. Hessian-vector products cost a few gradients. SGD uses cheap, noisy mini-batch gradients.

- A differentiable function is convex if and only if it lies above its tangent planes, and a twice differentiable one if and only if its Hessian is positive semidefinite everywhere. For convex functions, every critical point and every local minimum is a global minimum. Least squares, logistic regression and softmax regression have convex losses; neural networks do not.

- A μ-strongly convex function lies above the quadratic functions $f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y}-\mathbf{x}) + \frac{\mu}{2}\|\mathbf{y}-\mathbf{x}\|^2$ (for C^2 functions: all Hessian eigenvalues are at least μ). For a convex L-smooth function, gradient descent with $\eta = \frac{1}{L}$ reaches $f(\mathbf{x}_k) - f^* \le \frac{L\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{2k}$; with strong convexity, the error decreases geometrically, like $(1 - \frac{\mu}{L})^k$.

- Subgradients replace gradients for nonsmooth convex functions, and $\mathbf{x}^*$ minimizes f if and only if $\mathbf{0} \in \partial f(\mathbf{x}^*)$. The proximal gradient method $\mathbf{x}_{k+1} = \operatorname{prox}_{\eta h}(\mathbf{x}_k - \eta \nabla f(\mathbf{x}_k))$ minimizes $f + h$; for the lasso, the proximal step is soft thresholding, which sets small weights exactly to 0.

- Lagrange multipliers: at a constrained local minimum (with linearly independent constraint gradients), $\nabla f + \sum_i \lambda_i \nabla g_i = \mathbf{0}$. This gives eigenvectors for $\max_{\|\mathbf{x}\|=1} \mathbf{x}^\top A \mathbf{x}$ and the minimum norm solution $A^\top (A A^\top)^{-1} \mathbf{b}$. For inequality constraints, the KKT conditions add $\mu_i \ge 0$ and complementary slackness $\mu_i g_i(\mathbf{x}^*) = 0$, and for convex problems they are sufficient.

- The dual function $d(\boldsymbol{\mu}) = \inf_{\mathbf{x}} \mathcal{L}(\mathbf{x}, \boldsymbol{\mu})$ is concave, and $d(\boldsymbol{\mu}) \le p^*$ for all $\boldsymbol{\mu} \ge \mathbf{0}$ (weak duality). For convex problems in which the KKT conditions hold, the KKT multipliers maximize d and $d^* = p^*$ (strong duality). The dual of the support vector machine only depends on the inner products $\mathbf{x}_i^\top \mathbf{x}_j$, which leads to kernel methods. The convex conjugate $f^*(\mathbf{y}) = \sup_{\mathbf{x}} (\mathbf{y}^\top \mathbf{x} - f(\mathbf{x}))$ satisfies $f + f^* \ge \mathbf{x}^\top \mathbf{y}$ and $f^{**} = f$ for convex differentiable f; for linear constraints $A\mathbf{x} \le \mathbf{b}$, $d(\boldsymbol{\mu}) = -\mathbf{b}^\top \boldsymbol{\mu} - f^*(-A^\top \boldsymbol{\mu})$, so the dual of a linear program is a linear program, and the dual of a quadratic program has only sign constraints.

- A vector field $F : D \to \mathbb{R}^n$ has a square Jacobian, which splits into a symmetric part (stretching) and a skew-symmetric part (rotation). The divergence $\operatorname{div} F = \operatorname{tr} DF$ is the net outflow of F per unit volume, and the Laplacian is $\Delta f = \operatorname{div} \nabla f = \operatorname{tr} \nabla^2 f$. The curl ($\frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2}$ in the plane, $\nabla \times F$ in space) measures rotation. Gradient fields have zero curl, and on $\mathbb{R}^n$ the converse holds; the rotation field $(-y, x)$ of the game $\min_x \max_y xy$ is not the gradient of anything.

- Multiple integrals are computed as iterated one-dimensional integrals (Fubini), which needs $\int |f| < \infty$ for unbounded functions or regions. Under a change of variables, small volumes scale by $|\det D\Phi|$; for example, $\int e^{-\frac{1}{2}\mathbf{x}^\top \Sigma^{-1} \mathbf{x}} \, d\mathbf{x} = (2\pi)^{n/2} \sqrt{\det \Sigma}$. Line integrals of gradients depend only on the endpoints (integrated gradients). The divergence theorem $\int_V \operatorname{div} F \, d\mathbf{x} = \int_{\partial V} F^\top \mathbf{n} \, dS$ (Green's theorem in the plane) gives integration by parts, $\int g \operatorname{div} F = -\int \nabla g^\top F$, the basis of score matching.

- An ODE $\mathbf{x}' = F(\mathbf{x})$ with a Lipschitz continuous F has a unique solution for each initial value. Linear systems $\mathbf{x}' = A\mathbf{x}$ are solved with eigenvectors, $\mathbf{x}(t) = \sum_i c_i e^{\lambda_i t} \mathbf{v}_i = e^{At} \mathbf{x}_0$, and the signs of the real parts of the eigenvalues decide stability. Along the gradient flow $\mathbf{x}' = -\nabla f(\mathbf{x})$, f decreases; gradient descent is Euler's method for it, and its step size limit $\frac{2}{\lambda_{\max}}$ is the stability limit of Euler's method for stiff problems.

- The flow φ_t of a vector field F expands volumes at the rate $\operatorname{div} F$; for linear fields, $\det e^{At} = e^{t \operatorname{tr} A}$. A probability density moved by the flow satisfies the continuity equation $\frac{\partial p_t}{\partial t} = -\operatorname{div}(p_t F)$, and

along each solution, $\frac{d}{dt}\log p_t(\mathbf{x}(t)) = -\operatorname{div} F(\mathbf{x}(t))$, which continuous normalizing flows use instead of a Jacobian determinant.

3.18 Exercises

1. (a) Describe the level sets of $f(x,y) = x^2 + 4y^2$ and of $g(x,y) = xy$, and sketch a contour plot of each function.
   (b) Let $f(x,y) = \frac{x^2 y}{x^4+y^2}$ for $(x,y) \neq (0,0)$, and $f(0,0) = 0$. Show that $f(t\mathbf{v}) \to 0$ as $t \to 0$ for every vector $\mathbf{v} \in \mathbb{R}^2$, i.e., along every line through the origin, but that f is not continuous at the origin. (Hint: look at the parabola $y = x^2$.)

2. (a) Show that the derivative of the softplus function $s(x) = \log(1+e^x)$ is the sigmoid function σ, and that s is convex.
   (b) Show that $\tanh(x) = 2\sigma(2x) - 1$, and use $\sigma' = \sigma(1-\sigma)$ to show that $\tanh' = 1 - \tanh^2$.

3. Compute the Taylor polynomial T_3 of $f(x) = \log(1+x)$ at 0, and use it to approximate $\log(1.1)$. Use Taylor's theorem to show that the error is at most $2.5 \cdot 10^{-5}$.

4. Let $f(x,y,z) = xe^{yz} + y^2 z$. Compute ∇f and the directional derivative of f at $(1,0,2)$ in the direction of the unit vector $\frac{1}{3}(1,2,2)^\top$. In which direction does f decrease fastest at this point, and at which rate?

5. Let $f(x,y) = \sqrt{|xy|}$. Show that both partial derivatives of f at the origin exist, but that f is not differentiable at the origin. (Hint: Proposition 3.3.5.)

6. (a) Compute the Jacobian and the Jacobian determinant of $f(x,y) = (e^x \cos y,\, e^x \sin y)^\top$.
   (b) Show that spherical coordinates $f(r,\theta,\varphi) = (r\sin\theta\cos\varphi,\, r\sin\theta\sin\varphi,\, r\cos\theta)^\top$ have the Jacobian determinant $r^2\sin\theta$. Use the change of variables formula (Remark 3.4.8) to show that the ball of radius R has the volume $\frac{4}{3}\pi R^3$.

7. (a) Let $\mathbf{x}(t)$ be a differentiable curve in $\mathbb{R}^n$. Show that $\frac{d}{dt}\|\mathbf{x}(t)\|^2 = 2\mathbf{x}(t)^\top \mathbf{x}'(t)$, and conclude that if the curve lies on a sphere $\|\mathbf{x}\| = r$, then its velocity $\mathbf{x}'(t)$ is always orthogonal to $\mathbf{x}(t)$.
   (b) Let $f(u,v) = u^2 v$ and $g(x,y) = (x\cos y,\, x\sin y)^\top$. Compute $D(f\circ g)$ in two ways: as the product $Df(g(x,y))\, Dg(x,y)$, and by differentiating $f(g(x,y))$ directly.

8. (Ridge regression.) Let $X \in \mathbb{R}^{N\times d}$, $\mathbf{y} \in \mathbb{R}^N$ and $\lambda > 0$, and let $f(\mathbf{w}) = \|X\mathbf{w} - \mathbf{y}\|^2 + \lambda\|\mathbf{w}\|^2$. Compute ∇f and $\nabla^2 f$. Show that $X^\top X + \lambda I$ is invertible, and that the unique minimum of f is $\mathbf{w}^* = (X^\top X + \lambda I)^{-1} X^\top \mathbf{y}$.

9. Let A be a symmetric $n \times n$ matrix and $R(\mathbf{x}) = \frac{\mathbf{x}^\top A\mathbf{x}}{\mathbf{x}^\top\mathbf{x}}$ for $\mathbf{x} \neq \mathbf{0}$ (the Rayleigh quotient). Show that $\nabla R(\mathbf{x}) = \frac{2}{\mathbf{x}^\top\mathbf{x}}(A\mathbf{x} - R(\mathbf{x})\mathbf{x})$, and conclude that the critical points of R are exactly the eigenvectors of A.

10. (a) Show that $\nabla_X \operatorname{tr}(AXB) = A^\top B^\top$.
    (b) Let $X \in \mathbb{R}^{N\times d}$ and $Y \in \mathbb{R}^{N\times k}$. Compute $\nabla_W \|XW - Y\|_F^2$, and show that if X has linearly independent columns, the minimum is $W^* = (X^\top X)^{-1} X^\top Y$. (This is linear regression with k outputs.)
    (c) Let X be invertible. Show that $(X+H)^{-1} = X^{-1} - X^{-1}HX^{-1} + o(\|H\|_F)$, by multiplying the right side by $X+H$. Conclude that $\nabla_X \operatorname{tr}(AX^{-1}) = -X^{-\top}A^\top X^{-\top}$.

11. Let $\mathbf{y} \in \mathbb{R}^K$ be a probability vector (with $y_k \geq 0$ and $\sum_k y_k = 1$), and let

$$\ell(\mathbf{z}) = -\sum_{k=1}^{K} y_k \log \operatorname{softmax}(\mathbf{z})_k$$

be the cross-entropy with "soft labels". Show that $\nabla \ell(\mathbf{z}) = \operatorname{softmax}(\mathbf{z}) - \mathbf{y}$. Check that $\ell(\mathbf{z} + c\mathbf{1}) = \ell(\mathbf{z})$ for all $c \in \mathbb{R}$, and that $\nabla \ell(\mathbf{z})^\top \mathbf{1} = 0$. Why do these two facts fit together?

12. Let $f(x_1, x_2) = x_1^2 x_2 + e^{x_1 x_2}$. Draw the computation graph of f, and compute $\nabla f(1, 0)$ with forward mode (two passes) and with reverse mode (one pass), as in Example 3.7.1.

13. Consider the network $\hat{y} = w_2\, \sigma(w_1 x + b_1) + b_2$ with one input, one hidden unit with the sigmoid activation function, and one output, and the loss $L = \frac{1}{2}(\hat{y} - y)^2$. Derive formulas for the four partial derivatives of L with respect to w_1, b_1, w_2 and b_2, using backpropagation. Evaluate them for $x = 1$, $y = 0$, $w_1 = b_1 = b_2 = 0$ and $w_2 = 2$.

14. Compute the gradient and the Hessian of $f(x, y) = \log(1 + x + 2y)$, and its second-order Taylor approximation at the origin. Use it to approximate $f(0.1, 0.05)$, and compare with the exact value. Is the Hessian at the origin positive definite, negative definite, or neither?

15. Find and classify all critical points of $f(x, y) = x^4 + y^4 - 4xy$. Which of them are global minima?

16. Show that $f(x, y) = x^2 + y^4$ and $g(x, y) = x^2 - y^4$ have the same gradient and the same Hessian at the origin, but that the origin is a minimum of f and a saddle point of g.

17. Let $f(x_1, x_2) = x_1^2 + 9x_2^2$.

    (a) For which step sizes η does gradient descent converge for every starting point?

    (b) Find the best step size and the corresponding rate ρ. How many steps are needed to reduce the error by a factor 10^{-6}?

    (c) Substitute $x_2 = \frac{u}{3}$. How does the function look in the variables (x_1, u), and how many steps does gradient descent need now with the best step size?

18. (a) Show that Newton's method finds the minimum of $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top \mathbf{x}$ (with A symmetric positive definite) in one step, from every starting point.

    (b) Let B be an invertible matrix, $\mathbf{c}$ a vector, and $g(\mathbf{y}) = f(B\mathbf{y} + \mathbf{c})$. Show that if $\mathbf{x}_0 = B\mathbf{y}_0 + \mathbf{c}$, then the Newton iterates for f and g satisfy $\mathbf{x}_k = B\mathbf{y}_k + \mathbf{c}$ for all k. Show by an example that this is false for gradient descent.

    (c) Apply Newton's method to $f(x) = x^4$, starting at $x_0 = 1$. Show that $x_k = (\frac{2}{3})^k$. Why is the convergence not quadratic here?

19. (Line search and Gauss–Newton.)

    (a) Let f be L-smooth, and let $0 < c < 1$ and $0 < \beta < 1$. Show that $f(\mathbf{x} - \eta \nabla f(\mathbf{x})) \leq f(\mathbf{x}) - c\eta \|\nabla f(\mathbf{x})\|^2$ for all $\eta \leq \frac{2(1-c)}{L}$, and that backtracking that multiplies η by β accepts a step size $\eta \geq \min\left(\bar{\eta}, \frac{2\beta(1-c)}{L}\right)$.

    (b) Show that for a linear model $\mathbf{r}(\boldsymbol{\theta}) = A\boldsymbol{\theta} - \mathbf{y}$, where A has linearly independent columns, the Gauss–Newton method reaches the least squares solution in one step, from every starting point.

    (c) Show that the Levenberg–Marquardt step $\mathbf{h}_\lambda$ satisfies $\mathbf{h}_\lambda \to \mathbf{0}$ and $\lambda \mathbf{h}_\lambda \to -\nabla f(\boldsymbol{\theta})$ as $\lambda \to \infty$.

20. Which of the following functions are convex? Justify your answers. (a) $e^{x_1+x_2}$ on $\mathbb{R}^2$; (b) x_1x_2 on $\mathbb{R}^2$; (c) $\|\mathbf{x}\|_1 + \|\mathbf{x}\|^2$ on $\mathbb{R}^n$; (d) $\max(0, 1-x)$ on $\mathbb{R}$; (e) $\mathbf{x}^\top A\mathbf{x}$ on $\mathbb{R}^2$ with $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$; (f) $-\log(x_1x_2)$ on $\{\mathbf{x} \in \mathbb{R}^2 : x_1 > 0, x_2 > 0\}$.

21. (a) Show that if f is convex, then every sublevel set $\{\mathbf{x} : f(\mathbf{x}) \le c\}$ is convex. Give an example of a function on $\mathbb{R}$ that is not convex, although all its sublevel sets are convex.

    (b) Show that the gradient of a differentiable convex function satisfies $(\nabla f(\mathbf{x}) - \nabla f(\mathbf{y}))^\top(\mathbf{x} - \mathbf{y}) \ge 0$ for all $\mathbf{x}, \mathbf{y}$. (Hint: add two inequalities of Theorem 3.11.3.) What does this say for $n = 1$?

22. Let $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x}$ with $A = \operatorname{diag}(1, 4)$.

    (a) Find the largest μ and the smallest L for which f is μ-strongly convex and L-smooth.

    (b) How many steps of gradient descent with $\eta = \frac{1}{L}$ does Theorem 3.11.12 need to guarantee $f(\mathbf{x}_k) - f^* \le 10^{-6}(f(\mathbf{x}_0) - f^*)$? Compare with the exact behavior of gradient descent on this function (Theorem 3.10.7).

23. Compute the subdifferentials of $f(x) = \max(0, 1-x)$ (the hinge loss) and of $g(x) = |x| + x^2$ at every point $x \in \mathbb{R}$. Use the condition $0 \in \partial f(x^*)$ to minimize g and $h(x) = |x-1|+|x-2|+|x-6|$. What is the minimum of $\sum_{i=1}^N |x - a_i|$ in general?

24. (Weight decay.) Show that the proximal operator of $h(\mathbf{u}) = \frac{\lambda}{2}\|\mathbf{u}\|^2$ is $\operatorname{prox}_{\eta h}(\mathbf{v}) = \frac{1}{1+\eta\lambda}\mathbf{v}$. Write down the proximal gradient method for $f + h$, and compare it with gradient descent on $f + h$. Both shrink the weights towards $\mathbf{0}$ in every step, which is called weight decay.

25. (Maximum entropy.) Use a Lagrange multiplier to find the probability vector $\mathbf{p} = (p_1, \dots, p_n)$ with $p_i > 0$ and $\sum_i p_i = 1$ that maximizes the entropy $H(\mathbf{p}) = -\sum_i p_i \log p_i$. What is the maximal entropy?

26. (a) Use a Lagrange multiplier to find the point of the plane $x + 2y + 2z = 9$ that is closest to the origin. Compare with the formula for the distance to a hyperplane in Lecture 2.

    (b) Find the rectangle with sides parallel to the axes and the largest area that fits into the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} \le 1$.

27. Solve the problem: minimize $x^2 + y^2$ subject to $x + y \ge 2$, using the KKT conditions. Which constraint is active, and what is the multiplier? What changes for the constraint $x + y \ge -2$?

28. Find the dual function and solve the dual problem for (a) minimizing x^2+y^2 subject to $x+y \ge 2$, and (b) minimizing e^x subject to $x \ge 0$. Check that $d^* = p^*$ in both cases, and compare the dual solutions with the KKT multipliers.

29. (Convex conjugates.)

    (a) Compute the conjugates of $f(x) = ax^2 + bx$ (with $a > 0$), of $f(x) = |x|$, and of the softplus function $f(x) = \log(1 + e^x)$. (For the last one, the answer is a binary negative entropy.)

    (b) Show that $f^*(\mathbf{y}) \ge -f(\mathbf{0})$, and that $(cf)^*(\mathbf{y}) = cf^*(\mathbf{y}/c)$ for $c > 0$.

    (c) Write down the dual of the linear program: minimize $x_1 + 2x_2$ subject to $x_1 + x_2 \ge 1$, $x_1 \ge 0$ and $x_2 \ge 0$. Solve both problems, and check that their optimal values agree.

30. (a) Compute the divergence and the curl of $F(x, y, z) = (yz, xz, xy)^\top$ and of $G(x, y, z) = (-y, x, z)^\top$. Find a function f with $\nabla f = F$. Is G a gradient field?

    (b) Show that $\operatorname{div}(gF) = g \operatorname{div} F + \nabla g^\top F$ and $\Delta(fg) = g\Delta f + 2\nabla f^\top \nabla g + f\Delta g$.

(c) Show that $\Delta \log\|\mathbf{x}\| = 0$ on $\mathbb{R}^2 \setminus \{\mathbf{0}\}$, and that $\Delta\|\mathbf{x}\|^{2-n} = 0$ on $\mathbb{R}^n \setminus \{\mathbf{0}\}$ for $n \geq 3$. (Hint: $\nabla\|\mathbf{x}\| = \frac{\mathbf{x}}{\|\mathbf{x}\|}$, and use (b).)

31. Let $F(\mathbf{x}) = A\mathbf{x}$ with $A \in \mathbb{R}^{n\times n}$.

(a) Show that $\operatorname{div} F = \operatorname{tr} A$, that F is a gradient field if and only if A is symmetric, and that then $F = \nabla(\frac{1}{2}\mathbf{x}^\top A\mathbf{x})$.

(b) For $A = \begin{pmatrix} -1 & 2 \\ 0 & -1 \end{pmatrix}$, split A into its symmetric and skew-symmetric parts, and describe how the vector field stretches and rotates small neighborhoods of the origin. Compute $\operatorname{div} F$ and $\operatorname{curl} F$.

32. (Hutchinson's estimator.) Let $A \in \mathbb{R}^{n\times n}$ be symmetric, and let $\boldsymbol{\varepsilon} \in \{-1, 1\}^n$ be a vector of signs. Show that $\boldsymbol{\varepsilon}^\top A\boldsymbol{\varepsilon} = \operatorname{tr} A + 2\sum_{i<j} A_{ij}\varepsilon_i\varepsilon_j$, and conclude that the average of $\boldsymbol{\varepsilon}^\top A\boldsymbol{\varepsilon}$ over all 2^n sign vectors is $\operatorname{tr} A$. For which matrices is every single estimate exact?

33. (a) Compute $\iint_D (x + y)\,dx\,dy$ for $D = \{(x, y) : 0 \leq x \leq 1,\ x^2 \leq y \leq x\}$, integrating first over y and then over x, and in the other order.

(b) A random point $\mathbf{X}$ is uniformly distributed on the unit disk, i.e., it has the density $\frac{1}{\pi}$ on the disk and 0 outside. Use polar coordinates to compute its average distance $\int \frac{1}{\pi}\|\mathbf{x}\|\,d\mathbf{x}$ from the center. Find the density of the first coordinate X_1 by integrating out x_2.

34. (a) Differentiate $\int_{-\infty}^{\infty} e^{-ax^2}\,dx = \sqrt{\pi/a}$ (for $a > 0$) with respect to a under the integral sign, and conclude that $\int_{-\infty}^{\infty} x^2 e^{-x^2/2}\,dx = \sqrt{2\pi}$.

(b) (Stein's identity.) Let $\varphi(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$. Use integration by parts to show that for every polynomial f,

$$\int_{-\infty}^{\infty} f'(x)\varphi(x)\,dx = \int_{-\infty}^{\infty} xf(x)\varphi(x)\,dx,$$

and use it to compute $\int x^4\varphi(x)\,dx$.

35. (a) Verify the divergence theorem for $F(x, y) = (x^3, y^3)^\top$ on the unit disk.

(b) Use Green's theorem to show that the area enclosed by a closed curve $(x(t), y(t))$, $0 \leq t \leq T$, traversed counterclockwise, is $\frac{1}{2}\int_0^T (x(t)y'(t) - y(t)x'(t))\,dt$. Compute the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} \leq 1$ in this way.

36. Solve the ODE $\mathbf{x}' = A\mathbf{x}$ with $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$ and $\mathbf{x}(0) = (1, 0)^\top$, using the eigenvectors of A. Which initial values give solutions that stay bounded as $t \to \infty$? Compute e^{At}.

37. (a) Show that Euler's method for $x' = -\lambda x$ with $\lambda > 0$ gives $x_k = (1 - h\lambda)^k x_0$. For which step sizes h does the numerical solution tend to 0, like the exact solution? Relate the answer to Theorem 3.10.7.

(b) Let A be skew-symmetric. Show that $Q(t) = e^{At}$ is orthogonal for all t, by differentiating $Q(t)^\top Q(t)$. Show that Euler's method for $\mathbf{x}' = A\mathbf{x}$ increases $\|\mathbf{x}_k\|$ in every step (unless $A\mathbf{x}_k = \mathbf{0}$).

38. Let $A = \begin{pmatrix} -1 & 2 \\ 0 & -1 \end{pmatrix} = -I + N$ with $N = \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$.

(a) Show that A is not diagonalizable, and that $N^2 = 0$.

(b) Show that $e^{At} = e^{-t}(I + tN)$, by checking that it equals I at $t = 0$ and that its derivative is A times itself. Solve $\mathbf{x}' = A\mathbf{x}$ with $\mathbf{x}(0) = (0, 1)^\top$.

(c) Check Liouville's formula $\det e^{At} = e^{t \operatorname{tr} A}$ for this matrix. By which factor does the flow change areas in the time t?

39. Let $\mathbf{x}' = A\mathbf{x}$, and let $\mathbf{x}(0)$ be random with the density p_0. Use the change of variables formula (Remark 3.4.8) and Proposition 3.16.1 to show that $\log p_t(\mathbf{x}(t)) = \log p_0(\mathbf{x}(0)) - t \operatorname{tr} A$, and check that this agrees with Proposition 3.16.3. What is the distribution of $\mathbf{x}(t)$ if $\mathbf{x}(0) \sim \mathcal{N}(\mathbf{0}, I)$?

3.19 Programming Exercises

As in the previous lectures, implement the algorithms yourself with NumPy, and use library routines only to check your results. Most exercises also need Matplotlib for plots. Start each file with

```
import numpy as np
import matplotlib.pyplot as plt
rng = np.random.default_rng(0)  # random number generator for test data
```

P1. Numerical differentiation and its errors. Write functions `forward_diff(f, x, h)` and `central_diff(f, x, h)` for functions of one variable.

- For $f = \sin$ at $x = 1$, compute the errors of both approximations of $f'(1) = \cos 1$ for $h = 10^{-1}, 10^{-2}, \ldots, 10^{-15}$, and plot them against h with `plt.loglog`.
- Explain the V-shape of the curves: for large h, the error of Proposition 3.2.9 dominates; for small h, rounding errors of size about $\frac{10^{-16}}{h}$ dominate. Which h is best for each method, and how accurate is the result?
- Write a function `numerical_gradient(f, X, h=1e-5)` that computes the gradient of a scalar function `f` with respect to an array `X` of any shape by central differences, one entry at a time. (Hint: loop over `np.ndindex(X.shape)`.) You will use it in the following exercises.

P2. Gradient checking. For random vectors and matrices, compare every formula of Table 3.1 with `numerical_gradient`, using the relative error $\frac{\|\mathbf{g} - \tilde{\mathbf{g}}\|}{\|\mathbf{g}\| + \|\tilde{\mathbf{g}}\|}$.

- Compute $\operatorname{softmax}(\mathbf{z})$ for $\mathbf{z} = (1000, 0, -1000)^\top$ directly from the definition. What goes wrong? Since $\operatorname{softmax}(\mathbf{z} - c\mathbf{1}) = \operatorname{softmax}(\mathbf{z})$, you can subtract $c = \max_i z_i$ first. Implement a stable `softmax(z)` and a stable `logsumexp(z)` in this way, and check them against `scipy.special` if you have SciPy installed.
- Deliberately introduce a mistake into one of your gradient formulas (e.g., forget a transpose) and observe the relative error.

P3. Contour plots and gradient fields. Use `np.meshgrid`, `plt.contour` and `plt.quiver` to plot level curves and gradient vectors of $x^2 + y^2$, $x^2 - y^2$, $x^3 + y^3 - 3xy$ (mark the critical points of Example 3.9.4) and the Rosenbrock function $(1 - x)^2 + 100(y - x^2)^2$ (use logarithmically spaced levels). Use `plt.gca().set_aspect("equal")` so that right angles look right.

- Check Corollary 3.5.4 numerically: `cs = plt.contour(...)` returns an object, and `cs.get_paths()[i].vertices` contains the points of the i-th level curve. Compute the cosine of the angle between ∇f and the segments between consecutive points.
- Plot the graph of $x^2 - y^2$ as a surface. For this, create three-dimensional axes with the command `ax = plt.axes(projection="3d")`, and then use `ax.plot_surface`.

P4. Automatic differentiation. Implement reverse-mode automatic differentiation for scalar functions. Each `Var` stores its value, its parents with the local partial derivatives, and its adjoint `grad`:

```
class Var:
    def __init__(self, value, parents=()):
        self.value = value
        self.parents = parents  # pairs (parent Var, local partial derivative)
        self.grad = 0.0

    def __add__(self, other):
        other = other if isinstance(other, Var) else Var(other)
        return Var(self.value + other.value, [(self, 1.0), (other, 1.0)])

    def __mul__(self, other):
        other = other if isinstance(other, Var) else Var(other)
        return Var(self.value * other.value,
                   [(self, other.value), (other, self.value)])

def log(x):
    return Var(np.log(x.value), [(x, 1.0 / x.value)])
```

- Add subtraction, negation, division, powers with a constant exponent, `__radd__` and `__rmul__` (for expressions like `2 * x`), and the functions `exp`, `sin`, `cos` and `sigmoid`.
- Write a method `backward(self)`: list all variables that the output depends on in topological order (every variable after its parents, e.g., by a depth-first search), set the adjoint of the output to 1, and go through the list in reverse order, adding `child.grad * local` to `parent.grad` for every pair `(parent, local)`.
- Check your implementation on Example 3.7.1 (you should get 5.5 and 1.716), on the logistic regression loss of Example 3.5.6, and on random examples against `numerical_gradient`.
- (Forward mode.) Implement a class `Dual` that stores a value and a derivative $\dot{v}$, with the forward rules of Section 3.7, and check it on the same examples.

P5. Logistic regression. Generate two overlapping classes of points in the plane, and add a constant feature for the bias:

```
N = 200
X0 = rng.normal([-1, -1], 1.0, size=(N // 2, 2))  # class 0
X1 = rng.normal([1, 1], 1.0, size=(N // 2, 2))    # class 1
X = np.hstack([np.vstack([X0, X1]), np.ones((N, 1))])
y = np.r_[np.zeros(N // 2), np.ones(N // 2)]
```

- Implement the average cross-entropy loss, its gradient and its Hessian (Examples 3.5.6 and 3.8.4). For numerical stability, use $-y \log \sigma(z) - (1-y)\log(1-\sigma(z)) = \log(1+e^z) - yz$ and `np.logaddexp(0, z)`. Check the gradient with `numerical_gradient`.
- Run gradient descent with $\eta = 0.1$, 1 and 10, and Newton's method, until $\|\nabla L(\mathbf{w})\| < 10^{-6}$. Plot the loss against the number of iterations (with a logarithmic y-axis for $L(\mathbf{w}_k) - L(\mathbf{w}^*)$). How many iterations does each method need?
- Plot the data and the decision boundary $\mathbf{w}^\top \mathbf{x} = 0$.
- Move the class centers to $(\pm 3, \pm 3)$, so that the classes are (very likely) linearly separable, and run gradient descent again. What happens to $\|\mathbf{w}_k\|$ and to the loss, and why does the loss have no minimum? How does a ridge penalty $\lambda\|\mathbf{w}\|^2$ change this?

P6. Training a neural network. Fit $y = \sin x$ on $[-\pi, \pi]$ with a network with one hidden layer of 20 units with the tanh activation function, as in Section 3.7, with the loss $L = \frac{1}{2N}\sum_k (\hat{y}_k - y_k)^2$ over $N = 100$ equally spaced points.

- Implement the forward and backward passes for a whole batch, with the inputs as the rows of a matrix (Example 3.6.5). Check all four gradients with `numerical_gradient` on a small network.
- Initialize W_1 with standard normal entries, W_2 with normal entries with standard deviation $\frac{1}{\sqrt{20}}$, and the biases with zeros. Train with gradient descent with $\eta = 0.05$ for 20 000 steps. Plot the loss (log scale) and the fitted function.
- Build a deep network with 10 hidden layers of 20 sigmoid units (initialize the weights between hidden layers with standard deviation $\frac{1}{\sqrt{20}}$), and print the norms $\|\nabla_{W_l} L\|_F$ for all layers at initialization. What do you observe? Repeat with tanh and with ReLU.

P7. Gradient descent on quadratic functions. Let $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x}$ with $A = \mathrm{diag}(1, \kappa)$.

- For $\kappa = 10$, plot the level curves of f and the paths of gradient descent from $\mathbf{x}_0 = (8, 2)^\top$ for $\eta = 0.05$, $\frac{2}{11}$, 0.19 and 0.21, as in Figure 3.8.
- For $\kappa = 1, 10, 100, 1000$ and $\eta = \frac{2}{1+\kappa}$, count the steps needed to reach $\|\mathbf{x}_k\| \le 10^{-6}\|\mathbf{x}_0\|$ from $\mathbf{x}_0 = (1, 1)^\top$, and compare with Theorem 3.10.7.
- Implement the heavy ball method with $\eta = \frac{4}{(1+\sqrt{\kappa})^2}$ and $\beta = \left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}\right)^2$, and repeat the count. How does the number of steps grow with κ now?

P8. The Rosenbrock function. Let $f(x, y) = (1 - x)^2 + 100(y - x^2)^2$, whose minimum is $(1, 1)$, at the bottom of a long, curved valley.

- Compute ∇f and $\nabla^2 f$ by hand, and check them with `numerical_gradient`. Check that $(1, 1)$ is the only critical point, and that the Hessian there is positive definite. What is its condition number?
- Starting at $(-1.2, 1)$, run gradient descent with $\eta = 10^{-3}$ until $\|\mathbf{x}_k - (1, 1)^\top\| < 10^{-4}$. How many steps does it need? What happens for $\eta = 3 \cdot 10^{-3}$? (Look at consecutive iterates, and compare η with $\frac{2}{\lambda_{\max}}$ for the Hessian at $(1, 1)$.)
- Implement gradient descent with a backtracking line search: in each step, start with $\eta = 1$ and halve it until $f(\mathbf{x} - \eta\nabla f(\mathbf{x})) \le f(\mathbf{x}) - 10^{-4}\eta\|\nabla f(\mathbf{x})\|^2$.
- Run Newton's method from the same starting point. Plot the paths of all three methods on a contour plot, and compare the numbers of steps.

P9. Stochastic gradient descent. Generate a linear regression problem with many data points:

```
N, d = 10_000, 20
X = rng.normal(size=(N, d))
w_true = rng.normal(size=d)
y = X @ w_true + 0.5 * rng.normal(size=N)
```

and let $L(\mathbf{w}) = \frac{1}{N}\|X\mathbf{w} - \mathbf{y}\|^2$.

- Compute the minimum $\mathbf{w}^*$ with `np.linalg.lstsq`, and the eigenvalues of the Hessian $\frac{2}{N}X^\top X$. Which step sizes are safe for gradient descent?
- Compare gradient descent ($\eta = 0.1$) with SGD with mini-batches of size 1, 32 and 256 (draw the indices with `rng.integers(0, N, size=B)`), with step sizes 0.01, 0.05 and 0.1. Plot $L(\mathbf{w}_k) - L(\mathbf{w}^*)$ against the number of processed data points (one gradient step of full gradient descent processes N points).

- With a constant step size, SGD stops improving at some level. Try the decreasing step sizes $\eta_k = \frac{\eta_0}{1+k/1000}$ and compare.

P10. Constrained optimization.

- (Projected gradient ascent.) For a random symmetric positive semidefinite matrix $A = B^\top B$, maximize $\mathbf{x}^\top A\mathbf{x}$ on the unit sphere by repeating $\mathbf{x} \leftarrow \frac{\mathbf{x}+\eta A\mathbf{x}}{\|\mathbf{x}+\eta A\mathbf{x}\|}$. Compare the result with the top eigenvector from `np.linalg.eigh` (Example 3.12.3). What happens as $\eta \to \infty$? (Compare with power iteration from Lecture 1.)
- (Projected gradient descent.) Let `A = rng.normal(size=(30, 10))`, let `x_true` be a random vector of norm 3, and `b = A @ x_true + 0.1 * rng.normal(size=30)`. Minimize $\|A\mathbf{x}-\mathbf{b}\|^2$ subject to $\|\mathbf{x}\| \le 1$, by alternating gradient steps with step size $\frac{1}{L}$, where $L = 2\|A\|_2^2$, with the projection of Example 3.12.7. At the result, compute the multiplier μ from the stationarity condition, and check all four KKT conditions.
- (Minimum norm solutions.) For a random 3×6 matrix A and $\mathbf{b} \in \mathbb{R}^3$, compute $\mathbf{x}^* = A^\top(AA^\top)^{-1}\mathbf{b}$ (Example 3.12.4) and compare it with `np.linalg.pinv(A) @ b`. Get a basis of $\mathrm{null}(A)$ from the SVD (the last rows of $V^\top$), and check that adding a null space vector to $\mathbf{x}^*$ gives another solution of $A\mathbf{x} = \mathbf{b}$ with a larger norm.

P11. Vector fields.

- Plot the three vector fields of Figure 3.11 and the vortex field of Example 3.14.11 with `plt.quiver`. Compute their divergence and curl on a grid with `np.gradient`, which approximates partial derivatives by central differences, and compare with the exact values.
- (Hutchinson's estimator.) Let $F(\mathbf{x}) = \tanh(W\mathbf{x}+\mathbf{b})$ (entrywise) with a random 100×100 matrix W and a random vector $\mathbf{b}$. Show that $\operatorname{div} F(\mathbf{x}) = \sum_i (1-\tanh^2 z_i)W_{ii}$ with $\mathbf{z} = W\mathbf{x}+\mathbf{b}$, and that $\boldsymbol{\varepsilon}^\top DF(\mathbf{x}) = \left(\boldsymbol{\varepsilon} \odot (1-\tanh^2 \mathbf{z})\right)^\top W$. At a random point, average $\boldsymbol{\varepsilon}^\top DF(\mathbf{x})\boldsymbol{\varepsilon}$ over m random sign vectors, and plot the error against $m = 1, 10, 100, 1000$ on a log-log scale. What is the slope?
- (Game dynamics.) For $f(x, y) = xy$, start at $(1, 1)$ and run 1000 steps with $\eta = 0.1$ of (i) simultaneous gradient descent–ascent, $(x, y) \leftarrow (x - \eta y,\ y + \eta x)$; (ii) alternating updates, $x \leftarrow x - \eta y$ and then $y \leftarrow y + \eta x$ with the new x; (iii) the extragradient method, which computes the trial point $(\tilde{x}, \tilde{y}) = (x-\eta y,\ y+\eta x)$ and then updates $(x, y) \leftarrow (x-\eta\tilde{y},\ y+\eta\tilde{x})$ with the gradients at the trial point. Plot the paths and $\|(x_k, y_k)\|$ against k. Show that (i) multiplies the norm by $\sqrt{1+\eta^2}$ in each step, and (iii) by $\sqrt{1-\eta^2+\eta^4}$.

P12. Multiple integrals.

- By Fubini's theorem, $\int_{[0,1]^d} e^{-\|\mathbf{x}\|^2}\,d\mathbf{x} = \left(\int_0^1 e^{-t^2}\,dt\right)^d$. Compute the one-dimensional integral accurately with the midpoint rule (the Riemann sum with the midpoints of the small intervals). Then approximate the d-dimensional integral for $d = 1, 2, 3, 5, 8$ by the midpoint rule on a grid with k points per axis (k^d evaluations of the function), and by the average of the function over N uniformly random points (Monte Carlo, Lecture 4) with the same number of evaluations. Plot the errors against the number of evaluations. Which method is better in low dimensions, and which in high dimensions?
- Check the divergence theorem numerically for $F(x, y) = (x^3 + y,\ \sin x + y^2)^\top$ on the ellipse $\frac{x^2}{4} + y^2 \le 1$: compute the left side on a fine grid, using the indicator function of the ellipse, and the right side with the parametrization $\gamma(t) = (2\cos t, \sin t)$, for which $\mathbf{n}\|\gamma'(t)\| = (\gamma_2'(t), -\gamma_1'(t))^\top$.

- (Integrated gradients.) For the logistic regression model $f(\mathbf{x}) = \sigma(\mathbf{w}^\top \mathbf{x})$ of P5, compute the attributions of Example 3.14.9 for a few data points with the baseline $\mathbf{x}' = \mathbf{0}$ (approximate the integrals by the midpoint rule with 50 points), and check that they add up to $f(\mathbf{x}) - f(\mathbf{x}')$. Compare with the attributions $x_i \frac{\partial f}{\partial x_i}(\mathbf{x})$ ("gradient times input").

P13. ODEs and gradient flow.

- Implement Euler's method for $\mathbf{x}' = F(\mathbf{x})$. For $A = \begin{pmatrix} -2 & 1 \\ 1 & -2 \end{pmatrix}$ and $\mathbf{x}_0 = (2, 0)^\top$, compare the numerical solution at $t = 1$ with the exact solution of Example 3.15.5, for $h = 0.1, 0.05, 0.025, \ldots$. Plot the error against h on a log-log scale; what is the slope? If you have SciPy installed, also check your solver against `scipy.integrate.solve_ivp`.
- Implement the fourth-order Runge–Kutta method, $\mathbf{x}_{k+1} = \mathbf{x}_k + \frac{h}{6}(\mathbf{k}_1 + 2\mathbf{k}_2 + 2\mathbf{k}_3 + \mathbf{k}_4)$ with $\mathbf{k}_1 = F(\mathbf{x}_k)$, $\mathbf{k}_2 = F(\mathbf{x}_k + \frac{h}{2}\mathbf{k}_1)$, $\mathbf{k}_3 = F(\mathbf{x}_k + \frac{h}{2}\mathbf{k}_2)$, $\mathbf{k}_4 = F(\mathbf{x}_k + h\mathbf{k}_3)$, and repeat the error plot.
- Plot the vector field of Figure 3.12 with `plt.quiver`, together with numerical solutions. Do the same for the gradient flow of the Rosenbrock function, and compare the solution of the flow (computed with a small step) with the gradient descent iterates of P8 for $\eta = 10^{-3}$ and $\eta = 3 \cdot 10^{-3}$.

P14. Flows and volumes.

- Draw 10 000 samples from $\mathcal{N}(\mathbf{0}, I)$ in $\mathbb{R}^2$, and move them with the flow of $\mathbf{x}' = A\mathbf{x}$ for the matrix A of Figure 3.12 up to $t = 1$, using your Runge–Kutta method from P13. Compare the sample covariance matrix with $e^{At} e^{A^\top t}$, and its determinant with $e^{2t \operatorname{tr} A}$.
- The pendulum $x_1' = x_2$, $x_2' = -\sin x_1$ has $\operatorname{div} F = 0$, and with friction, $x_2' = -\sin x_1 - \gamma x_2$, it has $\operatorname{div} F = -\gamma$. Represent a small disk around $(1, 0)$ by 200 points on its boundary circle, move them with the flow, and compute the enclosed area over time with the shoelace formula $\frac{1}{2}\sum_i (x_i y_{i+1} - x_{i+1} y_i)$. Compare with the prediction of the divergence for $\gamma = 0$ and $\gamma = 0.5$, and plot the deformed disks.

P15. The lasso. Generate a regression problem in which only 5 of 50 features matter:

```
N, d = 100, 50
X = rng.normal(size=(N, d))
w_true = np.zeros(d)
w_true[:5] = [3, -2, 1.5, -1, 0.5]
y = X @ w_true + 0.3 * rng.normal(size=N)
```

- Implement soft thresholding (Proposition 3.11.17) and the iterative soft-thresholding algorithm for $F(\mathbf{w}) = \frac{1}{2}\|X\mathbf{w} - \mathbf{y}\|^2 + \lambda\|\mathbf{w}\|_1$, with the step size $\eta = 1/\|X\|_2^2$ (the smoothness constant of the first term). Check that $F(\mathbf{w}_k)$ decreases in every step (Theorem 3.11.18).
- For $\lambda = 1, 10, 50$, count the nonzero weights of the solution, and compare them with $\mathbf{w}_{\text{true}}$ and with the least squares solution. Check the optimality condition of Theorem 3.11.15: $|(X^\top(X\mathbf{w} - \mathbf{y}))_i| \le \lambda$ for the zero weights, and $= \lambda$ for the others.
- Plot the weights of the solution as functions of λ (the lasso path). At which values of λ do the true features enter the model?

P16. Line search, Gauss–Newton and Hessian-vector products.

- Implement gradient descent with backtracking line search (Proposition 3.10.6, with $\bar{\eta} = 1$), and apply it to the Rosenbrock function $f(x, y) = (1 - x)^2 + 100(y - x^2)^2$ from $(-1.2, 1)$. Compare with the constant step size $\eta = 10^{-3}$: how many iterations and how many evaluations of f does each method need to reach $\|\nabla f\| < 10^{-4}$?

- Generate 20 data points $y_i = 2e^{-0.5 t_i} + 0.05 \varepsilon_i$ with t_i equally spaced in $[0, 10]$ and standard normal ε_i, and fit the model $y = a e^{-bt}$ with Gauss–Newton and with Levenberg–Marquardt, starting from $(a, b) = (1, 2)$. Plot $f(\boldsymbol{\theta}_k)$ for both methods.
- For the Rosenbrock function, compute Hessian-vector products as $\frac{1}{2\varepsilon}\big(\nabla f(\mathbf{x} + \varepsilon \mathbf{v}) - \nabla f(\mathbf{x} - \varepsilon \mathbf{v})\big)$, compare them with the exact Hessian, and use them in power iteration to find the largest eigenvalue of the Hessian at the minimum $(1, 1)$. What does it say about the largest stable step size of gradient descent near the minimum?

Lecture 4

Probability and Statistics

Introduction

Machine learning is about learning from data, and data is noisy, incomplete and random: measurements contain errors, labels can be wrong, and the training set is only a sample of all the data we might see. Probability theory is the mathematical language for uncertainty, and statistics is the art of drawing conclusions from random data. Both are everywhere in machine learning:

- classifiers such as logistic regression and softmax regression (Lecture 3) output probabilities;
- the loss functions of Lecture 3 are negative log-likelihoods: least squares corresponds to Gaussian noise, and cross-entropy to categorical labels;
- regularization corresponds to prior beliefs about the parameters;
- stochastic gradient descent uses random mini-batches, and works because averages of many random terms concentrate around their mean;
- generative models, such as normalizing flows (Lecture 3), learn probability distributions.

In this lecture, we introduce probability spaces, conditional probability and Bayes' theorem, random variables, expectation and variance, the Poisson process, the variance of a neural network layer (which explains the Xavier and He initializations), conditional independence, graphical models and naive Bayes, moment generating functions, and random vectors with their covariance matrices, where the linear algebra of Lectures 1 and 2 enters. We then study the Gaussian distribution in detail, the law of large numbers and the central limit theorem, Monte Carlo sampling and the convergence of stochastic gradient descent, and Markov chains with Markov chain Monte Carlo (Metropolis and Gibbs sampling), the Bellman equations of reinforcement learning, and stationary time series with autoregressive models. The second half of the lecture is about learning from data: estimation (the method of moments, maximum likelihood, the Cramér–Rao bound, sufficiency and the Rao–Blackwell theorem, exponential families and generalized linear models, the empirical CDF and kernel density estimates, and survival analysis with censored data), hypothesis testing (p-values, power, the Neyman–Pearson lemma, t-tests and χ^2-tests, standard errors, t-tests and heteroscedasticity in linear regression, multiple testing, the bootstrap and permutation tests), the evaluation of models and experiments (cross-validation, classification metrics and ROC curves, generalization bounds, and A/B testing with Simpson's paradox), Bayesian inference (including Gaussian processes), information theory (entropy, cross-entropy and the Kullback–Leibler divergence), and latent variable models (Gaussian mixtures, the EM algorithm, k-means, probabilistic PCA and variational autoencoders). Finally, we add Gaussian noise to the ordinary differential equations of Lecture 3: Brownian motion, stochastic differential equations and the Ornstein–Uhlenbeck process, and Langevin dynamics, which samples from a distribution by noisy gradient descent and is behind

diffusion models. Appendix B builds probability theory on measure theory, which treats discrete and continuous distributions, and everything in between, in a single framework.

Notation. As in the previous lectures, vectors of $\mathbb{R}^n$ are bold. Random variables are denoted by capital letters such as X and Y, their values by small letters such as x and y, and random vectors by bold capital letters such as $\mathbf{X}$. We write $P(A)$ for the probability of an event A, $\mathbf{1}_A$ for its indicator function, which is 1 on A and 0 outside A, $\mathbb{E}[X]$ for the expectation of X, and log for the natural logarithm.

4.1 Probability Spaces

A random experiment has a set of possible outcomes, and we want to assign probabilities to sets of outcomes.

Definition 4.1.1 (Probability space)**.** A probability space consists of a sample space Ω, the set of all possible outcomes, and a probability P, which assigns a number $P(A)$ to every event $A \subseteq \Omega$, such that:

1. $P(A) \geq 0$ for every event A;
2. $P(\Omega) = 1$;
3. if $A_1, A_2, \ldots$ are pairwise disjoint events, then $P(A_1 \cup A_2 \cup \cdots) = P(A_1) + P(A_2) + \cdots$.

Remark 4.1.2 (Measure theory). If Ω is uncountable, such as $\Omega = \mathbb{R}$, it is impossible to assign a probability to every subset of Ω in a consistent way. A rigorous treatment specifies a collection of events (a σ-algebra) and uses measure theory. All sets that we meet in practice, such as intervals, balls and their unions and complements, are events, and we ignore these technicalities in this lecture; Appendix B treats them.

Proposition 4.1.3. *For all events A, B and $A_1, A_2, \ldots$:*

1. *$P(A^c) = 1 - P(A)$, where $A^c = \Omega \setminus A$, and $P(\emptyset) = 0$;*
2. *if $A \subseteq B$, then $P(A) \leq P(B)$;*
3. *$P(A \cup B) = P(A) + P(B) - P(A \cap B)$;*
4. *(union bound) $P(A_1 \cup A_2 \cup \cdots) \leq P(A_1) + P(A_2) + \cdots$.*

Proof. (1) A and A^c are disjoint with union Ω, so $P(A)+P(A^c) = 1$; for $A = \Omega$, this gives $P(\emptyset) = 0$. (2) B is the disjoint union of A and $B \setminus A$, so $P(B) = P(A) + P(B \setminus A) \geq P(A)$. (3) $A \cup B$ is the disjoint union of A and $B \setminus A$, and B is the disjoint union of $B \setminus A$ and $A \cap B$; so $P(A \cup B) = P(A) + P(B \setminus A) = P(A) + P(B) - P(A \cap B)$. (4) Let $B_1 = A_1$ and $B_k = A_k \setminus (A_1 \cup \cdots \cup A_{k-1})$. The B_k are disjoint, $B_k \subseteq A_k$, and they have the same union as the A_k; now use the third axiom and (2). □

Example 4.1.4 (Equally likely outcomes)**.** If Ω is finite and all outcomes are equally likely, then $P(A) = \frac{|A|}{|\Omega|}$. For two fair dice, $\Omega = \{1, \ldots, 6\}^2$ has 36 outcomes. The sum is 7 for the 6 outcomes $(1,6), (2,5), \ldots, (6,1)$, so $P(\text{sum} = 7) = \frac{6}{36} = \frac{1}{6}$. The event "at least one six" is the complement of "no six", which has $5 \cdot 5 = 25$ outcomes, so its probability is $1 - \frac{25}{36} = \frac{11}{36}$.

4.1.1 Conditional probability and Bayes' theorem

If we learn that an event B has happened, the probabilities of other events change: only the outcomes in B are still possible.

Definition 4.1.5 (Conditional probability)**.** If $P(B) > 0$, the conditional probability of A given B is

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}.$$

For fixed B, the function $A \mapsto P(A \mid B)$ is again a probability. Rearranging the definition gives the product rule $P(A \cap B) = P(A \mid B)\, P(B)$.

Theorem 4.1.6 (Total probability and Bayes' theorem)**.** *Let $B_1, \ldots, B_n$ be disjoint events with union Ω and $P(B_i) > 0$. Then for every event A,*

$$P(A) = \sum_{i=1}^{n} P(A \mid B_i)\, P(B_i),$$

and if $P(A) > 0$, then

$$P(B_j \mid A) = \frac{P(A \mid B_j)\, P(B_j)}{P(A)} = \frac{P(A \mid B_j)\, P(B_j)}{\sum_{i=1}^{n} P(A \mid B_i)\, P(B_i)}.$$

Proof. A is the disjoint union of the events $A \cap B_i$, so $P(A) = \sum_i P(A \cap B_i) = \sum_i P(A \mid B_i) P(B_i)$ by the product rule. For the second formula, use the product rule twice: $P(B_j \mid A) P(A) = P(A \cap B_j) = P(A \mid B_j) P(B_j)$. □

Bayes' theorem reverses conditional probabilities: from the probability of the evidence A given each hypothesis B_j, it computes the probability of each hypothesis given the evidence. The probabilities $P(B_j)$ are called prior probabilities, and the $P(B_j \mid A)$ posterior probabilities.

Example 4.1.7 (A medical test)**.** A disease affects 1% of a population. A test detects it in 99% of the sick people, but also gives a positive result for 5% of the healthy people. What is the probability that a person with a positive test is sick? With D = "sick" and $+$ = "positive test",

$$P(D \mid +) = \frac{P(+ \mid D) P(D)}{P(+ \mid D) P(D) + P(+ \mid D^c) P(D^c)} = \frac{0.99 \cdot 0.01}{0.99 \cdot 0.01 + 0.05 \cdot 0.99} = \frac{1}{6}.$$

Although the test is quite accurate, only one in six people with a positive test is sick, because the disease is rare: among 10 000 people, about 99 sick people and 495 healthy people test positive. Ignoring the prior probability is a common mistake, called the base rate fallacy. In machine learning, the same effect makes classifiers for rare classes (such as fraud detection) produce many false alarms.

4.1.2 Independence

Definition 4.1.8 (Independence)**.** Two events A and B are independent if $P(A \cap B) = P(A) P(B)$. Events $A_1, \ldots, A_n$ are (mutually) independent if $P(A_{i_1} \cap \cdots \cap A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$ for every choice of indices $i_1 < \cdots < i_k$.

If $P(B) > 0$, then A and B are independent if and only if $P(A \mid B) = P(A)$: learning that B happened does not change the probability of A. For example, for two fair dice, the events A = "the first die is even" and B = "the sum is 7" are independent: $A \cap B$ consists of the 3 outcomes $(2,5), (4,3), (6,1)$, so $P(A \cap B) = \frac{3}{36} = \frac{1}{2} \cdot \frac{1}{6} = P(A) P(B)$. Independence is not the same as disjointness: disjoint events with positive probabilities are never independent, since $P(A \cap B) = 0$.

4.2 Random Variables and Their Distributions

Definition 4.2.1 (Random variable)**.** A random variable is a function $X : \Omega \to \mathbb{R}$ on a probability space. Its distribution describes the probabilities $P(X \in I) = P(\{\omega : X(\omega) \in I\})$ for intervals I. Its cumulative distribution function (CDF) is $F(x) = P(X \le x)$.

The CDF is nondecreasing and right-continuous, tends to 0 as $x \to -\infty$ and to 1 as $x \to \infty$, and $P(a < X \le b) = F(b) - F(a)$. In practice, we rarely look at the sample space: a random variable is described by its distribution. Most distributions in machine learning are of one of two kinds.

Definition 4.2.2 (Discrete and continuous random variables)**.**

1. X is discrete if it takes only finitely or countably many values $x_1, x_2, \ldots$. Its probability mass function (pmf) is $p(x) = P(X = x)$, and $\sum_i p(x_i) = 1$.
2. X is continuous if there is a function $p \ge 0$, its probability density function (pdf), with

$$P(a \le X \le b) = \int_a^b p(x)\,dx \qquad \text{for all } a \le b, \qquad \text{and} \qquad \int_{-\infty}^{\infty} p(x)\,dx = 1.$$

For a continuous random variable, $P(X = x) = 0$ for every single value x, and the density is the derivative of the CDF, $p = F'$ (where p is continuous). A density is not a probability: it can be larger than 1. For a small interval, $P(x \le X \le x + h) \approx p(x)h$.

Example 4.2.3 (Discrete distributions)**.**

1. Bernoulli(p): $X \in \{0, 1\}$ with $P(X = 1) = p$; for example, a coin flip, or a binary label.
2. Binomial(n, p): the number of successes in n independent Bernoulli(p) trials,

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \qquad k = 0, 1, \ldots, n,$$

since each of the $\binom{n}{k}$ sequences with k successes has probability $p^k(1-p)^{n-k}$. For example, for $n = 10$ and $p = 0.3$, $P(X = 3) = 120 \cdot 0.3^3 \cdot 0.7^7 \approx 0.267$ (Figure 4.1).

3. Geometric(p): the number of independent Bernoulli(p) trials until the first success, $P(X = k) = (1-p)^{k-1}p$ for $k = 1, 2, \ldots$.
4. Poisson(λ): $P(X = k) = e^{-\lambda}\frac{\lambda^k}{k!}$ for $k = 0, 1, 2, \ldots$; it describes counts of rare events, such as the number of clicks on an ad in an hour. It is the limit of Binomial($n, \frac{\lambda}{n}$) as $n \to \infty$.
5. Categorical($\mathbf{p}$): $X \in \{1, \ldots, K\}$ with $P(X = k) = p_k$, where $p_k \ge 0$ and $\sum_k p_k = 1$; for example, the class predicted by a softmax classifier.

Example 4.2.4 (Continuous distributions)**.**

1. Uniform(a, b): $p(x) = \frac{1}{b-a}$ for $a \le x \le b$, and 0 otherwise.
2. Exponential(λ): $p(x) = \lambda e^{-\lambda x}$ for $x \ge 0$; then $P(X > t) = e^{-\lambda t}$. It describes waiting times, and is memoryless: $P(X > s + t \mid X > s) = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} = P(X > t)$.
3. Normal (Gaussian) $\mathcal{N}(\mu, \sigma^2)$, with mean μ and variance $\sigma^2 > 0$:

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\Big(-\frac{(x-\mu)^2}{2\sigma^2}\Big).$$

It integrates to 1 by the Gaussian integral of Lecture 3. The standard normal distribution $\mathcal{N}(0, 1)$ has the density $\varphi(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}$ and the CDF $\Phi(z) = \int_{-\infty}^{z} \varphi(t)\,dt$, which has no formula in terms of elementary functions.




Figure 4.1: Left: the pmf of the Binomial$(10, 0.3)$ distribution. Right: the densities of the standard normal distribution, the Exponential(1) distribution and the Uniform$(-1, 1)$ distribution.

4.2.1 Transformations of random variables

If X is a random variable and $g : \mathbb{R} \to \mathbb{R}$ is a function, then $Y = g(X)$ is again a random variable. For continuous random variables and invertible g, the density changes by the derivative of g, just as areas change by the Jacobian determinant in Lecture 3.

Proposition 4.2.5 (Change of variables)**.** *Let X have the density p_X, and let g be differentiable and strictly increasing (or strictly decreasing). Then $Y = g(X)$ has the density*

$$p_Y(y) = p_X(x)\,\frac{1}{|g'(x)|}, \qquad \text{where } x = g^{-1}(y).$$

Proof. For increasing g, $F_Y(y) = P(g(X) \le y) = P(X \le g^{-1}(y)) = F_X(g^{-1}(y))$. Differentiating with the chain rule gives $p_Y(y) = p_X(g^{-1}(y))\,(g^{-1})'(y) = \frac{p_X(x)}{g'(x)}$. For decreasing g, $F_Y(y) = 1 - F_X(g^{-1}(y))$, which gives the same formula with $|g'(x)|$. □

Example 4.2.6.

1. (Standardization.) If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Z = \frac{X-\mu}{\sigma}$ has the density $\sigma\, p_X(\mu + \sigma z) = \varphi(z)$, so $Z \sim \mathcal{N}(0, 1)$. Conversely, $X = \mu + \sigma Z$. Hence $P(|X - \mu| \le k\sigma) = P(|Z| \le k)$, which is 0.683, 0.954 and 0.997 for $k = 1, 2, 3$, and 0.95 for $k = 1.96$.

2. (Log-normal distribution.) If $X \sim \mathcal{N}(0, 1)$, then $Y = e^X$ has the density $\frac{\varphi(\log y)}{y}$ for $y > 0$.

Theorem 4.2.7 (Inverse transform sampling)**.** *Let F be nondecreasing and right-continuous with the limits 0 at $-\infty$ and 1 at ∞, let*

$$F^{-1}(u) = \min\{x \in \mathbb{R} : F(x) \ge u\} \qquad \text{for } 0 < u < 1$$

be its quantile function, and let $U \sim$ Uniform$(0, 1)$. Then $X = F^{-1}(U)$ has the CDF F.

Proof. For $0 < u < 1$, the set $\{x : F(x) \ge u\}$ is nonempty and bounded below, since F tends to 1 and to 0. Since F is nondecreasing, it is an interval that extends to ∞, and since F is right-continuous, it contains its left endpoint. So the minimum exists, and $F^{-1}(u) \le x$ if and only if $F(x) \ge u$. Hence $P(X \le x) = P(U \le F(x)) = F(x)$. □

If F is continuous and strictly increasing, F^{-1} is the inverse function of F. So a computer that can generate uniform random numbers can generate random numbers with any distribution on $\mathbb{R}$. For example, the Exponential(λ) distribution has $F(x) = 1 - e^{-\lambda x}$, so $X = -\frac{1}{\lambda}\log(1 - U)$ is exponentially distributed. For a Bernoulli(p) variable, F jumps from 0 to $1 - p$ at 0 and from $1 - p$ to 1 at 1, so $F^{-1}(u) = 0$ for $u \le 1 - p$ and $F^{-1}(u) = 1$ for $u > 1 - p$: we return 1 if $U > 1 - p$, which has the probability p. Normalizing flows (Lecture 3) generalize this idea to many dimensions: they transform simple random vectors by learned invertible maps.

4.3 Expectation and Variance

Definition 4.3.1 (Expectation)**.** The expectation (expected value, or mean) of a random variable X is

$$\mathbb{E}[X] = \sum_i x_i\, p(x_i) \qquad \text{or} \qquad \mathbb{E}[X] = \int_{-\infty}^{\infty} x\, p(x)\, dx,$$

for discrete and continuous X, respectively, provided that the sum or integral converges absolutely.

The expectation is the center of mass of the distribution: if we put the mass $p(x_i)$ at each point x_i, the balance point is $\mathbb{E}[X]$. To compute the expectation of a function of X, we do not need the distribution of $g(X)$; one can show that

$$\mathbb{E}[g(X)] = \sum_i g(x_i)\, p(x_i) \qquad \text{or} \qquad \mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)\, p(x)\, dx.$$

The expectation is linear: $\mathbb{E}[aX + bY + c] = a\mathbb{E}[X] + b\mathbb{E}[Y] + c$ for all random variables X, Y with finite expectations, and all numbers a, b, c (we prove this in Section 4.4 for discrete random variables).

Example 4.3.2 (Tail sums)**.** Let X take values in $\{0, 1, 2, \ldots\}$. Then

$$\mathbb{E}[X] = \sum_{k=0}^{\infty} P(X > k).$$

Indeed, $P(X > k) = \sum_{j>k} P(X = j)$, and in the double series $\sum_{k\geq 0} \sum_{j>k} P(X = j)$, whose terms are nonnegative and can therefore be rearranged, each $P(X = j)$ appears j times, for $k = 0, \ldots, j-1$; so the sum is $\sum_j j\, P(X = j) = \mathbb{E}[X]$. In other words, $X = \sum_{k\geq 0} \mathbf{1}_{\{X>k\}}$, since exactly the first X terms are 1, and we may take expectations term by term. For a Geometric(p) variable, $P(X > k) = (1-p)^k$, the probability that the first k trials fail, so $\mathbb{E}[X] = \sum_{k\geq 0}(1-p)^k = \frac{1}{p}$. Example 4.4.7 gives the analogous formula for continuous random variables.

Definition 4.3.3 (Variance)**.** The variance of X with mean $\mu = \mathbb{E}[X]$ is $\mathrm{Var}(X) = \mathbb{E}[(X-\mu)^2]$, and its standard deviation is $\sigma = \sqrt{\mathrm{Var}(X)}$.

The variance measures how far X typically is from its mean. Expanding the square and using linearity,

$$\mathrm{Var}(X) = \mathbb{E}[X^2 - 2\mu X + \mu^2] = \mathbb{E}[X^2] - 2\mu^2 + \mu^2 = \mathbb{E}[X^2] - \mathbb{E}[X]^2,$$

and $\mathrm{Var}(aX + b) = \mathbb{E}[(aX - a\mu)^2] = a^2\, \mathrm{Var}(X)$: shifting does not change the spread, and scaling by a scales the standard deviation by $|a|$.

Proposition 4.3.4. *The distributions of Section 4.2 have the following means and variances:*

distribution	$\mathbb{E}[X]$	$\mathrm{Var}(X)$
Bernoulli(p)	p	$p(1-p)$
Binomial(n,p)	np	$np(1-p)$
Geometric(p)	$\frac{1}{p}$	$\frac{1-p}{p^2}$
Poisson(λ)	λ	λ
Uniform(a,b)	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
Exponential(λ)	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
$\mathcal{N}(\mu, \sigma^2)$	μ	σ^2

Proof of some cases. Bernoulli: $\mathbb{E}[X] = 0 \cdot (1-p) + 1 \cdot p = p$, and $X^2 = X$, so $\mathrm{Var}(X) = p - p^2$. Uniform: $\mathbb{E}[X] = \frac{1}{b-a}\int_a^b x\,dx = \frac{b^2-a^2}{2(b-a)} = \frac{a+b}{2}$, and similarly $\mathbb{E}[X^2] = \frac{a^2+ab+b^2}{3}$, so $\mathrm{Var}(X) = \frac{(b-a)^2}{12}$. Exponential: integrating by parts, $\mathbb{E}[X] = \int_0^\infty x\lambda e^{-\lambda x}\,dx = \left[-xe^{-\lambda x}\right]_0^\infty + \int_0^\infty e^{-\lambda x}\,dx = \frac{1}{\lambda}$. Poisson: $\mathbb{E}[X] = \sum_{k\geq 1} k e^{-\lambda}\frac{\lambda^k}{k!} = \lambda \sum_{k\geq 1} e^{-\lambda}\frac{\lambda^{k-1}}{(k-1)!} = \lambda$. Normal: by Example 4.2.6, $X = \mu + \sigma Z$ with $Z \sim \mathcal{N}(0,1)$, so it suffices to show $\mathbb{E}[Z] = 0$ and $\mathrm{Var}(Z) = 1$. The first holds since $z\varphi(z)$ is an odd function, and for the second, integration by parts with $\varphi'(z) = -z\varphi(z)$ gives

$$\mathbb{E}[Z^2] = \int_{-\infty}^{\infty} z \cdot z\varphi(z)\,dz = \big[-z\varphi(z)\big]_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \varphi(z)\,dz = 1.$$

The binomial case follows from the Bernoulli case and Proposition 4.4.9 below. □

Example 4.3.5. For a fair die, $\mathbb{E}[X] = \frac{1+2+\cdots+6}{6} = 3.5$, which is not a possible value, and $\mathbb{E}[X^2] = \frac{91}{6}$, so $\mathrm{Var}(X) = \frac{91}{6} - \frac{49}{4} = \frac{35}{12} \approx 2.92$.

4.3.1 Inequalities

The variance controls how likely large deviations from the mean are.

Proposition 4.3.6 (Markov and Chebyshev)**.** *1. (Markov's inequality.) If $X \geq 0$ and $a > 0$, then $P(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$.*

2. (Chebyshev's inequality.) If X has mean μ and variance σ^2, then $P(|X-\mu| \geq k) \leq \frac{\sigma^2}{k^2}$ for every $k > 0$.

Proof. (1) Since $X \geq 0$, we have $X \geq a\mathbf{1}_{\{X\geq a\}}$. Taking expectations, $\mathbb{E}[X] \geq aP(X \geq a)$. (2) Apply (1) to $(X-\mu)^2$ and $a = k^2$: $P(|X-\mu| \geq k) = P((X-\mu)^2 \geq k^2) \leq \frac{\sigma^2}{k^2}$. □

For example, every random variable is at least 2 standard deviations from its mean with probability at most $\frac{1}{4}$. This bound holds for all distributions, so it is weak for specific ones: for a normal distribution, the probability is only 0.046.

Theorem 4.3.7 (Jensen's inequality)**.** *If f is convex, then $f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)]$. If f is concave, then $f(\mathbb{E}[X]) \geq \mathbb{E}[f(X)]$.*

Proof. We prove this for differentiable f. Let $\mu = \mathbb{E}[X]$. By the first-order characterization of convexity in Lecture 3, $f(x) \geq f(\mu) + f'(\mu)(x-\mu)$ for all x. So $f(X) \geq f(\mu) + f'(\mu)(X-\mu)$, and taking expectations gives $\mathbb{E}[f(X)] \geq f(\mu) + f'(\mu)\cdot 0$. For concave f, apply this to $-f$. □

For example, $f(x) = x^2$ gives $\mathbb{E}[X]^2 \leq \mathbb{E}[X^2]$, i.e., $\mathrm{Var}(X) \geq 0$, and the concave function log gives $\mathbb{E}[\log X] \leq \log \mathbb{E}[X]$ for $X > 0$. We will use the latter to show that the Kullback–Leibler divergence is nonnegative (Section 4.13).

Example 4.3.8 (The Chernoff bound)**.** Markov's inequality becomes much stronger when it is applied to an exponential: for every $t > 0$, $P(X \geq a) = P(e^{tX} \geq e^{ta}) \leq e^{-ta}\mathbb{E}[e^{tX}]$, and we may choose the best t. For $S \sim \mathrm{Binomial}(n, \frac{1}{2})$, the sum of n independent fair coin flips, $\mathbb{E}[e^{tS}] = \left(\frac{1+e^t}{2}\right)^n$ (by Proposition 4.4.4 below). For $n = 100$ and $a = 60$, the best choice is $e^t = \frac{3}{2}$, which gives $P(S \geq 60) \leq \left(\frac{5}{4}\right)^{100}\left(\frac{2}{3}\right)^{60} \approx 0.134$, while Chebyshev's inequality only gives $P(S \geq 60) \leq P(|S-50| \geq 10) \leq \frac{25}{100} = 0.25$; the exact value is 0.028. As n grows, Chernoff bounds decrease exponentially, while Chebyshev's bound only decreases like $\frac{1}{n}$. Hoeffding's inequality (Theorem 4.7.5) is proved in this way.

4.4 Joint Distributions and Independence

To describe how several random variables depend on each other, we need their joint distribution.

Definition 4.4.1 (Joint, marginal and conditional distributions)**.** The joint pmf of two discrete random variables X and Y is $p(x,y) = P(X = x, Y = y)$. Two continuous random variables have a joint density $p(x,y) \geq 0$ if $P((X,Y) \in A) = \iint_A p(x,y)\,dx\,dy$ for all (reasonable) sets $A \subseteq \mathbb{R}^2$. The marginal distributions of X and Y are

$$p_X(x) = \sum_y p(x,y) \qquad \text{or} \qquad p_X(x) = \int_{-\infty}^{\infty} p(x,y)\,dy,$$

and similarly for p_Y. The conditional distribution of Y given $X = x$ (for $p_X(x) > 0$) is

$$p(y \mid x) = \frac{p(x,y)}{p_X(x)}.$$

The formulas $p_X(x) = \sum_y p(x,y)$ (the sum rule) and $p(x,y) = p(y \mid x)\, p_X(x)$ (the product rule) are the two basic rules of probabilistic machine learning, and Bayes' theorem $p(x \mid y) = \frac{p(y|x)p_X(x)}{p_Y(y)}$ follows from them. The same definitions apply to more than two random variables.

Definition 4.4.2 (Independent random variables)**.** X and Y are independent if $p(x,y) = p_X(x)\, p_Y(y)$ for all x and y; equivalently, $p(y \mid x) = p_Y(y)$ whenever $p_X(x) > 0$. Random variables $X_1, \ldots, X_n$ are independent if their joint pmf or density is the product of the marginal ones. They are independent and identically distributed (i.i.d.) if moreover they all have the same distribution.

The standard assumption in machine learning is that the training examples are i.i.d. samples from an unknown distribution: each example is drawn from the same distribution, independently of the others.

Example 4.4.3. Let $X, Y \in \{0,1\}$ have the joint pmf

	$Y=0$	$Y=1$	p_X
$X=0$	0.3	0.2	0.5
$X=1$	0.1	0.4	0.5
p_Y	0.4	0.6	

The marginals are the row and column sums. The conditional probabilities are $P(Y = 1 \mid X = 1) = \frac{0.4}{0.5} = 0.8$ and $P(Y = 1 \mid X = 0) = \frac{0.2}{0.5} = 0.4$. Since $p(1,1) = 0.4 \neq 0.5 \cdot 0.6 = p_X(1)p_Y(1)$, X and Y are not independent: learning X changes the probabilities of Y.

For a function $g(X,Y)$, the expectation is $\mathbb{E}[g(X,Y)] = \sum_{x,y} g(x,y)\, p(x,y)$, or the corresponding integral. In particular, linearity of expectation follows: for discrete X and Y,

$$\mathbb{E}[X+Y] = \sum_{x,y} (x+y)\, p(x,y) = \sum_x x \sum_y p(x,y) + \sum_y y \sum_x p(x,y) = \mathbb{E}[X] + \mathbb{E}[Y].$$

Proposition 4.4.4. *If X and Y are independent, then $\mathbb{E}[XY] = \mathbb{E}[X]\,\mathbb{E}[Y]$, and more generally $\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\,\mathbb{E}[h(Y)]$.*

Proof. For discrete random variables,

$$\mathbb{E}[g(X)h(Y)] = \sum_{x,y} g(x)h(y)p_X(x)p_Y(y) = \Big(\sum_x g(x)p_X(x)\Big)\Big(\sum_y h(y)p_Y(y)\Big);$$

for densities, the same computation works with integrals. □

Example 4.4.5 (Sums of independent random variables)**.** If X and Y are independent with the densities p_X and p_Y, then $X+Y$ has the density

$$p_{X+Y}(z) = \int_{-\infty}^{\infty} p_X(x)\, p_Y(z-x)\, dx,$$

the convolution of p_X and p_Y. Indeed, substituting $y = z - x$ in the inner integral and exchanging the order of integration, which is allowed for nonnegative functions (Fubini's theorem, Lecture 3),

$$P(X+Y \le t) = \iint \mathbf{1}_{\{x+y\le t\}}\, p_X(x)\, p_Y(y)\, dy\, dx = \int_{-\infty}^{t}\int_{-\infty}^{\infty} p_X(x)\, p_Y(z-x)\, dx\, dz.$$

For two independent Uniform$(0,1)$ variables, the integrand is 1 for $\max(0, z-1) \le x \le \min(1, z)$ and 0 otherwise, so $p_{X+Y}(z) = z$ for $0 \le z \le 1$ and $p_{X+Y}(z) = 2 - z$ for $1 \le z \le 2$: a triangular density, and $P(X+Y \le \frac{1}{2}) = \frac{1}{8}$. For independent $X \sim \text{Poisson}(\lambda)$ and $Y \sim \text{Poisson}(\mu)$, the same computation with sums and the binomial theorem gives

$$P(X+Y=n) = \sum_{k=0}^{n} e^{-\lambda}\frac{\lambda^k}{k!}\, e^{-\mu}\frac{\mu^{n-k}}{(n-k)!} = e^{-(\lambda+\mu)}\frac{(\lambda+\mu)^n}{n!},$$

so $X+Y \sim \text{Poisson}(\lambda+\mu)$. For example, if a website gets independent Poisson numbers of clicks from search engines and from ads, with the means 3 and 2 per minute, then the total number of clicks per minute is Poisson(5).

Example 4.4.6 (The Poisson process)**.** Customers arrive at a shop, or requests at a server, with independent Exponential(λ) waiting times $T_1, T_2, \dots$ between consecutive arrivals. Let $S_n = T_1 + \cdots + T_n$ be the time of the n-th arrival, and $N(t)$ the number of arrivals up to time t. By the convolution formula and induction, S_n has the density

$$p_{S_n}(t) = \frac{\lambda^n t^{n-1}}{(n-1)!}e^{-\lambda t} \qquad (t \ge 0),$$

since $\int_0^t \frac{\lambda^n s^{n-1}}{(n-1)!}e^{-\lambda s}\, \lambda e^{-\lambda(t-s)}\, ds = \lambda^{n+1}e^{-\lambda t}\int_0^t \frac{s^{n-1}}{(n-1)!}\, ds = \frac{\lambda^{n+1}t^n}{n!}e^{-\lambda t}$. Integration by parts gives

$$P(S_{n+1} \le t) = \int_0^t \frac{\lambda^{n+1}s^n}{n!}e^{-\lambda s}\, ds = -\frac{(\lambda t)^n}{n!}e^{-\lambda t} + \int_0^t \frac{\lambda^n s^{n-1}}{(n-1)!}e^{-\lambda s}\, ds = P(S_n \le t) - \frac{(\lambda t)^n}{n!}e^{-\lambda t}.$$

Since $N(t) \ge n$ exactly when $S_n \le t$, we get $P(N(t)=n) = P(S_n \le t) - P(S_{n+1} \le t) = e^{-\lambda t}\frac{(\lambda t)^n}{n!}$ for $n \ge 1$, and $P(N(t)=0) = P(T_1 > t) = e^{-\lambda t}$: the number of arrivals in the time t is Poisson(λt). This is the Poisson process with rate λ. One can show more: by the memorylessness of the exponential distribution, the numbers of arrivals in disjoint time intervals are independent and Poisson distributed, with means λ times the lengths of the intervals. This explains why counts of many independent rare events, such as clicks, emails or radioactive decays, are approximately Poisson distributed.

Example 4.4.7 (Tail integrals)**.** For a random variable $X \ge 0$ with a density p, exchanging the order of integration (Fubini's theorem for nonnegative functions) gives

$$\int_0^\infty P(X > t)\, dt = \int_0^\infty\int_t^\infty p(x)\, dx\, dt = \int_0^\infty \Big(\int_0^x dt\Big)p(x)\, dx = \int_0^\infty x\, p(x)\, dx = \mathbb{E}[X],$$

the continuous version of Example 4.3.2; Appendix B shows that $\mathbb{E}[X] = \int_0^\infty P(X > t)\, dt$ for every random variable $X \ge 0$. For example, for $X \sim \text{Exponential}(\lambda)$, $\mathbb{E}[X] = \int_0^\infty e^{-\lambda t}\, dt = \frac{1}{\lambda}$, without integration by parts. Similarly, $\mathbb{E}[X^p] = \int_0^\infty pt^{p-1}P(X > t)\, dt$ for $p > 0$, which computes moments from tail bounds.

4.4.1 Covariance and correlation

Definition 4.4.8 (Covariance and correlation)**.** The covariance of X and Y, with means μ_X and μ_Y, is

$$\operatorname{Cov}(X,Y) = \mathbb{E}[(X-\mu_X)(Y-\mu_Y)] = \mathbb{E}[XY] - \mathbb{E}[X]\,\mathbb{E}[Y],$$

and if both have positive variances, their correlation is $\operatorname{Corr}(X,Y) = \frac{\operatorname{Cov}(X,Y)}{\sigma_X \sigma_Y}$. If $\operatorname{Cov}(X,Y) = 0$, they are uncorrelated.

The covariance is positive if X and Y tend to be larger than their means at the same time, and negative if one tends to be large when the other is small. Note that $\operatorname{Cov}(X,X) = \operatorname{Var}(X)$, and the covariance is bilinear and symmetric. By Proposition 4.4.4, independent random variables are uncorrelated. The converse is false: if X is uniform on $\{-1,0,1\}$ and $Y = X^2$, then $\operatorname{Cov}(X,Y) = \mathbb{E}[X^3] - \mathbb{E}[X]\mathbb{E}[X^2] = 0$, but Y is a function of X. Correlation only measures linear dependence.

Proposition 4.4.9. $\operatorname{Var}(X+Y) = \operatorname{Var}(X) + \operatorname{Var}(Y) + 2\operatorname{Cov}(X,Y)$. *In particular, if $X_1, \dots, X_n$ are pairwise uncorrelated (for example, independent), then* $\operatorname{Var}(X_1 + \cdots + X_n) = \operatorname{Var}(X_1) + \cdots + \operatorname{Var}(X_n)$.

Proof. With $\tilde{X} = X - \mu_X$ and $\tilde{Y} = Y - \mu_Y$: $\operatorname{Var}(X+Y) = \mathbb{E}[(\tilde{X}+\tilde{Y})^2] = \mathbb{E}[\tilde{X}^2] + \mathbb{E}[\tilde{Y}^2] + 2\mathbb{E}[\tilde{X}\tilde{Y}]$. The second statement follows by induction. □

For example, a Binomial(n,p) random variable is the sum of n independent Bernoulli(p) random variables, so its mean is np and its variance is $np(1-p)$, as claimed in Proposition 4.3.4.

The covariance behaves like an inner product. For random variables U and V with finite variances, $\langle U, V\rangle = \mathbb{E}[UV]$ is bilinear, symmetric, and $\langle U, U\rangle = \mathbb{E}[U^2] \geq 0$, with equality only if $U = 0$ with probability 1. So, if we do not distinguish random variables that are equal with probability 1, it is an inner product as in Lecture 2. For centered random variables, the inner product is the covariance, the norm is the standard deviation, and the correlation is the cosine of the angle between them.

Theorem 4.4.10. $|\operatorname{Cov}(X,Y)| \leq \sigma_X \sigma_Y$, *i.e.,* $-1 \leq \operatorname{Corr}(X,Y) \leq 1$. *Moreover,* $|\operatorname{Corr}(X,Y)| = 1$ *if and only if $Y = aX + b$ with probability* 1*, for some $a \neq 0$ and b; the sign of the correlation is the sign of a.*

Proof. This is the Cauchy–Schwarz inequality for the centered random variables $\tilde{X}$ and $\tilde{Y}$. Directly: for every $t \in \mathbb{R}$,

$$0 \leq \mathbb{E}[(\tilde{Y} - t\tilde{X})^2] = \sigma_Y^2 - 2t\operatorname{Cov}(X,Y) + t^2\sigma_X^2.$$

A quadratic polynomial in t that is never negative has a discriminant ≤ 0, so $4\operatorname{Cov}(X,Y)^2 - 4\sigma_X^2\sigma_Y^2 \leq 0$. Equality holds if and only if the polynomial has a zero $t = a$, i.e., $\mathbb{E}[(\tilde{Y} - a\tilde{X})^2] = 0$, i.e., $\tilde{Y} = a\tilde{X}$ with probability 1; then $Y = aX + (\mu_Y - a\mu_X)$, and $\operatorname{Cov}(X,Y) = a\sigma_X^2$. □

Example 4.4.11. In Example 4.4.3, $\mathbb{E}[XY] = P(X=1, Y=1) = 0.4$, so $\operatorname{Cov}(X,Y) = 0.4 - 0.5 \cdot 0.6 = 0.1$. With $\operatorname{Var}(X) = 0.25$ and $\operatorname{Var}(Y) = 0.24$, the correlation is $\frac{0.1}{\sqrt{0.06}} \approx 0.41$.

Initializing neural networks. Variances of sums explain how the weights of a neural network have to be scaled when training starts. A unit of a layer computes $z = W_1x_1 + \cdots + W_nx_n$ from the n outputs x_j of the previous layer, and passes $a(z)$ on, for an activation function a such as $\operatorname{ReLU}(z) = \max(z, 0)$ (Lecture 3). At initialization, the weights are random, and the following proposition tracks the size of the signal.

Proposition 4.4.12 (Variance of a layer)**.** *Let $x_1, \dots, x_n$ be random variables with $\mathbb{E}[x_j^2] = q$ for all j, and let $W_1, \dots, W_n$ have mean* 0 *and variance v, such that $W_1, \dots, W_n$ and the vector $(x_1, \dots, x_n)$ are independent. Then $z = W_1x_1 + \cdots + W_nx_n$ has*

$$\mathbb{E}[z] = 0 \qquad \textit{and} \qquad \operatorname{Var}(z) = nvq.$$

If moreover $-W_j$ has the same distribution as W_j for every j, then $-z$ has the same distribution as z, and $\mathbb{E}[\mathrm{ReLU}(z)^2] = \frac{1}{2}\operatorname{Var}(z)$.

Proof. By Proposition 4.4.4, $\mathbb{E}[W_j x_j] = \mathbb{E}[W_j]\,\mathbb{E}[x_j] = 0$, and for $j \neq k$, $\mathbb{E}[W_j x_j W_k x_k] = \mathbb{E}[W_j]\,\mathbb{E}[W_k x_j x_k] = 0$, since W_j is independent of (W_k, x_j, x_k). So the terms $W_j x_j$ are uncorrelated, with $\operatorname{Var}(W_j x_j) = \mathbb{E}[W_j^2]\,\mathbb{E}[x_j^2] = vq$, and $\operatorname{Var}(z) = nvq$ by Proposition 4.4.9. If every W_j is symmetric, then $(-W_1, \ldots, -W_n, x_1, \ldots, x_n)$ has the same joint distribution as $(W_1, \ldots, W_n, x_1, \ldots, x_n)$, by independence, so $-z$ has the same distribution as z. Then $\mathbb{E}[z^2 \mathbf{1}_{\{z>0\}}] = \mathbb{E}[z^2 \mathbf{1}_{\{z<0\}}]$, and the two add up to $\mathbb{E}[z^2]$, so $\mathbb{E}[\mathrm{ReLU}(z)^2] = \mathbb{E}[z^2 \mathbf{1}_{\{z>0\}}] = \frac{1}{2}\mathbb{E}[z^2] = \frac{1}{2}\operatorname{Var}(z)$. □

Note that $q = \mathbb{E}[x_j^2]$ is the second moment, not the variance: the outputs of a ReLU layer are nonnegative and do not have mean 0, but only their second moments matter.

Example 4.4.13 (Xavier and He initialization)**.** Consider a network of ReLU layers, where layer l has n_{l-1} inputs and weights with mean 0 and variance v_l, drawn independently and symmetrically. By Proposition 4.4.12, the second moments q_l of the outputs of layer l satisfy $q_l = \frac{1}{2} n_{l-1} v_l q_{l-1}$, so after L layers,

$$q_L = q_0 \prod_{l=1}^{L} \frac{n_{l-1} v_l}{2}.$$

With $v_l = \frac{2}{n_{l-1}}$ (He initialization), every factor is 1, and the signal keeps its size through any number of layers. With $v_l = \frac{1}{n_{l-1}}$, it is halved in every layer, and after 50 layers, it is multiplied by $2^{-50} \approx 9 \cdot 10^{-16}$: it vanishes. With standard normal weights ($v_l = 1$) and 512 units per layer, it is multiplied by 256 in every layer, and explodes. The same computation for backpropagation, which multiplies the gradients by $W_l^\top$ and sums over the n_l outputs of layer l (Lecture 3), asks for $n_l v_l \approx 2$; for activations such as tanh, which are linear near 0 (no factor $\frac{1}{2}$), the forward and backward conditions are $n_{l-1} v_l = 1$ and $n_l v_l = 1$, and Xavier (Glorot) initialization compromises with $v_l = \frac{2}{n_{l-1} + n_l}$. For weights that are uniform on $[-b, b]$, the variance is $\frac{b^2}{3}$, so the uniform He initialization uses $b = \sqrt{6/n_{l-1}}$. These are the default initializations of deep learning libraries, and normalization layers such as batch normalization keep the scale under control during training, by standardizing z.

4.4.2 Conditional expectation

Definition 4.4.14 (Conditional expectation)**.** The conditional expectation of Y given $X = x$ is the mean of the conditional distribution,

$$m(x) = \mathbb{E}[Y \mid X = x] = \sum_y y\, p(y \mid x) \qquad \text{or} \qquad \int_{-\infty}^{\infty} y\, p(y \mid x)\, dy.$$

The random variable $\mathbb{E}[Y \mid X] = m(X)$ is the conditional expectation of Y given X.

Theorem 4.4.15. *1. (Tower property.)* $\mathbb{E}\big[\mathbb{E}[Y \mid X]\big] = \mathbb{E}[Y]$.

2. (Best prediction.) For every function g, $\mathbb{E}[(Y - g(X))^2] \geq \mathbb{E}[(Y - m(X))^2]$, with $m(x) = \mathbb{E}[Y \mid X = x]$. More precisely,

$$\mathbb{E}[(Y - g(X))^2] = \mathbb{E}[(Y - m(X))^2] + \mathbb{E}[(m(X) - g(X))^2].$$

Proof. We give the proof for discrete random variables. (1) $\mathbb{E}[m(X)] = \sum_x p_X(x) \sum_y y\, p(y \mid x) = \sum_{x,y} y\, p(x, y) = \mathbb{E}[Y]$. (2) For every function h, the error $Y - m(X)$ is orthogonal to $h(X)$:

$$\mathbb{E}[(Y - m(X))h(X)] = \sum_x p_X(x) h(x) \sum_y (y - m(x)) p(y \mid x) = \sum_x p_X(x) h(x) \big(m(x) - m(x)\big) = 0.$$

Now write $Y - g(X) = (Y - m(X)) + (m(X) - g(X))$, expand the square, and use this with $h = m - g$; the mixed term vanishes, as in the theorem of Pythagoras. □

So $\mathbb{E}[Y \mid X]$ is the best prediction of Y from X in the sense of the mean squared error: it is the orthogonal projection of Y onto the space of functions of X, with respect to the inner product $\mathbb{E}[UV]$ (compare with orthogonal projections in Lecture 2). A regression model trained with the squared loss on enough data therefore estimates the conditional mean $\mathbb{E}[Y \mid X = \mathbf{x}]$. In Example 4.4.3, $\mathbb{E}[Y \mid X = 0] = 0.4$ and $\mathbb{E}[Y \mid X = 1] = 0.8$, and indeed $0.5 \cdot 0.4 + 0.5 \cdot 0.8 = 0.6 = \mathbb{E}[Y]$.

Example 4.4.16. 1. (Symmetry.) Let $X_1, \dots, X_n$ be i.i.d., and $S = X_1 + \cdots + X_n$. The pairs (X_i, S) all have the same joint distribution, so the $\mathbb{E}[X_i \mid S]$ are the same function of S, and they add up to $\mathbb{E}[S \mid S] = S$. Hence $\mathbb{E}[X_1 \mid S] = \frac{S}{n}$. For example, the expected value of the first of two dice, given that their sum is 8, is 4.

2. (Law of total variance.) By part (2) of Theorem 4.4.15, with the constant function $g = \mathbb{E}[Y]$,

$$\operatorname{Var}(Y) = \mathbb{E}\big[(Y - m(X))^2\big] + \mathbb{E}\big[(m(X) - \mathbb{E}[Y])^2\big].$$

By the tower property, $\mathbb{E}[m(X)] = \mathbb{E}[Y]$, so the second term is $\operatorname{Var}(\mathbb{E}[Y \mid X])$. Summing first over y, as in the proof of the tower property, shows that the first term is $\mathbb{E}[v(X)]$, where $v(x)$ is the variance of the conditional distribution of Y given $X = x$. With the conditional variance $\operatorname{Var}(Y \mid X) = v(X)$, this is the law of total variance

$$\operatorname{Var}(Y) = \mathbb{E}\big[\operatorname{Var}(Y \mid X)\big] + \operatorname{Var}\big(\mathbb{E}[Y \mid X]\big) :$$

the variance of Y is the mean of the variances within the groups $\{X = x\}$ plus the variance of the group means. For example, let the number of customers of a shop on a day be $N \sim \text{Poisson}(\lambda)$, and let customer i spend Y_i, where the Y_i are i.i.d. with mean μ and variance σ^2, and independent of N. The revenue $T = Y_1 + \cdots + Y_N$ has $\mathbb{E}[T \mid N] = N\mu$ and $\operatorname{Var}(T \mid N) = N\sigma^2$, so $\mathbb{E}[T] = \lambda\mu$ and $\operatorname{Var}(T) = \mathbb{E}[N]\sigma^2 + \operatorname{Var}(N)\mu^2 = \lambda(\sigma^2 + \mu^2)$. For $\lambda = 50$, $\mu = 20$ and $\sigma^2 = 100$, the revenue has the mean 1000 and the standard deviation $\sqrt{25\,000} \approx 158$.

4.4.3 Conditional independence and graphical models

Many models in machine learning are built from the assumption that some variables are independent once others are known.

Definition 4.4.17 (Conditional independence)**.** Random variables X and Y are conditionally independent given Z, written $X \perp\!\!\!\perp Y \mid Z$, if

$$p(x, y \mid z) = p(x \mid z)\, p(y \mid z) \qquad \text{for all } x, y \text{ and all } z \text{ with } p_Z(z) > 0.$$

Equivalently, $p(x \mid y, z) = p(x \mid z)$ whenever $p(y, z) > 0$: once Z is known, Y carries no further information about X. Indeed, $p(x \mid y, z) = \frac{p(x,y,z)}{p(y,z)} = \frac{p(x,y|z)}{p(y|z)}$, which equals $p(x \mid z)$ if and only if $p(x, y \mid z) = p(x \mid z)p(y \mid z)$. Conditional independence and independence are different properties.

- (A common cause.) Choose one of two coins with probability $\frac{1}{2}$ each, a fair one ($Z = 0$) and one with heads probability 0.9 ($Z = 1$), and let X and Y indicate heads in two flips of the chosen coin. Then $X \perp\!\!\!\perp Y \mid Z$. But $P(X = 1) = \frac{1}{2}(0.5 + 0.9) = 0.7$ and $P(X = 1, Y = 1) = \frac{1}{2}(0.25 + 0.81) = 0.53 \neq 0.7^2$, so X and Y are dependent: heads in the first flip makes the biased coin more likely, and $P(Y = 1 \mid X = 1) = \frac{0.53}{0.7} \approx 0.76$. Bayesian inference has this structure, with the parameter θ in the role of Z: the data are i.i.d. given θ, and we learn about future data from past data.

- (Explaining away.) Let X and Y be independent fair bits, and $Z = X + Y$. Then $X \perp\!\!\!\perp Y$, but not given Z: $P(X = 1 \mid Z = 1) = \frac{1}{2}$, while $P(X = 1 \mid Y = 1, Z = 1) = 0$. If an alarm ($Z$) can be set off by a burglary (X) or an earthquake (Y), then learning about the earthquake explains the alarm away and makes the burglary less likely.
- The defining property of a Markov chain (Section 4.8) is that the future is independent of the past given the present.

Applying the product rule repeatedly gives the chain rule of probability,

$$p(x_1, \dots, x_n) = p(x_1)\, p(x_2 \mid x_1)\, p(x_3 \mid x_1, x_2) \cdots p(x_n \mid x_1, \dots, x_{n-1}).$$

A directed graphical model, or Bayesian network, assumes that each variable only depends on a few of the earlier ones, its parents $\mathrm{pa}(i) \subseteq \{1, \dots, i-1\}$:

$$p(x_1, \dots, x_n) = \prod_{i=1}^{n} p(x_i \mid x_{\mathrm{pa}(i)}).$$

It is drawn as a graph with an arrow from each parent to its child. The factorization saves parameters: a general distribution of n binary variables has $2^n - 1$ free parameters, but if every variable has at most k parents, the conditional tables have at most $n2^k$ entries. Such factorizations are equivalent to conditional independence statements; the three basic patterns are the following.

Proposition 4.4.18. *Let X, Y, Z have a joint pmf or density of one of the following forms.*

1. *(Chain $X \to Z \to Y$.) $p(x, y, z) = p(x)\, p(z \mid x)\, p(y \mid z)$. Then $X \perp\!\!\!\perp Y \mid Z$.*
2. *(Fork $X \leftarrow Z \to Y$.) $p(x, y, z) = p(z)\, p(x \mid z)\, p(y \mid z)$. Then $X \perp\!\!\!\perp Y \mid Z$.*
3. *(Collider $X \to Z \leftarrow Y$.) $p(x, y, z) = p(x)\, p(y)\, p(z \mid x, y)$. Then $X \perp\!\!\!\perp Y$, but in general not given Z.*

Proof. (1) By Bayes' theorem, $p(x)p(z \mid x) = p_Z(z)p(x \mid z)$, so $p(x, y \mid z) = \frac{p(x,y,z)}{p_Z(z)} = p(x \mid z)p(y \mid z)$. (2) Divide by $p(z)$. (3) Summing over z gives $p(x, y) = p(x)p(y) \sum_z p(z \mid x, y) = p(x)p(y)$; the explaining-away example above shows that X and Y can be dependent given Z. □

Many models of this lecture are graphical models: a mixture model $z \to \mathbf{x}$ (Section 4.14), Bayesian inference $\theta \to x_1, \dots, x_N$ (a fork with N branches), a hidden Markov model, in which hidden states form a Markov chain $z_1 \to z_2 \to \cdots$ and each z_t emits an observation x_t, and the following classifier.

Example 4.4.19 (Naive Bayes)**.** A naive Bayes classifier predicts a class $y \in \{1, \dots, K\}$ from features $x_1, \dots, x_d$ under the assumption that the features are conditionally independent given the class: $p(\mathbf{x}, y) = p(y) \prod_{j=1}^{d} p(x_j \mid y)$. By Bayes' theorem,

$$p(y \mid \mathbf{x}) \propto p(y) \prod_{j=1}^{d} p(x_j \mid y),$$

and maximum likelihood estimates each factor by counting. For a spam filter, let $P(\text{spam}) = 0.4$, let the word "free" appear in 30% of spam and 2% of other emails, and the word "meeting" in 1% of spam and 10% of other emails. An email that contains "free" but not "meeting" has the posterior odds

$$\frac{P(\text{spam} \mid \mathbf{x})}{P(\text{other} \mid \mathbf{x})} = \frac{0.4}{0.6} \cdot \frac{0.3}{0.02} \cdot \frac{0.99}{0.9} = 11,$$

so $P(\text{spam} \mid \mathbf{x}) = \frac{11}{12} \approx 0.92$. With binary features $x_j \in \{0,1\}$ and two classes, the log-odds

$$\log \frac{p(y=1 \mid \mathbf{x})}{p(y=0 \mid \mathbf{x})} = \log \frac{p(y=1)}{p(y=0)} + \sum_{j=1}^{d} \Big(x_j \log \frac{p(x_j=1 \mid y=1)}{p(x_j=1 \mid y=0)} + (1-x_j) \log \frac{p(x_j=0 \mid y=1)}{p(x_j=0 \mid y=0)} \Big)$$

are an affine function of $\mathbf{x}$, as for logistic regression (Lecture 3). Naive Bayes is a generative model, which fits $p(\mathbf{x}, y)$ by counting, while logistic regression is the corresponding discriminative model, which fits $p(y \mid \mathbf{x})$ directly; naive Bayes needs less data, but its probabilities are overconfident when the features are dependent. A word that never appears in the spam of the training data would get $p(x_j = 1 \mid \text{spam}) = 0$ and veto the class; adding a pseudo-count to all counts (Laplace smoothing, a MAP estimate with a Beta prior, Section 4.12) avoids this.

4.4.4 Moment generating functions

The Chernoff bound (Example 4.3.8) used the expectation of e^{tX}. As a function of t, it contains all the moments of X, and it turns sums of independent random variables into products.

Definition 4.4.20 (Moment generating function)**.** The moment generating function (MGF) of a random variable X is $M_X(t) = \mathbb{E}[e^{tX}]$, for all $t \in \mathbb{R}$ for which this expectation is finite. Its logarithm $K_X(t) = \log M_X(t)$ is the cumulant generating function.

Always $M_X(0) = 1$. For a Bernoulli(p) variable, $M_X(t) = 1 - p + pe^t$; for $X \sim \text{Poisson}(\lambda)$, $M_X(t) = \sum_k e^{tk} e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)}$; for $X \sim \text{Exponential}(\lambda)$, $M_X(t) = \int_0^\infty e^{tx} \lambda e^{-\lambda x}\, dx = \frac{\lambda}{\lambda - t}$ for $t < \lambda$ (and ∞ for $t \geq \lambda$); and for $X \sim \mathcal{N}(\mu, \sigma^2)$, completing the square, $tx - \frac{(x-\mu)^2}{2\sigma^2} = \mu t + \frac{\sigma^2 t^2}{2} - \frac{(x - \mu - \sigma^2 t)^2}{2\sigma^2}$, so

$$M_X(t) = e^{\mu t + \sigma^2 t^2/2} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu-\sigma^2 t)^2/(2\sigma^2)}\, dx = e^{\mu t + \sigma^2 t^2/2}, \qquad K_X(t) = \mu t + \frac{\sigma^2 t^2}{2}.$$

Proposition 4.4.21. *1.* $M_{aX+b}(t) = e^{bt} M_X(at)$.

2. *If X and Y are independent, then $M_{X+Y}(t) = M_X(t) M_Y(t)$ and $K_{X+Y} = K_X + K_Y$.*

3. *If $M_X(t)$ is finite for all t in an interval $(-t_0, t_0)$ with $t_0 > 0$, then M_X has derivatives of all orders on $(-t_0, t_0)$, with $M_X^{(k)}(t) = \mathbb{E}[X^k e^{tX}]$. In particular, $\mathbb{E}[X^k] = M_X^{(k)}(0)$, and $K_X'(0) = \mathbb{E}[X]$, $K_X''(0) = \text{Var}(X)$.*

Proof. (1) $\mathbb{E}[e^{t(aX+b)}] = e^{bt}\mathbb{E}[e^{(at)X}]$. (2) By Proposition 4.4.4, $\mathbb{E}[e^{tX}e^{tY}] = \mathbb{E}[e^{tX}]\mathbb{E}[e^{tY}]$.

(3) Let $|t| < t_0$, and choose a with $|t| < a < t_0$; let $s = \frac{1}{2}(a - |t|)$, and $0 < |h| \leq s$. For every $k \geq 0$, the number $c_k = \max_{x \in \mathbb{R}} |x|^{k+2} e^{-s|x|}$ is finite. By Taylor's theorem (Lecture 3), $|e^u - 1 - u| \leq \frac{1}{2} u^2 e^{|u|}$, so for all x,

$$\begin{aligned} \big|x^k \big(e^{(t+h)x} - e^{tx} - hxe^{tx}\big)\big| = |x|^k e^{tx} \left|e^{hx} - 1 - hx\right| &\leq \frac{h^2}{2} |x|^{k+2} e^{(|t|+s)|x|} \\ &\leq \frac{h^2 c_k}{2}\, e^{a|x|} \leq \frac{h^2 c_k}{2} \big(e^{ax} + e^{-ax}\big). \end{aligned}$$

Similarly, $|x^k e^{tx}| \leq c_k'(e^{ax} + e^{-ax})$ with $c_k' = \max_x |x|^k e^{-(a-|t|)|x|}$, so $m_k(t) = \mathbb{E}[X^k e^{tX}]$ is finite. Taking expectations, $|m_k(t+h) - m_k(t) - h\, m_{k+1}(t)| \leq \frac{h^2 c_k}{2}\big(M_X(a) + M_X(-a)\big)$, so m_k is differentiable with $m_k' = m_{k+1}$. Since $m_0 = M_X$, induction gives $M_X^{(k)} = m_k$. Finally, $K_X' = \frac{M_X'}{M_X}$ and $K_X'' = \frac{M_X''}{M_X} - \big(\frac{M_X'}{M_X}\big)^2$, which at $t = 0$ give $\mathbb{E}[X]$ and $\mathbb{E}[X^2] - \mathbb{E}[X]^2$. □

For example, for $X \sim \text{Poisson}(\lambda)$, $K_X(t) = \lambda(e^t - 1)$ has $K'_X(0) = K''_X(0) = \lambda$: the mean and the variance are both λ. For $S \sim \text{Binomial}(n, p)$, a sum of n independent Bernoulli variables, $M_S(t) = (1 - p + pe^t)^n$, and $K_S(t) = n \log(1 - p + pe^t)$ gives $\mathbb{E}[S] = np$ and $\text{Var}(S) = np(1 - p)$. Cumulant generating functions of independent variables add, just like means and variances. This gives a heuristic explanation of the central limit theorem: for i.i.d. standardized X_i (with mean 0 and variance 1), $K_{(X_1+\cdots+X_N)/\sqrt{N}}(t) = NK_X(t/\sqrt{N}) \approx N\frac{t^2}{2N} = \frac{t^2}{2}$ by Taylor's theorem, since $K_X(0) = K'_X(0) = 0$ and $K''_X(0) = 1$, and $\frac{t^2}{2}$ is the cumulant generating function of $\mathcal{N}(0, 1)$.

Chernoff bounds. By Markov's inequality, $P(X \geq a) \leq e^{-ta} M_X(t) = e^{-(ta - K_X(t))}$ for every $t > 0$, so

$$P(X \geq a) \leq \exp\Big(- \sup_{t>0} \big(ta - K_X(t)\big)\Big).$$

For $X \sim \mathcal{N}(0, \sigma^2)$ and $a > 0$, the supremum of $ta - \frac{\sigma^2 t^2}{2}$ is $\frac{a^2}{2\sigma^2}$, attained at $t = \frac{a}{\sigma^2}$, so $P(X \geq a) \leq e^{-a^2/(2\sigma^2)}$. Random variables with $K_X(t) \leq \frac{\sigma^2 t^2}{2}$ for all t, i.e., with tails at least as light as those of $\mathcal{N}(0, \sigma^2)$, are called sub-Gaussian. Hoeffding's lemma (Lemma 4.7.6) says that a bounded variable with mean 0 and values in $[a, b]$ is sub-Gaussian with $\sigma^2 = \frac{(b-a)^2}{4}$, and the proof of Hoeffding's inequality adds the cumulant generating functions of the summands.

4.5 Random Vectors and Covariance Matrices

A random vector $\mathbf{X} = (X_1, \ldots, X_d)^\top$ is a vector of random variables, such as the features of a randomly chosen data point. Its expectation is taken entrywise.

Definition 4.5.1 (Mean and covariance matrix)**.** The mean of a random vector $\mathbf{X}$ is $\boldsymbol{\mu} = \mathbb{E}[\mathbf{X}] = (\mathbb{E}[X_1], \ldots, \mathbb{E}[X_d])^\top$, and its covariance matrix is the $d \times d$ matrix

$$\Sigma = \text{Cov}(\mathbf{X}) = \mathbb{E}\big[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top\big], \qquad \text{with entries } \Sigma_{ij} = \text{Cov}(X_i, X_j).$$

The diagonal entries of Σ are the variances $\text{Var}(X_i)$. Since $(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top$ is an outer product, linear algebra enters immediately.

Proposition 4.5.2. *Let $\mathbf{X}$ have mean $\boldsymbol{\mu}$ and covariance matrix Σ.*

1. $\Sigma = \mathbb{E}[\mathbf{X}\mathbf{X}^\top] - \boldsymbol{\mu}\boldsymbol{\mu}^\top$.
2. *For every matrix $A \in \mathbb{R}^{m \times d}$ and vector $\mathbf{b} \in \mathbb{R}^m$, $\mathbb{E}[A\mathbf{X} + \mathbf{b}] = A\boldsymbol{\mu} + \mathbf{b}$ and $\text{Cov}(A\mathbf{X} + \mathbf{b}) = A\Sigma A^\top$.*
3. *For every $\mathbf{u} \in \mathbb{R}^d$, $\text{Var}(\mathbf{u}^\top \mathbf{X}) = \mathbf{u}^\top \Sigma \mathbf{u}$. In particular, Σ is symmetric and positive semidefinite.*

Proof. (2) Each entry of $A\mathbf{X} + \mathbf{b}$ is a linear combination of the X_j plus a constant, so linearity of expectation gives $\mathbb{E}[A\mathbf{X} + \mathbf{b}] = A\boldsymbol{\mu} + \mathbf{b}$, and in the same way $\mathbb{E}[AMB] = A\,\mathbb{E}[M]\,B$ for a random matrix M and fixed matrices A and B. Hence

$$\text{Cov}(A\mathbf{X} + \mathbf{b}) = \mathbb{E}\big[A(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top A^\top\big] = A\Sigma A^\top.$$

(1) is proved like the formula $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$. (3) is (2) with $A = \mathbf{u}^\top$, and $\text{Var}(\mathbf{u}^\top \mathbf{X}) \geq 0$; Σ is symmetric since $\text{Cov}(X_i, X_j) = \text{Cov}(X_j, X_i)$. □

By (3), Σ fails to be positive definite exactly if $\text{Var}(\mathbf{u}^\top \mathbf{X}) = 0$ for some $\mathbf{u} \neq \mathbf{0}$, i.e., if $\mathbf{u}^\top \mathbf{X}$ is constant: then $\mathbf{X}$ lies in an affine hyperplane (with probability 1). Part (3) also describes the variance in every direction: for a unit vector $\mathbf{u}$, the variance of the projection $\mathbf{u}^\top \mathbf{X}$ is $\mathbf{u}^\top \Sigma \mathbf{u}$. By the

Rayleigh quotient (Lecture 2), it is largest for an eigenvector of Σ with the largest eigenvalue. This is the population version of principal component analysis, and the total variance $\sum_i \operatorname{Var}(X_i) = \operatorname{tr}\Sigma$ is the sum of the eigenvalues of Σ.

Example 4.5.3. Let $\Sigma = \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix}$. Then $\operatorname{Var}(X_1 + X_2) = (1,1)\Sigma(1,1)^\top = 4 + 3 + 2 \cdot 2 = 11$, $\operatorname{Var}(X_1 - X_2) = 4 + 3 - 4 = 3$, and $\operatorname{Corr}(X_1, X_2) = \frac{2}{\sqrt{12}} \approx 0.58$. The eigenvalues of Σ are $\frac{7 \pm \sqrt{17}}{2} \approx 5.56$ and 1.44: the variance in the direction of the first principal component is 5.56, out of the total variance $\operatorname{tr}\Sigma = 7$.

Remark 4.5.4 (Moments as tensors). In the language of the section on tensors in Lecture 1, $(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top = (\mathbf{X} - \boldsymbol{\mu}) \otimes (\mathbf{X} - \boldsymbol{\mu})$, so the covariance matrix is the expectation of a random tensor in $\mathbb{R}^d \otimes \mathbb{R}^d$. Part 2 of Proposition 4.5.2 says that it transforms like such a tensor: in Einstein notation, $\operatorname{Cov}(A\mathbf{X})_{ab} = A_{ai}A_{bj}\Sigma_{ij}$, which is $A\Sigma A^\top$. Higher moments are tensors of higher order. The third central moment $T = \mathbb{E}\big[(\mathbf{X} - \boldsymbol{\mu}) \otimes (\mathbf{X} - \boldsymbol{\mu}) \otimes (\mathbf{X} - \boldsymbol{\mu})\big]$ has the entries $T_{ijk} = \mathbb{E}[(X_i - \mu_i)(X_j - \mu_j)(X_k - \mu_k)]$, it changes to $A_{ai}A_{bj}A_{ck}T_{ijk}$ under $\mathbf{X} \mapsto A\mathbf{X}$, and it is 0 for every distribution that is symmetric about its mean, such as a Gaussian (since $\mathbf{X} - \boldsymbol{\mu}$ and $\boldsymbol{\mu} - \mathbf{X}$ have the same distribution, $T = -T$). Moment tensors are computed with `np.einsum`, and their decompositions into sums of pure tensors (the tensor rank of Lecture 1) are used to estimate the parameters of mixture models.

Definition 4.5.5 (Whitening and Mahalanobis distance)**.** Let Σ be symmetric positive definite, with spectral decomposition $\Sigma = Q\Lambda Q^\top$ (Lecture 2), and let $\Sigma^{-1/2} = Q\Lambda^{-1/2}Q^\top$, where $\Lambda^{-1/2} = \operatorname{diag}(\lambda_1^{-1/2}, \dots, \lambda_d^{-1/2})$. The whitened random vector is $\mathbf{Z} = \Sigma^{-1/2}(\mathbf{X} - \boldsymbol{\mu})$, and the Mahalanobis distance of $\mathbf{x}$ from $\boldsymbol{\mu}$ is

$$d_\Sigma(\mathbf{x}, \boldsymbol{\mu}) = \sqrt{(\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})} = \|\Sigma^{-1/2}(\mathbf{x} - \boldsymbol{\mu})\|.$$

The matrix $\Sigma^{-1/2}$ is symmetric with $(\Sigma^{-1/2})^2 = \Sigma^{-1}$, so by Proposition 4.5.2, $\operatorname{Cov}(\mathbf{Z}) = \Sigma^{-1/2}\Sigma\Sigma^{-1/2} = I$: whitening removes all correlations and scales all directions to unit variance. The Mahalanobis distance is the norm of the inner product $\langle \mathbf{x}, \mathbf{y} \rangle_{\Sigma^{-1}} = \mathbf{x}^\top \Sigma^{-1} \mathbf{y}$ (Lecture 2): it measures distances in units of standard deviations along each principal axis. Standardizing or whitening the features of a data set is a common preprocessing step; by Lecture 3, it improves the condition number and speeds up gradient descent.

Sample statistics. Given data points $\mathbf{x}_1, \dots, \mathbf{x}_N \in \mathbb{R}^d$, the sample mean and the sample covariance matrix are

$$\bar{\mathbf{x}} = \frac{1}{N} \sum_{i=1}^{N} \mathbf{x}_i \qquad \text{and} \qquad S = \frac{1}{N} \sum_{i=1}^{N} (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top = \frac{1}{N} X_c^\top X_c,$$

where X_c is the centered data matrix with rows $(\mathbf{x}_i - \bar{\mathbf{x}})^\top$. These are the mean and covariance matrix of the empirical distribution, which gives probability $\frac{1}{N}$ to each data point; principal component analysis in Lecture 2 used exactly this matrix. In Section 4.9, we will see why one often divides by $N - 1$ instead of N.

4.6 The Gaussian Distribution

The Gaussian (normal) distribution is the most important distribution in statistics and machine learning. It appears as the limit of sums of many small random effects (Section 4.7), it is completely described by its mean and covariance matrix, and its marginals and conditionals can be computed with linear algebra.

Definition 4.6.1 (Multivariate Gaussian distribution)**.** Let $\boldsymbol{\mu} \in \mathbb{R}^d$, and let Σ be a symmetric positive definite $d \times d$ matrix. A random vector $\mathbf{X}$ has the Gaussian distribution $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ if it has the density

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} \det(\Sigma)^{1/2}} \exp\Big(-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})\Big).$$

For $d = 1$ and $\Sigma = \sigma^2$, this is the density of Section 4.2. For $\boldsymbol{\mu} = \mathbf{0}$ and $\Sigma = I$, the density is $(2\pi)^{-d/2} e^{-\|\mathbf{z}\|^2/2} = \prod_i \varphi(z_i)$: a standard Gaussian vector $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$ has i.i.d. standard normal entries. All other Gaussians are obtained from it by affine maps.

Theorem 4.6.2. *Let $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$, let A be an invertible $d \times d$ matrix, and let $\boldsymbol{\mu} \in \mathbb{R}^d$. Then $\mathbf{X} = A\mathbf{Z} + \boldsymbol{\mu} \sim \mathcal{N}(\boldsymbol{\mu}, AA^\top)$. Consequently, the density of Definition 4.6.1 integrates to* 1*, and $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ has mean $\boldsymbol{\mu}$ and covariance matrix Σ.*

Proof. By the change of variables formula (Lecture 3), with $\mathbf{z} = A^{-1}(\mathbf{x} - \boldsymbol{\mu})$, whose Jacobian has the determinant $\det(A)^{-1}$,

$$p_{\mathbf{X}}(\mathbf{x}) = p_{\mathbf{Z}}(A^{-1}(\mathbf{x}-\boldsymbol{\mu}))\, \frac{1}{|\det A|} = \frac{1}{(2\pi)^{d/2}|\det A|} \exp\Big(-\frac{1}{2}\|A^{-1}(\mathbf{x}-\boldsymbol{\mu})\|^2\Big).$$

With $\Sigma = AA^\top$, we have $\|A^{-1}\mathbf{y}\|^2 = \mathbf{y}^\top A^{-\top} A^{-1}\mathbf{y} = \mathbf{y}^\top \Sigma^{-1}\mathbf{y}$ and $\det \Sigma = \det(A)^2$, which gives the density of $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$. Every symmetric positive definite Σ can be written as $AA^\top$, for example with $A = \Sigma^{1/2} = Q\Lambda^{1/2}Q^\top$. So every Gaussian density is the density of some random vector, and integrates to 1. By Proposition 4.5.2, $\mathbb{E}[\mathbf{X}] = A\mathbb{E}[\mathbf{Z}] + \boldsymbol{\mu} = \boldsymbol{\mu}$ and $\mathrm{Cov}(\mathbf{X}) = A\, I\, A^\top = \Sigma$. □

This is also how Gaussian vectors are sampled: compute a factorization $\Sigma = AA^\top$ (in practice, the Cholesky factorization $\Sigma = LL^\top$ with a lower triangular L, as in `np.linalg.cholesky`), draw $\mathbf{z}$ with i.i.d. standard normal entries, and return $\boldsymbol{\mu} + L\mathbf{z}$. If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ and B is invertible, then $B\mathbf{X} + \mathbf{b} = (BA)\mathbf{Z} + (B\boldsymbol{\mu} + \mathbf{b})$, so $B\mathbf{X} + \mathbf{b} \sim \mathcal{N}(B\boldsymbol{\mu} + \mathbf{b}, B\Sigma B^\top)$.

Remark 4.6.3 (The reparameterization trick). Writing a Gaussian random vector as $\boldsymbol{\mu} + \boldsymbol{\sigma} \odot \mathbf{Z}$ with a fixed source of randomness $\mathbf{Z}$ makes it a differentiable function of the parameters $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}$. Variational autoencoders use this to backpropagate (Lecture 3) through random sampling steps.

The density of $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ is constant on the ellipsoids $(\mathbf{x}-\boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu}) = r^2$, i.e., on the sets of points with Mahalanobis distance r from $\boldsymbol{\mu}$. By the spectral theorem (Lecture 2), their axes point in the directions of the eigenvectors $\mathbf{q}_i$ of Σ, with half-lengths $r\sqrt{\lambda_i}$, i.e., r standard deviations in the direction $\mathbf{q}_i$ (Figure 4.2).

Theorem 4.6.4 (Independence of blocks)**.** *Let $\mathbf{X} = (\mathbf{X}_a, \mathbf{X}_b) \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$, and let $\Sigma = \left(\begin{smallmatrix} \Sigma_{aa} & 0 \\ 0 & \Sigma_{bb} \end{smallmatrix}\right)$ be block diagonal. Then $\mathbf{X}_a$ and $\mathbf{X}_b$ are independent, with $\mathbf{X}_a \sim \mathcal{N}(\boldsymbol{\mu}_a, \Sigma_{aa})$ and $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b, \Sigma_{bb})$.*

Proof. Σ^{-1} is block diagonal with blocks Σ_{aa}^{-1} and Σ_{bb}^{-1}, and $\det \Sigma = \det \Sigma_{aa} \det \Sigma_{bb}$. Hence the quadratic form in the exponent splits into a sum of two terms, and the density is the product of the densities of $\mathcal{N}(\boldsymbol{\mu}_a, \Sigma_{aa})$ and $\mathcal{N}(\boldsymbol{\mu}_b, \Sigma_{bb})$. Integrating over $\mathbf{x}_b$ shows that the first factor is the marginal density of $\mathbf{X}_a$, and similarly for $\mathbf{X}_b$. □

So for jointly Gaussian random vectors, uncorrelated parts are independent. This is special to Gaussians, and it is the key to the following theorem, which is used constantly in Gaussian processes, Kalman filters and Bayesian linear regression.

Theorem 4.6.5 (Marginals and conditionals)**.** *Let $\mathbf{X} = (\mathbf{X}_a, \mathbf{X}_b) \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ with*

$$\boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_a \\ \boldsymbol{\mu}_b \end{pmatrix}, \qquad \Sigma = \begin{pmatrix} \Sigma_{aa} & \Sigma_{ab} \\ \Sigma_{ba} & \Sigma_{bb} \end{pmatrix}.$$

Then $\mathbf{X}_a \sim \mathcal{N}(\boldsymbol{\mu}_a, \Sigma_{aa})$, $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b, \Sigma_{bb})$, and the conditional distribution of $\mathbf{X}_a$ given $\mathbf{X}_b = \mathbf{x}_b$ is

$$\mathcal{N}(\boldsymbol{\mu}_a + \Sigma_{ab}\Sigma_{bb}^{-1}(\mathbf{x}_b - \boldsymbol{\mu}_b),\ \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}).$$

Proof. The block Σ_{bb} is positive definite (since $\mathbf{u}^\top \Sigma_{bb}\mathbf{u} = (\mathbf{0},\mathbf{u})^\top \Sigma (\mathbf{0},\mathbf{u}) > 0$ for $\mathbf{u} \neq \mathbf{0}$), so $K = \Sigma_{ab}\Sigma_{bb}^{-1}$ is defined. Let $\mathbf{W} = \mathbf{X}_a - K\mathbf{X}_b$, i.e.,

$$\begin{pmatrix} \mathbf{W} \\ \mathbf{X}_b \end{pmatrix} = B\mathbf{X}, \qquad B = \begin{pmatrix} I & -K \\ 0 & I \end{pmatrix}.$$

B is invertible, so $(\mathbf{W}, \mathbf{X}_b)$ is Gaussian with covariance matrix $B\Sigma B^\top$. Its off-diagonal block is $\mathrm{Cov}(\mathbf{W}, \mathbf{X}_b) = \Sigma_{ab} - K\Sigma_{bb} = 0$, and its upper left block is

$$\mathrm{Cov}(\mathbf{W}) = \Sigma_{aa} - K\Sigma_{ba} - \Sigma_{ab}K^\top + K\Sigma_{bb}K^\top = \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}.$$

By Theorem 4.6.4, $\mathbf{X}_b \sim \mathcal{N}(\boldsymbol{\mu}_b, \Sigma_{bb})$, and $\mathbf{W}$ is independent of $\mathbf{X}_b$, with $\mathbf{W} \sim \mathcal{N}(\boldsymbol{\mu}_a - K\boldsymbol{\mu}_b, \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba})$. Since $\det B = 1$, the joint density of $\mathbf{X}$ is $p(\mathbf{x}_a, \mathbf{x}_b) = p_{\mathbf{W}}(\mathbf{x}_a - K\mathbf{x}_b)\, p_{\mathbf{X}_b}(\mathbf{x}_b)$, so the conditional density is $p(\mathbf{x}_a \mid \mathbf{x}_b) = p_{\mathbf{W}}(\mathbf{x}_a - K\mathbf{x}_b)$: given $\mathbf{X}_b = \mathbf{x}_b$, $\mathbf{X}_a = \mathbf{W} + K\mathbf{x}_b$, which has the claimed distribution. The marginal of $\mathbf{X}_a$ follows by exchanging the roles of a and b. □

The conditional mean is an affine function of $\mathbf{x}_b$, and the conditional covariance $\Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}$ (the Schur complement of Σ_{bb}, as in the subsection on block matrices in Lecture 1, with the roles of the two blocks exchanged) does not depend on $\mathbf{x}_b$. It is never larger than Σ_{aa}, since $\Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}$ is positive semidefinite: observing $\mathbf{X}_b$ reduces the uncertainty about $\mathbf{X}_a$.

Example 4.6.6 (Regression to the mean)**.** Let (X, Y) be Gaussian with means 0, variances 1, and correlation ρ. By Theorem 4.6.5, given $X = x$, $Y \sim \mathcal{N}(\rho x, 1 - \rho^2)$. For example, for $\rho = 0.5$ and $x = 2$, $Y \sim \mathcal{N}(1, 0.75)$. If X and Y are the heights of parents and their children (measured in standard deviations from the mean), then children of very tall parents are on average tall, but less extreme than their parents. Francis Galton observed this "regression to the mean" in 1886, and it gave linear regression its name. Note that the regression line $y = \rho x$ is not the major axis of the ellipses of constant density, which is the line $y = x$ (Figure 4.2).




Figure 4.2: Level curves of the density of $\mathcal{N}(\mathbf{0}, \Sigma)$ with $\Sigma = \left(\begin{smallmatrix} 3 & 1 \\ 1 & 3 \end{smallmatrix}\right)$, for Mahalanobis distances $r = 0.5, 1, 1.5, 2$. The eigenvectors $\mathbf{q}_1 = \frac{1}{\sqrt{2}}(1,1)$ and $\mathbf{q}_2 = \frac{1}{\sqrt{2}}(1,-1)$ of Σ have the eigenvalues 4 and 2, so the axes have half-lengths $2r$ and $\sqrt{2}\,r$ (orange: one standard deviation in each direction). The conditional mean $\mathbb{E}[X_2 \mid X_1 = x_1] = \frac{\Sigma_{21}}{\Sigma_{11}}x_1 = \frac{1}{3}x_1$ (red) is flatter than the major axis $x_2 = x_1$ (dashed).

Corollary 4.6.7. *Let $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$, and let $A \in \mathbb{R}^{m \times d}$ have rank m. Then $A\mathbf{X} + \mathbf{b} \sim \mathcal{N}(A\boldsymbol{\mu} + \mathbf{b}, A\Sigma A^\top)$. In particular, if $X \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$ are independent, then $X + Y \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.*

Proof. The rows of A are linearly independent, so they can be extended to a basis of $\mathbb{R}^d$ (Lecture 1): there is a matrix C such that $B = \binom{A}{C}$ is invertible. Then $B\mathbf{X}+(\mathbf{b},\mathbf{0})$ is Gaussian, and $A\mathbf{X}+\mathbf{b}$ is its first block, which is Gaussian by Theorem 4.6.5, with the mean and covariance of Proposition 4.5.2. For the second statement, (X, Y) is Gaussian with the covariance matrix $\operatorname{diag}(\sigma_1^2, \sigma_2^2)$ (its density is the product of the two densities), and $X + Y = (1, 1)(X, Y)^\top$. □

4.7 Limit Theorems and Monte Carlo Methods

Machine learning relies on averages: the training loss is an average over the data, a mini-batch gradient is an average over a few examples, and the test accuracy is an average over the test set. The limit theorems explain why averages of many random quantities are reliable.

Proposition 4.7.1. *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and variance σ^2, and let $\bar{X}_N = \frac{1}{N}(X_1 + \cdots + X_N)$ be their sample mean. Then*

$$\mathbb{E}[\bar{X}_N] = \mu \qquad \text{and} \qquad \operatorname{Var}(\bar{X}_N) = \frac{\sigma^2}{N}.$$

Proof. By linearity, $\mathbb{E}[\bar{X}_N] = \frac{1}{N} \cdot N\mu$, and by Proposition 4.4.9, $\operatorname{Var}(\bar{X}_N) = \frac{1}{N^2} \cdot N\sigma^2$. □

So the standard deviation of the sample mean, its standard error, is $\frac{\sigma}{\sqrt{N}}$: to halve the error, we need four times as much data.

Theorem 4.7.2 (Weak law of large numbers)**.** *In the situation of Proposition 4.7.1, for every $\varepsilon > 0$,*

$$P(|\bar{X}_N - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{N\varepsilon^2} \to 0 \qquad \text{as } N \to \infty.$$

Proof. Chebyshev's inequality (Proposition 4.3.6) for $\bar{X}_N$, which has mean μ and variance $\frac{\sigma^2}{N}$. □

The law of large numbers says that the sample mean is very likely close to μ for large N. (The strong law of large numbers, which we discuss in Appendix B, even says that $\bar{X}_N \to \mu$ with probability 1.) The central limit theorem describes the size and shape of the remaining fluctuations: they are approximately Gaussian, whatever the distribution of the X_i.

Theorem 4.7.3 (Central limit theorem)**.** *In the situation of Proposition 4.7.1, with $\sigma > 0$, for every $z \in \mathbb{R}$,*

$$P\Big(\frac{\sqrt{N}(\bar{X}_N - \mu)}{\sigma} \leq z\Big) \to \Phi(z) \qquad \text{as } N \to \infty.$$

We prove the central limit theorem in Section 4.7.1, after introducing convergence in distribution. Informally, $\bar{X}_N$ is approximately $\mathcal{N}(\mu, \frac{\sigma^2}{N})$, and the sum $X_1 + \cdots + X_N$ is approximately $\mathcal{N}(N\mu, N\sigma^2)$. This explains why Gaussian distributions are so common: a measurement error that is the sum of many small independent effects is approximately Gaussian.

Example 4.7.4. We flip a fair coin 100 times. The number of heads S is Binomial$(100, \frac{1}{2})$, with mean 50 and standard deviation $\sqrt{100 \cdot \frac{1}{4}} = 5$. By the central limit theorem, $P(S \geq 60) = P(S \geq 59.5) \approx 1 - \Phi\big(\frac{59.5-50}{5}\big) = 1 - \Phi(1.9) \approx 0.0287$, while the exact value is 0.0284. (Replacing 60 by 59.5, the "continuity correction", improves the approximation of a discrete distribution by a continuous one.) Figure 4.3 compares a binomial distribution with its Gaussian approximation.




Figure 4.3: The pmf of the Binomial$(40, 0.3)$ distribution, the sum of 40 independent Bernoulli(0.3) variables, and the density of its Gaussian approximation $\mathcal{N}(12, 8.4)$ with the same mean and variance (red).

Monte Carlo estimation. To compute an expectation $\mathbb{E}[g(X)]$ that we cannot compute exactly, we can draw i.i.d. samples $x_1, \dots, x_N$ of X and use the average $\frac{1}{N}\sum_i g(x_i)$. By the central limit theorem, its error is approximately Gaussian with standard deviation $\frac{\sigma_g}{\sqrt{N}}$, where σ_g is the standard deviation of $g(X)$, no matter how high-dimensional X is. For example, the fraction of N uniformly random points of the square $[-1,1]^2$ that land in the unit disk estimates $\frac{\pi}{4}$, so 4 times this fraction estimates π, with standard error $4\sqrt{\frac{\pi}{4}(1-\frac{\pi}{4})/N}$, which is 0.016 for $N = 10\,000$. Each additional correct digit costs 100 times more samples.

Concentration inequalities. The central limit theorem is only an approximation for finite N. For bounded random variables, there is an exact bound that is much stronger than Chebyshev's inequality.

Theorem 4.7.5 (Hoeffding's inequality)**.** *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and values in $[a, b]$. Then for every $\varepsilon > 0$,*

$$P(|\bar{X}_N - \mu| \geq \varepsilon) \leq 2\exp\Big(-\frac{2N\varepsilon^2}{(b-a)^2}\Big).$$

The proof uses the Chernoff bound of Example 4.3.8, together with a bound for the expectation of an exponential.

Lemma 4.7.6 (Hoeffding's lemma)**.** *Let X have values in $[a, b]$ and $\mathbb{E}[X] = 0$. Then $\mathbb{E}[e^{tX}] \leq e^{t^2(b-a)^2/8}$ for every $t \in \mathbb{R}$.*

Proof. If $a = b$, then $X = 0$. Otherwise, $a \leq 0 \leq b$, since $\mathbb{E}[X] = 0$. The function $x \mapsto e^{tx}$ is convex, so on $[a, b]$ it lies below the chord between (a, e^{ta}) and (b, e^{tb}):

$$e^{tx} \leq \frac{b-x}{b-a}\,e^{ta} + \frac{x-a}{b-a}\,e^{tb} \qquad \text{for } a \leq x \leq b.$$

Taking expectations and using $\mathbb{E}[X] = 0$ gives, with $p = \frac{-a}{b-a} \in [0,1]$ and $u = t(b-a)$, so that $ta = -pu$,

$$\mathbb{E}[e^{tX}] \leq \frac{b\,e^{ta} - a\,e^{tb}}{b-a} = e^{ta}(1 - p + pe^u) = e^{L(u)}, \qquad L(u) = -pu + \log\,(1 - p + pe^u).$$

We have $L(0) = 0$ and $L'(u) = -p + q(u)$ with $q(u) = \frac{pe^u}{1-p+pe^u} \in [0,1]$, so $L'(0) = 0$, and $L''(u) = q(u)(1-q(u)) \leq \frac{1}{4}$. By Taylor's theorem (Lecture 3), $L(u) = \frac{1}{2}L''(\xi)u^2 \leq \frac{u^2}{8} = \frac{t^2(b-a)^2}{8}$ for some ξ. □

Proof of Theorem 4.7.5. Let $t > 0$ and $S = \sum_i (X_i - \mu)$. Each $X_i - \mu$ has mean 0 and values in the interval $[a - \mu, b - \mu]$ of length $b - a$. By Markov's inequality for e^{tS} (as in Example 4.3.8), independence and Lemma 4.7.6,

$$P(\bar{X}_N - \mu \geq \varepsilon) = P(S \geq N\varepsilon) \leq e^{-tN\varepsilon}\,\mathbb{E}[e^{tS}] = e^{-tN\varepsilon} \prod_{i=1}^{N} \mathbb{E}\big[e^{t(X_i-\mu)}\big] \leq \exp\Big(-tN\varepsilon + \frac{Nt^2(b-a)^2}{8}\Big).$$

The choice $t = \frac{4\varepsilon}{(b-a)^2}$ gives the bound $\exp\big(-\frac{2N\varepsilon^2}{(b-a)^2}\big)$. The variables $-X_i$, with values in $[-b, -a]$, give the same bound for $P(\bar{X}_N - \mu \leq -\varepsilon)$, and the union bound gives the factor 2. □

For example, let a classifier have the (unknown) error rate μ, and let $\bar{X}_N$ be its error rate on a test set of $N = 1000$ examples drawn independently of the training data, where $X_i \in \{0, 1\}$ indicates an error. Then $P(|\bar{X}_N - \mu| \geq 0.05) \leq 2e^{-5} \approx 0.013$, while Chebyshev's inequality with $\sigma^2 \leq \frac{1}{4}$ only gives 0.1.

Remark 4.7.7 (Limit theorems in machine learning).

1. The average loss on the training data, $\frac{1}{N}\sum_i \ell(\boldsymbol{\theta}; \mathbf{x}_i, y_i)$, is a sample mean. It estimates the expected loss $\mathbb{E}[\ell(\boldsymbol{\theta}; \mathbf{X}, Y)]$ on new data, which is the quantity we actually care about. For a fixed $\boldsymbol{\theta}$, the limit theorems apply. But the trained parameters depend on the training data, so the training loss of the trained model is a biased, usually too optimistic, estimate (overfitting). This is why we evaluate on separate test data, and why learning theory combines Hoeffding's inequality with the union bound (Proposition 4.1.3) to control all candidate models simultaneously (Section 4.11.3).

2. In stochastic gradient descent (Lecture 3), the mini-batch gradient of B randomly chosen examples is an unbiased estimate of the full gradient, and its covariance matrix decreases like $\frac{1}{B}$ (for sampling with replacement), by Proposition 4.7.1 applied to each coordinate. Larger batches give less noisy gradients, but each step costs more.

4.7.1 Slutsky's theorem and the delta method

The law of large numbers and the central limit theorem are statements about limits of probabilities. We give names to these two kinds of convergence.

Definition 4.7.8 (Convergence in probability and in distribution)**.** Let $Y, Y_1, Y_2, \ldots$ be random variables. We say that $Y_n \to Y$

1. in probability if $P(|Y_n - Y| > \varepsilon) \to 0$ for every $\varepsilon > 0$;
2. in distribution if $P(Y_n \leq y) \to P(Y \leq y)$ at every point y at which the CDF of Y is continuous.

Convergence in distribution only depends on the distributions, and we also write $Y_n \to \mathcal{N}(0, 1)$, for example. The law of large numbers (Theorem 4.7.2) says that $\bar{X}_N \to \mu$ in probability, and the central limit theorem (Theorem 4.7.3) says that $\frac{\sqrt{N}(\bar{X}_N - \mu)}{\sigma} \to \mathcal{N}(0, 1)$ in distribution; Appendix B discusses further modes of convergence. Convergence in distribution can also be tested with expectations of functions.

Lemma 4.7.9 (Test functions)**.** *Let $Y, Y_1, Y_2, \ldots$ be random variables.*

1. *If $\mathbb{E}[h(Y_n)] \to \mathbb{E}[h(Y)]$ for every three times continuously differentiable function $h : \mathbb{R} \to \mathbb{R}$ for which h, h', h'' and h''' are bounded, then $Y_n \to Y$ in distribution.*
2. *Conversely, if $Y_n \to Y$ in distribution, then $\mathbb{E}[h(Y_n)] \to \mathbb{E}[h(Y)]$ for every bounded continuous function h.*

Proof. Let F and F_n be the CDFs of Y and Y_n. A CDF has at most countably many points of discontinuity, since it has at most k jumps of size larger than $\frac{1}{k}$ (the jumps add up to at most 1). Hence every interval contains continuity points of F.

(1) Let $\psi(s) = 1$ for $s \le 0$, $\psi(s) = 0$ for $s \ge 1$, and $\psi(s) = 1 - 35s^4 + 84s^5 - 70s^6 + 20s^7$ for $0 < s < 1$. Then $\psi'(s) = -140s^3(1-s)^3 \le 0$ on $(0,1)$, so the derivatives of orders 1, 2 and 3 of the polynomial vanish at 0 and 1, and ψ is a nonincreasing, three times continuously differentiable function with values in $[0,1]$ and bounded derivatives. Let y be a continuity point of F, and let $\delta > 0$. The function $h(x) = \psi(\frac{x-y}{\delta})$ satisfies $\mathbf{1}_{\{x \le y\}} \le h(x) \le \mathbf{1}_{\{x \le y+\delta\}}$, so

$$P(Y_n \le y) \le \mathbb{E}[h(Y_n)] \to \mathbb{E}[h(Y)] \le P(Y \le y + \delta) = F(y+\delta).$$

Similarly, $\tilde{h}(x) = \psi(\frac{x-y+\delta}{\delta})$ satisfies $\mathbf{1}_{\{x \le y-\delta\}} \le \tilde{h}(x) \le \mathbf{1}_{\{x \le y\}}$, which gives $P(Y_n \le y) \ge \mathbb{E}[\tilde{h}(Y_n)] \to \mathbb{E}[\tilde{h}(Y)] \ge F(y-\delta)$. So all limit points of the sequence $P(Y_n \le y)$ lie between $F(y-\delta)$ and $F(y+\delta)$, and these tend to $F(y)$ as $\delta \to 0$, since F is continuous at y.

(2) Let $|h| \le C$, and let $\varepsilon > 0$. Choose continuity points $a < b$ of F with $F(a) < \varepsilon$ and $1 - F(b) < \varepsilon$. The function h is uniformly continuous on $[a,b]$ (by Proposition A.3.8), so there are continuity points $a = t_0 < t_1 < \cdots < t_K = b$ of F such that $|h(x) - h(t_k)| \le \varepsilon$ for $t_{k-1} \le x \le t_k$. Let s be the step function with $s(x) = h(t_k)$ for $t_{k-1} < x \le t_k$, and $s(x) = 0$ for $x \le a$ and for $x > b$. Then $|h - s| \le \varepsilon$ on $(a,b]$ and $|h - s| \le C$ elsewhere, so for every random variable V,

$$\big|\mathbb{E}[h(V)] - \mathbb{E}[s(V)]\big| \le \varepsilon + C\,P(V \le a) + C\,P(V > b).$$

Now $\mathbb{E}[s(Y_n)] = \sum_k h(t_k)\big(F_n(t_k) - F_n(t_{k-1})\big) \to \sum_k h(t_k)\big(F(t_k) - F(t_{k-1})\big) = \mathbb{E}[s(Y)]$, and $P(Y_n \le a) + P(Y_n > b) = F_n(a) + 1 - F_n(b) \to F(a) + 1 - F(b) < 2\varepsilon$. Hence all limit points of $\big|\mathbb{E}[h(Y_n)] - \mathbb{E}[h(Y)]\big|$ are at most $2(\varepsilon + 2C\varepsilon)$. Since ε was arbitrary, $\mathbb{E}[h(Y_n)] \to \mathbb{E}[h(Y)]$. □

Proof of the central limit theorem (Theorem 4.7.3). Let $Y_i = \frac{X_i - \mu}{\sigma}$. These are i.i.d. with mean 0 and variance 1, and $S_N = \frac{\sqrt{N}(\bar{X}_N - \mu)}{\sigma} = \frac{1}{\sqrt{N}}(Y_1 + \cdots + Y_N)$. Let $Z \sim \mathcal{N}(0,1)$. Since Φ is continuous, by Lemma 4.7.9 it suffices to show that $\mathbb{E}[h(S_N)] \to \mathbb{E}[h(Z)]$ for every h as in part 1 of the lemma; let C_2 and C_3 be bounds for $|h''|$ and $|h'''|$.

A Taylor bound. Let $r(w,x) = h(w+x) - h(w) - h'(w)x - \frac{1}{2}h''(w)x^2$. Taylor's theorem (Lecture 3) with $k = 2$ gives $r(w,x) = \frac{1}{6}h'''(\xi)x^3$, and with $k = 1$ it gives $r(w,x) = \frac{1}{2}\big(h''(\xi') - h''(w)\big)x^2$, for some ξ and ξ' between w and $w + x$. Hence

$$|r(w,x)| \le m(x) = \min\Big(\frac{C_3}{6}|x|^3,\, C_2 x^2\Big).$$

Replacing one variable at a time. Let $Z_1, \ldots, Z_N$ be i.i.d. $\mathcal{N}(0,1)$ variables, independent of the Y_i. By Corollary 4.6.7, $T_N = \frac{1}{\sqrt{N}}(Z_1 + \cdots + Z_N) \sim \mathcal{N}(0,1)$, so $\mathbb{E}[h(Z)] = \mathbb{E}[h(T_N)]$. For $k = 0, \ldots, N$, let $V_k = \frac{1}{\sqrt{N}}(Y_1 + \cdots + Y_k + Z_{k+1} + \cdots + Z_N)$, so that $V_N = S_N$ and $V_0 = T_N$, and let $W_k = V_k - \frac{Y_k}{\sqrt{N}} = V_{k-1} - \frac{Z_k}{\sqrt{N}}$ for $k \ge 1$. Then

$$\mathbb{E}[h(S_N)] - \mathbb{E}[h(T_N)] = \sum_{k=1}^{N} \big(\mathbb{E}[h(V_k)] - \mathbb{E}[h(V_{k-1})]\big) = \sum_{k=1}^{N} \Big(\mathbb{E}\Big[h\Big(W_k + \frac{Y_k}{\sqrt{N}}\Big)\Big] - \mathbb{E}\Big[h\Big(W_k + \frac{Z_k}{\sqrt{N}}\Big)\Big]\Big).$$

Expand $h(W_k + x) = h(W_k) + h'(W_k)x + \frac{1}{2}h''(W_k)x^2 + r(W_k, x)$ for $x = \frac{Y_k}{\sqrt{N}}$ and for $x = \frac{Z_k}{\sqrt{N}}$. The variable W_k is independent of Y_k and of Z_k, and Y_k and Z_k both have mean 0 and variance 1, so $\mathbb{E}[h'(W_k)Y_k] = \mathbb{E}[h'(W_k)]\mathbb{E}[Y_k] = 0 = \mathbb{E}[h'(W_k)Z_k]$ and $\mathbb{E}[h''(W_k)Y_k^2] = \mathbb{E}[h''(W_k)] = \mathbb{E}[h''(W_k)Z_k^2]$. So only the remainders remain:

$$\big|\mathbb{E}[h(S_N)] - \mathbb{E}[h(Z)]\big| \le \sum_{k=1}^{N} \Big(\mathbb{E}\Big[m\Big(\frac{Y_k}{\sqrt{N}}\Big)\Big] + \mathbb{E}\Big[m\Big(\frac{Z_k}{\sqrt{N}}\Big)\Big]\Big) = N\,\mathbb{E}\Big[m\Big(\frac{Y_1}{\sqrt{N}}\Big)\Big] + N\,\mathbb{E}\Big[m\Big(\frac{Z_1}{\sqrt{N}}\Big)\Big].$$

Bounding the remainders. Using the cubic bound, $N\,\mathbb{E}[m(Z_1/\sqrt{N})] \le \frac{C_3\mathbb{E}[|Z_1|^3]}{6\sqrt{N}} \to 0$. For the Y_1 term, let $\varepsilon > 0$, and use the cubic bound if $|Y_1| \le \varepsilon\sqrt{N}$ (then $|Y_1|^3 \le \varepsilon\sqrt{N}\,Y_1^2$) and the quadratic bound otherwise:

$$N\,\mathbb{E}\Big[m\Big(\frac{Y_1}{\sqrt{N}}\Big)\Big] \le \frac{C_3}{6}\,\varepsilon\,\mathbb{E}[Y_1^2] + C_2\,\mathbb{E}\big[Y_1^2\,\mathbf{1}_{\{|Y_1|>\varepsilon\sqrt{N}\}}\big].$$

Here $\mathbb{E}[Y_1^2] = 1$, and the last expectation tends to 0 as $N \to \infty$, since it is the tail of the convergent sum or integral $\mathbb{E}[Y_1^2]$. Hence all limit points of $\big|\mathbb{E}[h(S_N)] - \mathbb{E}[h(Z)]\big|$ are at most $\frac{C_3\varepsilon}{6}$, for every $\varepsilon > 0$, and $\mathbb{E}[h(S_N)] \to \mathbb{E}[h(Z)]$. □

The proof replaces the variables Y_k by Gaussian ones, one at a time; each replacement changes the expectation only by the remainder of order $N^{-3/2}$, because the first two moments match. This is Lindeberg's proof of the central limit theorem. Two rules allow us to combine limits in distribution.

Theorem 4.7.10 (Continuous mapping and Slutsky)**.** *Let $X_n \to X$ in distribution.*

1. *If g is continuous, then $g(X_n) \to g(X)$ in distribution.*
2. *(Slutsky) If $Y_n \to c$ in probability for a constant c, then $X_n + Y_n \to X + c$ and $X_nY_n \to cX$ in distribution.*
3. *If $Y_n \to c$ and $Z_n \to d$ in probability, and g is continuous at (c, d), then $g(Y_n, Z_n) \to g(c, d)$ in probability. The same holds for functions of one variable.*
4. *$Y_n \to c$ in distribution if and only if $Y_n \to c$ in probability.*

Proof. By Lemma 4.7.9, it suffices for parts 1 and 2 to show convergence of $\mathbb{E}[h(\cdot)]$ for every h as in part 1 of the lemma; let $|h| \le C_0$ and $|h'| \le C_1$.

(1) The function $h\circ g$ is bounded and continuous, so $\mathbb{E}[h(g(X_n))] \to \mathbb{E}[h(g(X))]$ by Lemma 4.7.9.

(2) Let $\varepsilon > 0$. By the mean value theorem, $|h(x+y) - h(x+c)| \le C_1|y-c|$, and the difference is at most $2C_0$ in any case. So

$$\big|\mathbb{E}[h(X_n + Y_n)] - \mathbb{E}[h(X_n + c)]\big| \le C_1\varepsilon + 2C_0\,P(|Y_n - c| > \varepsilon),$$

where the probability tends to 0. Since $x \mapsto h(x + c)$ is bounded and continuous, $\mathbb{E}[h(X_n + c)] \to \mathbb{E}[h(X + c)]$ by Lemma 4.7.9. So all limit points of $\big|\mathbb{E}[h(X_n + Y_n)] - \mathbb{E}[h(X + c)]\big|$ are at most $C_1\varepsilon$, for every $\varepsilon > 0$. For the product, let $\eta > 0$, choose $M > 0$ such that $\pm M$ are continuity points of the CDF F of X with $F(-M) + 1 - F(M) < \eta$ (see the proof of Lemma 4.7.9), and then $\varepsilon > 0$ with $C_1M\varepsilon < \eta$. If $|X_n| \le M$ and $|Y_n - c| \le \varepsilon$, then $|h(X_nY_n) - h(cX_n)| \le C_1|X_n||Y_n - c| \le \eta$, so

$$\big|\mathbb{E}[h(X_nY_n)] - \mathbb{E}[h(cX_n)]\big| \le \eta + 2C_0\big(P(|X_n| > M) + P(|Y_n - c| > \varepsilon)\big).$$

As $n \to \infty$, $P(|X_n| > M) \le P(X_n \le -M) + 1 - P(X_n \le M) \to F(-M) + 1 - F(M) < \eta$, and $\mathbb{E}[h(cX_n)] \to \mathbb{E}[h(cX)]$ by Lemma 4.7.9. So all limit points of $\big|\mathbb{E}[h(X_nY_n)] - \mathbb{E}[h(cX)]\big|$ are at most $\eta + 2C_0\eta$, for every $\eta > 0$.

(3) Let $\varepsilon > 0$. By continuity, there is a $\delta > 0$ with $|g(y, z) - g(c, d)| \le \varepsilon$ whenever $|y - c| \le \delta$ and $|z - d| \le \delta$. Hence $P(|g(Y_n, Z_n) - g(c, d)| > \varepsilon) \le P(|Y_n - c| > \delta) + P(|Z_n - d| > \delta) \to 0$.

(4) The CDF of the constant c is continuous everywhere except at c. If $Y_n \to c$ in distribution, then $P(|Y_n - c| > \varepsilon) \le P(Y_n \le c - \varepsilon) + 1 - P(Y_n \le c + \varepsilon) \to 0 + 1 - 1 = 0$. The converse is part 2, applied to the constant sequence 0 and to Y_n. □

Slutsky's theorem justifies the confidence intervals of Section 4.9, in which we replace σ by the sample standard deviation s_N. By the law of large numbers, applied to the X_i and the X_i^2 (if

$\mathbb{E}[X_i^4] < \infty$, so that Theorem 4.7.2 applies to the X_i^2), and by part 3, $s_N \to \sigma$ and $\frac{\sigma}{s_N} \to 1$ in probability, so by part 2,

$$\frac{\sqrt{N}(\bar{X}_N - \mu)}{s_N} = \frac{\sigma}{s_N} \cdot \frac{\sqrt{N}(\bar{X}_N - \mu)}{\sigma} \to \mathcal{N}(0,1) \qquad \text{in distribution,}$$

and $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$ is an asymptotic 95% confidence interval.

Proposition 4.7.11 (Delta method)**.** *Let $\sqrt{N}(T_N - \theta) \to \mathcal{N}(0, \tau^2)$ in distribution, and let g be continuously differentiable. Then $\sqrt{N}(g(T_N) - g(\theta)) \to \mathcal{N}(0, g'(\theta)^2\tau^2)$ in distribution.*

Proof. By the mean value theorem, $g(T_N) - g(\theta) = g'(\xi_N)(T_N - \theta)$ for some ξ_N between θ and T_N. By part 2 of Theorem 4.7.10, $T_N - \theta = \frac{1}{\sqrt{N}} \cdot \sqrt{N}(T_N - \theta) \to 0 \cdot Y = 0$ in distribution, where $Y \sim \mathcal{N}(0, \tau^2)$, hence in probability by part 4. Since $|\xi_N - \theta| \le |T_N - \theta|$, also $\xi_N \to \theta$ in probability, and $g'(\xi_N) \to g'(\theta)$ in probability by part 3, since g' is continuous. By part 2, $\sqrt{N}(g(T_N) - g(\theta)) = g'(\xi_N) \cdot \sqrt{N}(T_N - \theta) \to g'(\theta)Y$ in distribution, and $g'(\theta)Y \sim \mathcal{N}(0, g'(\theta)^2\tau^2)$. □

Example 4.7.12 (Confidence intervals on the log-odds scale)**.** For a proportion $\hat{p} = \bar{X}_N$ of N Bernoulli(p) observations, $\sqrt{N}(\hat{p} - p) \to \mathcal{N}(0, p(1-p))$ by the central limit theorem. The log-odds $g(p) = \log \frac{p}{1-p}$ have the derivative $g'(p) = \frac{1}{p(1-p)}$, so the estimated log-odds $g(\hat{p})$ have approximately the standard error $\frac{1}{\sqrt{Np(1-p)}}$. For 30 successes in 100 trials, the log-odds are $\log \frac{0.3}{0.7} \approx -0.847$, with the estimated standard error $\frac{1}{\sqrt{21}} \approx 0.218$. This gives the approximate 95% confidence interval $-0.847 \pm 1.96 \cdot 0.218 \approx [-1.275, -0.420]$ for the log-odds, and, transforming back with the sigmoid function, $[0.218, 0.397]$ for p. Unlike the interval $\hat{p} \pm 1.96\sqrt{\hat{p}(1-\hat{p})/N} = [0.210, 0.390]$, it is not symmetric around $\hat{p}$, and it never leaves $[0, 1]$. Logistic regression (Lecture 3) models the log-odds, and its standard errors are computed on this scale.

4.7.2 Monte Carlo sampling

Monte Carlo estimation needs samples from the distribution of interest. We have already seen inverse transform sampling (Theorem 4.2.7) and sampling Gaussian vectors by affine maps (Section 4.6), and we will see Markov chain Monte Carlo in Section 4.8. Two further general methods are rejection sampling, which produces exact samples, and importance sampling, which reweights samples from a different distribution.

Theorem 4.7.13 (Rejection sampling)**.** *Let p and q be densities, and let $M \ge 1$ be such that $p(x) \le Mq(x)$ for all x. Repeat the following trial until a value is accepted: draw $X \sim q$ and, independently, $U \sim$ Uniform$(0,1)$, and accept X if $U \le \frac{p(X)}{Mq(X)}$. Then the accepted value has the density p, and each trial is accepted with probability $\frac{1}{M}$.*

Proof. For every set A, $P(X \in A \text{ and accepted}) = \int_A q(x)\frac{p(x)}{Mq(x)}\,dx = \frac{1}{M}\int_A p(x)\,dx$. For $A = \mathbb{R}$, this gives $P(\text{accepted}) = \frac{1}{M}$, so $P(X \in A \mid \text{accepted}) = \int_A p(x)\,dx$. Since the trials are independent, the accepted value has this conditional distribution. □

Geometrically, the points $(X, UMq(X))$ are uniformly distributed under the graph of Mq, and we keep those that lie under the graph of p; their first coordinates have the density p. The method also works if p is only known up to a constant. For example, the Beta$(2,2)$ density $6x(1-x)$ is at most 1.5 on $[0,1]$, so with the uniform proposal and $M = 1.5$, two out of three trials are accepted. In high dimensions, however, M is usually enormous: for the target $\mathcal{N}(\mathbf{0}, I)$ in $\mathbb{R}^d$ and the proposal $\mathcal{N}(\mathbf{0}, 2I)$, the ratio $\frac{p}{q} = 2^{d/2}e^{-\|\mathbf{x}\|^2/4}$ forces $M \ge 2^{d/2}$, so only one in 1024 trials is accepted for $d = 20$. This is one reason why high-dimensional sampling relies on MCMC.

Proposition 4.7.14 (Importance sampling)**.** *Let p and q be densities such that $q(x) > 0$ for all x with $f(x)p(x) \neq 0$, and let $w = \frac{p}{q}$ (the importance weights). Then $\mathbb{E}_p[f(X)] = \mathbb{E}_q[f(Y)w(Y)]$. Hence, for i.i.d. samples $y_1, \dots, y_N$ from q, the estimator*

$$\hat{I} = \frac{1}{N} \sum_{i=1}^{N} f(y_i) w(y_i)$$

is unbiased, with variance $\frac{1}{N} \operatorname{Var}_q(f(Y)w(Y))$.

Proof. $\int f(x)p(x)\,dx = \int f(y)\frac{p(y)}{q(y)}q(y)\,dy$; the variance follows from Proposition 4.7.1. □

The variance is small if q puts its mass where $|f|p$ is large; for $f \geq 0$, the ideal (but impractical) choice $q \propto fp$ even gives variance 0.

Example 4.7.15 (A rare event)**.** The tail probability $P(Z > 4) = 3.17 \cdot 10^{-5}$ of $Z \sim \mathcal{N}(0,1)$ is hard to estimate by plain Monte Carlo: among $N = 10\,000$ samples, we expect only 0.32 samples above 4, and the relative standard error of the estimate is about 178%. With the proposal $q = \mathcal{N}(4,1)$, about half of the samples land above 4, and they are reweighted by $w(y) = \frac{\varphi(y)}{\varphi(y-4)} = e^{8-4y}$. With the same N, the relative standard error drops to about 2%. Probabilities of rare failures are estimated in this way.

If p is only known up to a constant, one uses the self-normalized estimator $\sum_i f(y_i)w_i / \sum_i w_i$, which is slightly biased but consistent. If a few weights dominate, the estimate is unreliable; the effective sample size $\frac{(\sum_i w_i)^2}{\sum_i w_i^2}$ measures how many samples effectively contribute. In machine learning, importance weights correct for differences between distributions: in off-policy evaluation in reinforcement learning (data collected with one policy, used to evaluate another), and under covariate shift, where training examples are weighted by the ratio of the test and training densities.

4.7.3 Stochastic gradient descent

Stochastic gradient descent (Remark 3.10.10) replaces the gradient of the training loss by the gradient on a random mini-batch. Its iterates are random vectors, and we can now analyze it. We write SGD as

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \eta_k \mathbf{g}_k,$$

where the stochastic gradient $\mathbf{g}_k$ is computed from a new random mini-batch in each step. We assume that it is unbiased and has bounded second moments: $\mathbb{E}[\mathbf{g}_k \mid \mathbf{x}_k] = \nabla f(\mathbf{x}_k)$ and $\mathbb{E}[\|\mathbf{g}_k\|^2 \mid \mathbf{x}_k] \leq G^2$. (For uniformly random mini-batches, the first condition holds as in Remark 3.10.10, and $\mathbf{x}_k$ takes finitely many values, so these conditional expectations are those of Section 4.4.2, applied entrywise.)

Theorem 4.7.16 (Convergence of SGD)**.** *Let f be differentiable and μ-strongly convex (Definition 3.11.10), with the minimum $\mathbf{x}^*$, and let $a_k = \mathbb{E}\big[\|\mathbf{x}_k - \mathbf{x}^*\|^2\big]$.*

1. *With the step sizes $\eta_k = \frac{1}{\mu(k+1)}$, $a_k \leq \dfrac{G^2}{\mu^2 k}$ for all $k \geq 1$.*
2. *With a constant step size $\eta \leq \frac{1}{2\mu}$, $a_k \leq (1 - 2\mu\eta)^k a_0 + \dfrac{\eta G^2}{2\mu}$.*

Proof. Expanding the square,

$$\|\mathbf{x}_{k+1} - \mathbf{x}^*\|^2 = \|\mathbf{x}_k - \mathbf{x}^*\|^2 - 2\eta_k \mathbf{g}_k^\top (\mathbf{x}_k - \mathbf{x}^*) + \eta_k^2 \|\mathbf{g}_k\|^2.$$

Summing over the values $\mathbf{x}$ of $\mathbf{x}_k$, as in the proof of the tower property (Theorem 4.4.15),

$$\mathbb{E}[\mathbf{g}_k^\top(\mathbf{x}_k - \mathbf{x}^*)] = \sum_{\mathbf{x}} P(\mathbf{x}_k = \mathbf{x})\,\mathbb{E}[\mathbf{g}_k \mid \mathbf{x}_k = \mathbf{x}]^\top(\mathbf{x} - \mathbf{x}^*) = \mathbb{E}\big[\nabla f(\mathbf{x}_k)^\top(\mathbf{x}_k - \mathbf{x}^*)\big],$$

which is at least μa_k, by Lemma 3.11.11. Similarly, $\mathbb{E}[\|\mathbf{g}_k\|^2] \le G^2$. Taking expectations,

$$a_{k+1} \le (1 - 2\mu\eta_k)\,a_k + \eta_k^2 G^2.$$

(1) With $\eta_k = \frac{1}{\mu(k+1)}$, the step $k = 0$ gives $a_1 \le -a_0 + \frac{G^2}{\mu^2} \le \frac{G^2}{\mu^2}$. If $a_k \le \frac{G^2}{\mu^2 k}$ for some $k \ge 1$, then

$$a_{k+1} \le \Big(1 - \frac{2}{k+1}\Big)\frac{G^2}{\mu^2 k} + \frac{G^2}{\mu^2(k+1)^2} = \frac{G^2}{\mu^2(k+1)}\Big(\frac{k-1}{k} + \frac{1}{k+1}\Big) \le \frac{G^2}{\mu^2(k+1)},$$

since $\frac{1}{k+1} \le \frac{1}{k}$. (2) With $0 \le 1 - 2\mu\eta < 1$, applying the recursion k times gives $a_k \le (1-2\mu\eta)^k a_0 + \eta^2 G^2 \sum_{j=0}^{k-1}(1-2\mu\eta)^j \le (1-2\mu\eta)^k a_0 + \frac{\eta^2 G^2}{2\mu\eta}$. □

So SGD with decreasing step sizes converges, but only at the rate $\frac{1}{k}$, much slower than the geometric rate of gradient descent (Theorem 3.11.12). With a constant step size, the error first decreases geometrically, but then fluctuates at a level proportional to η: the noise of the stochastic gradients keeps the iterates in a ball around $\mathbf{x}^*$. This is why step sizes are decreased during training. Each step of SGD costs only $|B|$ gradient evaluations instead of N; for large data sets and moderate accuracy, this more than compensates for the slower rate. Larger mini-batches reduce the noise: $\mathbb{E}[\|\mathbf{g}\|^2] = \|\nabla f\|^2 + \operatorname{tr}\operatorname{Cov}(\mathbf{g})$, and the covariance matrix of a mini-batch gradient decreases like $\frac{1}{|B|}$ (Proposition 4.7.1).

Example 4.7.17 (SGD computes running means)**.** Let $f(\theta) = \frac{1}{2N}\sum_{i=1}^N(\theta - y_i)^2$, which is 1-strongly convex with the minimum $\bar{y}$, the mean of the data. SGD with mini-batches of size 1 uses $g_k = \theta_k - y_{I_k}$ with a uniformly random index I_k. With $\eta_k = \frac{1}{k+1}$,

$$\theta_{k+1} = \theta_k - \frac{\theta_k - y_{I_k}}{k+1} = \frac{k\,\theta_k + y_{I_k}}{k+1},$$

so $\theta_1 = y_{I_0}$, and by induction $\theta_k = \frac{1}{k}(y_{I_0} + \cdots + y_{I_{k-1}})$ is the mean of k data points drawn with replacement. By Proposition 4.7.1, $\mathbb{E}[(\theta_k - \bar{y})^2] = \frac{s^2}{k}$, where $s^2 = \frac{1}{N}\sum_i (y_i - \bar{y})^2$. So the rate $\frac{1}{k}$ of Theorem 4.7.16 cannot be improved in general.

4.8 Markov Chains

So far, we mostly considered independent random variables. Many random sequences have memory: tomorrow's weather depends on today's, the next word of a sentence depends on the previous ones, and a user browsing the web chooses the next page depending on the current one. The simplest model with memory is a Markov chain, in which the future depends on the past only through the present.

Definition 4.8.1 (Markov chain)**.** A sequence of random variables $X_0, X_1, X_2, \ldots$ with values in a finite set of states $\{1, \ldots, n\}$ is a Markov chain with transition matrix $P \in \mathbb{R}^{n\times n}$ if

$$P(X_{t+1} = j \mid X_t = i, X_{t-1} = i_{t-1}, \ldots, X_0 = i_0) = P_{ij}$$

for all times t and all states $i, j, i_0, \ldots, i_{t-1}$ (whenever the condition has positive probability).

The entry P_{ij} is the probability of moving from state i to state j in one step. So P has nonnegative entries, and each row sums to 1, i.e., $P\mathbf{1} = \mathbf{1}$; such matrices are called stochastic matrices.

Example 4.8.2 (Weather)**.** Let state 1 be "sunny" and state 2 "rainy". After a sunny day, the next day is sunny with probability 0.9; after a rainy day, it is sunny with probability 0.5:

$$P = \begin{pmatrix} 0.9 & 0.1 \\ 0.5 & 0.5 \end{pmatrix}.$$

The chain is drawn as a graph with an arrow $i \to j$ labeled P_{ij} for each possible transition (Figure 4.4).

We write the distribution of X_t as a row vector $\boldsymbol{\pi}_t = \big(P(X_t = 1), \dots, P(X_t = n)\big)$. (The same weather model appears in the section on eigenvalues in Lecture 1, with the distributions written as columns and multiplied by the transposed matrix $P^\top$.)

Proposition 4.8.3. *$\boldsymbol{\pi}_{t+1} = \boldsymbol{\pi}_t P$, hence $\boldsymbol{\pi}_t = \boldsymbol{\pi}_0 P^t$. More generally, $P(X_{t+k} = j \mid X_t = i) = (P^k)_{ij}$.*

Proof. By the law of total probability (Theorem 4.1.6), $P(X_{t+1} = j) = \sum_i P(X_t = i)\, P(X_{t+1} = j \mid X_t = i) = \sum_i (\boldsymbol{\pi}_t)_i P_{ij}$. The second statement follows by induction on k, in the same way. □

So the distribution evolves by repeated multiplication with P, and the long-term behavior of the chain is a question about the powers of a matrix, i.e., about its eigenvalues (Lecture 1). For the weather chain, starting on a sunny day, $\boldsymbol{\pi}_0 = (1, 0)$, $\boldsymbol{\pi}_1 = (0.9, 0.1)$, $\boldsymbol{\pi}_2 = (0.86, 0.14)$, $\boldsymbol{\pi}_3 = (0.844, 0.156)$, and the distribution seems to approach a limit.

Definition 4.8.4 (Stationary distribution)**.** A probability distribution $\boldsymbol{\pi}$ (a row vector with $\pi_i \geq 0$ and $\sum_i \pi_i = 1$) is stationary for P if $\boldsymbol{\pi} P = \boldsymbol{\pi}$.

If X_0 has a stationary distribution, then so does every X_t. Transposing, $P^\top \boldsymbol{\pi}^\top = \boldsymbol{\pi}^\top$: a stationary distribution is an eigenvector of $P^\top$ with the eigenvalue 1 (a left eigenvector of P), scaled to have entries that sum to 1.

Proposition 4.8.5. *Let P be a stochastic matrix. Then 1 is an eigenvalue of P and of $P^\top$, and every (complex) eigenvalue λ of P satisfies $|\lambda| \leq 1$.*

Proof. $P\mathbf{1} = \mathbf{1}$, and $P^\top$ has the same characteristic polynomial as P, since $\det(A^\top) = \det(A)$ (Lecture 1). If $P\mathbf{v} = \lambda\mathbf{v}$ with $\mathbf{v} \neq \mathbf{0}$, choose i with $|v_i| = \max_j |v_j|$. Then $|\lambda||v_i| = |\sum_j P_{ij} v_j| \leq \sum_j P_{ij}|v_j| \leq \sum_j P_{ij}|v_i| = |v_i|$. □

Whether the distribution converges to a stationary distribution depends on two properties. A chain is irreducible if every state can be reached from every other state, i.e., for all i and j there is a t with $(P^t)_{ij} > 0$. It is aperiodic if for every state i, the numbers $t \geq 1$ with $(P^t)_{ii} > 0$ have greatest common divisor 1; for an irreducible chain, it suffices that some $P_{ii} > 0$.

Theorem 4.8.6 (Convergence of Markov chains)**.** *An irreducible Markov chain has a unique stationary distribution $\boldsymbol{\pi}$, and all $\pi_i > 0$. If the chain is also aperiodic, then $\boldsymbol{\pi}_0 P^t \to \boldsymbol{\pi}$ as $t \to \infty$, for every initial distribution $\boldsymbol{\pi}_0$.*

The key step of the proof is that some power of P has only positive entries.

Lemma 4.8.7. *If a Markov chain is irreducible and aperiodic, then there is an $m \geq 1$ such that all entries of P^m are positive.*

Proof. Fix a state i, and let $S_i = \{t \geq 1 : (P^t)_{ii} > 0\}$. Since $(P^{s+t})_{ii} = \sum_k (P^s)_{ik}(P^t)_{ki} \geq (P^s)_{ii}(P^t)_{ii}$, the set S_i is closed under addition. We show that it contains all large integers. Enumerate $S_i = \{s_1, s_2, \dots\}$. The greatest common divisors of $s_1, \dots, s_l$ form a nonincreasing sequence of positive integers, which is eventually constant; its final value divides every element of

S_i, so it is 1, and $s_1, \dots, s_l$ have the greatest common divisor 1 for some l. By Bézout's identity from elementary number theory, $1 = \sum_k a_k s_k$ for some integers a_k. Let u be the sum of the terms with $a_k > 0$, and v minus the sum of the terms with $a_k < 0$, so that $u - v = 1$. As sums of elements of S_i, $u \in S_i$, and $v \in S_i$ or $v = 0$. If $v = 0$, then $1 \in S_i$, so S_i contains every $t \geq 1$. Otherwise, write $t \geq v^2$ as $t = lv + r$ with $0 \leq r < v \leq l$. Then $t = (l - r)v + ru$ with $l - r \geq 1$, so $t \in S_i$.

So there is an N with $(P^t)_{ii} > 0$ for all states i and all $t \geq N$. By irreducibility, for $i \neq j$ there is a t_{ij} with $(P^{t_{ij}})_{ij} > 0$. Let $m = N + \max_{i \neq j} t_{ij}$. Then $(P^m)_{ii} > 0$, and for $i \neq j$, $(P^m)_{ij} \geq (P^{m - t_{ij}})_{ii} (P^{t_{ij}})_{ij} > 0$. □

Proof of Theorem 4.8.6. We measure the distance between distributions by $\|\mathbf{d}\|_1 = \sum_i |d_i|$.

Step 1: contraction. Let Q be a stochastic matrix whose entries are all at least $\delta > 0$; since the rows sum to 1, $n\delta \leq 1$. Let $\mathbf{d}$ be a row vector with $\sum_i d_i = 0$. Then $(\mathbf{d}Q)_j = \sum_i d_i Q_{ij} = \sum_i d_i (Q_{ij} - \delta)$, so

$$\|\mathbf{d}Q\|_1 \leq \sum_j \sum_i |d_i|(Q_{ij} - \delta) = (1 - n\delta)\|\mathbf{d}\|_1.$$

The same computation with $\delta = 0$ shows $\|\mathbf{d}P\|_1 \leq \|\mathbf{d}\|_1$ for every stochastic matrix P, and the entries of $\mathbf{d}P$ again sum to 0, since $(\mathbf{d}P)\mathbf{1} = \mathbf{d}(P\mathbf{1}) = \mathbf{d}\mathbf{1} = 0$.

Step 2: irreducible and aperiodic chains. By Lemma 4.8.7, the entries of $Q = P^m$ are at least some $\delta > 0$. By Proposition 4.8.5, 1 is an eigenvalue of $P^\top$, so there is a real row vector $\mathbf{v} \neq \mathbf{0}$ with $\mathbf{v}P = \mathbf{v}$, and then $\mathbf{v}Q = \mathbf{v}$. Now $|v_j| = |\sum_i v_i Q_{ij}| \leq \sum_i |v_i| Q_{ij}$, and summing over j gives $\sum_j |v_j| \leq \sum_i |v_i|$. So equality holds for every j, which is only possible if the nonzero numbers $v_i Q_{ij}$, and hence the nonzero entries v_i, all have the same sign. Thus $\boldsymbol{\pi} = \mathbf{v} / \sum_i v_i$ is a stationary distribution. For every initial distribution $\boldsymbol{\pi}_0$, the entries of $\mathbf{d} = \boldsymbol{\pi}_0 - \boldsymbol{\pi}$ sum to 0, and $\boldsymbol{\pi}_0 P^t - \boldsymbol{\pi} = \mathbf{d}P^t$. Writing $t = km + l$ with $0 \leq l < m$, Step 1 gives

$$\|\boldsymbol{\pi}_0 P^t - \boldsymbol{\pi}\|_1 = \|\mathbf{d}Q^k P^l\|_1 \leq (1 - n\delta)^k \|\mathbf{d}\|_1 \leq 2(1 - n\delta)^k \to 0.$$

Every stationary distribution $\boldsymbol{\pi}'$ satisfies $\boldsymbol{\pi}' = \boldsymbol{\pi}' P^t \to \boldsymbol{\pi}$, so $\boldsymbol{\pi}' = \boldsymbol{\pi}$. Moreover, the argument shows that every real row vector $\mathbf{v} \neq \mathbf{0}$ with $\mathbf{v}P = \mathbf{v}$ is a multiple of $\boldsymbol{\pi}$.

Step 3: irreducible chains. Let $P' = \frac{1}{2}(I + P)$, the "lazy" chain that stays where it is with probability $\frac{1}{2}$ and otherwise moves according to P. It is stochastic, and since I and P commute, $P'^t = 2^{-t} \sum_{k=0}^{t} \binom{t}{k} P^k$ has entries $(P'^t)_{ij} \geq 2^{-t}(P^t)_{ij}$ and $(P'^t)_{ii} \geq 2^{-t}$. So P' is irreducible and aperiodic, and by Step 2 it has a unique stationary distribution $\boldsymbol{\pi}$. Since $\boldsymbol{\pi}P' = \frac{1}{2}(\boldsymbol{\pi} + \boldsymbol{\pi}P)$, a distribution is stationary for P' if and only if it is stationary for P, so $\boldsymbol{\pi}$ is the unique stationary distribution of P; in the same way, every real row vector $\mathbf{v} \neq \mathbf{0}$ with $\mathbf{v}P = \mathbf{v}$ is a multiple of $\boldsymbol{\pi}$. Finally, choose i with $\pi_i > 0$. For every state j, there is a t with $(P^t)_{ij} > 0$ (with $t = 0$ if $j = i$), and $\pi_j = (\boldsymbol{\pi}P^t)_j \geq \pi_i (P^t)_{ij} > 0$. □

This is a special case of the Perron–Frobenius theorem on matrices with nonnegative entries. The proof gives the rate $(1 - n\delta)^{\lfloor t/m \rfloor}$; more precisely, the speed of convergence is governed by the second largest absolute value of an eigenvalue of P. Both assumptions are needed. For $P = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, the chain alternates between the two states: $(\frac{1}{2}, \frac{1}{2})$ is stationary, but $\boldsymbol{\pi}_0 = (1, 0)$ gives $\boldsymbol{\pi}_t = (1, 0), (0, 1), (1, 0), \dots$ (the chain has period 2, and -1 is an eigenvalue). For $P = I$, the chain never moves, and every distribution is stationary. For two states, everything can be computed by hand.

Proposition 4.8.8 (Two-state chains)**.** *Let* $P = \begin{pmatrix} 1 - a & a \\ b & 1 - b \end{pmatrix}$ *with* $a + b > 0$. *Then* $\boldsymbol{\pi} = \left(\frac{b}{a+b}, \frac{a}{a+b}\right)$ *is stationary, and for every initial distribution,*

$$\boldsymbol{\pi}_t - \boldsymbol{\pi} = (1 - a - b)^t (\boldsymbol{\pi}_0 - \boldsymbol{\pi}).$$

Proof. One checks $\boldsymbol{\pi}P = \boldsymbol{\pi}$ directly. The difference $\mathbf{d} = \boldsymbol{\pi}_0 - \boldsymbol{\pi}$ has entries that sum to 0, so $\mathbf{d} = c(1,-1)$, and $(1,-1)P = (1-a-b,\, a-1+b) = (1-a-b)(1,-1)$: the row vector $(1,-1)$ is a left eigenvector with the eigenvalue $1-a-b$. Hence $\boldsymbol{\pi}_t - \boldsymbol{\pi} = (\boldsymbol{\pi}_0 - \boldsymbol{\pi})P^t = (1-a-b)^t\,\mathbf{d}$. □

For the weather chain, $a = 0.1$ and $b = 0.5$, so the stationary distribution is $(\frac{5}{6}, \frac{1}{6})$, and the distance to it shrinks by the factor 0.4 (the second eigenvalue of P) every day. Starting on a sunny day, $P(X_t = \text{sunny}) = \frac{5}{6} + \frac{1}{6}\,0.4^t$; starting on a rainy day, it is $\frac{5}{6} - \frac{5}{6}\,0.4^t$ (Figure 4.4).




Figure 4.4: Left: the weather chain of Example 4.8.2. Right: the probability of a sunny day t days after a sunny day (orange) and after a rainy day (blue); both converge to the stationary probability $\frac{5}{6}$, and the distance shrinks by the factor 0.4 every day.

The ergodic theorem. For an irreducible chain, time averages converge to averages over the stationary distribution. This is a law of large numbers for dependent random variables.

Theorem 4.8.9 (Ergodic theorem)**.** *Let $X_0, X_1, \ldots$ be an irreducible Markov chain with an arbitrary initial distribution and the stationary distribution $\boldsymbol{\pi}$, and let f be a function on the states. Then $\frac{1}{T}\sum_{t=0}^{T-1} f(X_t) \to \sum_i \pi_i f(i)$ in probability as $T \to \infty$. In particular, the fraction of time that the chain spends in state i converges to π_i in probability.*

Proof. Let $\bar{f} = \sum_i \pi_i f(i)$, and let $\mathbf{g}$ be the column vector with the entries $g_i = f(i) - \bar{f}$, so that $\boldsymbol{\pi}\mathbf{g} = 0$. We first solve $\mathbf{g} = (I-P)\mathbf{h}$. By the proof of Theorem 4.8.6, the null space of $(I-P)^\top$ is spanned by $\boldsymbol{\pi}^\top$, so $\operatorname{rank}(I-P) = \operatorname{rank}\left((I-P)^\top\right) = n-1$ (Lecture 1). Every vector $(I-P)\mathbf{h}$ satisfies $\boldsymbol{\pi}(I-P)\mathbf{h} = (\boldsymbol{\pi} - \boldsymbol{\pi}P)\mathbf{h} = 0$, so the column space of $I-P$ is contained in the null space of the nonzero row vector $\boldsymbol{\pi}$; both have dimension $n-1$, so they are equal (Lecture 1), and $\mathbf{g} = (I-P)\mathbf{h}$ for some $\mathbf{h}$. Let $c = \max_i |h_i|$ and $G = \max_i |g_i|$. Since P^k is stochastic, the entries of $P^k\mathbf{h}$ are at most c in absolute value.

Let $S_T = \sum_{t=0}^{T-1} g_{X_t} = \sum_{t=0}^{T-1} f(X_t) - T\bar{f}$. For $s < t$, Proposition 4.8.3 gives $P(X_s = i, X_t = j) = P(X_s = i)(P^{t-s})_{ij}$, so $\mathbb{E}[g_{X_s} g_{X_t}] = \sum_i P(X_s = i)\, g_i\, (P^{t-s}\mathbf{g})_i$. The sum over t telescopes:

$$\sum_{t=s+1}^{T-1} P^{t-s}\mathbf{g} = \sum_{k=1}^{T-1-s} \left(P^k\mathbf{h} - P^{k+1}\mathbf{h}\right) = P\mathbf{h} - P^{T-s}\mathbf{h},$$

whose entries are at most $2c$ in absolute value. Hence $\left|\sum_{t>s} \mathbb{E}[g_{X_s} g_{X_t}]\right| \le 2cG$ for every s, and

$$\mathbb{E}[S_T^2] = \sum_{s=0}^{T-1} \mathbb{E}[g_{X_s}^2] + 2\sum_{s=0}^{T-1}\sum_{t=s+1}^{T-1} \mathbb{E}[g_{X_s} g_{X_t}] \le T(G^2 + 4cG).$$

By Markov's inequality for S_T^2 (Proposition 4.3.6), for every $\varepsilon > 0$,

$$P\Big(\Big|\frac{1}{T}\sum_{t=0}^{T-1} f(X_t) - \bar{f}\Big| > \varepsilon\Big) \le \frac{\mathbb{E}[S_T^2]}{T^2\varepsilon^2} \le \frac{G^2 + 4cG}{T\varepsilon^2} \to 0.$$

□

As for the strong law of large numbers in Appendix B, the convergence even holds with probability 1; we only prove the weak form. In the long run, $\frac{5}{6}$ of the days are sunny. Since successive states are correlated, these averages converge more slowly than averages of independent samples.

4.8.1 Detailed balance and Markov chain Monte Carlo

Finding a stationary distribution means solving a linear system. It is often easier to check a stronger, local condition.

Proposition 4.8.10 (Detailed balance)**.** *If a distribution $\boldsymbol{\pi}$ satisfies $\pi_i P_{ij} = \pi_j P_{ji}$ for all states i and j, then it is stationary.*

Proof. $(\boldsymbol{\pi}P)_j = \sum_i \pi_i P_{ij} = \sum_i \pi_j P_{ji} = \pi_j \sum_i P_{ji} = \pi_j$. □

Detailed balance says that in the stationary regime, the probability flow from i to j equals the flow from j to i; such chains are called reversible.

Example 4.8.11 (Random walks on graphs)**.** Let G be a connected undirected graph, and let a walker move from each vertex to a uniformly random neighbor, i.e., $P_{ij} = \frac{1}{\deg(i)}$ for each edge $\{i, j\}$. Then $\pi_i = \frac{\deg(i)}{2|E|}$, where $|E|$ is the number of edges, satisfies detailed balance: $\pi_i P_{ij} = \frac{1}{2|E|} = \pi_j P_{ji}$ for every edge. So in the long run, the walker visits each vertex in proportion to its degree. For the graph with the edges $\{1,2\}$, $\{1,3\}$, $\{2,3\}$, $\{3,4\}$, the degrees are $2, 2, 3, 1$, and $\boldsymbol{\pi} = (\frac{1}{4}, \frac{1}{4}, \frac{3}{8}, \frac{1}{8})$.

Remark 4.8.12 (PageRank). Google's original PageRank models a "random surfer" who, at each step, follows a uniformly random link on the current page with probability α (typically 0.85), and jumps to a uniformly random page with probability $1 - \alpha$. All entries of the resulting transition matrix are positive, so the chain is irreducible and aperiodic, and its unique stationary distribution ranks the pages. It is computed by the power iteration $\boldsymbol{\pi} \leftarrow \boldsymbol{\pi}P$ of the programming exercises of Lecture 1.

Detailed balance is also the key to Markov chain Monte Carlo (MCMC), the most important method for sampling from complicated distributions, such as the posterior distributions of Section 4.12. These are often only known up to a constant: $p(\theta \mid \mathcal{D}) \propto p(\mathcal{D} \mid \theta)\, p(\theta)$, where the evidence $p(\mathcal{D})$ is an intractable integral. The idea is to construct a Markov chain whose stationary distribution is the target distribution, run it for a long time, and use its states as (dependent) samples.

Theorem 4.8.13 (Metropolis algorithm)**.** *Let $\boldsymbol{\pi}$ be a distribution with all $\pi_i > 0$, and let Q be a symmetric stochastic matrix ($Q_{ij} = Q_{ji}$), the proposal. From the state i, propose a state j with probability Q_{ij}, and move to j with the acceptance probability* $\min\left(1, \frac{\pi_j}{\pi_i}\right)$*; otherwise, stay at i. This defines a Markov chain that satisfies detailed balance with respect to $\boldsymbol{\pi}$, so $\boldsymbol{\pi}$ is stationary.*

Proof. For $i \neq j$, $P_{ij} = Q_{ij} \min\left(1, \frac{\pi_j}{\pi_i}\right)$, so $\pi_i P_{ij} = Q_{ij} \min(\pi_i, \pi_j)$, which is symmetric in i and j because Q is. For $i = j$, detailed balance holds trivially. □

The algorithm only uses the ratios $\frac{\pi_j}{\pi_i}$, so the unknown normalizing constant cancels. Moves to more probable states are always accepted, and moves to less probable states sometimes. For densities, the same algorithm works with a symmetric proposal such as $\theta' = \theta + \eta\,\varepsilon$ with $\varepsilon \sim \mathcal{N}(0,1)$, and the ratio of the (unnormalized) densities. If the resulting chain is irreducible and aperiodic, Theorem 4.8.6 and the ergodic theorem justify using averages along the chain as estimates of expectations under $\boldsymbol{\pi}$.

Example 4.8.14. Let the states $1, 2, 3, 4$ lie on a cycle, let Q propose each of the two neighbors with probability $\frac{1}{2}$, and let the target be $\boldsymbol{\pi} = (0.1, 0.2, 0.3, 0.4)$. The Metropolis chain has the transition matrix

$$P = \begin{pmatrix} 0 & 0.5 & 0 & 0.5 \\ 0.25 & 0.25 & 0.5 & 0 \\ 0 & 1/3 & 1/6 & 0.5 \\ 0.125 & 0 & 0.375 & 0.5 \end{pmatrix};$$

for example, from state 3, the move to state 2 is proposed with probability $\frac{1}{2}$ and accepted with probability $\frac{0.2}{0.3}$, so $P_{32} = \frac{1}{3}$. One checks that $\boldsymbol{\pi} P = \boldsymbol{\pi}$.

Gibbs sampling. If the state is a vector $\mathbf{x} = (x_1, \dots, x_d)$, for example the parameters of a Bayesian model or the pixels of an image, it is often easy to sample a single entry from its conditional distribution given all the others, even if the joint distribution π is complicated. The Gibbs sampler updates the entries one at a time: in the update of entry i, it replaces x_i by a sample from $\pi(x_i \mid \mathbf{x}_{-i})$, the conditional distribution of the i-th entry given the other entries $\mathbf{x}_{-i}$, and keeps $\mathbf{x}_{-i}$.

Proposition 4.8.15 (Gibbs sampling)**.** *Let π be a distribution on a finite set of vectors $\mathbf{x} = (x_1, \dots, x_d)$. The update of entry i is a Markov chain that satisfies detailed balance with respect to π. Hence every sequence of such updates, for example a sweep through $i = 1, \dots, d$, leaves π stationary. (The same holds for densities.)*

Proof. The update of entry i moves from $\mathbf{x}$ to $\mathbf{x}'$ only if $\mathbf{x}'_{-i} = \mathbf{x}_{-i}$, and then with the probability $P_{\mathbf{x}\mathbf{x}'} = \pi(x'_i \mid \mathbf{x}_{-i}) = \frac{\pi(\mathbf{x}')}{\pi_{-i}(\mathbf{x}_{-i})}$, where π_{-i} is the marginal distribution of $\mathbf{x}_{-i}$. So $\pi(\mathbf{x})P_{\mathbf{x}\mathbf{x}'} = \frac{\pi(\mathbf{x})\pi(\mathbf{x}')}{\pi_{-i}(\mathbf{x}_{-i})}$ is symmetric in $\mathbf{x}$ and $\mathbf{x}'$, and π is stationary by Proposition 4.8.10. If $\pi P_1 = \pi$ and $\pi P_2 = \pi$, then also $\pi P_1 P_2 = \pi P_2 = \pi$. □

Gibbs sampling needs no proposal and no rejections, but it moves only along the coordinate axes, which is slow if the entries are strongly dependent. For a standard bivariate Gaussian with correlation ρ, the conditional distributions are $X_1 \mid X_2 = x_2 \sim \mathcal{N}(\rho x_2, 1 - \rho^2)$ and vice versa (Theorem 4.6.5). A sweep maps X_1 to $\rho^2 X_1$ plus independent Gaussian noise, so successive values of X_1 have the correlation ρ^2. For $\rho = 0.99$, it is 0.98, and it takes about 35 sweeps until the correlation drops below $\frac{1}{2}$ (since $0.98^{35} \approx 0.5$). Gibbs sampling is the method of choice for models with conditionally conjugate priors, such as mixture models and topic models, and for Markov random fields such as the Ising model.

4.8.2 Markov reward processes and the Bellman equation

Reinforcement learning studies agents that collect rewards while they move through the states of a Markov chain. Before the agent makes any decisions, the basic question is how much reward to expect in the long run.

Definition 4.8.16 (Markov reward process)**.** A Markov reward process consists of a Markov chain $X_0, X_1, \dots$ with the transition matrix P on the states $\{1, \dots, n\}$, a reward vector $\mathbf{r} \in \mathbb{R}^n$ (the reward r_i is collected in every step that the chain spends in state i), and a discount factor $\gamma \in [0, 1)$. The value of the state i is the expected discounted total reward

$$V_i = \mathbb{E}\Big[\sum_{t=0}^{\infty} \gamma^t r_{X_t} \;\Big|\; X_0 = i\Big],$$

and $\mathbf{V} = (V_1, \dots, V_n)^\top$ is the value vector.

The discount factor makes rewards in the far future count less. One can also read $1-\gamma$ as the probability that the process stops after each step, so that an episode lasts $\frac{1}{1-\gamma}$ steps on average. Since $|r_{X_t}| \le \|\mathbf{r}\|_\infty$, the series converges absolutely, uniformly in the outcomes, so we may take expectations term by term, and $|V_i| \le \frac{\|\mathbf{r}\|_\infty}{1-\gamma}$. We use the maximum norm $\|\mathbf{x}\|_\infty = \max_i |x_i|$ of Lecture 2. For a stochastic matrix P, each entry $(P\mathbf{x})_i = \sum_j P_{ij}x_j$ is a weighted average of the entries of $\mathbf{x}$, so

$$\|P\mathbf{x}\|_\infty \le \|\mathbf{x}\|_\infty \qquad \text{for all } \mathbf{x} \in \mathbb{R}^n.$$

Theorem 4.8.17 (Bellman equation)**.** *The value vector of a Markov reward process satisfies the Bellman equation*

$$\mathbf{V} = \mathbf{r} + \gamma P\mathbf{V}, \qquad \textit{i.e.,} \qquad V_i = r_i + \gamma \sum_{j=1}^{n} P_{ij}V_j.$$

The matrix $I-\gamma P$ is invertible, with $(I-\gamma P)^{-1} = \sum_{k=0}^{\infty} \gamma^k P^k$, and $\mathbf{V} = (I-\gamma P)^{-1}\mathbf{r}$ is the only solution of the Bellman equation.

Proof. By Proposition 4.8.3, $\mathbb{E}[r_{X_k} \mid X_0 = i] = \sum_j (P^k)_{ij} r_j = (P^k\mathbf{r})_i$, so $\mathbf{V} = \sum_{k\ge 0} \gamma^k P^k \mathbf{r}$. For every $\mathbf{x}$, $\|\gamma^k P^k \mathbf{x}\|_\infty \le \gamma^k \|\mathbf{x}\|_\infty$, so the series $S = \sum_{k\ge 0} \gamma^k P^k$ converges (apply it to $\mathbf{x} = \mathbf{e}_j$ to get the columns). Since

$$(I-\gamma P)\sum_{k=0}^{K-1} \gamma^k P^k = I - \gamma^K P^K \to I \qquad (K \to \infty),$$

and in the same way with the factors in the other order, $(I-\gamma P)S = S(I-\gamma P) = I$. So $I-\gamma P$ is invertible with the inverse S, and $\mathbf{V} = S\mathbf{r}$ is the only solution of $(I-\gamma P)\mathbf{V} = \mathbf{r}$, which is the Bellman equation. □

The Bellman equation says that the value of a state is the reward now plus the discounted expected value of the next state; this "first-step analysis" is how it is usually derived.

Example 4.8.18 (The value of sunny days)**.** In the weather chain of Example 4.8.2, let every sunny day give the reward 1, i.e., $\mathbf{r} = (1,0)^\top$, and let $\gamma = 0.9$. Then

$$I - \gamma P = \begin{pmatrix} 0.19 & -0.09 \\ -0.45 & 0.55 \end{pmatrix}, \qquad \mathbf{V} = (I-\gamma P)^{-1}\mathbf{r} = \frac{1}{0.064}\begin{pmatrix} 0.55 \\ 0.45 \end{pmatrix} = \begin{pmatrix} 8.59375 \\ 7.03125 \end{pmatrix}.$$

Check: $1 + 0.9(0.9 \cdot 8.59375 + 0.1 \cdot 7.03125) = 8.59375$. With $\frac{1}{1-\gamma} = 10$ days as the effective horizon, the normalized values $(1-\gamma)\mathbf{V} = (0.859, 0.703)$ are discounted averages of the fraction of sunny days. A sunny start is worth more, but the advantage fades: as $\gamma \to 1$, both entries of $(1-\gamma)\mathbf{V}$ tend to the long-run fraction $\pi_1 = \frac{5}{6}$ of sunny days (Exercise 25).

For large state spaces, solving the linear system is too expensive, and P may not even be known explicitly. The Bellman equation then suggests an iteration.

Proposition 4.8.19 (Iterative policy evaluation)**.** *Let $\mathbf{V}_0 \in \mathbb{R}^n$ be arbitrary, and let $\mathbf{V}_{k+1} = \mathbf{r} + \gamma P\mathbf{V}_k$. Then $\|\mathbf{V}_k - \mathbf{V}\|_\infty \le \gamma^k \|\mathbf{V}_0 - \mathbf{V}\|_\infty$.*

Proof. Subtracting the Bellman equation, $\mathbf{V}_{k+1} - \mathbf{V} = \gamma P(\mathbf{V}_k - \mathbf{V})$, so $\|\mathbf{V}_{k+1} - \mathbf{V}\|_\infty \le \gamma \|\mathbf{V}_k - \mathbf{V}\|_\infty$; use induction. □

In other words, the map $B(\mathbf{V}) = \mathbf{r} + \gamma P\mathbf{V}$ is a contraction, $\|B(\mathbf{U}) - B(\mathbf{V})\|_\infty \le \gamma \|\mathbf{U} - \mathbf{V}\|_\infty$, and $\mathbf{V}$ is its fixed point. In Example 4.8.18, starting from $\mathbf{V}_0 = \mathbf{0}$, the iterates are $(1,0)$, $(1.81, 0.45)$, $(2.51, 1.02)$, …, and the error bound $0.9^k \cdot 8.6$ drops below 0.01 after 65 steps. The convergence is slow if γ is close to 1, i.e., if the horizon is long.

Markov decision processes. In reinforcement learning, an agent chooses an action a from a finite set A in every state s. It then receives the reward $r(s,a)$, and the next state is s' with the probability $P(s' \mid s,a)$. A (deterministic) policy π chooses the action $\pi(s)$ in the state s. For a fixed policy, the states form a Markov reward process with the transition probabilities $P^\pi_{ss'} = P(s' \mid s,\pi(s))$ and the rewards $r^\pi_s = r(s,\pi(s))$, and its value vector $\mathbf{V}^\pi$ solves the Bellman equation $\mathbf{V}^\pi = \mathbf{r}^\pi + \gamma P^\pi \mathbf{V}^\pi$ (policy evaluation). The agent wants a policy with the largest value. The Bellman optimality operator is

$$(T\mathbf{V})_s = \max_{a\in A} \Big(r(s,a) + \gamma \sum_{s'} P(s' \mid s,a) V_{s'} \Big).$$

Theorem 4.8.20 (Value iteration)**.** *1. $\|T\mathbf{U} - T\mathbf{V}\|_\infty \le \gamma \|\mathbf{U} - \mathbf{V}\|_\infty$ for all $\mathbf{U}, \mathbf{V} \in \mathbb{R}^n$. Hence T has exactly one fixed point $\mathbf{V}^*$, the solution of the Bellman optimality equation $\mathbf{V}^* = T\mathbf{V}^*$, and the iterates $\mathbf{V}_{k+1} = T\mathbf{V}_k$ of value iteration satisfy $\|\mathbf{V}_k - \mathbf{V}^*\|_\infty \le \gamma^k \|\mathbf{V}_0 - \mathbf{V}^*\|_\infty$.*

2. $\mathbf{V}^\pi \le \mathbf{V}^*$ (entrywise) for every policy π.

3. The greedy policy π^*, which chooses in every state s an action that attains the maximum in $(T\mathbf{V}^*)_s$, has $\mathbf{V}^{\pi^*} = \mathbf{V}^*$. So it is an optimal policy, and $\mathbf{V}^*$ is the optimal value.

Proof. 1. For functions f, g on A, $|\max_a f(a) - \max_a g(a)| \le \max_a |f(a) - g(a)|$: if $\max f$ is attained at a^*, then $\max f - \max g \le f(a^*) - g(a^*)$, and symmetrically. Apply this with $f(a)$ and $g(a)$ the brackets in $(T\mathbf{U})_s$ and $(T\mathbf{V})_s$, whose difference is $\gamma \sum_{s'} P(s' \mid s,a)(U_{s'} - V_{s'})$, of absolute value at most $\gamma\|\mathbf{U} - \mathbf{V}\|_\infty$. Now let $\mathbf{V}_0$ be arbitrary and $\mathbf{V}_{k+1} = T\mathbf{V}_k$. Then $\|\mathbf{V}_{k+1} - \mathbf{V}_k\|_\infty \le \gamma^k \|\mathbf{V}_1 - \mathbf{V}_0\|_\infty$, so for $m > k$, $\|\mathbf{V}_m - \mathbf{V}_k\|_\infty \le (\gamma^k + \cdots + \gamma^{m-1})\|\mathbf{V}_1 - \mathbf{V}_0\|_\infty \le \frac{\gamma^k}{1-\gamma}\|\mathbf{V}_1 - \mathbf{V}_0\|_\infty$. Each entry of $\mathbf{V}_k$ is a Cauchy sequence of real numbers, so $\mathbf{V}_k$ converges to some $\mathbf{V}^*$, and since T is continuous, $T\mathbf{V}^* = \lim T\mathbf{V}_k = \lim \mathbf{V}_{k+1} = \mathbf{V}^*$. If also $T\mathbf{U} = \mathbf{U}$, then $\|\mathbf{U} - \mathbf{V}^*\|_\infty \le \gamma\|\mathbf{U} - \mathbf{V}^*\|_\infty$, so $\mathbf{U} = \mathbf{V}^*$. The error bound follows as in Proposition 4.8.19. (This is the Banach fixed-point theorem, Theorem A.5.3, for the maximum norm.)

2. T is monotone: if $\mathbf{U} \le \mathbf{V}$, then $T\mathbf{U} \le T\mathbf{V}$, since the probabilities $P(s' \mid s,a)$ are nonnegative. Moreover, $\mathbf{V}^\pi = \mathbf{r}^\pi + \gamma P^\pi \mathbf{V}^\pi \le T\mathbf{V}^\pi$, since the action $\pi(s)$ is one of the candidates in the maximum. By monotonicity and induction, $\mathbf{V}^\pi \le T^k \mathbf{V}^\pi$ for all k, and $T^k\mathbf{V}^\pi \to \mathbf{V}^*$ by part 1.

3. By the choice of π^*, $\mathbf{r}^{\pi^*} + \gamma P^{\pi^*}\mathbf{V}^* = T\mathbf{V}^* = \mathbf{V}^*$. So $\mathbf{V}^*$ solves the Bellman equation of the Markov reward process of π^*, and by the uniqueness in Theorem 4.8.17, $\mathbf{V}^{\pi^*} = \mathbf{V}^*$. □

The expression in the bracket is the action value $Q^*(s,a) = r(s,a) + \gamma \sum_{s'} P(s' \mid s,a) V^*_{s'}$, the value of taking the action a in the state s and acting optimally afterwards, and $V^*_s = \max_a Q^*(s,a)$. If the transition probabilities are unknown, Q-learning observes single transitions (s,a,r,s') of the agent and updates

$$Q(s,a) \leftarrow Q(s,a) + \eta\big(r + \gamma \max_{a'} Q(s',a') - Q(s,a)\big) :$$

it replaces the expectation over s' in value iteration by a sample, in the same way as SGD replaces the full gradient by a stochastic gradient (Section 4.7.3). Its convergence, for suitable step sizes η and if every pair (s,a) is tried infinitely often, is proved with the same kind of stochastic approximation argument as for SGD. Deep Q-networks approximate Q by a neural network.

4.8.3 Stationary time series and autoregressive models

A time series is a sequence of random variables $X_0, X_1, X_2, \ldots$ indexed by time: daily sales, temperatures, server loads, or the training loss of a neural network. Unlike i.i.d. data, neighboring values are dependent, and the simplest description of this dependence is by covariances.

Definition 4.8.21 (Stationary time series)**.** A sequence $X_0, X_1, \ldots$ of random variables with finite variances is (weakly) stationary if the mean $\mathbb{E}[X_t] = \mu$ and the autocovariances $c(h) = \operatorname{Cov}(X_t, X_{t+h})$ do not depend on t, for all $t, h \geq 0$. Its autocorrelation function is $\rho(h) = \frac{c(h)}{c(0)}$.

So a stationary series has the same mean and variance at all times, and the dependence between two values depends only on their distance in time. (We write $c(h)$ rather than the common $\gamma(h)$, to keep γ for discount factors.) Three basic examples, with uncorrelated noise terms ε_t with mean 0 and variance σ^2 (white noise):

- White noise itself is stationary, with $c(0) = \sigma^2$ and $c(h) = 0$ for $h \geq 1$.
- The moving average $X_t = \varepsilon_t + \theta\varepsilon_{t-1}$, an MA(1) process, is stationary, with $c(0) = (1+\theta^2)\sigma^2$, $c(1) = \theta\sigma^2$ and $c(h) = 0$ for $h \geq 2$: values more than one step apart are uncorrelated. Its autocorrelation $\rho(1) = \frac{\theta}{1+\theta^2}$ is at most $\frac{1}{2}$ in absolute value.
- The random walk $S_t = \varepsilon_1 + \cdots + \varepsilon_t$ (with $S_0 = 0$) has $\operatorname{Var}(S_t) = t\sigma^2$, so it is not stationary. Its differences $S_t - S_{t-1} = \varepsilon_t$ are. Trends of this kind are removed by differencing before a stationary model is fitted; this is the "I" (integrated) in ARIMA models.

The most important model is the autoregressive process, in which each value is a multiple of the previous one plus fresh noise.

Proposition 4.8.22 (The AR(1) process)**.** *Let $\varepsilon_1, \varepsilon_2, \ldots$ be uncorrelated, with mean 0 and variance $\sigma^2 > 0$, let X_0 have a finite variance and be uncorrelated with all ε_t, and let $X_t = \phi X_{t-1} + \varepsilon_t$ for $t \geq 1$.*

1. *$X_t = \phi^t X_0 + \sum_{k=0}^{t-1} \phi^k \varepsilon_{t-k}$. Hence $\mathbb{E}[X_t] = \phi^t \mathbb{E}[X_0]$, $\operatorname{Var}(X_t) = \phi^{2t} \operatorname{Var}(X_0) + \sigma^2 \sum_{k=0}^{t-1} \phi^{2k}$, and $\operatorname{Cov}(X_t, X_{t+h}) = \phi^h \operatorname{Var}(X_t)$.*
2. *Let $|\phi| < 1$. If $\mathbb{E}[X_0] = 0$ and $\operatorname{Var}(X_0) = \frac{\sigma^2}{1-\phi^2}$, then the series is stationary, with*

$$c(h) = \frac{\sigma^2 \phi^h}{1-\phi^2} \qquad \text{and} \qquad \rho(h) = \phi^h.$$

   For every X_0, $\mathbb{E}[X_t] \to 0$ and $\operatorname{Var}(X_t) \to \frac{\sigma^2}{1-\phi^2}$ as $t \to \infty$.
3. *If $|\phi| \geq 1$, then $\operatorname{Var}(X_t) \geq \operatorname{Var}(X_0) + t\sigma^2 \to \infty$, so the series is not stationary, whatever X_0 is.*

Proof. 1. The formula for X_t follows by induction. The terms $\phi^t X_0, \phi^0 \varepsilon_t, \ldots, \phi^{t-1}\varepsilon_1$ are uncorrelated, which gives the variance (Proposition 4.4.9). In the same way, $X_{t+h} = \phi^h X_t + \sum_{k=0}^{h-1} \phi^k \varepsilon_{t+h-k}$, and the noise terms $\varepsilon_{t+1}, \ldots, \varepsilon_{t+h}$ are uncorrelated with X_t, which is a linear combination of $X_0, \varepsilon_1, \ldots, \varepsilon_t$. So $\operatorname{Cov}(X_t, X_{t+h}) = \phi^h \operatorname{Var}(X_t)$.

2. By part 1 and the geometric sum, $\operatorname{Var}(X_t) = \phi^{2t} \operatorname{Var}(X_0) + \sigma^2 \frac{1-\phi^{2t}}{1-\phi^2}$, which is $\frac{\sigma^2}{1-\phi^2}$ for all t if $\operatorname{Var}(X_0) = \frac{\sigma^2}{1-\phi^2}$. Then $c(h) = \phi^h \frac{\sigma^2}{1-\phi^2}$, and the mean is $\phi^t \cdot 0 = 0$. For a general X_0, $\phi^t \to 0$ and $\phi^{2t} \to 0$.

3. $\operatorname{Var}(X_t) = \phi^2 \operatorname{Var}(X_{t-1}) + \sigma^2 \geq \operatorname{Var}(X_{t-1}) + \sigma^2$; use induction. □

So for $|\phi| < 1$, the process forgets its starting point and settles into a stationary regime, whose correlations decay geometrically. If the noise is Gaussian and independent, the AR(1) process is a Markov chain with values in $\mathbb{R}$, its stationary distribution is $\mathcal{N}(0, \frac{\sigma^2}{1-\phi^2})$, and the best prediction of X_{t+h} given the past is $\mathbb{E}[X_{t+h} \mid X_t] = \phi^h X_t$ (by the formula for X_{t+h} in the proof), with the

error variance $\sigma^2 \frac{1-\phi^{2h}}{1-\phi^2}$. Forecasts revert to the mean, and their uncertainty grows to the stationary variance. For example, for $\phi = 0.8$ and $\sigma^2 = 1$, the stationary variance is $\frac{1}{0.36} \approx 2.78$, and $\rho(3) = 0.512$. The AR(1) process is the time discretization of the Ornstein–Uhlenbeck process of Section 4.15.

To fit the model to data $x_0, \dots, x_T$, one regresses x_t on x_{t-1}: the least squares estimate, which is the maximum likelihood estimate for Gaussian noise given x_0, is

$$\hat{\phi} = \frac{\sum_{t=1}^{T} x_t x_{t-1}}{\sum_{t=1}^{T} x_{t-1}^2},$$

close to the sample autocorrelation at lag 1.

AR(p) processes and the companion matrix. An AR(p) process $X_t = \phi_1 X_{t-1} + \cdots + \phi_p X_{t-p} + \varepsilon_t$ uses p past values. The vector $\mathbf{Y}_t = (X_t, X_{t-1}, \dots, X_{t-p+1})^\top$ follows the AR(1)-type recursion

$$\mathbf{Y}_t = A\mathbf{Y}_{t-1} + \varepsilon_t \mathbf{e}_1, \qquad A = \begin{pmatrix} \phi_1 & \phi_2 & \cdots & \phi_{p-1} & \phi_p \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix},$$

with the companion matrix A. As in part 1 of Proposition 4.8.22, $\mathbf{Y}_t = A^t \mathbf{Y}_0 + \sum_{k=0}^{t-1} A^k \mathbf{e}_1 \varepsilon_{t-k}$, so the process forgets its start and becomes stationary if $A^k \to 0$. This happens if and only if all complex eigenvalues of A have absolute value less than 1; for a diagonalizable $A = QDQ^{-1}$ (with complex entries), this follows from $A^k = QD^kQ^{-1}$ (Lecture 1). The eigenvalues are the roots of $\lambda^p - \phi_1\lambda^{p-1} - \cdots - \phi_p$ (Exercise 26), and this condition is what time series software checks when it fits AR, ARMA and ARIMA models. For example, $X_t = 0.5X_{t-1} + 0.3X_{t-2} + \varepsilon_t$ has the eigenvalues 0.85 and -0.35 and is stationary, while for $X_t = 0.5X_{t-1} + 0.6X_{t-2} + \varepsilon_t$, the eigenvalue 1.06 makes the variance explode. The same condition on the eigenvalues of a matrix that is applied again and again decides whether the gradients of a recurrent neural network vanish or explode (Lecture 3).

Remark 4.8.23 (Markov models in machine learning). An n-gram language model is a Markov chain on words (or on tuples of $n-1$ words), whose transition probabilities are estimated by counting in a large text, which is maximum likelihood estimation (Section 4.9). Hidden Markov models, in which only a noisy function of the state is observed, were long the standard model in speech recognition. Reinforcement learning is based on Markov decision processes, in which the transition probabilities depend on the actions of an agent (Section 4.8.2). Diffusion models for image generation learn to reverse a Markov chain that gradually adds Gaussian noise to an image (Section 4.15.1).

4.9 Parameter Estimation

Statistics turns the questions of probability around: instead of computing probabilities from a known distribution, we infer an unknown distribution from data. We assume that the data $x_1, \dots, x_N$ are i.i.d. samples from a distribution $p(x \mid \theta)$ that belongs to a known family, with an unknown parameter θ (for example, the mean and variance of a Gaussian, or the weights of a model).

Definition 4.9.1 (Estimator, bias and variance)**.** An estimator is a function $\hat{\theta} = \hat{\theta}(X_1, \dots, X_N)$ of the data; as a function of random data, it is a random variable. Its bias is $\mathbb{E}[\hat{\theta}] - \theta$, and it is unbiased if the bias is 0 for all θ. Its mean squared error is $\mathrm{MSE}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \theta)^2]$.

Proposition 4.9.2 (Bias–variance decomposition)**.** $\mathrm{MSE}(\hat{\theta}) = \big(\mathbb{E}[\hat{\theta}] - \theta\big)^2 + \mathrm{Var}(\hat{\theta})$.

Proof. Let $m = \mathbb{E}[\hat{\theta}]$. Then $\mathbb{E}[(\hat{\theta}-\theta)^2] = \mathbb{E}[(\hat{\theta}-m)^2] + 2(m-\theta)\mathbb{E}[\hat{\theta}-m] + (m-\theta)^2$, and the middle term is 0. □

A good estimator has a small bias and a small variance, and sometimes we accept some bias for a much smaller variance. This is the idea behind regularization: ridge regression shrinks the weights towards 0, which introduces a bias but reduces the variance. For a prediction model, a similar decomposition says that the expected squared test error is the sum of the squared bias, the variance and the irreducible noise; very flexible models have a small bias but a large variance (they overfit), and very simple ones the opposite.

Example 4.9.3 (Shrinkage)**.** Let $X_1, \dots, X_N$ be i.i.d. $\mathcal{N}(\mu, \sigma^2)$, and consider the estimators $\hat{\mu}_c = c\bar{X}_N$ with $0 \le c \le 1$, which shrink the sample mean towards 0. Their bias is $(c-1)\mu$ and their variance $c^2\frac{\sigma^2}{N}$, so

$$\mathrm{MSE}(\hat{\mu}_c) = (1-c)^2\mu^2 + c^2\frac{\sigma^2}{N},$$

which is minimal for $c^* = \frac{\mu^2}{\mu^2+\sigma^2/N}$. For $\mu = 2$, $\sigma^2 = 16$ and $N = 4$, the unbiased sample mean has the MSE 4, while $\hat{\mu}_{1/2}$ has the MSE $1 + 1 = 2$. The best c depends on the unknown μ, so this estimator is not available in practice, but it shows that some bias can pay off. Remarkably, the James–Stein estimator achieves this in a data-driven way: for the means of $d \ge 3$ independent Gaussians, it shrinks the sample means towards 0, and its total MSE is smaller than that of the sample means, whatever the true means are.

By Proposition 4.7.1, the sample mean is an unbiased estimator of μ, with variance $\frac{\sigma^2}{N}$. The natural estimator of the variance, however, is biased.

Proposition 4.9.4 (Sample variance)**.** *Let $X_1, \dots, X_N$ be i.i.d. with mean μ and variance σ^2, and $N \ge 2$. Then*

$$\mathbb{E}\Big[\frac{1}{N}\sum_{i=1}^{N}(X_i - \bar{X}_N)^2\Big] = \frac{N-1}{N}\sigma^2, \qquad \textit{so} \qquad s^2 = \frac{1}{N-1}\sum_{i=1}^{N}(X_i - \bar{X}_N)^2$$

is an unbiased estimator of σ^2.

Proof. Expanding $X_i - \mu = (X_i - \bar{X}_N) + (\bar{X}_N - \mu)$ and using $\sum_i (X_i - \bar{X}_N) = 0$ gives

$$\sum_i (X_i - \mu)^2 = \sum_i (X_i - \bar{X}_N)^2 + N(\bar{X}_N - \mu)^2.$$

Taking expectations, $N\sigma^2 = \mathbb{E}\big[\sum_i (X_i - \bar{X}_N)^2\big] + N \cdot \frac{\sigma^2}{N}$. □

Intuitively, the data are on average closer to their own mean $\bar{X}_N$ than to the true mean μ, so the version with $\frac{1}{N}$ underestimates the variance. The factor $\frac{N}{N-1}$ (Bessel's correction) only matters for small N; `np.var` uses $\frac{1}{N}$ by default, and `np.var(x, ddof=1)` uses $\frac{1}{N-1}$.

4.9.1 The method of moments

The oldest general method of estimation chooses the parameters so that the model reproduces the moments of the data.

Definition 4.9.5 (Method of moments)**.** Let $p(x \mid \boldsymbol{\theta})$ be a family of distributions with k parameters. The method of moments estimate $\hat{\boldsymbol{\theta}}$ solves the k equations

$$\mathbb{E}_{\boldsymbol{\theta}}[X^j] = \frac{1}{N}\sum_{i=1}^{N} x_i^j, \qquad j = 1, \dots, k,$$

(or equivalent equations for the mean, the variance, and so on).

By the law of large numbers, the sample moments converge to the true moments, so the method of moments gives consistent estimates whenever the moment equations determine $\boldsymbol{\theta}$ in a continuous way.

Example 4.9.6. 1. Exponential(λ): $\mathbb{E}[X] = \frac{1}{\lambda}$ gives $\hat{\lambda} = \frac{1}{\bar{x}}$.

2. Uniform$(0, \theta)$: $\mathbb{E}[X] = \frac{\theta}{2}$ gives $\hat{\theta} = 2\bar{x}$. This estimator is unbiased, but it can be smaller than the largest observation, which is impossible for the true θ (compare with the MLE in the exercises).

3. The Gamma distribution with shape $k > 0$ and scale $s > 0$ has the density $\frac{x^{k-1}e^{-x/s}}{\Gamma(k)s^k}$ for $x > 0$, the mean ks and the variance ks^2. Matching the sample mean $\bar{x}$ and the sample variance $\hat{\sigma}^2$ gives $\hat{s} = \frac{\hat{\sigma}^2}{\bar{x}}$ and $\hat{k} = \frac{\bar{x}^2}{\hat{\sigma}^2}$; for example, $\bar{x} = 6$ and $\hat{\sigma}^2 = 12$ give $\hat{k} = 3$ and $\hat{s} = 2$. The maximum likelihood estimate of k has no closed form, and the method of moments provides a good starting point for its numerical computation (for example with Newton's method, Lecture 3).

The method of moments is simple, but its estimates usually have a larger variance than maximum likelihood estimates, which we discuss next.

4.9.2 Maximum likelihood estimation

The most important general method to construct estimators is to choose the parameter under which the observed data are most probable.

Definition 4.9.7 (Maximum likelihood)**.** The likelihood of a parameter θ for i.i.d. data $x_1, \dots, x_N$ is $L(\theta) = \prod_{i=1}^{N} p(x_i \mid \theta)$, and the log-likelihood is $\ell(\theta) = \log L(\theta) = \sum_{i=1}^{N} \log p(x_i \mid \theta)$. A maximum likelihood estimate (MLE) is a parameter $\hat{\theta}$ that maximizes L, or equivalently ℓ.

Taking logarithms turns the product into a sum, which is easier to differentiate and numerically more stable. Maximizing ℓ is the same as minimizing the negative log-likelihood $-\ell(\theta)$, which is a loss function in the sense of Lecture 3.

Example 4.9.8 (Bernoulli)**.** Let $x_i \in \{0, 1\}$ with k ones among N observations, and $p(x \mid \theta) = \theta^x(1-\theta)^{1-x}$. Then $\ell(\theta) = k \log\theta + (N-k)\log(1-\theta)$, and $\ell'(\theta) = \frac{k}{\theta} - \frac{N-k}{1-\theta} = 0$ gives $\hat{\theta} = \frac{k}{N}$, the relative frequency. Since $\ell'' < 0$, this is the maximum. Note that after three heads in three flips, $\hat{\theta} = 1$: the MLE claims that tails are impossible. With little data, maximum likelihood overfits; we return to this example in Section 4.12.

Example 4.9.9 (Gaussian)**.** For $x_i \sim \mathcal{N}(\mu, \sigma^2)$,

$$\ell(\mu, \sigma^2) = -\frac{N}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}\sum_{i=1}^{N}(x_i - \mu)^2.$$

Setting the partial derivatives to zero gives $\hat{\mu} = \bar{x}$ and $\hat{\sigma}^2 = \frac{1}{N}\sum_i (x_i - \bar{x})^2$: the MLE of the variance is the biased version of Proposition 4.9.4. For the data $2.1, 3.4, 1.9, 2.8, 3.3$, we get $\hat{\mu} = 2.7$, $\hat{\sigma}^2 = 0.372$, while $s^2 = 0.465$.

Theorem 4.9.10 (MLE of a multivariate Gaussian)**.** *For i.i.d. data $\mathbf{x}_1, \dots, \mathbf{x}_N$ from $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$, whose sample covariance matrix S is invertible, the maximum likelihood estimates are the sample mean $\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}}$ and the sample covariance matrix $\hat{\Sigma} = S = \frac{1}{N}\sum_i (\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top$.*

Proof. Up to a constant, $\ell = -\frac{N}{2}\log\det\Sigma - \frac{1}{2}\sum_i (\mathbf{x}_i - \boldsymbol{\mu})^\top \Sigma^{-1}(\mathbf{x}_i - \boldsymbol{\mu})$. For fixed Σ, this is a concave quadratic function of $\boldsymbol{\mu}$ with gradient $\Sigma^{-1}\sum_i (\mathbf{x}_i - \boldsymbol{\mu})$ (Lecture 3), which vanishes for

$\boldsymbol{\mu} = \bar{\mathbf{x}}$, whatever Σ is. With $\boldsymbol{\mu} = \bar{\mathbf{x}}$, we write ℓ in terms of the precision matrix $\Lambda = \Sigma^{-1}$. Since $\mathbf{a}^\top \Lambda \mathbf{a} = \operatorname{tr}(\Lambda \mathbf{a}\mathbf{a}^\top)$,

$$\ell = \frac{N}{2} \log\det\Lambda - \frac{1}{2}\sum_i \operatorname{tr}\left(\Lambda(\mathbf{x}_i - \bar{\mathbf{x}})(\mathbf{x}_i - \bar{\mathbf{x}})^\top\right) = \frac{N}{2}\big(\log\det\Lambda - \operatorname{tr}(\Lambda S)\big).$$

By the matrix gradients of Lecture 3 ($\nabla \log\det\Lambda = \Lambda^{-\top}$ and $\nabla \operatorname{tr}(\Lambda S) = S^\top$), the gradient with respect to Λ is $\frac{N}{2}(\Lambda^{-1} - S)$, which vanishes for $\Lambda = S^{-1}$, i.e., $\Sigma = S$. (One can show that $\log\det$ is concave on the symmetric positive definite matrices, so this critical point is the maximum.) □

4.9.3 Consistency and efficiency

How good can an estimator be? Two properties are central: an estimator should converge to the true parameter as the amount of data grows, and its variance should be as small as possible.

Definition 4.9.11 (Consistency)**.** A sequence of estimators $\hat{\theta}_N$, computed from N data points, is consistent if $\hat{\theta}_N \to \theta$ in probability (Definition 4.7.8) for every true parameter θ, i.e., if $P(|\hat{\theta}_N - \theta| \geq \varepsilon) \to 0$ as $N \to \infty$ for every $\varepsilon > 0$.

Proposition 4.9.12. *If the bias and the variance of $\hat{\theta}_N$ tend to 0 as $N \to \infty$, then $\hat{\theta}_N$ is consistent.*

Proof. By Markov's inequality applied to $(\hat{\theta}_N - \theta)^2$ and Proposition 4.9.2, $P(|\hat{\theta}_N - \theta| \geq \varepsilon) \leq \frac{\operatorname{MSE}(\hat{\theta}_N)}{\varepsilon^2} = \frac{\operatorname{bias}^2 + \operatorname{Var}(\hat{\theta}_N)}{\varepsilon^2} \to 0$. □

For example, the sample mean and both versions of the sample variance are consistent, and so are the method of moments estimators. A lower bound for the variance comes from the curvature of the log-likelihood. For a family $p(x \mid \theta)$ with a single parameter, the score is $s(x;\theta) = \frac{\partial}{\partial\theta}\log p(x \mid \theta)$. Differentiating $\int p(x \mid \theta)\,dx = 1$ under the integral sign gives $\int \frac{\partial p}{\partial\theta}\,dx = \int s\,p\,dx = 0$, i.e., $\mathbb{E}_\theta[s(X;\theta)] = 0$. The Fisher information is

$$I(\theta) = \mathbb{E}_\theta\big[s(X;\theta)^2\big] = \operatorname{Var}_\theta\left(s(X;\theta)\right),$$

and one can show that also $I(\theta) = -\mathbb{E}_\theta\big[\frac{\partial^2}{\partial\theta^2}\log p(X \mid \theta)\big]$: it measures how sharply the log-likelihood is curved around the true parameter, i.e., how much information one observation carries about θ.

Theorem 4.9.13 (Cramér–Rao bound)**.** *Let $\hat{\theta}$ be an unbiased estimator of θ from N i.i.d. observations, and assume that we may differentiate under the integral sign (Lecture 3; this excludes, e.g., families whose support depends on θ). Then*

$$\operatorname{Var}_\theta(\hat{\theta}) \geq \frac{1}{N\,I(\theta)}.$$

Proof. Let $S = \sum_{i=1}^N s(X_i;\theta)$ be the score of the whole sample. It has mean 0 and, by independence, variance $N\,I(\theta)$. Write $p_N(\mathbf{x} \mid \theta) = \prod_i p(x_i \mid \theta)$, so that $\frac{\partial p_N}{\partial\theta} = p_N S$. Differentiating the unbiasedness condition $\theta = \int \hat{\theta}(\mathbf{x})\,p_N(\mathbf{x} \mid \theta)\,d\mathbf{x}$ gives

$$1 = \int \hat{\theta}(\mathbf{x})\,S(\mathbf{x})\,p_N(\mathbf{x} \mid \theta)\,d\mathbf{x} = \mathbb{E}[\hat{\theta}S] = \operatorname{Cov}(\hat{\theta}, S),$$

since $\mathbb{E}[S] = 0$. By Theorem 4.4.10 (the Cauchy–Schwarz inequality),

$$1 = \operatorname{Cov}(\hat{\theta}, S)^2 \leq \operatorname{Var}(\hat{\theta})\operatorname{Var}(S) = \operatorname{Var}(\hat{\theta})\,N\,I(\theta).$$ □

An unbiased estimator that attains the bound is called efficient.

Example 4.9.14. 1. Bernoulli(θ): $s(x;\theta) = \frac{x}{\theta} - \frac{1-x}{1-\theta}$ and $I(\theta) = \frac{1}{\theta(1-\theta)}$. The bound $\frac{\theta(1-\theta)}{N}$ is the variance of the relative frequency $\bar{X}_N$, which is therefore efficient.

2. $\mathcal{N}(\mu, \sigma^2)$ with known σ^2: $s = \frac{x-\mu}{\sigma^2}$ and $I(\mu) = \frac{1}{\sigma^2}$, so the sample mean, with variance $\frac{\sigma^2}{N}$, is efficient.

3. Uniform$(0, \theta)$ does not satisfy the assumptions, since its support depends on θ. Indeed, the unbiased estimator $\frac{N+1}{N} \max_i X_i$ has the variance $\frac{\theta^2}{N(N+2)}$, which decreases like $\frac{1}{N^2}$, faster than any bound of the form $\frac{c}{N}$.

Theorem 4.9.15 (Asymptotic normality of the MLE)**.** *Under regularity conditions, the maximum likelihood estimator is consistent, and $\sqrt{N}(\hat{\theta}_{MLE} - \theta)$ converges in distribution to $\mathcal{N}(0, \frac{1}{I(\theta)})$.*

We do not prove this theorem. It says that for large N, the MLE is approximately unbiased and attains the Cramér–Rao bound: maximum likelihood is asymptotically efficient, which is the main theoretical justification for it. In practice, the standard error of the MLE is estimated from the curvature of the log-likelihood at its maximum, $\mathrm{SE} \approx \left(-\ell''(\hat{\theta})\right)^{-1/2}$ (the observed information). For example, for k successes in N Bernoulli trials, $-\ell''(\hat{\theta}) = \frac{k}{\hat{\theta}^2} + \frac{N-k}{(1-\hat{\theta})^2} = \frac{N}{\hat{\theta}(1-\hat{\theta})}$, which gives the familiar $\mathrm{SE} = \sqrt{\hat{\theta}(1-\hat{\theta})/N}$, e.g., 0.046 for 30 successes in 100 trials. With several parameters, the Fisher information becomes a matrix, and the approximate covariance matrix of the MLE is the inverse of the Hessian of $-\ell$ (Lecture 3). The Fisher information matrix also defines the inner product of the natural gradient method mentioned in Lecture 3.

4.9.4 Sufficiency and the Rao–Blackwell theorem

Often, a few numbers computed from the data contain all the information about the parameter. A statistic $T = T(X_1, \ldots, X_N)$ of discrete data is sufficient for θ if the conditional distribution of the data given T does not depend on θ: once we know T, the remaining details of the data are noise that tells us nothing about θ. We write $\mathbf{x} = (x_1, \ldots, x_N)$.

Theorem 4.9.16 (Fisher–Neyman factorization)**.** *A statistic T of discrete data is sufficient for θ if and only if the joint pmf of the data can be written as $p(\mathbf{x} \mid \theta) = g(T(\mathbf{x}), \theta)\, h(\mathbf{x})$ for some functions g and h.*

Proof. If T is sufficient, let $g(t, \theta) = P_\theta(T = t)$, and let $h(\mathbf{x})$ be the conditional probability of $\mathbf{x}$ given $T = T(\mathbf{x})$, which does not depend on θ. Then $p(\mathbf{x} \mid \theta) = P_\theta\big(T = T(\mathbf{x})\big)\, h(\mathbf{x})$ (if $P_\theta(T = T(\mathbf{x})) = 0$, both sides are 0). Conversely, if $p(\mathbf{x} \mid \theta) = g(T(\mathbf{x}), \theta) h(\mathbf{x})$, then for $\mathbf{x}$ with $T(\mathbf{x}) = t$ and $P_\theta(T = t) > 0$,

$$P_\theta(\mathbf{X} = \mathbf{x} \mid T = t) = \frac{p(\mathbf{x} \mid \theta)}{\sum_{\mathbf{x}' : T(\mathbf{x}') = t} p(\mathbf{x}' \mid \theta)} = \frac{g(t, \theta) h(\mathbf{x})}{g(t, \theta) \sum_{\mathbf{x}' : T(\mathbf{x}') = t} h(\mathbf{x}')} = \frac{h(\mathbf{x})}{\sum_{\mathbf{x}' : T(\mathbf{x}') = t} h(\mathbf{x}')},$$

which does not depend on θ. □

For data with densities, conditional distributions given T are more delicate, and we take the factorization $p(\mathbf{x} \mid \theta) = g(T(\mathbf{x}), \theta) h(\mathbf{x})$ of the joint density as the definition of sufficiency. For Bernoulli data, $\prod_i \theta^{x_i}(1-\theta)^{1-x_i} = \theta^k (1-\theta)^{N-k}$ only depends on the number k of ones; for Poisson data, $\prod_i e^{-\lambda} \frac{\lambda^{x_i}}{x_i!} = e^{-N\lambda} \lambda^{\sum_i x_i} \cdot \prod_i \frac{1}{x_i!}$, so $\sum_i x_i$ is sufficient; and for Gaussian data, $\sum_i x_i$ and $\sum_i x_i^2$ together are sufficient for (μ, σ^2). Since the likelihood only depends on the data through $g(T, \theta)$, so does the MLE.

Theorem 4.9.17 (Rao–Blackwell)**.** *Let T be sufficient for θ, and let $\hat{\theta}$ be an estimator with $\mathbb{E}[\hat{\theta}^2] < \infty$. Then $\tilde{\theta} = \mathbb{E}[\hat{\theta} \mid T]$ is an estimator (it does not depend on θ), it has the same bias as $\hat{\theta}$, and $\mathrm{MSE}(\tilde{\theta}) \le \mathrm{MSE}(\hat{\theta})$.*

Proof. By sufficiency, the conditional distribution of the data given T, and hence $\mathbb{E}[\hat{\theta} \mid T]$, does not depend on θ. By the tower property, $\mathbb{E}[\tilde{\theta}] = \mathbb{E}[\hat{\theta}]$, so the bias is the same. By the law of total variance (Example 4.4.16), $\operatorname{Var}(\hat{\theta}) = \mathbb{E}\big[\operatorname{Var}(\hat{\theta} \mid T)\big] + \operatorname{Var}(\tilde{\theta}) \geq \operatorname{Var}(\tilde{\theta})$, and the bias–variance decomposition gives the claim. □

So averaging an estimator over all data sets with the same value of a sufficient statistic never hurts.

Example 4.9.18 (Estimating the probability of a quiet day)**.** Let $X_1, \dots, X_N$ be the numbers of server failures on N days, i.i.d. Poisson(λ), and suppose that we want to estimate the probability $e^{-\lambda} = P(X_1 = 0)$ of a day without failures. The indicator $\hat{\theta} = \mathbf{1}_{\{X_1 = 0\}}$ is unbiased, but it only uses the first day. Given $S = X_1 + \cdots + X_N = s$, a short computation with the Poisson pmf shows that X_1 is Binomial($s, \frac{1}{N}$): each of the s failures happened on day 1 with probability $\frac{1}{N}$, independently. Hence

$$\tilde{\theta} = \mathbb{E}\big[\mathbf{1}_{\{X_1 = 0\}} \mid S\big] = \Big(1 - \frac{1}{N}\Big)^S.$$

It is unbiased, since $\mathbb{E}[t^S] = e^{N\lambda(t-1)}$ for $S \sim \text{Poisson}(N\lambda)$ (Example 4.4.5). For $\lambda = 1$ and $N = 10$, its variance is $\mathbb{E}\big[(0.9)^{2S}\big] - e^{-2} = e^{-1.9} - e^{-2} \approx 0.014$, compared with $e^{-1}(1 - e^{-1}) \approx 0.233$ for $\hat{\theta}$. The plug-in estimator $e^{-\bar{X}_N}$, which is the MLE, is slightly biased; one can show that $\tilde{\theta}$ has the smallest variance among all unbiased estimators of $e^{-\lambda}$ (the Lehmann–Scheffé theorem).

4.9.5 Loss functions as negative log-likelihoods

Maximum likelihood explains the loss functions of Lectures 2 and 3.

Proposition 4.9.19 (Least squares as maximum likelihood)**.** *Assume the linear regression model $y_i = \mathbf{w}^\top \mathbf{x}_i + \varepsilon_i$, where the noise terms $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$ are i.i.d. Then the negative log-likelihood of $\mathbf{w}$ is*

$$-\ell(\mathbf{w}) = \frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 + \frac{N}{2}\log(2\pi\sigma^2),$$

where X has the rows $\mathbf{x}_i^\top$. So the MLE of $\mathbf{w}$ is the least squares solution, and the MLE of the noise variance is the mean squared residual $\frac{1}{N}\|\mathbf{y} - X\hat{\mathbf{w}}\|^2$.

Proof. Given $\mathbf{x}_i$, $y_i \sim \mathcal{N}(\mathbf{w}^\top \mathbf{x}_i, \sigma^2)$, so $-\log p(y_i \mid \mathbf{x}_i, \mathbf{w}) = \frac{(y_i - \mathbf{w}^\top \mathbf{x}_i)^2}{2\sigma^2} + \frac{1}{2}\log(2\pi\sigma^2)$; sum over i. The second statement is proved as in Example 4.9.9. □

Similarly, if $y_i \in \{0, 1\}$ is Bernoulli with $P(y_i = 1) = \sigma(\mathbf{w}^\top \mathbf{x}_i)$, the negative log-likelihood is $\sum_i \big(-y_i \log p_i - (1 - y_i)\log(1 - p_i)\big)$ with $p_i = \sigma(\mathbf{w}^\top \mathbf{x}_i)$: the cross-entropy loss of logistic regression in Lecture 3. For K classes with $P(y_i = k) = \operatorname{softmax}(\mathbf{z}_i)_k$, it is the softmax cross-entropy loss $-\sum_i \log \operatorname{softmax}(\mathbf{z}_i)_{y_i}$. So choosing a loss function amounts to choosing a noise model. For example, Laplace-distributed noise with density $\frac{1}{2b}e^{-|\varepsilon|/b}$ leads to the absolute loss $\sum_i |y_i - \mathbf{w}^\top \mathbf{x}_i|$, which is less sensitive to outliers than the squared loss.

4.9.6 Exponential families

The Bernoulli, Poisson, exponential and Gaussian distributions, and the categorical distributions behind softmax, have the same structure, which explains why their maximum likelihood estimates are so simple, and why logistic, softmax and least squares regression have the same gradient.

Definition 4.9.20 (Exponential family)**.** An exponential family is a family of pmfs or densities of the form

$$p_{\boldsymbol{\eta}}(x) = h(x)\exp\big(\boldsymbol{\eta}^\top T(x) - A(\boldsymbol{\eta})\big), \qquad A(\boldsymbol{\eta}) = \log \sum_x h(x) e^{\boldsymbol{\eta}^\top T(x)} \ \Big(\text{or } \log \int h(x) e^{\boldsymbol{\eta}^\top T(x)}\, dx\Big),$$

where $T(x) \in \mathbb{R}^m$ is the sufficient statistic, $h(x) \geq 0$, the natural parameter $\boldsymbol{\eta} \in \mathbb{R}^m$ ranges over the vectors for which the sum or integral is finite, and the log-partition function A makes the total probability 1.

Example 4.9.21. 1. Bernoulli(p): $p^x(1-p)^{1-x} = \exp\left(x \log \frac{p}{1-p} + \log(1-p)\right)$, so $T(x) = x$, $\eta = \log \frac{p}{1-p}$ (the log-odds) and $A(\eta) = \log(1+e^\eta)$, since $1 - p = \frac{1}{1+e^\eta}$. The inverse map $p = \frac{1}{1+e^{-\eta}} = \sigma(\eta)$ is the sigmoid function.

2. Poisson(λ): $\frac{\lambda^x}{x!}e^{-\lambda} = \frac{1}{x!}\exp(x \log \lambda - \lambda)$, so $T(x) = x$, $\eta = \log \lambda$, $h(x) = \frac{1}{x!}$ and $A(\eta) = e^\eta$.

3. $\mathcal{N}(\mu, \sigma^2)$: $\frac{1}{\sqrt{2\pi\sigma^2}}e^{-(x-\mu)^2/(2\sigma^2)} = \frac{1}{\sqrt{2\pi}}\exp\left(\frac{\mu}{\sigma^2}x - \frac{1}{2\sigma^2}x^2 - \frac{\mu^2}{2\sigma^2} - \log\sigma\right)$, so $T(x) = (x, x^2)$ and $\boldsymbol{\eta} = \left(\frac{\mu}{\sigma^2}, -\frac{1}{2\sigma^2}\right)$.

4. Categorical on $\{1, \ldots, K\}$ with probabilities softmax($\boldsymbol{\eta}$): $p(k) = \exp\left(\eta_k - \log \sum_j e^{\eta_j}\right)$, so $T(k) = \mathbf{e}_k$ (one-hot) and $A(\boldsymbol{\eta}) = \log \sum_j e^{\eta_j}$, the log-sum-exp function.

Proposition 4.9.22. *Let $\boldsymbol{\eta}$ lie in the interior of the set of natural parameters, and let $X \sim p_{\boldsymbol{\eta}}$. For every $\mathbf{v} \in \mathbb{R}^m$, the random variable $Y = \mathbf{v}^\top T(X)$ has the cumulant generating function $K_Y(r) = A(\boldsymbol{\eta} + r\mathbf{v}) - A(\boldsymbol{\eta})$ for small $|r|$. Hence*

$$\frac{d}{dr}A(\boldsymbol{\eta} + r\mathbf{v})\Big|_{r=0} = \mathbf{v}^\top \mathbb{E}[T(X)], \qquad \frac{d^2}{dr^2}A(\boldsymbol{\eta} + r\mathbf{v})\Big|_{r=0} = \operatorname{Var}\left(\mathbf{v}^\top T(X)\right) \geq 0,$$

In particular, the partial derivatives of A are $\frac{\partial A}{\partial \eta_j}(\boldsymbol{\eta}) = \mathbb{E}[T_j(X)]$, and A is convex on the interior of the set of natural parameters.

Proof. In the discrete case, $\mathbb{E}[e^{rY}] = \sum_x h(x)e^{(\boldsymbol{\eta}+r\mathbf{v})^\top T(x) - A(\boldsymbol{\eta})} = e^{A(\boldsymbol{\eta}+r\mathbf{v}) - A(\boldsymbol{\eta})}$, and similarly with integrals. This is finite for small $|r|$, since $\boldsymbol{\eta}$ is an interior point. The derivatives follow from Proposition 4.4.21, with $\mathbf{v} = \mathbf{e}_j$ for the partial derivatives. The set of natural parameters is convex, since $e^{(\theta\boldsymbol{\eta}_1 + (1-\theta)\boldsymbol{\eta}_2)^\top T(x)} \leq \theta e^{\boldsymbol{\eta}_1^\top T(x)} + (1-\theta)e^{\boldsymbol{\eta}_2^\top T(x)}$ by the convexity of exp, and so is its interior. Along every segment in the interior, A has a nonnegative second derivative, so it is convex there (Theorem 3.11.6, in one variable), and a function that is convex on every segment is convex. □

For example, for Poisson(λ), $A'(\eta) = A''(\eta) = e^\eta = \lambda$ is the mean and the variance, and for Bernoulli, $A'(\eta) = \sigma(\eta) = p$ and $A''(\eta) = \sigma(\eta)(1 - \sigma(\eta)) = p(1-p)$.

Theorem 4.9.23 (Maximum likelihood in exponential families)**.** *For i.i.d. data $x_1, \ldots, x_N$ from an exponential family, the log-likelihood $\ell(\boldsymbol{\eta}) = \boldsymbol{\eta}^\top \sum_i T(x_i) - NA(\boldsymbol{\eta}) + \sum_i \log h(x_i)$ is concave on the interior of the set of natural parameters. If an interior parameter $\hat{\boldsymbol{\eta}}$ satisfies*

$$\nabla A(\hat{\boldsymbol{\eta}}) = \mathbb{E}_{\hat{\boldsymbol{\eta}}}[T(X)] = \frac{1}{N}\sum_{i=1}^N T(x_i),$$

then it maximizes the likelihood over the interior. Moreover, $\sum_i T(x_i)$ is a sufficient statistic.

Proof. Let $\boldsymbol{\eta}$ be another interior parameter, $\mathbf{v} = \boldsymbol{\eta} - \hat{\boldsymbol{\eta}}$, and $\phi(r) = -\ell(\hat{\boldsymbol{\eta}} + r\mathbf{v})$ for $0 \leq r \leq 1$. By Proposition 4.9.22, ϕ is convex, and $\phi'(0) = N\mathbf{v}^\top \mathbb{E}_{\hat{\boldsymbol{\eta}}}[T(X)] - \mathbf{v}^\top \sum_i T(x_i) = 0$. By the first-order characterization of convexity (Theorem 3.11.3, in one variable), $\phi(1) \geq \phi(0) + \phi'(0) = \phi(0)$, i.e., $\ell(\boldsymbol{\eta}) \leq \ell(\hat{\boldsymbol{\eta}})$. The likelihood $\prod_i h(x_i) \cdot \exp\left(\boldsymbol{\eta}^\top \sum_i T(x_i) - NA(\boldsymbol{\eta})\right)$ has the form of the factorization theorem (Theorem 4.9.16). □

So the maximum likelihood estimate matches the expected sufficient statistics with their averages over the data (moment matching). For Poisson data, $\hat{\lambda} = \bar{x}$; for Gaussian data, the mean and the second moment are matched, which gives the estimates of Example 4.9.9; for categorical data, the predicted probabilities are the observed frequencies.

Generalized linear models. A generalized linear model predicts a response y from features $\mathbf{x}$ by an exponential family with the natural parameter $\eta = \mathbf{w}^\top \mathbf{x}$: the Gaussian with known variance gives linear regression, Bernoulli gives logistic regression, and Poisson gives Poisson regression for counts, with $\mathbb{E}[y \mid \mathbf{x}] = e^{\mathbf{w}^\top \mathbf{x}}$. For data $(\mathbf{x}_i, y_i)$ with $T(y) = y$, the negative log-likelihood is $\sum_i \left(A(\mathbf{w}^\top \mathbf{x}_i) - y_i \mathbf{w}^\top \mathbf{x}_i\right)$ plus a constant, and by the chain rule and Proposition 4.9.22,

$$\nabla_{\mathbf{w}}\big(-\ell(\mathbf{w})\big) = \sum_{i=1}^{N} \left(A'(\mathbf{w}^\top \mathbf{x}_i) - y_i\right)\mathbf{x}_i = \sum_{i=1}^{N} \left(\mathbb{E}[y \mid \mathbf{x}_i] - y_i\right)\mathbf{x}_i :$$

the gradient is always "prediction minus target" times the input, as we found for linear and logistic regression in Lecture 3. Its Hessian $\sum_i A''(\mathbf{w}^\top \mathbf{x}_i)\mathbf{x}_i \mathbf{x}_i^\top$ is positive semidefinite, so all these losses are convex.

4.9.7 Confidence intervals

An estimate is more useful with an error bar. By the central limit theorem, $\bar{X}_N$ is approximately $\mathcal{N}(\mu, \frac{\sigma^2}{N})$, so

$$P\Big(|\bar{X}_N - \mu| \le 1.96 \frac{\sigma}{\sqrt{N}}\Big) \approx 0.95.$$

Replacing the unknown σ by the sample standard deviation s (which is justified for large N by Slutsky's theorem, Theorem 4.7.10), the interval $\bar{x} \pm 1.96 \frac{s}{\sqrt{N}}$ is an approximate 95% confidence interval for μ. For example, for $N = 100$ data points with $\bar{x} = 5.2$ and $s = 1.5$, it is $[4.91, 5.49]$. Note that μ is fixed and the interval is random: if we repeated the experiment many times, about 95% of the intervals would contain μ. For a classifier with accuracy 0.9 on 1000 independent test examples, the standard error is $\sqrt{0.9 \cdot 0.1/1000} \approx 0.0095$, so the accuracy is known up to about ± 0.019, and a difference of one percentage point between two models on such a test set may well be due to chance.

4.9.8 Nonparametric estimation

So far, we assumed a parametric family of distributions and estimated its parameters. Nonparametric methods estimate the distribution itself. For i.i.d. data $X_1, \dots, X_N$ with the CDF F, the empirical CDF

$$\hat{F}_N(t) = \frac{1}{N} \#\{i : X_i \le t\} = \frac{1}{N} \sum_{i=1}^{N} \mathbf{1}_{\{X_i \le t\}}$$

is the CDF of the empirical distribution, which puts the probability $\frac{1}{N}$ on each data point. For fixed t, $N\hat{F}_N(t)$ is Binomial$(N, F(t))$, so $\hat{F}_N(t)$ is unbiased with the variance $\frac{F(t)(1-F(t))}{N}$, and Hoeffding's inequality (Theorem 4.7.5) gives $P(|\hat{F}_N(t) - F(t)| \ge \varepsilon) \le 2e^{-2N\varepsilon^2}$. In fact, the approximation is good for all t at once.

Theorem 4.9.24 (Uniform convergence of the empirical CDF)**.** *Let F be continuous. For every $\varepsilon \in (0, 1]$,*

$$P\Big(\sup_{t \in \mathbb{R}} |\hat{F}_N(t) - F(t)| > \varepsilon\Big) \le \frac{4}{\varepsilon} e^{-N\varepsilon^2/2}.$$

Proof. Let $m = \lceil 2/\varepsilon \rceil$. Since F is continuous and increases from 0 to 1, the intermediate value theorem gives points $t_1 < \cdots < t_{m-1}$ with $F(t_j) = \frac{j}{m}$; let $t_0 = -\infty$ and $t_m = \infty$, with $\hat{F}_N = F = 0$ at t_0 and $= 1$ at t_m. For $t_{j-1} \le t \le t_j$, both functions are nondecreasing, so

$$\hat{F}_N(t) - F(t) \le \hat{F}_N(t_j) - F(t_{j-1}) = \hat{F}_N(t_j) - F(t_j) + \tfrac{1}{m}, \qquad \hat{F}_N(t) - F(t) \ge \hat{F}_N(t_{j-1}) - F(t_{j-1}) - \tfrac{1}{m}.$$

Hence $\sup_t |\hat{F}_N(t) - F(t)| \le \max_{1\le j\le m-1} |\hat{F}_N(t_j) - F(t_j)| + \frac{\varepsilon}{2}$, and if the supremum exceeds ε, then $|\hat{F}_N(t_j) - F(t_j)| > \frac{\varepsilon}{2}$ for some j. By the union bound and Hoeffding's inequality, this has probability at most $(m-1)\cdot 2e^{-2N(\varepsilon/2)^2} \le \frac{4}{\varepsilon}e^{-N\varepsilon^2/2}$, since $m - 1 < \frac{2}{\varepsilon}$. □

So $\sup_t |\hat{F}_N(t) - F(t)| \to 0$ in probability (the Glivenko–Cantelli theorem; since the bounds are summable over N, the Borel–Cantelli lemma in Appendix B even gives almost sure convergence). The bound is crude; the Dvoretzky–Kiefer–Wolfowitz inequality, which we do not prove, improves it to $2e^{-2N\varepsilon^2}$ for every F. With it, $\hat{F}_N(t) \pm \sqrt{\log(40)/(2N)}$ is a 95% confidence band for the whole function F; for $N = 1000$, its half-width is 0.043. The largest distance $\sup_t |\hat{F}_N(t) - F_0(t)|$ to a hypothesized CDF F_0 is the statistic of the Kolmogorov–Smirnov test of H_0: $F = F_0$.

Kernel density estimation. A density cannot be estimated by the empirical distribution, which has no density. A histogram counts the data in bins; a smoother estimate replaces each data point by a small bump. With a kernel $K \ge 0$ with $\int K = 1$ and $K(-u) = K(u)$, such as the standard normal density φ, and a bandwidth $h > 0$, the kernel density estimate is

$$\hat{f}_h(x) = \frac{1}{Nh}\sum_{i=1}^{N} K\Big(\frac{x - X_i}{h}\Big),$$

a mixture of N copies of K, scaled by h and centered at the data points.

Proposition 4.9.25 (Bias and variance of kernel density estimates)**.** *Let the data have a density f with bounded continuous derivatives f', f'' and f''', and let $\kappa_2 = \int u^2 K(u)\,du$, $\kappa_3 = \int |u|^3 K(u)\,du$, $R = \int K(u)^2\,du$ and $R_1 = \int |u| K(u)^2\,du$ be finite. Then for every x,*

$$\Big|\mathbb{E}[\hat{f}_h(x)] - f(x) - \frac{h^2}{2}\kappa_2 f''(x)\Big| \le \frac{h^3}{6}\kappa_3 \sup|f'''|, \qquad \operatorname{Var}\big(\hat{f}_h(x)\big) \le \frac{f(x)R + hR_1 \sup|f'|}{Nh}.$$

Proof. Each term $Y_i = \frac{1}{h}K(\frac{x-X_i}{h})$ has the mean $\int \frac{1}{h}K(\frac{x-y}{h})f(y)\,dy = \int K(u)f(x-hu)\,du$, with the substitution $y = x - hu$. By Taylor's theorem (Lecture 3), $f(x-hu) = f(x) - huf'(x) + \frac{h^2u^2}{2}f''(x) - \frac{h^3u^3}{6}f'''(\xi)$ for some ξ. Integrating against K, with $\int K = 1$ and $\int uK(u)\,du = 0$ by symmetry, gives the bias. Since $\hat{f}_h(x)$ is the average of the i.i.d. Y_i, $\operatorname{Var}(\hat{f}_h(x)) = \frac{1}{N}\operatorname{Var}(Y_1) \le \frac{1}{N}\mathbb{E}[Y_1^2] = \frac{1}{Nh}\int K(u)^2 f(x-hu)\,du$, and $f(x-hu) \le f(x) + h|u|\sup|f'|$ by the mean value theorem. □

This is a bias–variance tradeoff, controlled by h: a small bandwidth gives a spiky estimate with a large variance, of order $\frac{1}{Nh}$, and a large bandwidth oversmooths, with a bias of order h^2. Minimizing the leading terms $\frac{h^4}{4}\kappa_2^2 f''(x)^2 + \frac{f(x)R}{Nh}$ of the mean squared error over h gives $h \propto N^{-1/5}$ and a mean squared error of order $N^{-4/5}$, slower than the rate $\frac{1}{N}$ of parametric estimates; in practice, h is chosen by cross-validation or by rules of thumb such as $h = 1.06\,\hat{\sigma}N^{-1/5}$ for the Gaussian kernel. In d dimensions, the same computation gives the rate $N^{-4/(4+d)}$, which is very slow for large d: the curse of dimensionality, and a reason why high-dimensional densities are modeled with parametric or neural network models instead.

4.9.9 Survival analysis and censored data

How long until a customer cancels a subscription, a machine fails, or a patient relapses? Such waiting times are positive random variables T, and their analysis has two peculiarities: they are naturally described by the risk of the event at each moment, and they are often only partially observed, because the study ends before the event happens. Let $T \ge 0$ have a continuous density p, the CDF F, and $F(0) = 0$.

Definition 4.9.26 (Survival function and hazard rate)**.** The survival function of T is $S(t) = P(T > t) = 1 - F(t)$. For t with $S(t) > 0$, the hazard rate is $h(t) = \frac{p(t)}{S(t)}$, and the cumulative hazard is $H(t) = \int_0^t h(u)\,du$.

The hazard rate is the rate of events among the survivors: $P(t < T \le t + \delta \mid T > t) = \frac{F(t+\delta)-F(t)}{S(t)} \approx h(t)\delta$ for small δ. A constant hazard describes events that strike at random; a decreasing hazard, "infant mortality" (defective units fail early); and an increasing one, wear-out.

Proposition 4.9.27. *1. $S(t) = e^{-H(t)}$ and $p(t) = h(t)e^{-H(t)}$, so the hazard rate determines the distribution. Moreover, $\mathbb{E}[T] = \int_0^\infty S(t)\,dt$.*

2. The hazard rate is constant, $h(t) = \lambda$, if and only if $T \sim$ Exponential(λ).

Proof. 1. $S' = -p$, so $(\log S)' = -\frac{p}{S} = -h$. Since $S(0) = 1$, integration gives $\log S(t) = -H(t)$, and $p = hS$. The formula for $\mathbb{E}[T]$ is the tail integral of Example 4.4.7.

2. If $h = \lambda$, then $S(t) = e^{-\lambda t}$, the survival function of Exponential(λ), and conversely $h(t) = \frac{\lambda e^{-\lambda t}}{e^{-\lambda t}} = \lambda$. □

A constant hazard is the memorylessness of the exponential distribution (Section 4.2). The Weibull distribution with the shape $k > 0$ and the scale $s > 0$ has the hazard $h(t) = \frac{k}{s}\left(\frac{t}{s}\right)^{k-1}$, so $H(t) = \left(\frac{t}{s}\right)^k$ and $S(t) = e^{-(t/s)^k}$; it has a decreasing hazard for $k < 1$, a constant one for $k = 1$ (the exponential distribution), and an increasing one for $k > 1$.

Censoring. In a study that runs for a limited time, some units have not experienced the event when the study ends, or they leave the study early. For each unit i, we then observe $t_i = \min(T_i, C_i)$, where C_i is the censoring time, and the indicator $\delta_i = 1$ if the event was observed ($T_i \le C_i$) and $\delta_i = 0$ if the observation was censored. For a censored unit, we only know that $T_i > t_i$. Assume that the censoring times are independent of the event times, with a distribution that does not depend on the parameter θ of the event times (non-informative censoring). Then an observed event contributes the density $p_\theta(t_i)$ to the likelihood, a censored observation contributes the probability $S_\theta(t_i)$, and the factors that come from the censoring times do not depend on θ. So the likelihood is

$$L(\theta) = \prod_{i=1}^{N} p_\theta(t_i)^{\delta_i} S_\theta(t_i)^{1-\delta_i} = \prod_{i=1}^{N} h_\theta(t_i)^{\delta_i} e^{-H_\theta(t_i)}.$$

Proposition 4.9.28 (Exponential survival times with censoring)**.** *For Exponential(λ) event times with non-informative censoring, the maximum likelihood estimate is*

$$\hat{\lambda} = \frac{D}{\sum_{i=1}^{N} t_i},$$

where $D = \sum_i \delta_i$ is the number of observed events: the number of events divided by the total time at risk.

Proof. With $h = \lambda$ and $H(t) = \lambda t$, $\log L(\lambda) = D \log \lambda - \lambda \sum_i t_i$, whose derivative $\frac{D}{\lambda} - \sum_i t_i$ vanishes at $\hat{\lambda}$; the second derivative $-\frac{D}{\lambda^2}$ is negative (for $D > 0$). □

Example 4.9.29 (Censored waiting times)**.** Six customers are observed until they cancel their subscription or until the study ends. The times (in months) are $2, 3^+, 4, 5^+, 7, 8^+$, where $^+$ marks a censored time: these customers had not canceled yet. With $D = 3$ and the total time 29, $\hat{\lambda} = \frac{3}{29} \approx 0.103$ per month, and the estimated mean lifetime is $\frac{1}{\hat{\lambda}} \approx 9.7$ months. Two naive approaches are biased: dropping the censored customers gives $\frac{3}{2+4+7} \approx 0.23$, and treating the censoring times as cancellations gives $\frac{6}{29} \approx 0.21$. Both ignore that the censored customers stayed at least as long as observed, and both overestimate the risk of canceling by a factor of about 2.

The Kaplan–Meier estimator. Without a parametric model, the survival function is estimated as a product of conditional survival probabilities. Let $\tau_1 < \tau_2 < \cdots$ be the distinct times at which events are observed, let d_j be the number of events at τ_j, and let n_j be the number of units at risk just before τ_j, i.e., with $t_i \geq \tau_j$. Since surviving past t means surviving each event time $\tau_j \leq t$, given survival until then, the Kaplan–Meier estimator is

$$\hat{S}(t) = \prod_{j:\tau_j \leq t} \Big(1 - \frac{d_j}{n_j}\Big).$$

A censored unit counts as at risk until its censoring time, and then leaves the risk set without causing an event. Without censoring, $n_{j+1} = n_j - d_j$, the product telescopes to $\frac{n_{j+1}}{N}$ for $\tau_j \leq t < \tau_{j+1}$, and $\hat{S}(t) = 1 - \hat{F}_N(t)$ is one minus the empirical CDF of Section 4.9.8. In Example 4.9.29, $\hat{S}$ drops to $\frac{5}{6}$ at $t = 2$, to $\frac{5}{6} \cdot \frac{3}{4} = \frac{5}{8}$ at $t = 4$ (only 4 customers are at risk then), and to $\frac{5}{8} \cdot \frac{1}{2} = \frac{5}{16}$ at $t = 7$. To compare groups, or to include features $\mathbf{x}$, the Cox proportional hazards model assumes $h(t \mid \mathbf{x}) = h_0(t)e^{\mathbf{w}^\top \mathbf{x}}$, with a baseline hazard h_0 that is common to all units.

4.10 Hypothesis Testing

Often we want to decide a yes-or-no question from data: is a coin fair? Does a new model have a lower error rate than the old one? Does a new website design increase sales? A hypothesis test makes such decisions while controlling the probability of a false alarm.

Definition 4.10.1 (Hypothesis test)**.** A test compares a null hypothesis H_0 (usually "no effect") with an alternative hypothesis H_1. It computes a test statistic T from the data and rejects H_0 if T falls into a rejection region. Rejecting a true H_0 is a type I error (a false positive), and keeping a false H_0 is a type II error (a false negative). A test has significance level α if the probability of a type I error is at most α; its power is the probability of rejecting H_0 when H_1 is true.

Definition 4.10.2 (p-value)**.** If large values of T speak against H_0, the p-value of an observed value t is $p = P_{H_0}(T \geq t)$: the probability, computed under H_0, of a test statistic at least as extreme as the observed one. The test that rejects H_0 if $p \leq \alpha$ has significance level α.

Proposition 4.10.3. *If T has a continuous, strictly increasing CDF F under H_0, then the p-value $1 - F(T)$ is Uniform$(0,1)$-distributed under H_0. In particular, $P_{H_0}(p \leq \alpha) = \alpha$.*

Proof. $P(F(T) \leq u) = P(T \leq F^{-1}(u)) = F(F^{-1}(u)) = u$, so $F(T)$ is uniform on $(0,1)$, as in Theorem 4.2.7, and so is $1 - F(T)$. □

Example 4.10.4 (Testing a coin)**.** A coin shows 60 heads in 100 flips. Is it fair? With H_0: $\theta = \frac{1}{2}$ and the number of heads S as the test statistic, the one-sided p-value is $P_{H_0}(S \geq 60) = 0.028$ (Example 4.7.4). If we had no reason to expect more heads than tails, a two-sided test is appropriate: it counts deviations in both directions, and its p-value is $P_{H_0}(|S - 50| \geq 10) = 0.057$. At the level $\alpha = 0.05$, the one-sided test rejects H_0, and the two-sided test does not. The test must be chosen before looking at the data; otherwise its significance level is not what it claims.

Example 4.10.5 (Power)**.** For the coin of Example 4.10.4, the one-sided test at the level 0.05 rejects H_0: $\theta = \frac{1}{2}$ if $S \geq 59$, since $P_{H_0}(S \geq 59) \approx 0.044 \leq 0.05 < P_{H_0}(S \geq 58) \approx 0.067$. If the coin actually has $\theta = 0.6$, the test rejects H_0 with the probability $P_{\theta=0.6}(S \geq 59) \approx 0.62$: it misses this bias in more than a third of the experiments. With 400 flips, the rejection region is $S \geq 217$, and the power for $\theta = 0.6$ is 0.99. How to plan the sample size for a desired power is explained in Section 4.11.

The most powerful test. Which test has the largest power at a given level? For two simple hypotheses, which each specify one distribution, the answer is the likelihood ratio test.

Theorem 4.10.6 (Neyman–Pearson lemma)**.** *Let the data* $\mathbf{x}$ *have the pmf or density* p_0 *under* H_0 *and* p_1 *under* H_1*. For* $k > 0$*, the likelihood ratio test rejects* H_0 *if* $p_1(\mathbf{x}) > kp_0(\mathbf{x})$*; let* α *be its level,* $\alpha = P_0(p_1(\mathbf{X}) > kp_0(\mathbf{X}))$*. Then every test with level at most* α *has at most the power of the likelihood ratio test.*

Proof. Let $\phi^*(\mathbf{x}) = 1$ if $p_1(\mathbf{x}) > kp_0(\mathbf{x})$ and 0 otherwise, and let $\phi(\mathbf{x}) \in \{0, 1\}$ indicate the rejection region of another test, with $\mathbb{E}_0[\phi] \le \alpha$. Then

$$\big(\phi^*(\mathbf{x}) - \phi(\mathbf{x})\big)\big(p_1(\mathbf{x}) - kp_0(\mathbf{x})\big) \ge 0 \qquad \text{for all } \mathbf{x}:$$

if $p_1(\mathbf{x}) > kp_0(\mathbf{x})$, then $\phi^*(\mathbf{x}) = 1 \ge \phi(\mathbf{x})$, and otherwise $\phi^*(\mathbf{x}) = 0 \le \phi(\mathbf{x})$. Summing or integrating over $\mathbf{x}$ gives $\mathbb{E}_1[\phi^*] - \mathbb{E}_1[\phi] \ge k\big(\mathbb{E}_0[\phi^*] - \mathbb{E}_0[\phi]\big) = k(\alpha - \mathbb{E}_0[\phi]) \ge 0$. Here $\mathbb{E}_1[\phi^*]$ and $\mathbb{E}_1[\phi]$ are the powers of the two tests. □

Example 4.10.7 (The z-test is most powerful)**.** Let $X_1, \dots, X_N$ be i.i.d. $\mathcal{N}(\mu, 1)$, and test H_0: $\mu = 0$ against H_1: $\mu = \mu_1$ with $\mu_1 > 0$. The likelihood ratio is

$$\frac{p_1(\mathbf{x})}{p_0(\mathbf{x})} = \prod_{i=1}^{N} \frac{e^{-(x_i-\mu_1)^2/2}}{e^{-x_i^2/2}} = \exp\Big(N\mu_1\bar{x} - \frac{N\mu_1^2}{2}\Big),$$

an increasing function of $\bar{x}$. So the likelihood ratio tests reject for large $\bar{x}$, and at the level 0.05, the most powerful test rejects if $\bar{x} > \frac{1.645}{\sqrt{N}}$, since $\bar{X}_N \sim \mathcal{N}(0, \frac{1}{N})$ under H_0. This is the one-sided z-test, and it does not depend on μ_1: it is the most powerful test against every $\mu_1 > 0$ at once. For $N = 25$ and $\mu_1 = 0.5$, it rejects if $\bar{x} > 0.329$, and its power is $P(\bar{X}_N > 0.329) = 1 - \Phi(\frac{0.329-0.5}{0.2}) \approx 0.80$.

In classification, the two hypotheses are the two classes, a test is a classifier, its level is the false positive rate, and its power is the true positive rate (Section 4.11.2). So a classifier that thresholds the likelihood ratio $\frac{p(\mathbf{x}|y=1)}{p(\mathbf{x}|y=0)}$, or equivalently the posterior probability $P(y = 1 \mid \mathbf{x})$, has at each threshold the largest true positive rate among all classifiers whose false positive rate is not larger. Training a classifier to predict $P(y = 1 \mid \mathbf{x})$ well is therefore the right goal for every tradeoff between the two kinds of errors.

Tests for a mean. To test H_0: $\mu = \mu_0$ for i.i.d. data, we use the standardized statistic $Z = \frac{\bar{X}_N - \mu_0}{s/\sqrt{N}}$, which is approximately $\mathcal{N}(0, 1)$ under H_0 by the central limit theorem (the z-test). The two-sided p-value is $2(1 - \Phi(|z|))$. For small samples from a Gaussian distribution, Z has a Student t-distribution with $N - 1$ degrees of freedom (Theorem 4.10.12 below, with $X = \mathbf{1}$), whose quantiles are larger than 1.96 (for example, 2.78, 2.26 and 2.05 for $N = 5$, 10 and 30); this is the t-test. Tests and confidence intervals are two sides of the same coin: the two-sided test at level 0.05 rejects $\mu = \mu_0$ exactly if μ_0 lies outside the 95% confidence interval $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$.

Example 4.10.8 (A paired t-test)**.** A new model B is compared with a model A on 8 benchmark data sets, and the differences of the accuracies (B minus A, in percentage points) are

$$1.2, \quad 0.4, \quad 2.1, \quad -0.3, \quad 1.5, \quad 0.8, \quad 1.1, \quad 0.6.$$

Their mean is $\bar{d} = 0.925$ and their standard deviation is $s = 0.729$, so $t = \frac{\bar{d}}{s/\sqrt{8}} \approx 3.59$. Under H_0: "on average, there is no difference", and if the differences are approximately Gaussian, t has the t-distribution with 7 degrees of freedom, and the two-sided p-value is 0.009. So B is significantly better at the level 0.05, and even at 0.01. Analyzing the differences, rather than the two lists of accuracies, removes the large variation between the data sets (Section 4.11).

Goodness of fit. To test whether categorical data with the counts $O_1, \dots, O_K$ in N observations follow given probabilities $p_1, \dots, p_K$, Pearson's statistic compares the observed counts with the expected counts $E_k = Np_k$:

$$\chi^2 = \sum_{k=1}^{K} \frac{(O_k - E_k)^2}{E_k}.$$

Under H_0, for large N, it has approximately the χ^2 distribution with $K-1$ degrees of freedom, the distribution of $Z_1^2 + \cdots + Z_{K-1}^2$ for independent standard normal Z_i; one degree of freedom is lost because the counts add up to N. Large values speak against H_0.

Example 4.10.9 (Is the die fair?)**.** A die is rolled 120 times, with the counts $18, 25, 16, 21, 13, 27$ for the faces $1, \dots, 6$. The expected counts are 20, so

$$\chi^2 = \frac{4 + 25 + 16 + 1 + 49 + 49}{20} = 7.2.$$

With 5 degrees of freedom, the p-value is $P(\chi_5^2 \geq 7.2) \approx 0.21$: the data are consistent with a fair die, although the counts 13 and 27 may look suspicious at first sight. The same test checks, for example, whether the class frequencies predicted by a classifier match the observed ones, or whether a random number generator is uniform.

Remark 4.10.10 (What a p-value is not)*.* A p-value is not the probability that H_0 is true. That probability depends on the prior plausibility of H_0, as in the medical test of Example 4.1.7, and only Bayesian inference can compute it. A small p-value can also come from a tiny effect that is irrelevant in practice, if N is huge (statistical versus practical significance), so effect sizes and confidence intervals should be reported as well. Conversely, a large p-value does not show that there is no effect: the test may just have too little power.

4.10.1 Inference for linear regression

A fitted regression model reports a coefficient for every feature. Is the effect of a feature real, or could the coefficient be 0? Standard errors, confidence intervals and t-tests answer this, and they are the numbers in the output tables of statistical software. We consider the model $\mathbf{y} = X\mathbf{w} + \boldsymbol{\varepsilon}$ with a fixed $N \times n$ matrix X with linearly independent columns (usually the first column is $\mathbf{1}$, for the intercept), an unknown $\mathbf{w} \in \mathbb{R}^n$, and noise with

$$\mathbb{E}[\boldsymbol{\varepsilon}] = \mathbf{0} \qquad \text{and} \qquad \mathrm{Cov}(\boldsymbol{\varepsilon}) = \sigma^2 I_N :$$

the noise terms are uncorrelated and have the same variance (homoscedasticity). Let $\hat{\mathbf{w}} = (X^\top X)^{-1} X^\top \mathbf{y}$ be the least squares estimate (Lecture 2), $\mathbf{e} = \mathbf{y} - X\hat{\mathbf{w}}$ the residuals, and $\mathrm{RSS} = \|\mathbf{e}\|^2$ the residual sum of squares. The exact distributions of the test statistics are the following.

Definition 4.10.11 (χ^2- and t-distributions)**.** For independent $Z_1, \dots, Z_k \sim \mathcal{N}(0,1)$, the distribution of $Z_1^2 + \cdots + Z_k^2$ is the χ^2-distribution with k degrees of freedom, χ_k^2. If $Z \sim \mathcal{N}(0,1)$ and $V \sim \chi_k^2$ are independent, the distribution of $\frac{Z}{\sqrt{V/k}}$ is the t-distribution with k degrees of freedom, t_k.

The χ_k^2-distribution has the mean k. The t_k-distribution is symmetric and has heavier tails than $\mathcal{N}(0,1)$, because the denominator is random; as $k \to \infty$, $\frac{V}{k} \to 1$ in probability by the law of large numbers, and t_k tends to $\mathcal{N}(0,1)$ by Slutsky's theorem. Its 97.5% quantiles are $t_{1,0.975} = 12.71$, $t_{2,0.975} = 4.30$, $t_{3,0.975} = 3.18$, $t_{10,0.975} = 2.23$ and $t_{30,0.975} = 2.04$.

Theorem 4.10.12 (Sampling distribution of least squares)**.** *In the model above:*

1. *$\hat{\mathbf{w}}$ is unbiased, $\mathbb{E}[\hat{\mathbf{w}}] = \mathbf{w}$, with the covariance matrix $\mathrm{Cov}(\hat{\mathbf{w}}) = \sigma^2(X^\top X)^{-1}$.*
2. *$\mathbb{E}[\mathrm{RSS}] = (N-n)\sigma^2$, so $s^2 = \frac{\mathrm{RSS}}{N-n}$ is an unbiased estimate of σ^2.*
3. *If $\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 I_N)$, then $\hat{\mathbf{w}} \sim \mathcal{N}\big(\mathbf{w}, \sigma^2(X^\top X)^{-1}\big)$, $\frac{\mathrm{RSS}}{\sigma^2} \sim \chi^2_{N-n}$, and $\hat{\mathbf{w}}$ and RSS are independent. Consequently, with the standard errors $\mathrm{SE}_j = s\sqrt{\big((X^\top X)^{-1}\big)_{jj}}$,*

$$\frac{\hat{w}_j - w_j}{\mathrm{SE}_j} \sim t_{N-n} \qquad \text{for } j = 1, \dots, n.$$

Proof. 1. $\hat{\mathbf{w}} = (X^\top X)^{-1}X^\top(X\mathbf{w} + \boldsymbol{\varepsilon}) = \mathbf{w} + (X^\top X)^{-1}X^\top\boldsymbol{\varepsilon}$. So $\mathbb{E}[\hat{\mathbf{w}}] = \mathbf{w}$, and by Proposition 4.5.2, $\mathrm{Cov}(\hat{\mathbf{w}}) = (X^\top X)^{-1}X^\top(\sigma^2 I)X(X^\top X)^{-1} = \sigma^2(X^\top X)^{-1}$.

2. Let $H = X(X^\top X)^{-1}X^\top$ be the matrix of the orthogonal projection onto $\mathrm{col}(X)$ (Lecture 2). Then $(I-H)X = 0$, so $\mathbf{e} = (I-H)\mathbf{y} = (I-H)\boldsymbol{\varepsilon}$, and since $I-H$ is symmetric with $(I-H)^2 = I-H$, $\mathrm{RSS} = \boldsymbol{\varepsilon}^\top(I-H)\boldsymbol{\varepsilon}$. For every matrix M, $\mathbb{E}[\boldsymbol{\varepsilon}^\top M\boldsymbol{\varepsilon}] = \sum_{i,j} M_{ij}\mathbb{E}[\varepsilon_i\varepsilon_j] = \sigma^2 \operatorname{tr}(M)$, and $\operatorname{tr}(I-H) = N - \operatorname{tr}\big((X^\top X)^{-1}X^\top X\big) = N-n$, since $\operatorname{tr}(AB) = \operatorname{tr}(BA)$.

3. Choose an orthonormal basis $\mathbf{u}_1, \dots, \mathbf{u}_n$ of $\mathrm{col}(X)$ and an orthonormal basis $\mathbf{u}_{n+1}, \dots, \mathbf{u}_N$ of $\mathrm{col}(X)^\perp$ (Lecture 2), and let Q be the orthogonal matrix with the columns $\mathbf{u}_1, \dots, \mathbf{u}_N$. By Corollary 4.6.7, $Q^\top\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 Q^\top Q) = \mathcal{N}(\mathbf{0}, \sigma^2 I)$, so the numbers $Z_i = \frac{1}{\sigma}\mathbf{u}_i^\top\boldsymbol{\varepsilon}$ are independent and $\mathcal{N}(0,1)$ (Theorem 4.6.4). Now $(I-H)\boldsymbol{\varepsilon} = \sum_{i>n}(\mathbf{u}_i^\top\boldsymbol{\varepsilon})\mathbf{u}_i$, so $\mathrm{RSS} = \sigma^2\sum_{i>n} Z_i^2$, and $\frac{\mathrm{RSS}}{\sigma^2} \sim \chi^2_{N-n}$. On the other hand, $X^\top = X^\top H$ (since $HX = X$), so $\hat{\mathbf{w}} - \mathbf{w} = (X^\top X)^{-1}X^\top H\boldsymbol{\varepsilon}$, and $H\boldsymbol{\varepsilon} = \sigma\sum_{i\le n} Z_i\mathbf{u}_i$ only depends on $Z_1, \dots, Z_n$. Functions of the independent groups $Z_1, \dots, Z_n$ and $Z_{n+1}, \dots, Z_N$ are independent, so $\hat{\mathbf{w}}$ and RSS are independent. The matrix $(X^\top X)^{-1}X^\top$ has rank n, so $\hat{\mathbf{w}}$ is Gaussian by Corollary 4.6.7, with the mean and covariance matrix of part 1. Finally, with $c_j = ((X^\top X)^{-1})_{jj}$, the number $\frac{\hat{w}_j - w_j}{\sigma\sqrt{c_j}}$ is $\mathcal{N}(0,1)$, it is independent of $V = \frac{\mathrm{RSS}}{\sigma^2} \sim \chi^2_{N-n}$, and dividing it by $\sqrt{V/(N-n)} = \frac{s}{\sigma}$ gives $\frac{\hat{w}_j - w_j}{\mathrm{SE}_j}$. □

So $\hat{w}_j \pm t_{N-n,0.975}\,\mathrm{SE}_j$ is an exact 95% confidence interval for w_j under Gaussian noise (and an approximate one in general, for large N, by the central limit theorem), and H_0: $w_j = 0$ ("the feature has no effect") is tested with the t-statistic $\frac{\hat{w}_j}{\mathrm{SE}_j}$. The one-sample t-test of a mean is the special case $X = \mathbf{1}$, with $\hat{w} = \bar{x}$ and $\mathrm{SE} = \frac{s}{\sqrt{N}}$. Two degrees of freedom are lost in a regression with an intercept and a slope, one for each estimated coefficient, as in the factor $\frac{1}{N-1}$ of the sample variance.

Example 4.10.13 (Standard errors of a regression line)**.** For the five points $x = -2, -1, 0, 1, 2$ with $y = 0.8, 2.3, 2.9, 4.4, 5.6$, the design matrix has the columns $\mathbf{1}$ and $\mathbf{x}$, and $X^\top X = \mathrm{diag}(5, 10)$, since $\sum_i x_i = 0$. So $\hat{w}_0 = \bar{y} = 3.2$, $\hat{w}_1 = \frac{\sum_i x_i y_i}{10} = 1.17$, the residuals are $-0.06, 0.27, -0.30, 0.03, 0.06$, $\mathrm{RSS} = 0.171$, and $s^2 = \frac{0.171}{5-2} = 0.057$. The standard error of the slope is $\sqrt{0.057/10} \approx 0.0755$, its t-statistic is 15.5, with a two-sided p-value of 0.0006 (3 degrees of freedom), and the 95% confidence interval is $1.17 \pm 3.18 \cdot 0.0755 = [0.93, 1.41]$.

Least squares is not only unbiased, it is also the most precise among the unbiased estimators that are linear in $\mathbf{y}$.

Proposition 4.10.14 (Gauss–Markov)**.** *Let $\tilde{\mathbf{w}} = C\mathbf{y}$, with an $n \times N$ matrix C, be unbiased for every $\mathbf{w}$. Then $\mathrm{Cov}(\tilde{\mathbf{w}}) - \mathrm{Cov}(\hat{\mathbf{w}})$ is positive semidefinite; in particular, $\mathrm{Var}(\tilde{w}_j) \ge \mathrm{Var}(\hat{w}_j)$ for every j.*

Proof. $\mathbb{E}[C\mathbf{y}] = CX\mathbf{w} = \mathbf{w}$ for all $\mathbf{w}$ means $CX = I$. Let $D = C - (X^\top X)^{-1}X^\top$, so that $DX = 0$. By Proposition 4.5.2, $\mathrm{Cov}(\tilde{\mathbf{w}}) = \sigma^2 CC^\top = \sigma^2\big((X^\top X)^{-1} + DD^\top\big)$, since the cross terms $(X^\top X)^{-1}X^\top D^\top = (X^\top X)^{-1}(DX)^\top$ vanish. And $DD^\top$ is positive semidefinite. □

Biased estimators such as ridge regression can still have a smaller mean squared error (Section 4.9): Gauss–Markov only compares unbiased ones.

Heteroscedasticity. Often the noise is larger for some data points than for others: prices of expensive houses vary more than those of cheap ones, and measurements of large quantities are less precise. If $\mathrm{Cov}(\boldsymbol{\varepsilon}) = \Sigma = \mathrm{diag}(\sigma_1^2, \dots, \sigma_N^2)$ with different variances (heteroscedasticity), the estimate $\hat{\mathbf{w}}$ is still unbiased, but by Proposition 4.5.2, its covariance matrix is the "sandwich"

$$\mathrm{Cov}(\hat{\mathbf{w}}) = (X^\top X)^{-1} X^\top \Sigma X (X^\top X)^{-1},$$

and the formula $s^2(X^\top X)^{-1}$ is wrong, together with the standard errors, confidence intervals and t-tests based on it. For example, in the model $y_i = wx_i + \varepsilon_i$ without intercept, with the noise standard deviations $\sigma_i = \sigma|x_i|$ that grow with $|x_i|$, the true variance is $\mathrm{Var}(\hat{w}) = \frac{\sigma^2 \sum_i x_i^4}{(\sum_i x_i^2)^2}$. The homoscedastic formula, with the average noise variance $\bar{\sigma}^2 = \frac{\sigma^2}{N}\sum_i x_i^2$, gives $\frac{\bar{\sigma}^2}{\sum_i x_i^2} = \frac{\sigma^2}{N}$. Their ratio is $\frac{N \sum_i x_i^4}{(\sum_i x_i^2)^2} \geq 1$, by the Cauchy–Schwarz inequality for the vectors $(x_i^2)_i$ and $\mathbf{1}$, with equality only if all $|x_i|$ are equal. For $x = 1, 2, \dots, 10$, the ratio is 1.71: the usual standard error is too small by a factor of 1.31, and the test finds effects too often. Heteroscedasticity is detected by plotting the residuals against the fitted values or against a feature, which shows a funnel shape. The remedies are robust (White) standard errors, which replace Σ in the sandwich by the estimate $\mathrm{diag}(e_1^2, \dots, e_N^2)$, and, if the variances are known up to a factor, weighted least squares, which minimizes $\sum_i \frac{(y_i - \mathbf{w}^\top \mathbf{x}_i)^2}{\sigma_i^2}$ and is the maximum likelihood estimate for Gaussian noise with these variances.

4.10.2 Multiple testing

If we perform many tests, some of them will reject by chance. For m independent tests of true null hypotheses at the level 0.05, the probability of at least one false positive is $1 - 0.95^m$, which is 0.64 for $m = 20$. This happens when we compare many models on the same test set, screen thousands of genes or features, or analyze many metrics of an experiment, and it is a main source of irreproducible results ("p-hacking"). The probability of at least one false positive among all tests is called the family-wise error rate (FWER).

Proposition 4.10.15 (Bonferroni correction)**.** *If each of m tests rejects its null hypothesis when its p-value is at most $\frac{\alpha}{m}$, then the family-wise error rate is at most α, whatever the dependence between the tests.*

Proof. Let I_0 be the set of true null hypotheses. By the union bound (Proposition 4.1.3) and Proposition 4.10.3, $P\big(\bigcup_{i \in I_0} \{p_i \leq \frac{\alpha}{m}\}\big) \leq \sum_{i \in I_0} \frac{\alpha}{m} \leq \alpha$. □

The Bonferroni correction is very conservative when m is large. When we screen many hypotheses, it is often more useful to control the false discovery rate (FDR), the expected fraction of false positives among all rejected hypotheses: if R hypotheses are rejected, V of them true, then $\mathrm{FDR} = \mathbb{E}\big[\frac{V}{\max(R,1)}\big]$. The Benjamini–Hochberg procedure sorts the p-values $p_{(1)} \leq \cdots \leq p_{(m)}$, finds the largest k with $p_{(k)} \leq \frac{k}{m} q$ ($k = 0$ if there is none), and rejects the k hypotheses with the smallest p-values.

Theorem 4.10.16 (Benjamini–Hochberg)**.** *Suppose that the m p-values are independent, and that the p-value p_i of every true null hypothesis satisfies $P(p_i \leq \alpha) \leq \alpha$ for all $\alpha \in [0, 1]$ (as in Proposition 4.10.3). Then the false discovery rate of the Benjamini–Hochberg procedure is at most $\frac{m_0}{m} q \leq q$, where m_0 is the number of true null hypotheses.*

Proof. Let $N(k)$ be the number of p-values that are at most $\frac{k}{m}q$. Since $p_{(k)} \le \frac{k}{m}q$ if and only if $N(k) \ge k$, the number R of rejections is the largest $k \in \{0, \dots, m\}$ with $N(k) \ge k$. Exactly R p-values are at most $\frac{R}{m}q$: at least R by definition, and if $R+1$ of them were, then $N(R+1) \ge R+1$. So hypothesis i is rejected if and only if $p_i \le \frac{R}{m}q$.

Let i be a true null hypothesis, and let $N^{(i)}(k)$ and $R^{(i)}$ be defined in the same way after replacing p_i by 0; then $R^{(i)}$ is a function of the other p-values, so it is independent of p_i. Let $k \ge 1$, and suppose that $p_i \le \frac{k}{m}q$. Then $N(k') = N^{(i)}(k')$ for all $k' \ge k$. If $R = k$, then $N^{(i)}(k) = N(k) \ge k$ and $N^{(i)}(k') = N(k') < k'$ for $k' > k$, so $R^{(i)} = k$. Conversely, if $R^{(i)} = k$, then $N(k) = N^{(i)}(k) \ge k$ and $N(k') = N^{(i)}(k') < k'$ for $k' > k$, so $R = k$. Hence the events $\{p_i \le \frac{k}{m}q,\, R = k\}$ and $\{p_i \le \frac{k}{m}q,\, R^{(i)} = k\}$ are equal, and

$$\begin{aligned}\mathbb{E}\Big[\frac{\mathbf{1}_{\{i \text{ rejected}\}}}{\max(R,1)}\Big] &= \sum_{k=1}^{m} \frac{1}{k} P\Big(p_i \le \frac{k}{m}q,\, R = k\Big) = \sum_{k=1}^{m} \frac{1}{k} P\Big(p_i \le \frac{k}{m}q\Big) P\big(R^{(i)} = k\big) \\ &\le \sum_{k=1}^{m} \frac{q}{m} P\big(R^{(i)} = k\big) \le \frac{q}{m}.\end{aligned}$$

Since V is the number of rejected true null hypotheses, adding this over the m_0 true null hypotheses gives $\mathrm{FDR} \le \frac{m_0}{m}q$. □

Example 4.10.17. Ten tests give the p-values

$$0.001,\ 0.008,\ 0.012,\ 0.030,\ 0.041,\ 0.10,\ 0.25,\ 0.40,\ 0.62,\ 0.90.$$

Without correction, five are below 0.05. Bonferroni with $\alpha = 0.05$ uses the threshold 0.005 and rejects only the first. Benjamini–Hochberg with $q = 0.05$ compares $p_{(k)}$ with $0.005k$: $0.001 \le 0.005$, $0.008 \le 0.010$ and $0.012 \le 0.015$, but $p_{(k)} > 0.005k$ for all $k \ge 4$, so it rejects three hypotheses.

4.10.3 The bootstrap and permutation tests

The formulas for standard errors and p-values so far required either a simple statistic, such as a mean, or a known distribution. Two resampling methods work without such formulas, at the cost of computation.

The bootstrap. To estimate the uncertainty of a statistic $\hat{\theta} = s(x_1, \dots, x_N)$, such as a median, a correlation, or the AUC of a classifier, the bootstrap replaces the unknown distribution of the data by the empirical distribution (the plug-in principle). It draws B bootstrap samples $x_1^*, \dots, x_N^*$, each consisting of N draws with replacement from the data, and computes $\hat{\theta}_b^* = s(x_1^*, \dots, x_N^*)$ for $b = 1, \dots, B$. The standard deviation of the $\hat{\theta}_b^*$ estimates the standard error of $\hat{\theta}$, and the 2.5% and 97.5% quantiles of the $\hat{\theta}_b^*$ give an approximate 95% confidence interval (the percentile interval). For the sample mean, everything can be computed exactly: a bootstrap sample consists of N i.i.d. draws from the empirical distribution, whose variance is $\hat{\sigma}^2 = \frac{1}{N}\sum_i (x_i - \bar{x})^2$, so the bootstrap standard error of the mean is $\frac{\hat{\sigma}}{\sqrt{N}}$, the familiar formula. The bootstrap can fail for very small samples, for statistics such as the maximum, and for dependent data.

Permutation tests. Suppose we observe $x_1, \dots, x_n$ in group A and $y_1, \dots, y_m$ in group B, and want to test H_0: both groups come from the same distribution. Under H_0, the group labels carry no information: given the $n + m$ observed values, each of the $\binom{n+m}{n}$ ways to assign n of them to group A is equally likely. So the p-value of a statistic such as the difference of the group means is the fraction of all relabelings whose statistic is at least as large as the observed one. For large samples, we use a few thousand random relabelings instead of all of them. The test has exactly the claimed level, and it requires no assumptions about the distribution.

Example 4.10.18. Let group A have the values $12, 15, 14$ and group B the values $9, 10, 13$. The observed difference of the means is $13\frac{2}{3} - 10\frac{2}{3} = 3$. Of the $\binom{6}{3} = 20$ ways to split the six values into two groups of three, only 2 give a difference of at least 3: the observed split, and the split $\{13, 14, 15\}$ versus $\{9, 10, 12\}$, with the difference $3\frac{2}{3}$. So the one-sided p-value is $\frac{2}{20} = 0.1$: with so few observations, even a clear difference is not significant.

4.11 Evaluating Models and Experiments

4.11.1 Cross-validation

To choose between models or hyperparameters, we need to estimate how well a learning algorithm generalizes, and we often cannot afford to set aside a large test set. In k-fold cross-validation, the data are split into k folds of about equal size; for $j = 1, \dots, k$, the model is trained on all folds except the j-th one and evaluated on the j-th fold, which gives an error e_j. The cross-validation estimate is the average $\bar{e} = \frac{1}{k}\sum_j e_j$. Typical choices are $k = 5$ or 10; for $k = N$, it is leave-one-out cross-validation. The statistical properties of $\bar{e}$ are subtle.

- Bias. Each model is trained on only a fraction $\frac{k-1}{k}$ of the data, so $\bar{e}$ slightly overestimates the error of a model trained on all data; this bias is smaller for larger k.
- Variance. The fold errors are not independent, since the training sets overlap. So the naive standard error $\frac{s_e}{\sqrt{k}}$, with the standard deviation s_e of the fold errors, usually underestimates the true uncertainty; one can show that no estimator of the variance of $\bar{e}$ is unbiased for all distributions. For example, the fold errors $0.12, 0.15, 0.10, 0.14, 0.13$ have the mean 0.128, the standard deviation 0.019 and the naive standard error 0.009; the true uncertainty is larger. Repeating cross-validation with different random splits reduces the variance that comes from the split, but not the dependence.
- Comparing two models. Comparing two models on the same folds (or the same test examples) and analyzing the differences is much more precise than comparing their separate error estimates, since $\mathrm{Var}(A - B) = \mathrm{Var}(A) + \mathrm{Var}(B) - 2\,\mathrm{Cov}(A, B)$ and the errors of two models on the same data are positively correlated. This is the idea of paired tests.
- Selection bias. If we evaluate many models or hyperparameter settings and choose the one with the smallest cross-validation error, the selected error estimate is biased downward: by Jensen's inequality, the expected minimum of noisy estimates is smaller than the minimum of their expectations. For example, if 20 models all have the true error 0.10 and their estimates have independent errors with standard deviation 0.01, the smallest estimate is about $0.10 - 1.87 \cdot 0.01 \approx 0.081$ on average (the expected maximum of 20 independent standard normal variables is about 1.87). The remedy is to estimate the error of the selected model on data that played no role in the selection: a separate test set, used once, or nested cross-validation, which repeats the whole selection procedure inside each training fold.

4.11.2 Classification metrics

The accuracy of a binary classifier hides which errors it makes. A classifier usually computes a score $s(\mathbf{x})$, such as an estimate of $P(y = 1 \mid \mathbf{x})$, and predicts $\hat{y} = 1$ if $s(\mathbf{x}) \geq t$ for a threshold t. On a test set, the confusion matrix counts the true positives (TP: $\hat{y} = 1$, $y = 1$), false positives (FP: $\hat{y} = 1$, $y = 0$), false negatives (FN: $\hat{y} = 0$, $y = 1$) and true negatives (TN: $\hat{y} = 0$, $y = 0$). From these counts, one computes

$$\text{precision} = \frac{\mathrm{TP}}{\mathrm{TP} + \mathrm{FP}}, \qquad \text{recall} = \mathrm{TPR} = \frac{\mathrm{TP}}{\mathrm{TP} + \mathrm{FN}}, \qquad \mathrm{FPR} = \frac{\mathrm{FP}}{\mathrm{FP} + \mathrm{TN}},$$

and the F_1 score, the harmonic mean $\frac{2}{1/\text{precision}+1/\text{recall}} = \frac{2\text{TP}}{2\text{TP}+\text{FP}+\text{FN}}$ of precision and recall. The recall (true positive rate) is the fraction of the positives that are found, the precision is the fraction of the positive predictions that are correct, and the false positive rate is the fraction of the negatives that raise a false alarm. In the language of hypothesis testing, the FPR is the level of the test "H_0: $y = 0$", and the recall is its power.

When positives are rare, accuracy is misleading: if 1% of the transactions are fraudulent, the classifier that never predicts fraud has an accuracy of 99%, and recall 0. Precision depends on the base rate, as in the medical test of Example 4.1.7: by Bayes' theorem, precision is $P(y = 1 \mid \hat{y} = 1) = \frac{\text{TPR}\cdot\pi_1}{\text{TPR}\cdot\pi_1+\text{FPR}\cdot(1-\pi_1)}$ for the fraction π_1 of positives, so the same classifier has a much lower precision on data with fewer positives.

ROC curves and the AUC. Lowering the threshold t finds more positives, but also raises more false alarms. The ROC curve (receiver operating characteristic) shows the pairs $(\text{FPR}(t), \text{TPR}(t))$ for all thresholds; it goes from $(0, 0)$ (nothing predicted positive) to $(1, 1)$ (everything predicted positive). A classifier with random scores has the diagonal as its ROC curve, and a perfect one goes through $(0, 1)$. The area under the ROC curve (AUC) summarizes the curve in one number, which has a simple meaning.

Theorem 4.11.1. *Let a test set have n_+ positives and n_- negatives, with distinct scores. Then the area under the ROC curve is the fraction of the n_+n_- pairs of a positive and a negative example in which the positive example has the higher score.*

Proof. Sort the examples by decreasing score, and lower the threshold past them one at a time. Passing a positive example raises the TPR by $\frac{1}{n_+}$, and passing a negative example raises the FPR by $\frac{1}{n_-}$. So the ROC curve is a staircase, and its area is the sum of the areas of the strips of width $\frac{1}{n_-}$ below its horizontal steps, one for each negative example. The strip of the negative example j has the height $\frac{1}{n_+}$ times the number of positives with a higher score than j. Summing over j gives the number of correctly ordered pairs, divided by n_+n_-. □

So the AUC estimates the probability that a random positive example gets a higher score than a random negative one. It only depends on the order of the scores, not on a threshold, and it does not change if the classes are rebalanced. An AUC of 0.5 is no better than random guessing.

Example 4.11.2. Ten test examples have the scores $0.95, 0.85, 0.8, 0.7, 0.6, 0.45, 0.4, 0.3, 0.2, 0.1$, and the labels $1, 1, 0, 1, 1, 0, 1, 0, 0, 0$. With the threshold 0.5, TP = 4, FP = 1 (the score 0.8), FN = 1 (the score 0.4) and TN = 4, so the accuracy, precision, recall and F_1 are all 0.8, and FPR = 0.2. Of the 25 positive–negative pairs, 21 are ordered correctly (the positive with the score 0.7 is below one negative, 0.8, and so are the positive 0.6, and the positive 0.4, which is also below 0.45), so the AUC is 0.84.

A classifier is calibrated if, among the examples with the predicted probability q, a fraction q are positive, so that its scores can be read as probabilities. The cross-entropy loss rewards calibration: if the true probability of $y = 1$ is p, the expected loss $-p \log q - (1 - p) \log(1 - q) = H(p, q)$ of predicting q is smallest for $q = p$, by Gibbs' inequality (Theorem 4.13.2). Losses with this property are called proper scoring rules; the accuracy is not one, since it only rewards being on the right side of $\frac{1}{2}$.

4.11.3 Generalization bounds

A model chosen because it fits the training data well will usually do worse on new data. How much worse can be bounded with Hoeffding's inequality and the union bound. Let $\ell(h; \mathbf{x}, y) \in [0, 1]$ be the loss of a model h on the example $(\mathbf{x}, y)$, for example the indicator of a misclassification.

The risk $R(h) = \mathbb{E}[\ell(h; \mathbf{X}, Y)]$ is the expected loss on new data, and the empirical risk $\hat{R}(h) = \frac{1}{N}\sum_{i=1}^{N} \ell(h; \mathbf{x}_i, y_i)$ is the average loss on N i.i.d. training examples.

Theorem 4.11.3 (Generalization bound for finitely many models)**.** *Let $\mathcal{H}$ be a finite set of models, and let $0 < \delta < 1$. With probability at least $1 - \delta$ over the training data,*

$$|R(h) - \hat{R}(h)| \le \varepsilon = \sqrt{\frac{\log(2|\mathcal{H}|/\delta)}{2N}} \qquad \text{for all } h \in \mathcal{H} \text{ simultaneously.}$$

In particular, the empirical risk minimizer $\hat{h}$, the model in $\mathcal{H}$ with the smallest $\hat{R}$, satisfies $R(\hat{h}) \le \min_{h\in\mathcal{H}} R(h) + 2\varepsilon$ with probability at least $1 - \delta$.

Proof. For a fixed h, $\hat{R}(h)$ is a mean of N i.i.d. variables with values in $[0, 1]$ and mean $R(h)$, so by Hoeffding's inequality (Theorem 4.7.5), $P\big(|R(h) - \hat{R}(h)| > \varepsilon\big) \le 2e^{-2N\varepsilon^2} = \frac{\delta}{|\mathcal{H}|}$. By the union bound, the probability that this happens for at least one $h \in \mathcal{H}$ is at most δ. If it happens for none, then for every $h \in \mathcal{H}$, $R(\hat{h}) \le \hat{R}(\hat{h}) + \varepsilon \le \hat{R}(h) + \varepsilon \le R(h) + 2\varepsilon$. □

The price for choosing among many models is the factor $\log|\mathcal{H}|$, which grows slowly. For a single model evaluated on $N = 10\,000$ independent test examples and $\delta = 0.05$, the bound is $\varepsilon = 0.014$; for the best of $|\mathcal{H}| = 10^6$ models, it is 0.030. This is why a model must be evaluated on data that played no role in choosing it (Section 4.11): on the training data, the factor $\log|\mathcal{H}|$ is not paid, and the error estimate is too optimistic. A model with d parameters stored as 32-bit numbers is one of at most 2^{32d} models, so $\log|\mathcal{H}| \le 32d \log 2$, and the bound is of the order $\sqrt{d/N}$: we need more data points than parameters. Neural networks often generalize much better than this; statistical learning theory replaces $\log|\mathcal{H}|$ by finer measures of complexity, such as the VC dimension, and explaining the generalization of large networks is an active research area.

4.11.4 A/B testing and experiment design

To find out whether a change causes an improvement, for example a new recommendation model, the most reliable method is a randomized controlled experiment, called an A/B test in industry: each user is randomly assigned to version A (the control) or version B (the treatment), and the outcomes are compared. Randomization makes the two groups comparable in all respects, observed or not, on average. So a significant difference can be attributed to the treatment. Observational comparisons, such as "users who chose the new feature buy more", can be confounded by other differences between the groups.

Confounding and Simpson's paradox. A classic medical study compared two treatments for kidney stones:

	small stones	large stones	all
treatment A	81/87 = 93%	192/263 = 73%	273/350 = 78%
treatment B	234/270 = 87%	55/80 = 69%	289/350 = 83%

Treatment A is more successful for small stones and for large stones, but less successful overall. The reason is that A was mostly given for large stones (263 of 350 patients), which are harder to treat, and B mostly for small ones. By the law of total probability, $P(\text{success} \mid A) = \sum_s P(\text{success} \mid A, s)\, P(s \mid A)$ averages over the stone sizes s with weights $P(s \mid A)$ that differ between the treatments; the stone size is a confounder. Weighting both treatments with the same distribution of stone sizes, that of all 700 patients (51% small), gives $0.51 \cdot 0.93 + 0.49 \cdot 0.73 = 0.83$ for A and $0.51 \cdot 0.87 + 0.49 \cdot 0.69 = 0.78$ for B, the right comparison if the size is the only confounder. Randomization makes $P(s \mid A) = P(s \mid B)$ on average for every confounder s, including the ones we do not know about.

Analyzing an A/B test. Let $\hat{p}_A$ and $\hat{p}_B$ be the conversion rates (for example, the fractions of users who buy something) of n_A and n_B users. Under H_0: $p_A = p_B$, the pooled rate $\hat{p}$ of all users estimates the common rate, and by the central limit theorem,

$$Z = \frac{\hat{p}_B - \hat{p}_A}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{n_A} + \frac{1}{n_B}\right)}}$$

is approximately $\mathcal{N}(0,1)$ (the two-proportion z-test). For example, if 200 of 5000 users convert with version A (4%) and 250 of 5000 with version B (5%), then $\hat{p} = 0.045$, $Z \approx 2.41$, and the two-sided p-value is 0.016. An approximate 95% confidence interval for the improvement $p_B - p_A$ is $0.01 \pm 1.96\sqrt{\frac{\hat{p}_A(1-\hat{p}_A)}{n_A} + \frac{\hat{p}_B(1-\hat{p}_B)}{n_B}} = [0.002, 0.018]$.

Sample size and power. How many users do we need to detect an improvement $\delta = p_B - p_A$? With n users per group, the standard error of $\hat{p}_B - \hat{p}_A$ is about $\sqrt{2\bar{p}(1-\bar{p})/n}$, where $\bar{p}$ is the average rate, and Z is approximately $\mathcal{N}(\delta/\mathrm{SE}, 1)$. The two-sided test at level α rejects if $|Z| > z_{1-\alpha/2}$, so it has power $1-\beta$ approximately when $\frac{\delta}{\mathrm{SE}} = z_{1-\alpha/2} + z_{1-\beta}$, i.e.,

$$n \approx \frac{(z_{1-\alpha/2} + z_{1-\beta})^2 \cdot 2\bar{p}(1-\bar{p})}{\delta^2}$$

per group, where $z_u = \Phi^{-1}(u)$. For $\alpha = 0.05$ and power 0.8, $z_{0.975} + z_{0.8} \approx 1.96 + 0.84$; to detect an increase from 4% to 5%, we need about 6750 users per group. Halving the effect that we want to detect quadruples the sample size.

Design principles and pitfalls.

- Fix the primary metric, the test and the sample size in advance. Peeking at the p-value repeatedly and stopping as soon as it is below 0.05 inflates the false positive rate: in a simulation with two identical versions (conversion rate 5%, up to 10 000 users per group), checking after every 200 users per group declared a significant difference in about one third of the runs. If early stopping is needed, use sequential tests designed for it, such as the likelihood ratio test of Example B.8.2.
- Many metrics or user segments mean many tests: correct for multiple testing, or treat secondary results as hypotheses for future experiments.
- Randomize at the right level. If users influence each other (for example in a social network or a marketplace), or one user is counted several times, the observations are not independent, and the standard errors above are too small.
- Reduce variance by design. Paired or blocked designs compare similar units (for example, the same users or the same queries under both versions), and adjusting for measurements taken before the experiment can reduce the variance considerably.
- Watch for novelty effects, which make new versions look better at first, and check that the groups have the planned sizes; a mismatch indicates a bug in the assignment.

The same principles apply to offline comparisons of machine learning models: whether model B is really better than model A is a hypothesis testing question, and many reported improvements are within the noise of the evaluation.

4.12 Bayesian Inference

Maximum likelihood treats the parameter θ as a fixed unknown number. Bayesian inference treats it as a random variable, which expresses our uncertainty about it. Before seeing data, our beliefs are described by a prior distribution $p(\theta)$; after seeing the data $\mathcal{D}$, Bayes' theorem gives the posterior distribution

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}, \qquad p(\mathcal{D}) = \int p(\mathcal{D} \mid \theta)\, p(\theta)\, d\theta.$$

In words: the posterior is proportional to the likelihood times the prior. The normalizing constant $p(\mathcal{D})$ is called the evidence; it does not depend on θ.

4.12.1 Coin flips and the Beta distribution

Definition 4.12.1 (Beta distribution)**.** For $a, b > 0$, the Beta(a, b) distribution on $[0, 1]$ has the density

$$p(\theta) = \frac{\theta^{a-1}(1-\theta)^{b-1}}{B(a,b)}, \qquad \text{where} \quad B(a,b) = \int_0^1 \theta^{a-1}(1-\theta)^{b-1}\, d\theta.$$

For positive integers, $B(a,b) = \frac{(a-1)!\,(b-1)!}{(a+b-1)!}$; for example, Beta$(1,1)$ is the uniform distribution, and Beta$(2,2)$ has the density $6\theta(1-\theta)$. The mean of Beta(a,b) is $\frac{a}{a+b}$, and for $a, b > 1$ its mode (the maximum of the density) is $\frac{a-1}{a+b-2}$.

Theorem 4.12.2 (Beta–Bernoulli conjugacy)**.** *Let the prior of θ be Beta(a,b), and let the data be N independent Bernoulli(θ) observations with k ones. Then the posterior is Beta$(a+k,\ b+N-k)$.*

Proof. $p(\theta \mid \mathcal{D}) \propto \theta^k(1-\theta)^{N-k} \cdot \theta^{a-1}(1-\theta)^{b-1} = \theta^{a+k-1}(1-\theta)^{b+N-k-1}$. The only density proportional to this function is that of Beta$(a+k, b+N-k)$. □

A prior is called conjugate to a likelihood if the posterior belongs to the same family as the prior; then updating our beliefs amounts to updating a few parameters. Here, the prior acts like a previously seen ones and b previously seen zeros.

Example 4.12.3. With the prior Beta$(2,2)$, which expresses a mild belief that the coin is fair, and three heads in three flips, the posterior is Beta$(5,2)$. Its mean is $\frac{5}{7} \approx 0.71$ and its mode is $\frac{4}{5} = 0.8$, whereas the MLE was 1 (Example 4.9.8). After 30 flips with 20 heads, the posterior is Beta$(22,12)$, with mean 0.65 and standard deviation 0.08: with more data, the posterior concentrates, and the prior matters less (Figure 4.5).

To predict the next flip, Bayesian inference averages over all values of θ, weighted by the posterior:

$$P(X_{N+1} = 1 \mid \mathcal{D}) = \int_0^1 \theta\, p(\theta \mid \mathcal{D})\, d\theta = \mathbb{E}[\theta \mid \mathcal{D}] = \frac{a+k}{a+b+N}.$$

For the uniform prior ($a = b = 1$), this is Laplace's rule of succession $\frac{k+1}{N+2}$: after three heads in three flips, the probability of heads is $\frac{4}{5}$, not 1. The same "add pseudo-counts" idea (Laplace smoothing) prevents zero probabilities for unseen words in language models.

Example 4.12.4 (Credible intervals)**.** A 95% credible interval contains θ with posterior probability 0.95, for example the interval between the 2.5% and the 97.5% quantiles of the posterior. For the posterior Beta$(22,12)$ after 20 heads in 30 flips (Example 4.12.3), it is $[0.48, 0.80]$ (computed numerically), close to the approximate confidence interval $\frac{2}{3} \pm 1.96\sqrt{\frac{2}{3} \cdot \frac{1}{3}/30} = [0.50, 0.84]$ of Section 4.9. The interpretations differ: the credible interval makes a probability statement about θ, given the data, while the confidence interval makes a probability statement about the random interval, for a fixed θ. As the amount of data grows, the two intervals agree more and more (the Bernstein–von Mises theorem).




Figure 4.5: Bayesian updating for the probability θ of heads: the prior Beta$(2,2)$ (blue), the posterior Beta$(5,2)$ after 3 heads in 3 flips (orange), and the posterior Beta$(22,12)$ after 20 heads in 30 flips (red). With more data, the posterior concentrates around the relative frequency of heads.

Example 4.12.5 (Gamma–Poisson conjugacy)**.** Let the numbers of clicks on an ad in N hours be i.i.d. Poisson(λ), and let the prior of λ be the Gamma distribution with the shape a and the rate b, i.e., with the density $\frac{b^a}{\Gamma(a)}\lambda^{a-1}e^{-b\lambda}$ for $\lambda > 0$ and the mean $\frac{a}{b}$ (the Gamma distribution of Example 4.9.6 with the scale $\frac{1}{b}$). If S is the total number of clicks, the posterior is

$$p(\lambda \mid \mathcal{D}) \propto e^{-N\lambda}\lambda^{S} \cdot \lambda^{a-1}e^{-b\lambda} = \lambda^{a+S-1}e^{-(b+N)\lambda},$$

i.e., Gamma$(a+S, b+N)$: the prior acts like b previous hours with a clicks in total. The posterior mean $\frac{a+S}{b+N}$ lies between the prior mean $\frac{a}{b}$ and the MLE $\frac{S}{N}$. For the prior Gamma$(2,1)$ and the counts $3,5,4,6,2$ in five hours, the posterior is Gamma$(22,6)$, with the mean $\frac{22}{6} \approx 3.67$ (the MLE is 4) and the standard deviation $\frac{\sqrt{22}}{6} \approx 0.78$. The predictive probability of an hour without clicks is $\int e^{-\lambda}\, p(\lambda \mid \mathcal{D})\, d\lambda = \left(\frac{6}{7}\right)^{22} \approx 0.034$, while plugging in the MLE gives $e^{-4} \approx 0.018$: the posterior accounts for the uncertainty about λ.

4.12.2 Maximum a posteriori estimation and regularization

Often we only want a single parameter value. The maximum a posteriori (MAP) estimate is the mode of the posterior,

$$\hat{\theta}_{\mathrm{MAP}} = \arg\max_{\theta} p(\theta \mid \mathcal{D}) = \arg\min_{\theta} \big(-\log p(\mathcal{D} \mid \theta) - \log p(\theta) \big),$$

since $p(\mathcal{D})$ does not depend on θ. So MAP estimation minimizes the negative log-likelihood plus a penalty $-\log p(\theta)$ coming from the prior: a prior is a regularizer.

Proposition 4.12.6 (Ridge regression as MAP estimation)**.** *In the linear regression model of Proposition 4.9.19, with the prior* $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1}I)$*, the MAP estimate minimizes* $\|\mathbf{y} - X\mathbf{w}\|^2 + \lambda\|\mathbf{w}\|^2$ *with* $\lambda = \alpha\sigma^2$*, i.e., it is the ridge regression solution* $\hat{\mathbf{w}} = (X^\top X + \lambda I)^{-1}X^\top \mathbf{y}$*.*

Proof. Up to constants, $-\log p(\mathcal{D} \mid \mathbf{w}) - \log p(\mathbf{w}) = \frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 + \frac{\alpha}{2}\|\mathbf{w}\|^2$; multiply by $2\sigma^2$. The minimum was computed in the exercises of Lecture 3. □

A strong prior belief that the weights are small (large α) means strong regularization. Similarly, the Laplace prior $p(\mathbf{w}) \propto e^{-\|\mathbf{w}\|_1/b}$ leads to ℓ_1 regularization (the lasso), which favors sparse weight vectors.

4.12.3 Bayesian linear regression

The MAP estimate discards the uncertainty that the posterior expresses. In linear regression with a Gaussian prior, the full posterior can be computed with the tools of Section 4.6.

Theorem 4.12.7 (Posterior of Bayesian linear regression)**.** *In the model of Proposition 4.9.19 with the prior* $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1} I)$*, the posterior is* $\mathcal{N}(\mathbf{m}, S)$ *with*

$$S = \left(\alpha I + \sigma^{-2} X^\top X\right)^{-1}, \qquad \mathbf{m} = \sigma^{-2} S X^\top \mathbf{y}.$$

For a new input $\mathbf{x}_*$*, the predictive distribution of* $y_* = \mathbf{w}^\top \mathbf{x}_* + \varepsilon_*$ *is* $\mathcal{N}(\mathbf{m}^\top \mathbf{x}_*,\ \mathbf{x}_*^\top S \mathbf{x}_* + \sigma^2)$*.*

Proof. Up to an additive constant that does not depend on $\mathbf{w}$,

$$\log p(\mathbf{w} \mid \mathcal{D}) = -\frac{1}{2\sigma^2}\|\mathbf{y} - X\mathbf{w}\|^2 - \frac{\alpha}{2}\|\mathbf{w}\|^2 = -\frac{1}{2}\mathbf{w}^\top S^{-1}\mathbf{w} + \sigma^{-2}\mathbf{y}^\top X \mathbf{w} + \text{const}.$$

Completing the square, this is $-\frac{1}{2}(\mathbf{w}-\mathbf{m})^\top S^{-1}(\mathbf{w}-\mathbf{m}) + \text{const}$, since the linear term of the latter is $\mathbf{m}^\top S^{-1}\mathbf{w} = \sigma^{-2}\mathbf{y}^\top X\mathbf{w}$. So the posterior density is proportional to the density of $\mathcal{N}(\mathbf{m}, S)$, hence equal to it. Given the data, $\mathbf{w}^\top \mathbf{x}_*$ is $\mathcal{N}(\mathbf{m}^\top \mathbf{x}_*, \mathbf{x}_*^\top S \mathbf{x}_*)$ by Corollary 4.6.7, and adding the independent noise $\varepsilon_* \sim \mathcal{N}(0, \sigma^2)$ adds σ^2 to the variance. □

The posterior mean $\mathbf{m} = (X^\top X + \alpha\sigma^2 I)^{-1} X^\top \mathbf{y}$ is the ridge regression solution, which is also the MAP estimate (the mode and the mean of a Gaussian coincide). But the posterior also gives error bars: the predictive variance $\mathbf{x}_*^\top S \mathbf{x}_* + \sigma^2$ consists of the uncertainty about the weights, which shrinks as we collect more data, and the noise, which does not.

Example 4.12.8. For the data $(1,2), (2,3), (3,5)$, the model $y = wx + \varepsilon$ without intercept, $\alpha = 1$ and $\sigma^2 = 1$: $S = (1 + 1 + 4 + 9)^{-1} = \frac{1}{15}$ and $m = \frac{1}{15}(2 + 6 + 15) = \frac{23}{15} \approx 1.53$. At $x_* = 4$, the predictive distribution is $\mathcal{N}(6.13,\ \frac{16}{15} + 1) \approx \mathcal{N}(6.13, 2.07)$.

4.12.4 Gaussian processes

Bayesian linear regression puts a prior on the weights of a fixed set of features. A Gaussian process puts a prior directly on the unknown function.

Definition 4.12.9 (Gaussian process)**.** A Gaussian process with mean function m and covariance function (or kernel) k is a random function f such that for all points $\mathbf{x}_1, \dots, \mathbf{x}_n$, the vector $(f(\mathbf{x}_1), \dots, f(\mathbf{x}_n))$ is Gaussian with the mean vector $(m(\mathbf{x}_i))_i$ and the covariance matrix K with entries $K_{ij} = k(\mathbf{x}_i, \mathbf{x}_j)$. We write $f \sim \mathcal{GP}(m, k)$.

For this to make sense, every such matrix K must be symmetric and positive semidefinite. This holds, for example, for $k(\mathbf{x}, \mathbf{x}') = \boldsymbol{\phi}(\mathbf{x})^\top \boldsymbol{\phi}(\mathbf{x}')$ with any feature map $\boldsymbol{\phi}$, since K is then a Gram matrix (compare with the remark on kernels in Lecture 2).

Example 4.12.10.

1. (Linear kernel.) If $f(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$ with $\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \alpha^{-1} I)$, then

$$\operatorname{Cov}(f(\mathbf{x}), f(\mathbf{x}')) = \mathbf{x}^\top \operatorname{Cov}(\mathbf{w}) \mathbf{x}' = \alpha^{-1} \mathbf{x}^\top \mathbf{x}'.$$

   So the prior of Bayesian linear regression is the Gaussian process with the kernel $k(\mathbf{x}, \mathbf{x}') = \alpha^{-1}\mathbf{x}^\top \mathbf{x}'$.

2. (Squared exponential kernel.) $k(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2\ell^2}\right)$ gives smooth random functions, whose values at points closer than the length scale ℓ are strongly correlated. It corresponds to infinitely many features.

3. (Brownian motion.) $k(s,t) = \min(s,t)$ for $s,t \geq 0$ is the covariance function of Brownian motion, a random function that we study in Section 4.15.

Proposition 4.12.11 (Gaussian process regression)**.** *Let $f \sim \mathcal{GP}(0,k)$, and let $y_i = f(\mathbf{x}_i) + \varepsilon_i$ for $i = 1,\dots,N$, with i.i.d. noise $\varepsilon_i \sim \mathcal{N}(0,\sigma^2)$ independent of f. Let K be the kernel matrix of the training inputs, and for a new input $\mathbf{x}_*$, let $\mathbf{k}_* = (k(\mathbf{x}_1,\mathbf{x}_*),\dots,k(\mathbf{x}_N,\mathbf{x}_*))^\top$. Then, given the data,*

$$f(\mathbf{x}_*) \sim \mathcal{N}\big(\mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{y},\ k(\mathbf{x}_*,\mathbf{x}_*) - \mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{k}_*\big).$$

Proof. The vector $(\mathbf{y}, f(\mathbf{x}_*))$ is Gaussian with mean $\mathbf{0}$ and the covariance blocks $\operatorname{Cov}(\mathbf{y}) = K+\sigma^2 I$, $\operatorname{Cov}(\mathbf{y}, f(\mathbf{x}_*)) = \mathbf{k}_*$ and $\operatorname{Var}(f(\mathbf{x}_*)) = k(\mathbf{x}_*,\mathbf{x}_*)$; now apply Theorem 4.6.5. □

The posterior mean $\sum_i \alpha_i k(\mathbf{x}_i,\mathbf{x}_*)$ with $\boldsymbol{\alpha} = (K+\sigma^2 I)^{-1}\mathbf{y}$ is a combination of kernel functions centered at the data points; it is also the prediction of kernel ridge regression. The posterior variance does not depend on $\mathbf{y}$: it is small near the data, and it returns to the prior variance far away from them (Figure 4.6). For the linear kernel, the proposition reproduces the predictive distribution of Theorem 4.12.7.

Example 4.12.12. With the squared exponential kernel ($\ell = 1$), $\sigma^2 = 0.1$, and the data $(0,1)$ and $(1,0)$: $K + \sigma^2 I = \begin{pmatrix} 1.1 & 0.607 \\ 0.607 & 1.1 \end{pmatrix}$, and at $x_* = 0.5$, $\mathbf{k}_* = (0.882, 0.882)^\top$. The posterior mean is 0.517, and the posterior variance is $1 - \mathbf{k}_*^\top (K+\sigma^2 I)^{-1}\mathbf{k}_* \approx 0.087$; a new noisy observation at x_* has the predictive variance $0.087 + 0.1 = 0.187$.




Figure 4.6: Gaussian process regression with the squared exponential kernel ($\ell = 0.7$, $\sigma^2 = 0.01$) for six data points (red): the posterior mean (blue) and a band of ± 2 posterior standard deviations. The uncertainty is small near the data and grows back to the prior standard deviation 1 between and beyond them.

The hyperparameters, such as ℓ and σ^2, are usually chosen by maximizing the marginal likelihood $\log p(\mathbf{y}) = -\frac{1}{2}\mathbf{y}^\top (K+\sigma^2 I)^{-1}\mathbf{y} - \frac{1}{2}\log\det(K+\sigma^2 I) - \frac{N}{2}\log(2\pi)$, since $\mathbf{y} \sim \mathcal{N}(\mathbf{0}, K+\sigma^2 I)$; its gradient can be computed with the matrix calculus of Lecture 3. The cost of solving with the $N \times N$ matrix $K+\sigma^2 I$ grows like N^3, so exact Gaussian processes are limited to moderate data sets. Remarkably, one can show that neural networks with random weights become Gaussian processes in the limit of infinitely wide layers.

Remark 4.12.13 (Beyond conjugate models). For most models, such as neural networks, the posterior has no closed form, and the evidence $p(\mathcal{D})$ is an intractable integral. Approximate methods include Markov chain Monte Carlo sampling (Section 4.8) and variational inference, which approximates the posterior by a simpler distribution by minimizing a Kullback–Leibler divergence (Section 4.13).

4.13 Information Theory

Information theory measures uncertainty and the difference between distributions. Its quantities appear in machine learning as loss functions (cross-entropy), as regularizers (the KL term of variational autoencoders), and in the analysis of learning algorithms.

Definition 4.13.1 (Entropy, cross-entropy and KL divergence)**.** Let $p = (p_1, \dots, p_K)$ and $q = (q_1, \dots, q_K)$ be probability distributions on $\{1, \dots, K\}$. The entropy of p, the cross-entropy of q relative to p, and the Kullback–Leibler (KL) divergence of q from p are

$$H(p) = -\sum_k p_k \log p_k, \qquad H(p, q) = -\sum_k p_k \log q_k, \qquad \mathrm{KL}(p \,\|\, q) = \sum_k p_k \log \frac{p_k}{q_k},$$

with the conventions $0 \log 0 = 0$ and $\mathrm{KL}(p \,\|\, q) = \infty$ if $q_k = 0 < p_k$ for some k.

The number $-\log p_k$ measures the surprise of seeing the outcome k: rare outcomes are surprising. The entropy is the expected surprise. With $\log_2$ instead of log, entropy is measured in bits, and it is the minimal average number of bits per symbol needed to encode outcomes of p (Shannon's source coding theorem). A fair coin has entropy 1 bit, a coin with $P(\text{heads}) = 0.9$ only 0.47 bits, and a deterministic outcome 0 bits. Clearly $H(p) \geq 0$, and

$$H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q) :$$

the cross-entropy is the average number of bits if we encode outcomes of p with a code designed for q, and the KL divergence is the number of extra bits we pay for using the wrong distribution.

Theorem 4.13.2 (Gibbs' inequality)**.** $\mathrm{KL}(p \,\|\, q) \geq 0$*, with equality if and only if* $p = q$*.*

Proof. We may assume $q_k > 0$ whenever $p_k > 0$. Let T be the set of k with $p_k > 0$, and let R be the random variable that takes the value $\frac{q_k}{p_k}$ with probability p_k, for $k \in T$. By Jensen's inequality for the concave function log (Theorem 4.3.7),

$$-\mathrm{KL}(p \,\|\, q) = \sum_{k \in T} p_k \log \frac{q_k}{p_k} = \mathbb{E}[\log R] \leq \log \mathbb{E}[R] = \log \sum_{k \in T} q_k \leq \log 1 = 0.$$

If equality holds, then $\sum_{k \in T} q_k = 1$, and equality holds in Jensen's inequality, which for the strictly concave log happens only if R is constant; so $q_k = c\, p_k$ for $k \in T$ and $q_k = 0$ otherwise, and $c = 1$ since both sum to 1. □

Two consequences: the cross-entropy $H(p, q)$ is minimal for $q = p$, and for the uniform distribution u on K outcomes, $0 \leq \mathrm{KL}(p \,\|\, u) = \log K - H(p)$, so $H(p) \leq \log K$: the uniform distribution has maximal entropy (compare with the exercise on maximum entropy in Lecture 3). The KL divergence is not a distance in the sense of Lecture 2, since it is not symmetric. For $p = (\frac{1}{2}, \frac{1}{2})$ and $q = (0.9, 0.1)$, $\mathrm{KL}(p \,\|\, q) \approx 0.511$ but $\mathrm{KL}(q \,\|\, p) \approx 0.368$.

Theorem 4.13.3 (Maximum likelihood minimizes the KL divergence)**.** *Let* $x_1, \dots, x_N \in \{1, \dots, K\}$ *be data with the empirical distribution* $\hat{p}$*, i.e.,* $\hat{p}_k$ *is the fraction of data points equal to* k*, and let* q_θ *be a model distribution. Then*

$$\frac{1}{N} \sum_{i=1}^{N} \log q_\theta(x_i) = -H(\hat{p}, q_\theta) = -H(\hat{p}) - \mathrm{KL}(\hat{p} \,\|\, q_\theta).$$

So maximizing the likelihood is the same as minimizing the cross-entropy $H(\hat{p}, q_\theta)$*, and the same as minimizing* $\mathrm{KL}(\hat{p} \,\|\, q_\theta)$*.*

Proof. Grouping the data points by their values, $\frac{1}{N} \sum_i \log q_\theta(x_i) = \sum_k \hat{p}_k \log q_\theta(k) = -H(\hat{p}, q_\theta)$. The term $H(\hat{p})$ does not depend on θ. □

This is why the loss of classifiers is called the cross-entropy loss: for each training example, it is the cross-entropy between the one-hot distribution of the true label and the predicted distribution, and training fits the model distribution to the data in the sense of the KL divergence.

Continuous distributions. For densities, the sums become integrals: the differential entropy is $h(p) = -\int p(x)\log p(x)\,dx$, and $\mathrm{KL}(p\,\|\,q) = \int p(x)\log\frac{p(x)}{q(x)}\,dx \geq 0$. The differential entropy of $\mathcal{N}(\mu,\sigma^2)$ is $\frac{1}{2}\log(2\pi e\sigma^2)$, and among all densities with variance σ^2, the Gaussian has the largest entropy.

Example 4.13.4 (Maximum entropy)**.** Among the distributions on $\mathbb{R}$ with the variance 1, the uniform distribution on $[-\sqrt{3},\sqrt{3}]$ has the differential entropy $\log(2\sqrt{3}) \approx 1.24$, the Laplace distribution with the density $\frac{1}{\sqrt{2}}e^{-\sqrt{2}|x|}$ has $1+\frac{1}{2}\log 2 \approx 1.35$, and $\mathcal{N}(0,1)$ has the largest value, $\frac{1}{2}\log(2\pi e) \approx 1.42$ nats. In fact, for every density p with mean 0 and variance 1, $\mathrm{KL}(p\,\|\,\mathcal{N}(0,1)) = -h(p) + \mathbb{E}_p[\frac{1}{2}\log(2\pi) + \frac{X^2}{2}] = \frac{1}{2}\log(2\pi e) - h(p)$, which is nonnegative by Gibbs' inequality. This proves the maximum entropy property of the Gaussian, which is one justification for Gaussian noise models: they assume nothing beyond the variance.

Proposition 4.13.5 (KL divergence between Gaussians)**.** *For $p=\mathcal{N}(\mu_1,\sigma_1^2)$ and $q=\mathcal{N}(\mu_2,\sigma_2^2)$,*

$$\mathrm{KL}(p\,\|\,q) = \log\frac{\sigma_2}{\sigma_1} + \frac{\sigma_1^2+(\mu_1-\mu_2)^2}{2\sigma_2^2} - \frac{1}{2}.$$

Proof. $\log\frac{p(x)}{q(x)} = \log\frac{\sigma_2}{\sigma_1} - \frac{(x-\mu_1)^2}{2\sigma_1^2} + \frac{(x-\mu_2)^2}{2\sigma_2^2}$. Take the expectation for $X\sim p$, using $\mathbb{E}[(X-\mu_1)^2]=\sigma_1^2$ and $\mathbb{E}[(X-\mu_2)^2] = \sigma_1^2+(\mu_1-\mu_2)^2$. □

In a variational autoencoder, each data point is encoded as a Gaussian $\mathcal{N}(\mu,\sigma^2)$ (in each latent dimension), and the loss contains the term $\mathrm{KL}\left(\mathcal{N}(\mu,\sigma^2)\,\|\,\mathcal{N}(0,1)\right) = \frac{1}{2}(\mu^2+\sigma^2-1-\log\sigma^2)$, which keeps the codes close to a standard Gaussian. For example, for $\mu=1$ and $\sigma=2$, it is $\frac{1}{2}(4-\log 4) \approx 1.31$.

Mutual information. The mutual information of two random variables is the KL divergence between their joint distribution and the product of their marginals,

$$I(X;Y) = \mathrm{KL}\left(p(x,y)\,\|\,p_X(x)p_Y(y)\right) = \sum_{x,y} p(x,y)\log\frac{p(x,y)}{p_X(x)p_Y(y)}.$$

By Gibbs' inequality, $I(X;Y)\geq 0$, with equality if and only if X and Y are independent. Unlike the correlation, it detects every kind of dependence. For Example 4.4.3, $I(X;Y)\approx 0.086$ nats.

4.14 Latent Variable Models and the EM Algorithm

Many data sets are explained by quantities that we do not observe: the cluster to which a customer belongs, the topic of a document, or the object in an image. A latent variable model describes the joint distribution $p(\mathbf{x}, z \mid \theta)$ of an observation $\mathbf{x}$ and a hidden variable z. Only $\mathbf{x}$ is observed, so the likelihood is the marginal

$$p(\mathbf{x}\mid\theta) = \sum_z p(\mathbf{x},z\mid\theta) \qquad \text{or} \qquad \int p(\mathbf{x},z\mid\theta)\,dz,$$

and the logarithm of a sum is hard to maximize directly. This section derives a general method for this problem from the KL divergence of Section 4.13.

4.14.1 Gaussian mixture models

Definition 4.14.1 (Gaussian mixture model)**.** A Gaussian mixture model with K components draws a component $z \in \{1, \dots, K\}$ with probabilities $\pi_1, \dots, \pi_K$, and then $\mathbf{x} \sim \mathcal{N}(\boldsymbol{\mu}_z, \Sigma_z)$. So the density of $\mathbf{x}$ is

$$p(\mathbf{x} \mid \theta) = \sum_{k=1}^{K} \pi_k \, \varphi(\mathbf{x}; \boldsymbol{\mu}_k, \Sigma_k),$$

where $\varphi(\cdot\,; \boldsymbol{\mu}, \Sigma)$ is the density of $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$, and θ consists of the weights π_k, the means $\boldsymbol{\mu}_k$ and the covariance matrices Σ_k.

Given an observation, Bayes' theorem gives the posterior probabilities of the components,

$$r_k(\mathbf{x}) = P(z = k \mid \mathbf{x}, \theta) = \frac{\pi_k \, \varphi(\mathbf{x}; \boldsymbol{\mu}_k, \Sigma_k)}{\sum_j \pi_j \, \varphi(\mathbf{x}; \boldsymbol{\mu}_j, \Sigma_j)},$$

called the responsibilities: a soft assignment of $\mathbf{x}$ to the clusters. If we knew the components z_i of the data points, maximum likelihood would be easy: $\hat{\pi}_k$ would be the fraction of points in cluster k, and $\hat{\boldsymbol{\mu}}_k$ and $\hat{\Sigma}_k$ the mean and covariance matrix of these points (Theorem 4.9.10). The EM algorithm alternates between guessing the components with the current parameters and fitting the parameters to the guess.

4.14.2 The evidence lower bound

Theorem 4.14.2 (Evidence lower bound)**.** *For every distribution q on the values of z (with $q(z) > 0$ only where $p(\mathbf{x}, z \mid \theta) > 0$),*

$$\log p(\mathbf{x} \mid \theta) = \underbrace{\sum_z q(z) \log \frac{p(\mathbf{x}, z \mid \theta)}{q(z)}}_{\mathrm{ELBO}(q,\theta)} + \mathrm{KL}\left(q \,\|\, p(\cdot \mid \mathbf{x}, \theta)\right).$$

Hence $\log p(\mathbf{x} \mid \theta) \geq \mathrm{ELBO}(q, \theta)$, with equality if and only if q is the posterior $p(z \mid \mathbf{x}, \theta)$.

Proof. Since $p(\mathbf{x}, z \mid \theta) = p(z \mid \mathbf{x}, \theta)\, p(\mathbf{x} \mid \theta)$,

$$\begin{aligned} \sum_z q(z) \log \frac{p(\mathbf{x}, z \mid \theta)}{q(z)} &= \sum_z q(z) \log p(\mathbf{x} \mid \theta) + \sum_z q(z) \log \frac{p(z \mid \mathbf{x}, \theta)}{q(z)} \\ &= \log p(\mathbf{x} \mid \theta) - \mathrm{KL}\left(q \,\|\, p(\cdot \mid \mathbf{x}, \theta)\right). \end{aligned}$$

The rest follows from Gibbs' inequality (Theorem 4.13.2). □

The ELBO is the evidence lower bound; the marginal likelihood $p(\mathbf{x} \mid \theta)$ is also called the evidence. For data $\mathbf{x}_1, \dots, \mathbf{x}_N$ with independent latent variables, we choose one distribution q_i for each data point, and then $\ell(\theta) = \sum_i \log p(\mathbf{x}_i \mid \theta) \geq \sum_i \mathrm{ELBO}(q_i, \theta)$.

Definition 4.14.3 (EM algorithm)**.** Starting from parameters θ_0, the expectation-maximization (EM) algorithm repeats two steps:

1. (E-step) $q_i = p(\cdot \mid \mathbf{x}_i, \theta_t)$, the posterior of the latent variable of each data point;
2. (M-step) θ_{t+1} maximizes $\sum_i \mathrm{ELBO}(q_i, \theta)$. Since the terms $-q_i(z) \log q_i(z)$ do not depend on θ, this is the same as maximizing the expected complete-data log-likelihood

$$\sum_{i=1}^{N} \sum_z q_i(z) \log p(\mathbf{x}_i, z \mid \theta).$$

Theorem 4.14.4. *The EM algorithm never decreases the log-likelihood: $\ell(\theta_{t+1}) \geq \ell(\theta_t)$.*

Proof. By Theorem 4.14.2, the choice of q_i in the E-step, and the M-step,

$$\ell(\theta_{t+1}) \geq \sum_i \mathrm{ELBO}(q_i, \theta_{t+1}) \geq \sum_i \mathrm{ELBO}(q_i, \theta_t) = \ell(\theta_t). \qquad \square$$

So EM is coordinate ascent on the ELBO: the E-step maximizes it over the q_i, which makes the bound tight, and the M-step maximizes it over θ. Like gradient ascent, it finds a local maximum (or a saddle point) of the likelihood, not necessarily the global one, so it is run from several starting points.

Theorem 4.14.5 (EM for Gaussian mixtures)**.** *For a Gaussian mixture model, let $r_{ik} = r_k(\mathbf{x}_i)$ be the responsibilities computed with θ_t in the E-step, and $N_k = \sum_i r_{ik}$. The M-step gives*

$$\pi_k = \frac{N_k}{N}, \qquad \boldsymbol{\mu}_k = \frac{1}{N_k} \sum_{i=1}^{N} r_{ik} \mathbf{x}_i, \qquad \Sigma_k = \frac{1}{N_k} \sum_{i=1}^{N} r_{ik} (\mathbf{x}_i - \boldsymbol{\mu}_k)(\mathbf{x}_i - \boldsymbol{\mu}_k)^\top.$$

Proof. The M-step maximizes

$$\sum_{i=1}^{N} \sum_{k=1}^{K} r_{ik} \big(\log \pi_k + \log \varphi(\mathbf{x}_i; \boldsymbol{\mu}_k, \Sigma_k) \big) = \sum_k N_k \log \pi_k + \sum_k \sum_i r_{ik} \log \varphi(\mathbf{x}_i; \boldsymbol{\mu}_k, \Sigma_k).$$

The weights only appear in the first sum, which is $-N\, H(\hat{\mathbf{p}}, \boldsymbol{\pi})$ for the distribution $\hat{p}_k = \frac{N_k}{N}$ (note $\sum_k N_k = N$); by Gibbs' inequality, it is largest for $\boldsymbol{\pi} = \hat{\mathbf{p}}$. The parameters of component k only appear in $\sum_i r_{ik} \log \varphi(\mathbf{x}_i; \boldsymbol{\mu}_k, \Sigma_k)$, the log-likelihood of a Gaussian in which the data point $\mathbf{x}_i$ has the weight r_{ik}. The computation of Theorem 4.9.10, with these weights, gives the weighted mean and covariance matrix. $\square$

So EM for Gaussian mixtures alternates between computing the responsibilities and refitting each Gaussian to all data points, weighted by their responsibilities. Each step costs $O(NK)$ density evaluations.

Example 4.14.6. Fit a mixture of two Gaussians with variance 1 and equal weights to the four numbers $1, 2, 4, 5$, starting with the means $\mu_1 = 0$ and $\mu_2 = 3$; only the means are fitted. The responsibilities of the first component are $r_{i1} = \frac{e^{-(x_i-\mu_1)^2/2}}{e^{-(x_i-\mu_1)^2/2} + e^{-(x_i-\mu_2)^2/2}}$: for $x = 1$, $\frac{1}{1+e^{-1.5}} = 0.818$; for $x = 2$, $\frac{1}{1+e^{1.5}} = 0.182$; for $x = 4$ and $x = 5$, 0.0006 and 0.00003. The M-step gives the new means $\mu_1 = \frac{0.818 \cdot 1 + 0.182 \cdot 2 + \cdots}{0.818 + 0.182 + \cdots} \approx 1.18$ and $\mu_2 \approx 3.61$, and the log-likelihood increases from -9.54 to -7.47. Further steps converge to $\mu_1 \approx 1.56$ and $\mu_2 \approx 4.44$ (not exactly 1.5 and 4.5, since the two Gaussians overlap), with the log-likelihood -6.84.

4.14.3 k-means clustering

The k-means algorithm is the "hard" version of EM. It looks for K centers $\boldsymbol{\mu}_1, \ldots, \boldsymbol{\mu}_K$ and an assignment $c(i) \in \{1, \ldots, K\}$ of the data points to the centers that minimize

$$J = \sum_{i=1}^{N} \|\mathbf{x}_i - \boldsymbol{\mu}_{c(i)}\|^2.$$

Lloyd's algorithm alternates two steps: assign each point to its nearest center, and move each center to the mean of its points.

Proposition 4.14.7. *Neither step of Lloyd's algorithm increases J. Hence the values of J after the update steps never increase, and they are eventually constant.*

Proof. For fixed centers, the assignment step minimizes each term of J separately. For a fixed assignment, the terms of cluster k are $\sum_{i:c(i)=k}\|\mathbf{x}_i-\boldsymbol{\mu}_k\|^2=\sum_{i:c(i)=k}\|\mathbf{x}_i-\bar{\mathbf{x}}_k\|^2+n_k\|\bar{\mathbf{x}}_k-\boldsymbol{\mu}_k\|^2$, where $\bar{\mathbf{x}}_k$ is the mean of the n_k points of the cluster (expand the squares; the mixed terms add up to 0). This is smallest for $\boldsymbol{\mu}_k=\bar{\mathbf{x}}_k$. After an update step, J only depends on the assignment, which can take finitely many values, so a nonincreasing sequence of these values is eventually constant. □

k-means is EM for a Gaussian mixture with equal weights and covariance matrices $\sigma^2 I$, in the limit $\sigma\to 0$: the responsibilities are proportional to $e^{-\|\mathbf{x}_i-\boldsymbol{\mu}_k\|^2/(2\sigma^2)}$, and as $\sigma\to 0$, the ratio of the responsibilities of two components k and j is $e^{(\|\mathbf{x}_i-\boldsymbol{\mu}_j\|^2-\|\mathbf{x}_i-\boldsymbol{\mu}_k\|^2)/(2\sigma^2)}$, which tends to ∞ if $\boldsymbol{\mu}_k$ is closer to $\mathbf{x}_i$. So the responsibility of the nearest center tends to 1, and the M-step becomes the update of the centers. Like EM, k-means only finds a local minimum and depends on the initial centers; the method k-means++ chooses them at random, far apart from each other. Since it measures distances with the Euclidean norm, k-means finds round clusters of similar size; for other shapes, Gaussian mixtures with general covariance matrices, or spectral clustering (Lecture 2), work better.

4.14.4 Probabilistic PCA

Principal component analysis (Lecture 2) can also be derived from a latent variable model, with a continuous latent variable: each data point is a linear function of k hidden factors plus noise.

Definition 4.14.8 (Probabilistic PCA)**.** Let $k<d$. In probabilistic PCA, the latent vector is $\mathbf{z}\sim\mathcal{N}(\mathbf{0},I_k)$, and the observation is $\mathbf{x}=W\mathbf{z}+\boldsymbol{\mu}+\boldsymbol{\varepsilon}$ with $\boldsymbol{\varepsilon}\sim\mathcal{N}(\mathbf{0},\sigma^2 I_d)$ independent of $\mathbf{z}$. The parameters are $W\in\mathbb{R}^{d\times k}$, $\boldsymbol{\mu}\in\mathbb{R}^d$ and $\sigma^2>0$.

The vector $(\mathbf{z},\boldsymbol{\varepsilon})$ is $\mathcal{N}(\mathbf{0},\operatorname{diag}(I_k,\sigma^2 I_d))$, and $(\mathbf{z},\mathbf{x})$ is an affine function of it, so it is Gaussian (Theorem 4.6.2), with

$$\mathbf{x}\sim\mathcal{N}(\boldsymbol{\mu},C),\qquad C=WW^\top+\sigma^2 I_d,\qquad \operatorname{Cov}(\mathbf{z},\mathbf{x})=\mathbb{E}[\mathbf{z}(W\mathbf{z}+\boldsymbol{\varepsilon})^\top]=W^\top.$$

So the model describes data whose covariance matrix is a rank-k matrix plus isotropic noise. By Theorem 4.6.5, the posterior of $\mathbf{z}$ is Gaussian with the mean $W^\top C^{-1}(\mathbf{x}-\boldsymbol{\mu})$ and the covariance $I_k-W^\top C^{-1}W$. With $M=W^\top W+\sigma^2 I_k$, the identity $W^\top(WW^\top+\sigma^2 I_d)=MW^\top$ gives $W^\top C^{-1}=M^{-1}W^\top$, so

$$\mathbb{E}[\mathbf{z}\mid\mathbf{x}]=M^{-1}W^\top(\mathbf{x}-\boldsymbol{\mu}),\qquad \operatorname{Cov}(\mathbf{z}\mid\mathbf{x})=I_k-M^{-1}W^\top W=\sigma^2 M^{-1},$$

which only needs the inverse of a $k\times k$ matrix. As $\sigma^2\to 0$, the posterior mean tends to $(W^\top W)^{-1}W^\top(\mathbf{x}-\boldsymbol{\mu})$, the coordinates of the orthogonal projection of $\mathbf{x}-\boldsymbol{\mu}$ onto the column space of W (Lecture 2).

Theorem 4.14.9 (Maximum likelihood for probabilistic PCA)**.** *Let $\mathbf{x}_1,\dots,\mathbf{x}_N\in\mathbb{R}^d$ have the mean $\bar{\mathbf{x}}$ and the sample covariance matrix $S=\frac{1}{N}\sum_i(\mathbf{x}_i-\bar{\mathbf{x}})(\mathbf{x}_i-\bar{\mathbf{x}})^\top$, with eigenvalues $\lambda_1\ge\cdots\ge\lambda_d$ and orthonormal eigenvectors $\mathbf{u}_1,\dots,\mathbf{u}_d$, and assume $\lambda_{k+1}>0$. Then the likelihood is maximized by*

$$\hat{\boldsymbol{\mu}}=\bar{\mathbf{x}},\qquad \hat{\sigma}^2=\frac{1}{d-k}\sum_{j=k+1}^{d}\lambda_j,\qquad \hat{W}=U_k(\Lambda_k-\hat{\sigma}^2 I_k)^{1/2},$$

where U_k has the columns $\mathbf{u}_1,\dots,\mathbf{u}_k$ and $\Lambda_k=\operatorname{diag}(\lambda_1,\dots,\lambda_k)$, and also by $\hat{W}R$ for every orthogonal $k\times k$ matrix R.

Proof. As in the proof of Theorem 4.9.10, $\boldsymbol{\mu}=\bar{\mathbf{x}}$ is optimal for every C, and then $\ell=\frac{N}{2}(-\log\det C-\operatorname{tr}(C^{-1}S))$ plus a constant. By the spectral theorem, $WW^\top=\sum_{j=1}^{k}s_j\mathbf{v}_j\mathbf{v}_j^\top$ with

$s_1 \geq \cdots \geq s_k \geq 0$ and orthonormal vectors $\mathbf{v}_1, \ldots, \mathbf{v}_d$ (its rank is at most k). So C has the eigenvalues $c_j = \sigma^2 + s_j$ for $j \leq k$ and σ^2 for $j > k$, $\log\det C = \sum_{j\leq k} \log c_j + (d-k)\log\sigma^2$, and

$$C^{-1} = \frac{1}{\sigma^2} I_d - \sum_{j=1}^{k} a_j \mathbf{v}_j \mathbf{v}_j^\top, \qquad a_j = \frac{1}{\sigma^2} - \frac{1}{c_j}, \qquad a_1 \geq \cdots \geq a_k \geq 0.$$

With $q_j = \mathbf{v}_j^\top S \mathbf{v}_j$ and $a_{k+1} = 0$, we have $\sum_{j\leq k} a_j q_j = \sum_{j\leq k}(a_j - a_{j+1})(q_1 + \cdots + q_j) \leq \sum_{j\leq k}(a_j - a_{j+1})(\lambda_1 + \cdots + \lambda_j) = \sum_{j\leq k} a_j \lambda_j$, by Ky Fan's inequality (Lecture 2) and since $a_j - a_{j+1} \geq 0$. Hence

$$\operatorname{tr}(C^{-1}S) = \frac{\operatorname{tr} S}{\sigma^2} - \sum_{j\leq k} a_j q_j \geq \sum_{j\leq k} \frac{\lambda_j}{c_j} + \sum_{j>k} \frac{\lambda_j}{\sigma^2},$$

and with $\bar\sigma^2 = \frac{1}{d-k}\sum_{j>k}\lambda_j$,

$$-\log\det C - \operatorname{tr}(C^{-1}S) \leq -\sum_{j\leq k}\Big(\log c_j + \frac{\lambda_j}{c_j}\Big) - (d-k)\Big(\log\sigma^2 + \frac{\bar\sigma^2}{\sigma^2}\Big).$$

For $\lambda > 0$, the function $c \mapsto \log c + \frac{\lambda}{c}$ on $(0,\infty)$ has the derivative $\frac{c-\lambda}{c^2}$, so its minimum is $\log\lambda + 1$, at $c = \lambda$. So the right side is at most $-\sum_{j\leq k}(\log\lambda_j + 1) - (d-k)(\log\bar\sigma^2 + 1)$, for all parameters. For $\sigma^2 = \bar\sigma^2$ and $W = \hat W$ (note $\lambda_j \geq \lambda_k \geq \bar\sigma^2$ for $j \leq k$), we get $WW^\top = \sum_{j\leq k}(\lambda_j - \bar\sigma^2)\mathbf{u}_j\mathbf{u}_j^\top$, so $\mathbf{v}_j = \mathbf{u}_j$ and $c_j = \lambda_j$, and all inequalities are equalities. Finally, $(\hat W R)(\hat W R)^\top = \hat W \hat W^\top$. □

So the maximum likelihood estimate spans the principal subspace $\operatorname{span}\{\mathbf{u}_1, \ldots, \mathbf{u}_k\}$ of PCA, reproduces the variances $\lambda_1, \ldots, \lambda_k$ in these directions exactly, and replaces the variances in the remaining $d-k$ directions by their average $\hat\sigma^2$, the noise level. For example, for $S = \operatorname{diag}(4,2,1)$ and $k = 1$, $\hat\sigma^2 = 1.5$ and $\hat W = (\sqrt{2.5}, 0, 0)^\top$, so the fitted covariance matrix is $\hat C = \operatorname{diag}(4, 1.5, 1.5)$. Unlike PCA, probabilistic PCA has a likelihood, which can be compared across different k or used in mixtures of such models, and it can be fitted with EM when some entries of the data are missing. Its decoder $p(\mathbf{x} \mid \mathbf{z}) = \mathcal{N}(W\mathbf{z} + \boldsymbol{\mu}, \sigma^2 I)$ is linear; a variational autoencoder replaces it by a neural network.

4.14.5 Variational inference and variational autoencoders

In more complex models, such as deep generative models, the posterior $p(z \mid \mathbf{x}, \theta)$ cannot be computed, and the E-step is impossible. Variational inference maximizes the ELBO over a family of tractable distributions q instead, for example Gaussians $q(z) = \mathcal{N}(\boldsymbol{\mu}, \operatorname{diag}(\boldsymbol{\sigma}^2))$; by Theorem 4.14.2, this finds the member of the family that is closest to the posterior in the sense of $\mathrm{KL}(q \,\|\, p(\cdot \mid \mathbf{x}, \theta))$. Writing $p(\mathbf{x}, z \mid \theta) = p(\mathbf{x} \mid z, \theta)p(z)$ gives the form

$$\mathrm{ELBO}(q, \theta) = \mathbb{E}_{z\sim q}\big[\log p(\mathbf{x} \mid z, \theta)\big] - \mathrm{KL}\left(q \,\|\, p(z)\right) :$$

a reconstruction term, which rewards latent codes that explain $\mathbf{x}$, and a KL term, which keeps q close to the prior $p(z)$. A variational autoencoder trains an encoder network that outputs the parameters of $q(z \mid \mathbf{x})$ for each data point and a decoder network that defines $p(\mathbf{x} \mid z, \theta)$, both by stochastic gradient ascent on the ELBO. For a Gaussian q and the prior $\mathcal{N}(\mathbf{0}, I)$, the KL term has the closed form of Proposition 4.13.5, and the gradient of the reconstruction term is estimated with the reparameterization trick, $z = \boldsymbol{\mu} + \boldsymbol{\sigma} \odot \boldsymbol{\varepsilon}$ with $\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, I)$ (Section 4.6).

4.15 Stochastic Differential Equations

Markov chains (Section 4.8) move in discrete steps. Many random processes change continuously in time: the position of a particle in a fluid, the price of a stock, the parameters of a network during

training with stochastic gradients, or the gradually noised images of a diffusion model. Stochastic differential equations (SDEs) describe such processes by adding random noise to ordinary differential equations (Section 3.15). The source of the noise is Brownian motion.

Definition 4.15.1 (Brownian motion)**.** A Brownian motion (or Wiener process) is a family of random variables $B(t)$, $t \geq 0$, with $B(0) = 0$, such that

1. for $0 \leq t_0 < t_1 < \cdots < t_k$, the increments $B(t_1) - B(t_0), \ldots, B(t_k) - B(t_{k-1})$ are independent;
2. $B(t) - B(s) \sim \mathcal{N}(0, t-s)$ for $s < t$;
3. the paths $t \mapsto B(t)$ are continuous.

That such a process exists is a theorem of Norbert Wiener, which we do not prove. On the grid $t_k = kh$, Brownian motion is simulated exactly by the random walk $B(t_{k+1}) = B(t_k) + \sqrt{h}\, Z_k$ with i.i.d. $Z_k \sim \mathcal{N}(0,1)$ (Figure 4.7). Conversely, by the central limit theorem, every random walk with many small independent steps looks like a Brownian motion from far away.

Proposition 4.15.2. $\mathbb{E}[B(t)] = 0$, $\mathrm{Var}(B(t)) = t$, *and* $\mathrm{Cov}(B(s), B(t)) = \min(s,t)$.

Proof. The first two statements are property (2) with $s = 0$. For $s \leq t$, $\mathrm{Cov}(B(s), B(t)) = \mathrm{Cov}\big(B(s), B(s) + (B(t) - B(s))\big) = \mathrm{Var}(B(s)) + 0 = s$, since the increment $B(t) - B(s)$ is independent of $B(s) = B(s) - B(0)$. □

So Brownian motion is a Gaussian process with mean 0 and covariance function $\min(s,t)$ (Section 4.12.4). Its typical displacement in a time h is $\sqrt{h}$, which is much larger than h for small h: the difference quotient $\frac{B(t+h)-B(t)}{h}$ has the variance $\frac{1}{h} \to \infty$. In fact, the paths of Brownian motion are continuous but nowhere differentiable, and ordinary calculus does not apply to them.

Definition 4.15.3 (Stochastic differential equation)**.** An SDE $dX = \mu(X)\, dt + \sigma(X)\, dB$, with drift μ and diffusion coefficient σ, describes a process with $X(t) = X(0) + \int_0^t \mu(X(s))\, ds + \int_0^t \sigma(X(s))\, dB(s)$, where the last integral is a stochastic (Itô) integral. Its Euler–Maruyama approximation with step h is

$$X_{k+1} = X_k + \mu(X_k)\, h + \sigma(X_k)\sqrt{h}\, Z_k, \qquad Z_k \sim \mathcal{N}(0,1) \text{ i.i.d.}$$

We do not define the stochastic integral; for our purposes, the Euler–Maruyama scheme, which converges to the solution as $h \to 0$, can serve as the definition. It is Euler's method for the ODE $x' = \mu(x)$ (Section 3.15.2), plus Gaussian noise whose standard deviation is proportional to $\sqrt{h}$, and it is a Markov chain with continuous states. For $\sigma = 0$, the SDE is an ODE.

Example 4.15.4 (The Ornstein–Uhlenbeck process)**.** The SDE $dX = -\theta X\, dt + \sigma\, dB$ with $\theta > 0$ pulls X towards 0 while the noise pushes it around. Its Euler–Maruyama approximation is $X_{k+1} = (1 - \theta h)X_k + \sigma\sqrt{h}\, Z_k$. Its mean satisfies $\mathbb{E}[X_{k+1}] = (1 - \theta h)\mathbb{E}[X_k]$, so it follows Euler's method for the ODE $x' = -\theta x$, with the solution $e^{-\theta t}X(0)$. If $X_k \sim \mathcal{N}(0, v)$ is independent of Z_k, then $X_{k+1} \sim \mathcal{N}\big(0, (1-\theta h)^2 v + \sigma^2 h\big)$ by Corollary 4.6.7. So for $0 < \theta h < 2$, the chain has the stationary distribution $\mathcal{N}(0, v_h)$ with

$$v_h = \frac{\sigma^2 h}{1 - (1 - \theta h)^2} = \frac{\sigma^2}{\theta(2 - \theta h)} \to \frac{\sigma^2}{2\theta} \qquad \text{as } h \to 0.$$

For $\theta = 1$, $\sigma = \sqrt{2}$, the stationary distribution of the SDE is $\mathcal{N}(0,1)$ (Figure 4.7); for $\theta = \sigma = 1$ and $h = 0.01$, $v_h \approx 0.5025$ instead of 0.5.




Figure 4.7: Left: three paths of Brownian motion, simulated with $h = 0.02$. Right: three paths of the Ornstein–Uhlenbeck process $dX = -X\,dt + \sqrt{2}\,dB$, started at 2.5, -2 and 0, together with the solutions $X(0)e^{-t}$ of the ODE without noise (dashed). The process forgets its starting point and fluctuates around 0 with the stationary distribution $\mathcal{N}(0,1)$; about 95% of the time, it stays between -2 and 2.

4.15.1 Langevin dynamics and diffusion models

The Ornstein–Uhlenbeck process is an example of a general principle: a gradient flow with the right amount of noise samples from a probability distribution.

Theorem 4.15.5 (Langevin dynamics)**.** *Let $U : \mathbb{R}^d \to \mathbb{R}$ be smooth with $\int e^{-U(\mathbf{x})}\,d\mathbf{x} < \infty$. Then the SDE $d\mathbf{X} = -\nabla U(\mathbf{X})\,dt + \sqrt{2}\,d\mathbf{B}$, where $\mathbf{B}$ is a vector of d independent Brownian motions, has the stationary density $p(\mathbf{x}) \propto e^{-U(\mathbf{x})}$, and under mild conditions, the distribution of $\mathbf{X}(t)$ converges to it.*

We do not prove the convergence, but we can see why e^{-U} is stationary. Without noise, the density p_t of $\mathbf{X}(t)$ under the flow of the vector field $-\nabla U$ satisfies the continuity equation $\frac{\partial p_t}{\partial t} = -\operatorname{div}(-p_t \nabla U)$ (Proposition 3.16.3). The noise adds a diffusion term, and one can show that p_t satisfies the Fokker–Planck equation

$$\frac{\partial p_t}{\partial t} = \operatorname{div}(p_t \nabla U) + \Delta p_t = \operatorname{div}(p_t \nabla U + \nabla p_t),$$

with the Laplacian $\Delta p_t = \operatorname{div} \nabla p_t$ (Lecture 3). For $p \propto e^{-U}$, we have $\nabla p = -p \nabla U$, so the right side vanishes, and p does not change in time. For $U = 0$, the Fokker–Planck equation is the heat equation $\frac{\partial p_t}{\partial t} = \Delta p_t$, which describes how the density of $\sqrt{2}\,\mathbf{B}(t)$ spreads out. For $U(x) = \frac{x^2}{2s^2}$, the SDE is the Ornstein–Uhlenbeck process with $\theta = \frac{1}{s^2}$ and $\sigma = \sqrt{2}$, whose stationary variance $\frac{\sigma^2}{2\theta} = s^2$ agrees with the density $e^{-U} \propto e^{-x^2/(2s^2)}$ of $\mathcal{N}(0, s^2)$. The Euler–Maruyama scheme gives the unadjusted Langevin algorithm

$$\mathbf{x}_{k+1} = \mathbf{x}_k - h \nabla U(\mathbf{x}_k) + \sqrt{2h}\,\mathbf{z}_k, \qquad \mathbf{z}_k \sim \mathcal{N}(\mathbf{0}, I),$$

which is gradient descent on U plus Gaussian noise. To sample from a posterior $p(\boldsymbol{\theta} \mid \mathcal{D}) \propto e^{-U(\boldsymbol{\theta})}$, we take $U(\boldsymbol{\theta}) = -\log p(\mathcal{D} \mid \boldsymbol{\theta}) - \log p(\boldsymbol{\theta})$, the training loss plus the regularizer. Without the noise, the iteration converges to the MAP estimate; with the noise, it explores the posterior. The discretization introduces a small bias: for the target $\mathcal{N}(2, 0.25)$ and $h = 0.01$, the stationary variance of the algorithm is 0.2551 instead of 0.25, by Example 4.15.4. The Metropolis-adjusted Langevin algorithm removes this bias by accepting or rejecting each step as in Theorem 4.8.13, and stochastic gradient Langevin dynamics uses mini-batch gradients to make the method feasible for large models.

Remark 4.15.6 (SDEs in machine learning). 1. Diffusion models for image generation use an SDE of Ornstein–Uhlenbeck type that gradually turns images into Gaussian noise. The time-reversed process is again an SDE, whose drift involves the score $\nabla \log p_t(\mathbf{x})$ of the noised data distribution at time t. A neural network learns this score, with a variant of the score matching loss of Lecture 3, and new images are generated by simulating the reverse SDE, starting from pure noise.

2. The noise of mini-batch gradients makes SGD with a small step size behave approximately like the Euler–Maruyama scheme for an SDE $d\boldsymbol{\theta} = -\nabla L(\boldsymbol{\theta})\,dt + \sqrt{\eta\,\Sigma(\boldsymbol{\theta})}\,d\mathbf{B}$, where Σ is the covariance matrix of the mini-batch gradients. This heuristic is used to study which minima SGD prefers.

4.16 Key Takeaways

- A probability assigns numbers in $[0,1]$ to events, additively for disjoint events. Conditional probability $P(A \mid B) = \frac{P(A \cap B)}{P(B)}$ and Bayes' theorem $P(B \mid A) = \frac{P(A|B)P(B)}{P(A)}$ update beliefs with evidence; ignoring the prior $P(B)$ is the base rate fallacy.

- Random variables are described by a pmf or a density. The sum rule $p(x) = \sum_y p(x,y)$ and the product rule $p(x,y) = p(y \mid x)p(x)$ are the basic rules of probabilistic modeling; independence means that the joint distribution factorizes, and the density of a sum of independent random variables is the convolution of their densities.

- The expectation is linear; $\mathrm{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$; Markov's, Chebyshev's, Chernoff's and Jensen's inequalities relate them. For $X \geq 0$, $\mathbb{E}[X] = \sum_{k \geq 0} P(X > k)$ for integer values and $\mathbb{E}[X] = \int_0^\infty P(X > t)\,dt$ for densities. For densities, $p_Y(y) = \frac{p_X(x)}{|g'(x)|}$ under $y = g(x)$, and $F^{-1}(U)$, with the quantile function $F^{-1}(u) = \min\{x : F(x) \geq u\}$, samples from the CDF F.

- The moment generating function $M_X(t) = \mathbb{E}[e^{tX}]$ has the derivatives $M_X^{(k)}(0) = \mathbb{E}[X^k]$, and $M_{X+Y} = M_X M_Y$ for independent X and Y; for $\mathcal{N}(\mu, \sigma^2)$, it is $e^{\mu t + \sigma^2 t^2/2}$. Optimizing the Chernoff bound $e^{-ta} M_X(t)$ over t gives exponential tail bounds.

- Covariance is an inner product on centered random variables, and correlation is the cosine of their angle, so $|\,\mathrm{Corr}\,| \leq 1$. Uncorrelated does not imply independent. $\mathbb{E}[Y \mid X]$ is the best mean-squared-error predictor of Y, the orthogonal projection of Y onto functions of X, and $\mathrm{Var}(Y) = \mathbb{E}[\mathrm{Var}(Y \mid X)] + \mathrm{Var}(\mathbb{E}[Y \mid X])$ (law of total variance).

- A unit $z = \sum_{j=1}^n W_j x_j$ with independent weights of mean 0 and variance v has $\mathrm{Var}(z) = nv\mathbb{E}[x_j^2]$, and a ReLU halves the second moment of a symmetric z. So signals keep their size through deep ReLU networks with $v = \frac{2}{n_{\text{in}}}$ (He initialization), and vanish or explode geometrically otherwise; Xavier initialization $v = \frac{2}{n_{\text{in}} + n_{\text{out}}}$ balances the forward and the backward pass.

- $X \perp\!\!\!\perp Y \mid Z$ means $p(x, y \mid z) = p(x \mid z)p(y \mid z)$; it neither implies nor follows from independence (common causes, explaining away). Directed graphical models factor $p(\mathbf{x}) = \prod_i p(x_i \mid x_{\mathrm{pa}(i)})$; in chains and forks, the middle variable makes the ends conditionally independent, in a collider, it makes independent causes dependent. Naive Bayes assumes conditionally independent features, $p(y \mid \mathbf{x}) \propto p(y) \prod_j p(x_j \mid y)$, and has linear log-odds like logistic regression.

- In a Poisson process with rate λ, the waiting times are i.i.d. Exponential(λ), the n-th arrival time has the density $\frac{\lambda^n t^{n-1}}{(n-1)!} e^{-\lambda t}$, and the number of arrivals in the time t is Poisson(λt).

- The covariance matrix $\Sigma = \mathbb{E}[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^\top]$ is symmetric positive semidefinite, $\mathrm{Cov}(A\mathbf{X} + \mathbf{b}) = A\Sigma A^\top$, and $\mathrm{Var}(\mathbf{u}^\top \mathbf{X}) = \mathbf{u}^\top \Sigma \mathbf{u}$ is maximal for the top eigenvector (PCA). Whitening $\Sigma^{-1/2}(\mathbf{X} - \boldsymbol{\mu})$ gives identity covariance.

- $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ is the distribution of $\boldsymbol{\mu} + A\mathbf{Z}$ with $AA^\top = \Sigma$ and $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$; its level sets are ellipsoids along the eigenvectors of Σ. Affine maps, marginals and conditionals of Gaussians are Gaussian, with $\mathbf{X}_a \mid \mathbf{x}_b \sim \mathcal{N}\big(\boldsymbol{\mu}_a + \Sigma_{ab}\Sigma_{bb}^{-1}(\mathbf{x}_b - \boldsymbol{\mu}_b), \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}\big)$; for jointly Gaussian vectors, uncorrelated means independent.

- The sample mean of N i.i.d. variables has standard error $\frac{\sigma}{\sqrt{N}}$; it converges to μ in probability (law of large numbers), its fluctuations are approximately Gaussian (central limit theorem), and for bounded variables Hoeffding's inequality bounds large deviations exponentially. Slutsky's theorem allows us to replace σ by s in confidence intervals, and the delta method gives the approximate standard error $|g'(\theta)|\,\mathrm{SE}$ of a smooth function $g(\hat{\theta})$ of an estimator.

- Averages of samples underlie Monte Carlo methods, test set evaluation, and stochastic gradients. Rejection sampling draws exact samples using an envelope $p \le Mq$ (acceptance rate $\frac{1}{M}$); importance sampling reweights samples from q by the likelihood ratio $\frac{p}{q}$, which can reduce the variance dramatically for rare events.

- SGD with unbiased stochastic gradients converges for strongly convex functions with step sizes $\eta_k = \frac{1}{\mu(k+1)}$, at the rate $\mathbb{E}\|\mathbf{x}_k - \mathbf{x}^*\|^2 \le \frac{G^2}{\mu^2 k}$; with a constant step size, it converges to a noise ball whose size is proportional to the step size.

- A Markov chain with transition matrix P evolves by $\boldsymbol{\pi}_{t+1} = \boldsymbol{\pi}_t P$. Stationary distributions are left eigenvectors of P with eigenvalue 1; an irreducible, aperiodic chain converges to its unique stationary distribution, at a speed governed by the second largest eigenvalue in absolute value. Detailed balance $\pi_i P_{ij} = \pi_j P_{ji}$ implies stationarity; the Metropolis algorithm uses it to sample from distributions that are only known up to a constant (MCMC).

- Gibbs sampling updates one coordinate at a time from its conditional distribution; each update satisfies detailed balance, so the target distribution is stationary. It is slow when the coordinates are strongly correlated.

- The value $\mathbf{V}$ of a Markov reward process, the expected discounted sum of rewards, solves the Bellman equation $\mathbf{V} = \mathbf{r} + \gamma P\mathbf{V}$, so $\mathbf{V} = (I - \gamma P)^{-1}\mathbf{r} = \sum_k \gamma^k P^k \mathbf{r}$. Since $\|P\mathbf{x}\|_\infty \le \|\mathbf{x}\|_\infty$, the Bellman maps are γ-contractions in the maximum norm: policy evaluation and value iteration converge like γ^k, the Bellman optimality equation has exactly one solution $\mathbf{V}^*$, and the greedy policy with respect to $\mathbf{V}^*$ is optimal. Q-learning is a stochastic version of value iteration.

- A time series is (weakly) stationary if its mean and its autocovariances $c(h) = \mathrm{Cov}(X_t, X_{t+h})$ do not depend on t. The AR(1) process $X_t = \phi X_{t-1} + \varepsilon_t$ is stationary for $|\phi| < 1$, with the variance $\frac{\sigma^2}{1-\phi^2}$ and the autocorrelations ϕ^h; for $|\phi| \ge 1$, its variance grows without bound, like that of a random walk, which differencing makes stationary. An AR(p) process is stationary if the eigenvalues of its companion matrix lie inside the unit circle.

- Estimators have a bias and a variance, and MSE = bias^2 + variance, so some bias (shrinkage) can pay off; if both tend to 0, the estimator is consistent. The sample variance needs the factor $\frac{1}{N-1}$ to be unbiased. No unbiased estimator has a variance below the Cramér–Rao bound $\frac{1}{N\,I(\theta)}$, where the Fisher information $I(\theta)$ is the variance of the score $\frac{\partial}{\partial\theta}\log p(x \mid \theta)$; the MLE attains it asymptotically, and its standard error is estimated from the curvature of the log-likelihood. A sufficient statistic carries all the information about θ, and conditioning an estimator on it never increases the MSE (Rao–Blackwell).

- Maximum likelihood maximizes $\sum_i \log p(x_i \mid \theta)$. Least squares is maximum likelihood with Gaussian noise, and the cross-entropy loss is maximum likelihood for categorical labels; the MLE of a Gaussian is the sample mean and the (biased) sample covariance. The method of moments matches sample moments instead; it is simpler but usually less efficient. Confidence intervals $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$ quantify uncertainty.

- Exponential families $h(x)e^{\boldsymbol{\eta}^\top T(x) - A(\boldsymbol{\eta})}$ include the Bernoulli, Poisson, Gaussian and categorical distributions. The log-partition function A is convex, with $\nabla A = \mathbb{E}[T]$, so the log-likelihood is concave and the MLE matches $\mathbb{E}[T]$ with the average of the $T(x_i)$. Generalized linear models have the gradient "prediction minus target" times the input.

- The empirical CDF converges to F uniformly in t, with exponentially small failure probabilities, which gives confidence bands and the Kolmogorov–Smirnov test. Kernel density estimates $\frac{1}{Nh}\sum_i K(\frac{x - X_i}{h})$ have a bias of order h^2 and a variance of order $\frac{1}{Nh}$; the best bandwidth is of order $N^{-1/5}$, and the rate $N^{-4/(4+d)}$ in d dimensions shows the curse of dimensionality.

- A waiting time T is described by its survival function $S(t) = P(T > t) = e^{-H(t)}$ and its hazard rate $h = \frac{p}{S}$, the rate of events among the survivors; a constant hazard means an exponential distribution. Censored observations, of which we only know $T > t_i$, contribute $S(t_i)$ to the likelihood; for exponential times, $\hat{\lambda}$ is the number of events divided by the total time at risk. Ignoring censoring biases the estimates. The Kaplan–Meier estimator $\prod_{\tau_j \le t}(1 - \frac{d_j}{n_j})$ generalizes the empirical survival function to censored data.

- A test rejects H_0 if the p-value, the probability under H_0 of data at least as extreme as observed, is at most α; under H_0, p-values are uniform, so the false positive rate is α, and the power is the probability of detecting a real effect. A p-value is not the probability that H_0 is true. Paired t-tests compare two models on the same data sets, and Pearson's χ^2 test compares observed counts with expected counts. Many tests need corrections: Bonferroni ($\frac{\alpha}{m}$) controls the family-wise error rate, Benjamini–Hochberg the false discovery rate. The bootstrap (resampling the data with replacement) estimates standard errors of any statistic, and permutation tests (relabeling the groups) give exact p-values without distributional assumptions.

- Neyman–Pearson: for two simple hypotheses, the likelihood ratio test is the most powerful test at its level; for a Gaussian mean, it is the one-sided z-test.

- In linear regression with uncorrelated noise of constant variance σ^2, least squares is unbiased with $\mathrm{Cov}(\hat{\mathbf{w}}) = \sigma^2(X^\top X)^{-1}$, and it is the best linear unbiased estimator (Gauss–Markov); $s^2 = \frac{\mathrm{RSS}}{N-n}$ estimates σ^2 without bias. For Gaussian noise, $\frac{\hat{w}_j - w_j}{\mathrm{SE}_j}$ has the t_{N-n}-distribution, which gives exact confidence intervals and t-tests for the coefficients. With heteroscedastic noise, $\hat{\mathbf{w}}$ is still unbiased, but its covariance is the sandwich $(X^\top X)^{-1}X^\top \Sigma X (X^\top X)^{-1}$, and the usual standard errors are wrong; robust standard errors or weighted least squares fix this.

- Cross-validation estimates the generalization error, but its naive standard error is too small, and the best of many cross-validated models is optimistically biased; compare models with paired differences and evaluate the final choice on untouched data. A/B tests randomize to measure causal effects, since a confounder can reverse a comparison in every group (Simpson's paradox); plan the sample size $n \approx (z_{1-\alpha/2} + z_{1-\beta})^2\, 2\bar{p}(1-\bar{p})/\delta^2$ per group in advance, and do not stop at the first significant peek.

- Precision, recall and the false positive rate describe the errors of a classifier at a threshold; the ROC curve shows all thresholds, and the AUC is the fraction of correctly ordered positive–negative pairs. The cross-entropy loss is a proper scoring rule. For $|\mathcal{H}|$ models and N i.i.d. examples, all empirical risks are within $\sqrt{\log(2|\mathcal{H}|/\delta)/(2N)}$ of the risks with probability $1 - \delta$ (Hoeffding and the union bound).

- Bayesian inference updates a prior to the posterior $p(\theta \mid \mathcal{D}) \propto p(\mathcal{D} \mid \theta)p(\theta)$. Conjugate priors (Beta for Bernoulli, Gamma for Poisson, Gaussian for linear regression) give closed-form posteriors, and credible intervals are quantiles of the posterior. MAP estimation adds the regularizer $-\log p(\theta)$: a Gaussian prior gives ridge regression, a Laplace prior the lasso. The Bayesian posterior also gives predictive uncertainty. A Gaussian process is a prior on functions given by a kernel; its posterior follows from Gaussian conditioning, with the mean $\mathbf{k}_*^\top (K + \sigma^2 I)^{-1}\mathbf{y}$ and a variance that is small near the data.

- Entropy measures uncertainty, cross-entropy $H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q)$, and $\mathrm{KL}(p \,\|\, q) \geq 0$ with equality only for $p = q$ (Gibbs). Maximum likelihood minimizes the cross-entropy and the KL divergence from the empirical distribution to the model. Among all densities with a given variance, the Gaussian has the largest entropy.

- In latent variable models, $\log p(\mathbf{x} \mid \theta) = \mathrm{ELBO}(q, \theta) + \mathrm{KL}(q \,\|\, p(z \mid \mathbf{x}, \theta))$. EM alternates between setting q to the posterior (E-step) and maximizing the ELBO over θ (M-step), and never decreases the likelihood; for Gaussian mixtures, it alternates responsibilities and weighted means and covariances. k-means is its hard-assignment limit. Probabilistic PCA, $\mathbf{x} = W\mathbf{z} + \boldsymbol{\mu} + \boldsymbol{\varepsilon}$, has the maximum likelihood solution $\hat{W} = U_k(\Lambda_k - \hat{\sigma}^2 I)^{1/2}$, which spans the principal subspace, with $\hat{\sigma}^2$ the average of the remaining eigenvalues. Variational autoencoders maximize the ELBO over an encoder and a decoder.

- Brownian motion has independent Gaussian increments $B(t) - B(s) \sim \mathcal{N}(0, t - s)$ and continuous, nowhere differentiable paths. An SDE $dX = \mu(X)\,dt + \sigma(X)\,dB$ is simulated by Euler–Maruyama, $X_{k+1} = X_k + \mu(X_k)h + \sigma(X_k)\sqrt{h}\,Z_k$: noise of size $\sqrt{h}$, not h. Langevin dynamics $d\mathbf{X} = -\nabla U\,dt + \sqrt{2}\,d\mathbf{B}$ samples from $p \propto e^{-U}$, the stationary solution of the Fokker–Planck equation; its discretization is gradient descent plus noise, and diffusion models generate data by reversing a noising SDE.

4.17 Exercises

1. (The birthday problem.) Assume that birthdays are independent and uniformly distributed over 365 days. Show that the probability that among n people at least two share a birthday is $1 - \frac{365 \cdot 364 \cdots (365 - n + 1)}{365^n}$, and that it exceeds $\frac{1}{2}$ for $n = 23$. How many people are needed for 99%?

2. (a) Prove the inclusion–exclusion formula $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$.

   (b) Two fair coins are flipped. Let A = "the first coin shows heads", B = "the second coin shows heads", and C = "exactly one coin shows heads". Show that any two of these events are independent, but the three events are not independent.

3. A spam filter knows that 40% of all emails are spam, that the word "free" appears in 30% of the spam emails and in 2% of the other emails. What is the probability that an email containing "free" is spam?

4. Let X be Geometric(p). Show that $\mathbb{E}[X] = \frac{1}{p}$ and that $P(X > s + t \mid X > s) = P(X > t)$ for all positive integers s and t. How many times do you expect to roll a fair die until you see a six?

5. Let U and V be independent and Uniform$(0, 1)$. Find the densities of (a) U^2, (b) $-\log U$, (c) $\max(U, V)$ (Hint: compute the CDF first.)

6. Let X be Poisson(λ). Compute $\mathbb{E}[X(X - 1)]$, and use it to show that $\mathrm{Var}(X) = \lambda$.

7. (a) A nonnegative random variable has mean 2. Bound $P(X \geq 10)$ with Markov's inequality, and compare with the exact value for $X \sim$ Exponential($\frac{1}{2}$).

(b) Use Jensen's inequality to show that $\mathbb{E}[\frac{1}{X}] \geq \frac{1}{\mathbb{E}[X]}$ for every positive random variable X, and that $\mathbb{E}[e^X] \geq e^{\mathbb{E}[X]}$.

8. Let (X, Y) have the joint pmf $p(0,0) = 0.1$, $p(0,1) = 0.2$, $p(0,2) = 0.1$, $p(1,0) = 0.2$, $p(1,1) = 0.1$, $p(1,2) = 0.3$. Compute the marginal distributions, $\mathbb{E}[Y \mid X = 0]$ and $\mathbb{E}[Y \mid X = 1]$, and check the tower property. Compute $\mathrm{Cov}(X, Y)$. Are X and Y independent?

9. Let $X \sim \mathcal{N}(0,1)$. Show that X and X^2 are uncorrelated, but not independent. Show that $\mathrm{Corr}(X, aX + b) = \mathrm{sign}(a)$ for $a \neq 0$.

10. (Initialization.)

(a) Let $z \sim \mathcal{N}(0, s^2)$. Show that $\mathbb{E}[\mathrm{ReLU}(z)] = \frac{s}{\sqrt{2\pi}}$, $\mathbb{E}[\mathrm{ReLU}(z)^2] = \frac{s^2}{2}$ and $\mathrm{Var}(\mathrm{ReLU}(z)) = s^2(\frac{1}{2} - \frac{1}{2\pi})$.

(b) (The backward pass.) Let W be an $m \times n$ matrix with independent entries of mean 0 and variance v, and let $\boldsymbol{\delta} \in \mathbb{R}^m$ be a random vector independent of W, with $\mathbb{E}[\delta_i^2] = q$ for all i. Show that every entry of $W^\top \boldsymbol{\delta}$, the gradient that backpropagation passes to the inputs of a linear layer $\mathbf{y} = W\mathbf{x}$, has mean 0 and variance mvq. Which condition on v keeps the size of the gradients constant?

(c) Find the standard deviations of the He and Xavier normal initializations for a layer with 256 inputs and 128 outputs, and the bound b of the uniform He initialization on $[-b, b]$.

11. Let X and Y be independent and Exponential(λ). Show that $X + Y$ has the density $\lambda^2 z e^{-\lambda z}$ for $z > 0$.

12. (The Poisson process.)

(a) Show that the minimum of independent $X \sim$ Exponential(λ) and $Y \sim$ Exponential(μ) is Exponential($\lambda + \mu$), and that $P(X < Y) = \frac{\lambda}{\lambda+\mu}$.

(b) Requests arrive at a server as a Poisson process with the rate 2 per second. Find the probability of no request in one second, the probability of at least 3 requests in two seconds, and the expected waiting time until the 10th request.

13. (Conditional independence.)

(a) Show that $X \perp\!\!\!\perp Y \mid Z$ if and only if $p(x, y, z) = g(x, z)h(y, z)$ for some functions g and h.

(b) In a chain $X \to Z \to Y$ of binary variables, let $P(Z = 1 \mid X = 1) = 0.9$, $P(Z = 1 \mid X = 0) = 0.2$, $P(Y = 1 \mid Z = 1) = 0.7$ and $P(Y = 1 \mid Z = 0) = 0.1$. Compute $P(Y = 1 \mid X = 1)$, $P(Y = 1 \mid X = 0)$ and $P(Y = 1 \mid X = 1, Z = 1)$. Are X and Y independent? Are they independent given Z?

(c) A naive Bayes classifier has seen the word "meeting" in 0 of 50 spam emails. What is the maximum likelihood estimate of $P(\text{meeting} \mid \text{spam})$, and what happens to an email that contains "meeting" and many typical spam words? What does Laplace smoothing, adding 1 to each of the counts of "contains the word" and "does not contain the word", estimate instead?

14. (Moment generating functions.) Let $X \sim$ Exponential(λ). Use the cumulant generating function $K_X(t) = \log \frac{\lambda}{\lambda - t}$ to find $\mathbb{E}[X]$ and $\mathrm{Var}(X)$. Find the moment generating function of the sum of n independent Exponential(λ) variables, and its mean and variance. Show that $\mathbb{E}[X^k] = \frac{k!}{\lambda^k}$ by expanding $M_X(t) = \frac{1}{1 - t/\lambda}$ as a geometric series.

15. Let $\mathbf{X}$ have the covariance matrix $\Sigma = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$. Compute $\operatorname{Var}(X_1 + X_2 + X_3)$ and $\operatorname{Var}(X_1 - X_3)$. In which direction $\mathbf{u}$ (with $\|\mathbf{u}\| = 1$) is $\operatorname{Var}(\mathbf{u}^\top \mathbf{X})$ largest, and what is it? Find a matrix A such that $A\mathbf{X}$ has covariance matrix I.

16. Let $(X, Y) \sim \mathcal{N}\Big(\mathbf{0}, \begin{pmatrix} 4 & 2 \\ 2 & 2 \end{pmatrix}\Big)$. Find the conditional distributions of Y given $X = x$ and of X given $Y = y$, the distribution of $X - Y$, and $P(X > Y)$. Describe the level curves of the density.

17. Let $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ in $\mathbb{R}^d$. Show that the squared Mahalanobis distance $(\mathbf{X} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{X} - \boldsymbol{\mu})$ has the same distribution as $\|\mathbf{Z}\|^2$ with $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I)$, and that its expectation is d. What does this say about where the probability mass of a high-dimensional Gaussian lies?

18. How many times do we have to flip a fair coin so that the fraction of heads is within 0.01 of $\frac{1}{2}$ with probability at least 95%? Answer with Chebyshev's inequality, with Hoeffding's inequality, and with the central limit theorem, and compare.

19. (Delta method.) Let $X_1, \dots, X_N$ be i.i.d. Exponential(λ). Show that $\sqrt{N}\big(\frac{1}{\bar{X}_N} - \lambda\big) \to \mathcal{N}(0, \lambda^2)$ in distribution, and compare with the Fisher information of the exponential distribution (Exercise 29).

20. Let $p(x) = \sqrt{2/\pi}\, e^{-x^2/2}$ for $x > 0$ be the density of $|Z|$ for $Z \sim \mathcal{N}(0,1)$, and let $q(x) = e^{-x}$ be the Exponential(1) density.

    (a) Find the smallest M with $p \le Mq$, and the acceptance rate of rejection sampling with this proposal.

    (b) Write down the importance sampling estimator of $\mathbb{E}_p[X^2]$ with samples from q, and explain why its variance is finite.

21. (SGD with a constant step size.) For $f(\theta) = \frac{1}{2N}\sum_i (\theta - y_i)^2$, SGD with mini-batches of size 1 and a constant step size $\eta \in (0,1)$ is $\theta_{k+1} = \theta_k - \eta(\theta_k - y_{I_k})$. Show that $e_k = \theta_k - \bar{y}$ satisfies $e_{k+1} = (1-\eta)e_k + \eta(y_{I_k} - \bar{y})$, and that $\mathbb{E}[e_k^2] \to \frac{\eta s^2}{2-\eta}$, where $s^2 = \frac{1}{N}\sum_i (y_i - \bar{y})^2$. Compare with Theorem 4.7.16.

22. Consider the Markov chain with the transition matrix

$$P = \begin{pmatrix} 0.7 & 0.2 & 0.1 \\ 0.3 & 0.5 & 0.2 \\ 0.2 & 0.3 & 0.5 \end{pmatrix}.$$

    Draw its transition graph, compute $P(X_2 = 3 \mid X_0 = 1)$, and find the stationary distribution. Is the chain irreducible and aperiodic? In the long run, what fraction of the time does it spend in state 1?

23. (Random walks on graphs.)

    (a) Let the walker of Example 4.8.11 move on the cycle with the four vertices $1, 2, 3, 4$ (and the edges $\{1,2\}, \{2,3\}, \{3,4\}, \{4,1\}$). Find the stationary distribution and the eigenvalues of P. Show that the distribution of X_t does not converge if $X_0 = 1$.

    (b) The lazy walk stays at the current vertex with probability $\frac{1}{2}$, i.e., it has the transition matrix $\frac{1}{2}(I + P)$. Show that it has the same stationary distribution, find its eigenvalues, and explain why it converges.

24. (Gibbs sampling.) Let (X_1, X_2) be uniformly distributed on the four points $(0,0)$, $(0,1)$, $(1,1)$ and $(1,2)$. Find the conditional distributions of X_1 given X_2 and of X_2 given X_1, write down the 4×4 transition matrix of one sweep of the Gibbs sampler (first update X_1, then X_2), and check that the uniform distribution is stationary. Is the chain irreducible?

25. (Markov reward processes and decision processes.)

    (a) For the weather chain of Example 4.8.18, compute $\mathbf{V}$ as a function of γ, and show that $(1-\gamma)\mathbf{V} \to (\frac{5}{6}, \frac{5}{6})^\top$ as $\gamma \to 1$. For a general irreducible, aperiodic chain with the stationary distribution $\boldsymbol{\pi}$, explain with Theorem 4.8.6 why $(1-\gamma)V_i \to \boldsymbol{\pi}\mathbf{r}$ for every state i: the long-run average reward does not depend on the start.

    (b) Show that the Bellman optimality operator satisfies $T(\mathbf{V}+c\mathbf{1}) = T\mathbf{V}+\gamma c\mathbf{1}$ for every $c \in \mathbb{R}$, and that value iteration started at $\mathbf{V}_0 = \mathbf{0}$ produces nondecreasing iterates if all rewards are nonnegative.

    (c) (Policy iteration.) Starting from a policy π_0, compute $\mathbf{V}^{\pi_k}$ by solving the Bellman equation, and let π_{k+1} be greedy with respect to $\mathbf{V}^{\pi_k}$, i.e., let it maximize $r(s,a)+\gamma \sum_{s'} P(s' \mid s,a) V_{s'}^{\pi_k}$ over a in every state s. Show that $\mathbf{V}^{\pi_{k+1}} \geq \mathbf{V}^{\pi_k}$, and that if $\mathbf{V}^{\pi_{k+1}} = \mathbf{V}^{\pi_k}$, then π_k is optimal. Conclude that policy iteration finds an optimal policy after finitely many steps.

26. (Time series.)

    (a) Compute the autocovariances of the MA(2) process $X_t = \varepsilon_t + \theta_1 \varepsilon_{t-1} + \theta_2 \varepsilon_{t-2}$, with white noise ε_t of variance σ^2.

    (b) Show that the companion matrix A of an AR(p) process has the characteristic polynomial $\pm(\lambda^p - \phi_1 \lambda^{p-1} - \cdots - \phi_p)$. (Hint: show that $(\lambda^{p-1}, \ldots, \lambda, 1)^\top$ is an eigenvector of A if λ is a root of this polynomial, or expand the determinant along the last column.)

    (c) Show that the AR(2) process $X_t = 1.2X_{t-1} - 0.5X_{t-2} + \varepsilon_t$ satisfies the stationarity condition. Its eigenvalues are complex; what does this suggest about the shape of its autocorrelation function?

    (d) For a stationary AR(1) process, show that $\operatorname{Cov}(X_t, X_{t-1}) = \phi \operatorname{Var}(X_{t-1})$, so that $\phi = \rho(1)$ (the Yule–Walker equation), and compare with the least squares estimate $\hat{\phi}$.

27. Let $X_1, \ldots, X_N$ be i.i.d. Uniform$(0, \theta)$.

    (a) Show that the MLE of θ is $M = \max_i X_i$, and that $\mathbb{E}[M] = \frac{N}{N+1}\theta$. (Hint: $P(M \leq t) = (t/\theta)^N$.)

    (b) Show that $\frac{N+1}{N}M$ and $2\bar{X}_N$ are unbiased, and compare their variances, $\frac{\theta^2}{N(N+2)}$ and $\frac{\theta^2}{3N}$. Which estimator is better?

28. Find the maximum likelihood estimates for i.i.d. data from (a) Poisson(λ), (b) Exponential(λ), (c) Geometric(p).

29. (Fisher information.)

    (a) Show that the Fisher information of Poisson(λ) is $I(\lambda) = \frac{1}{\lambda}$, and that the MLE $\bar{X}_N$ is efficient.

    (b) Show that $I(\lambda) = \frac{1}{\lambda^2}$ for Exponential(λ). The sum S of $N \geq 3$ i.i.d. Exponential(λ) variables has the Gamma density $\frac{\lambda^N s^{N-1} e^{-\lambda s}}{(N-1)!}$ for $s > 0$. Use it to show that the MLE $\frac{1}{\bar{X}_N}$ has the expectation $\frac{N\lambda}{N-1}$, that $\frac{N-1}{N\bar{X}_N}$ is unbiased, and that its variance $\frac{\lambda^2}{N-2}$ is larger than the Cramér–Rao bound. Is the MLE asymptotically efficient?

(c) Counts of $N = 50$ independent Poisson observations have the sample mean 3. Estimate the standard error of $\hat{\lambda}$ from the observed information $-\ell''(\hat{\lambda})$, and give an approximate 95% confidence interval for λ.

30. (Rao–Blackwell.) Let $X_1, \dots, X_N$ be i.i.d. Uniform$(0, \theta)$. Use the factorization criterion to show that $M = \max_i X_i$ is sufficient for θ. Show that $\mathbb{E}[2X_1 \mid M] = \frac{N+1}{N} M$: Rao–Blackwellizing the unbiased estimator $2X_1$ gives the unbiased estimator of Exercise 27. (Hint: given $M = m$, $X_1 = m$ with probability $\frac{1}{N}$, and otherwise X_1 is uniform on $[0, m]$.)

31. Assume the regression model $y_i = \mathbf{w}^\top \mathbf{x}_i + \varepsilon_i$ with i.i.d. Laplace noise, with density $\frac{1}{2b} e^{-|\varepsilon|/b}$. Show that the MLE of $\mathbf{w}$ minimizes $\sum_i |y_i - \mathbf{w}^\top \mathbf{x}_i|$. In the special case $\mathbf{x}_i = 1$ (estimating a single number w), show that every median of the y_i is a minimizer.

32. Find the method of moments estimators of (a) a and b for Uniform(a, b), using the mean and the variance; (b) the shape k and the scale s of a Gamma distribution, for data with the sample mean 4 and the sample variance 8.

33. (Exponential families.) Write the Geometric(p) distribution, $P(X = x) = p(1-p)^{x-1}$ for $x = 1, 2, \dots$, as an exponential family, and find T, η and $A(\eta)$. Use Proposition 4.9.22 to find the mean and the variance, and Theorem 4.9.23 to find the MLE of p. Show also that the Exponential(λ) and the Gamma distributions with a known shape are exponential families.

34. (Nonparametric estimation.)

(a) Show that a kernel density estimate $\hat{f}_h$ is a probability density.

(b) For the uniform kernel $K = \frac{1}{2}\mathbf{1}_{[-1,1]}$, show that $\hat{f}_h(x) = \frac{1}{2Nh} \#\{i : |x - X_i| \le h\}$, a histogram with a moving bin centered at x.

(c) For $\mathcal{N}(0,1)$ data and the Gaussian kernel, show that $\mathbb{E}[\hat{f}_h(x)]$ is the density of $\mathcal{N}(0, 1+h^2)$ at x. (Hint: it is the density of $X + hU$ with independent $X, U \sim \mathcal{N}(0,1)$.) Compare $\mathbb{E}[\hat{f}_h(0)] - \varphi(0)$ with the bias $\frac{h^2}{2}\kappa_2 f''(0)$ of Proposition 4.9.25 for small h.

35. (Survival analysis.)

(a) For the Weibull distribution with the shape k and the scale s, find the density, the median $s(\log 2)^{1/k}$, and the mean $s\,\Gamma(1 + \frac{1}{k})$, using $\mathbb{E}[T] = \int_0^\infty S(t)\,dt$ and the Gamma function $\Gamma(a) = \int_0^\infty u^{a-1} e^{-u}\,du$.

(b) (Discrete time.) A customer who is still subscribed at the beginning of month j cancels during that month with probability h_j. Show that the probability to be still subscribed after t months is $\prod_{j=1}^{t}(1 - h_j)$, and compare with the Kaplan–Meier estimator.

(c) For Weibull event times with a known shape k and non-informative censoring, show that the maximum likelihood estimate of $\theta = s^{-k}$ is $\hat{\theta} = \frac{D}{\sum_i t_i^k}$, where D is the number of observed events.

(d) Show that without censoring, the Kaplan–Meier estimator equals $1 - \hat{F}_N$.

36. A die is rolled 120 times and shows 30 sixes. Test at the level 0.05 whether the die favors sixes: state H_0 and H_1, and compute the exact one-sided p-value $P(S \ge 30)$ for $S \sim$ Binomial$(120, \frac{1}{6})$ with a computer, and its normal approximation.

37. (Neyman–Pearson.) Let $X_1, \dots, X_N$ be i.i.d. Exponential(λ). Find the most powerful test at the level α of H_0: $\lambda = 1$ against H_1: $\lambda = 2$, and show that it rejects if $\sum_i X_i$ is small. For $N = 1$ and $\alpha = 0.05$, find the rejection region and the power.

38. (Inference for linear regression.)

(a) For the simple regression $y_i = w_0 + w_1 x_i + \varepsilon_i$ with homoscedastic noise, show that $\operatorname{Var}(\hat{w}_1) = \frac{\sigma^2}{\sum_i (x_i - \bar{x})^2}$. How should the inputs x_i be chosen in an experiment to estimate the slope precisely?

(b) For heteroscedastic noise with $\operatorname{Cov}(\boldsymbol{\varepsilon}) = \operatorname{diag}(\sigma_1^2, \dots, \sigma_N^2)$, show that $\mathbb{E}[\mathrm{RSS}] = \sum_i \sigma_i^2 (1 - H_{ii})$, where $H = X(X^\top X)^{-1} X^\top$.

(c) In the model $y_i = w x_i + \varepsilon_i$ with $\sigma_i = \sigma |x_i|$ and all $x_i \neq 0$, show that weighted least squares gives $\tilde{w} = \frac{1}{N} \sum_i \frac{y_i}{x_i}$, with the variance $\frac{\sigma^2}{N}$, and that this is at most the variance $\frac{\sigma^2 \sum_i x_i^4}{(\sum_i x_i^2)^2}$ of ordinary least squares. Which theorem explains this, applied to the rescaled data $\frac{y_i}{x_i}$?

39. Out of 200 users, 70, 80 and 50 prefer the layouts A, B and C. Use Pearson's χ^2 test to test at the level 0.05 whether the three layouts are equally popular. (The χ^2_2 distribution has the tail $P(\chi^2_2 \geq c) = e^{-c/2}$.)

40. (a) Show that for m independent tests of true null hypotheses at the level $\frac{\alpha}{m}$, the probability of at least one rejection is $1 - (1 - \frac{\alpha}{m})^m \leq \alpha$.

(b) In a permutation test with two groups of three observations each, what is the smallest possible one-sided p-value? What does this mean for very small experiments?

(c) Show that the bootstrap standard error of the sample mean (for $B \to \infty$) is $\frac{\hat{\sigma}}{\sqrt{N}}$ with $\hat{\sigma}^2 = \frac{1}{N} \sum_i (x_i - \bar{x})^2$.

41. An online shop wants to test whether a new design increases the click-through rate from 10% to 11%. How many users per group are needed for a two-sided test at the level 0.05 with power 0.8? After the experiment, the two groups of 15 000 users have 1500 and 1650 clicks. Compute the z-statistic, the p-value, and a 95% confidence interval for the difference.

42. (Classification metrics.) A test for a disease with a prevalence of 2% has a true positive rate of 90% and a false positive rate of 5%. Compute its precision. Which threshold-independent number does not change when the prevalence changes, and why? For the scores $0.9, 0.7, 0.6, 0.4, 0.2$ of five examples with the labels $1, 0, 1, 0, 0$, draw the ROC curve and compute the AUC in two ways.

43. (Generalization.) How many i.i.d. test examples are needed to estimate the error rate of one classifier within ± 0.01 with probability at least 0.95, using Hoeffding's inequality? How many are needed to estimate the error rates of 1000 classifiers simultaneously with the same accuracy?

44. (Simpson's paradox.) Two treatments are compared in groups $s = 1, \dots, m$, with the success rates $a_s = P(\text{success} \mid A, s)$ and $b_s = P(\text{success} \mid B, s)$, and $a_s > b_s$ for every s. Show that if $P(s \mid A) = P(s \mid B)$ for all s, as in a randomized experiment, then $P(\text{success} \mid A) > P(\text{success} \mid B)$. Then construct an example with two groups in which $a_s > b_s$ for both groups, but $P(\text{success} \mid A) < P(\text{success} \mid B)$.

45. With the uniform prior $\mathrm{Beta}(1, 1)$, a coin shows 7 heads in 10 flips. Find the posterior distribution, its mean, the MAP estimate, the MLE, and the predictive probability of heads in the next flip.

46. (Gaussian–Gaussian conjugacy.) Let $x_1, \dots, x_N$ be i.i.d. $\mathcal{N}(\mu, \sigma^2)$ with known σ^2, and let the prior be $\mu \sim \mathcal{N}(\mu_0, \tau^2)$. Show that the posterior is Gaussian with precision $\frac{1}{\tau^2} + \frac{N}{\sigma^2}$ and mean

$$\frac{\mu_0/\tau^2 + N\bar{x}/\sigma^2}{1/\tau^2 + N/\sigma^2},$$

a weighted average of the prior mean and the sample mean. What happens as $N \to \infty$?

47. (Gaussian processes and Bayesian linear regression.)

    (a) Show that $(X^\top X + \lambda I)X^\top = X^\top(XX^\top + \lambda I)$ for every matrix X and $\lambda > 0$, and conclude that $(X^\top X + \lambda I)^{-1}X^\top = X^\top(XX^\top + \lambda I)^{-1}$.

    (b) Use (a) to show that for the linear kernel $k(\mathbf{x}, \mathbf{x}') = \alpha^{-1}\mathbf{x}^\top\mathbf{x}'$, the posterior mean of Proposition 4.12.11 equals $\mathbf{m}^\top\mathbf{x}_*$ from Theorem 4.12.7. Which of the two formulas is cheaper to evaluate when $N \gg d$, and which when $d \gg N$?

    (c) Let $f \sim \mathcal{GP}(0, k)$ with $k(s,t) = \min(s,t)$, and observe $f(1) = 1$ without noise. Find the posterior mean and variance of $f(t)$ for $0 \le t \le 1$ and for $t > 1$, and sketch them.

48. (a) Compute the entropy of a fair die in bits.

    (b) Compute $\mathrm{KL}(\mathrm{Bernoulli}(p) \,\|\, \mathrm{Bernoulli}(q))$, and show that

$$\mathrm{KL}\left(\mathcal{N}(\mu_1, \sigma^2) \,\|\, \mathcal{N}(\mu_2, \sigma^2)\right) = \frac{(\mu_1 - \mu_2)^2}{2\sigma^2}.$$

    (c) Show that $I(X;Y) = H(X) + H(Y) - H(X,Y)$ for discrete random variables, where $H(X,Y)$ is the entropy of the joint distribution. Compute $I(X;Y)$ for Exercise 8.

49. (EM for a mixture of two coins.) In each of 5 experiments, one of two coins with the unknown head probabilities θ_A and θ_B is chosen with probability $\frac{1}{2}$ and flipped 10 times. The numbers of heads are $5, 9, 8, 4, 7$, and we do not know which coin was used. Starting from $\theta_A = 0.6$ and $\theta_B = 0.5$, carry out one step of EM: compute the posterior probabilities that coin A was used in each experiment, and the new estimates. Show that the log-likelihood increased.

50. (k-means.) Show that every assignment step and every update step of Lloyd's algorithm can only decrease J, and find an example with $N = 4$ points in $\mathbb{R}$ and $K = 2$ for which Lloyd's algorithm, started from a bad pair of centers, stops at an assignment that is not optimal.

51. (Probabilistic PCA.)

    (a) Show that the maximal log-likelihood of probabilistic PCA is $-\frac{N}{2}\big(d\log(2\pi) + \sum_{j\le k}\log\lambda_j + (d-k)\log\hat\sigma^2 + d\big)$.

    (b) For data with $S = \mathrm{diag}(4, 2, 1)$, compute it (per data point) for $k = 0$, 1 and 2. Why does it always increase with k, and why is it nevertheless not a good idea to choose k by the likelihood alone?

    (c) Show that the posterior covariance $\sigma^2 M^{-1}$ of $\mathbf{z}$ is $\sigma^2 I_k$ minus a positive semidefinite matrix, so that observing $\mathbf{x}$ reduces the uncertainty about $\mathbf{z}$ in every direction.

52. (Brownian motion and the Ornstein–Uhlenbeck process.)

    (a) Compute $\mathrm{Var}(B(2) - B(0.5))$, $\mathrm{Cov}(B(1), B(3))$ and $\mathrm{Corr}(B(1), B(3))$.

    (b) Let $c > 0$. Show that $W(t) = \frac{1}{\sqrt{c}}B(ct)$ has the same mean and covariance functions as $B(t)$. (In fact, W is again a Brownian motion: zooming into a Brownian path by the factor c in time and $\sqrt{c}$ in space gives a Brownian path.)

    (c) Let $X_{k+1} = (1 - \theta h)X_k + \sigma\sqrt{h}\,Z_k$ with $X_0 = x_0$ and i.i.d. $Z_k \sim \mathcal{N}(0,1)$. Show that $\mathbb{E}[X_k] = (1-\theta h)^k x_0$ and $\mathrm{Var}(X_k) = v_h\big(1 - (1-\theta h)^{2k}\big)$, with v_h from Example 4.15.4. What happens for $\theta h > 2$?

4.18 Programming Exercises

As in the previous lectures, implement the methods yourself with NumPy, and use library routines (such as `rng.normal` or `scipy.stats`) only to check your results. Start each file with

```
import numpy as np
import matplotlib.pyplot as plt
rng = np.random.default_rng(0)
```

P1. Monte Carlo estimation. Estimate π from N uniform random points in $[-1,1]^2$, for $N = 10^2, 10^3, \dots, 10^7$. Plot the absolute error against N on a log-log scale, together with the standard error $4\sqrt{\frac{\pi}{4}(1-\frac{\pi}{4})/N}$. What is the slope? Then estimate the volume of the unit ball in $\mathbb{R}^d$ for $d = 2, 5, 10, 20$ in the same way. What goes wrong in high dimensions?

P2. Sampling. Generate 10^5 samples from each of the following distributions, using only `rng.random()` (uniform samples), and compare histograms with the densities.

- Exponential(λ) by inverse transform sampling (Theorem 4.2.7).
- A discrete distribution with the values $x_1 < \cdots < x_K$ and the probabilities $p_1, \dots, p_K$, by its quantile function $F^{-1}(u) = \min\{x : F(x) \geq u\}$ (Theorem 4.2.7), using the cumulative sums and `np.searchsorted`. Compare the frequencies with the probabilities.
- $\mathcal{N}(0,1)$ with the Box–Muller method: for independent $U_1, U_2 \sim \text{Uniform}(0,1)$, the numbers $\sqrt{-2\log U_1}\cos(2\pi U_2)$ and $\sqrt{-2\log U_1}\sin(2\pi U_2)$ are independent standard normal.
- $\mathcal{N}(\boldsymbol{\mu}, \Sigma)$ in $\mathbb{R}^2$ with $\Sigma = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$, via $\boldsymbol{\mu} + L\mathbf{z}$ with the Cholesky factor L. Plot the samples together with the ellipses of Figure 4.2, and check the sample mean and covariance matrix. What fraction of the samples lies inside the ellipse with $r = 1$, and with $r = 2$? (Compare with $1 - e^{-r^2/2}$.)

P3. The law of large numbers and the central limit theorem. For $X_i \sim \text{Exponential}(1)$, simulate 10^4 sample means $\bar{X}_N$ for $N = 1, 2, 5, 30$, and plot histograms of the standardized means $\sqrt{N}(\bar{X}_N - 1)$ together with the standard normal density. Repeat with the Cauchy distribution (the ratio of two independent standard normal variables), which has no mean: plot $\bar{X}_N$ against N for one long sequence. What happens?

P4. Gaussian classifiers. Generate two classes in $\mathbb{R}^2$ from Gaussians with means $(0,0)$ and $(2,1)$ and the same covariance matrix $\Sigma = \begin{pmatrix} 1 & 0.5 \\ 0.5 & 1 \end{pmatrix}$, with 200 points each.

- Estimate the means and the shared covariance matrix by maximum likelihood (Theorem 4.9.10).
- Show (on paper) that with equal class priors, Bayes' theorem gives $P(y = 1 \mid \mathbf{x}) = \sigma(\mathbf{w}^\top \mathbf{x} + b)$ with $\mathbf{w} = \Sigma^{-1}(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_0)$ and $b = -\frac{1}{2}(\boldsymbol{\mu}_1^\top \Sigma^{-1} \boldsymbol{\mu}_1 - \boldsymbol{\mu}_0^\top \Sigma^{-1} \boldsymbol{\mu}_0)$: a logistic regression model.
- Plot the data and the decision boundary, and compare with logistic regression trained by gradient descent (Lecture 3).

P5. Maximum likelihood and bias. Draw 10^5 data sets of size $N = 5$ from $\mathcal{N}(2, 9)$. Compute the average of the MLE $\hat{\sigma}^2$ and of s^2 over the data sets, and compare with $\frac{N-1}{N}\sigma^2 = 7.2$ and $\sigma^2 = 9$. Then estimate the mean squared errors of both estimators. Which one has the smaller MSE?

P6. Rao–Blackwell and the delta method.

- For 10^5 data sets of $N = 10$ Poisson(1) counts, compute the three estimators $\mathbf{1}_{\{X_1=0\}}$, $(1-\frac{1}{N})^S$ and $e^{-\bar{X}_N}$ of e^{-1} (Example 4.9.18), and estimate their biases, variances and mean squared errors.
- For $N = 30$ Bernoulli(p) observations with $p = 0.1$, estimate the coverage probabilities of the interval $\hat{p} \pm 1.96\sqrt{\hat{p}(1-\hat{p})/N}$ and of the log-odds interval of Example 4.7.12 by simulation (skip the data sets with $\hat{p} \in \{0, 1\}$, and report how often this happens). Which interval keeps its promise of 95% better?

P7. The bootstrap. For $N = 50$ samples from Exponential(1), estimate the standard error of the sample median by resampling the data 2000 times with replacement and computing the median of each resample. Compare with the standard error of the median over many independent data sets. Use the same method to compute a 95% interval for the mean (the 2.5% and 97.5% quantiles of the bootstrapped means), and compare with $\bar{x} \pm 1.96\frac{s}{\sqrt{N}}$.

P8. Bayesian updating and Thompson sampling.

- Simulate 100 flips of a coin with $\theta = 0.3$, and plot the Beta posterior (starting from Beta$(1, 1)$) after $0, 1, 5, 20$ and 100 flips, as in Figure 4.5.
- A slot machine has three arms with unknown success probabilities 0.3, 0.5 and 0.6. Thompson sampling keeps a Beta posterior for each arm, and in each round draws one sample from each posterior and plays the arm with the largest sample. Simulate 1000 rounds, and plot how often each arm is played. Compare the total reward with always playing a random arm and with always playing the best arm.

P9. Bayesian linear regression. Generate 20 points $y = \sin(3x)+\varepsilon$ with x uniform in $[-1, 1]$ and $\varepsilon \sim \mathcal{N}(0, 0.1^2)$, and use the features $1, x, \dots, x^7$. Compute the posterior of Theorem 4.12.7 with $\alpha = 1$ and $\sigma = 0.1$. Plot the data, the posterior mean, the band of ± 2 predictive standard deviations, and a few functions with weights sampled from the posterior. Where is the uncertainty largest? Check that the posterior mean equals the ridge solution with $\lambda = \alpha\sigma^2$.

P10. Cross-entropy and KL divergence.

- Write functions for $H(p)$, $H(p, q)$ and $\mathrm{KL}(p \,\|\, q)$, and check $H(p, q) = H(p) + \mathrm{KL}(p \,\|\, q)$ and $\mathrm{KL} \geq 0$ for random distributions (e.g., `rng.dirichlet(np.ones(5))`).
- Let p be the mixture $\frac{1}{2}\mathcal{N}(-2, 0.5^2) + \frac{1}{2}\mathcal{N}(2, 0.5^2)$. On a grid, find the Gaussian $q = \mathcal{N}(\mu, s^2)$ that minimizes $\mathrm{KL}(p \,\|\, q)$, and the one that minimizes $\mathrm{KL}(q \,\|\, p)$ (try several starting points). Plot both. Why does the first cover both modes, while the second picks one of them?

P11. Gaussian process regression (Section 4.12.4), with the squared exponential kernel $k(x, x') = \exp(-\frac{(x-x')^2}{2\ell^2})$.

- Sample and plot a few functions from the prior $\mathcal{GP}(0, k)$ on a grid of 200 points in $[-3, 3]$, for $\ell = 0.1, 0.3, 1$. (Add $10^{-8}I$ to the kernel matrix before the Cholesky factorization.) Do the same for the Brownian motion kernel $\min(s, t)$ (Section 4.15) on $[0, 3]$.
- For 8 noisy observations $y_i = \sin(3x_i) + \varepsilon_i$ ($\varepsilon_i \sim \mathcal{N}(0, 0.1^2)$), compute the posterior mean and variance of Proposition 4.12.11 on the grid, and plot them with a band of ± 2 standard deviations, as in Figure 4.6. Use `np.linalg.solve` (or a Cholesky factorization) rather than the inverse. Sample a few functions from the posterior.

- Compute the log marginal likelihood $\log p(\mathbf{y})$ for ℓ on a grid from 0.05 to 3, and plot it. Which length scale does it select? Compare the posterior with the one for $\ell = 0.1$ and $\ell = 1$.

P12. Markov chains and MCMC.

- (A bigram language model.) Take a text of a few thousand words, split it into words, and estimate the transition probabilities $P(\text{next word} = j \mid \text{current word} = i)$ by counting. Generate random sentences from the chain. Compute the stationary distribution by power iteration, and compare it with the word frequencies in the text.
- (Metropolis sampling.) For the probability θ of heads, use the prior density $p(\theta) \propto e^{-10(\theta-0.5)^2}$ on $[0,1]$, which is not a Beta distribution, and the data "7 heads in 10 flips". Sample from the posterior $p(\theta \mid \mathcal{D}) \propto \theta^7(1-\theta)^3 e^{-10(\theta-0.5)^2}$ with the Metropolis algorithm, using the proposals $\theta' = \theta + \eta\varepsilon$ with $\varepsilon \sim \mathcal{N}(0,1)$ (and target 0 outside $[0,1]$). Compare the histogram of 50 000 samples with the posterior density, normalized numerically on a grid. Plot the chain for $\eta = 0.01$, 0.1 and 1, and report the acceptance rates. Which step size works best, and why?

P13. Rejection and importance sampling.

- Sample from the Beta$(2,5)$ distribution by rejection sampling with the uniform proposal. Find the best M numerically, compare the acceptance rate with $\frac{1}{M}$, and compare a histogram with the density.
- Estimate $P(Z > 4)$ and $P(Z > 6)$ for $Z \sim \mathcal{N}(0,1)$ with plain Monte Carlo and with importance sampling from $\mathcal{N}(a,1)$ for $a = 4$ (or 6), using $N = 10^4$ samples. Repeat 100 times to estimate the relative standard errors, and compare with the exact values $1-\Phi(4)$ and $1-\Phi(6)$. What happens with the proposal $\mathcal{N}(0,1)$ shifted to $a = 8$? Compute the effective sample size $\frac{(\sum_i w_i)^2}{\sum_i w_i^2}$ in each case.

P14. Hypothesis tests by simulation.

- Simulate 10 000 data sets of size $N = 30$ from $\mathcal{N}(0,1)$, compute the two-sided z-test p-value for H_0: $\mu = 0$ in each, and plot the histogram of the p-values. Repeat with data from $\mathcal{N}(0.5,1)$, and estimate the power of the test at the level 0.05.
- Generate 1000 tests, of which 50 have a real effect (data from $\mathcal{N}(0.8,1)$) and 950 do not (data from $\mathcal{N}(0,1)$), each with $N = 20$. Count the true and false discoveries without correction, with Bonferroni, and with Benjamini–Hochberg at the level 0.05.
- Implement a permutation test for the difference of the means of two groups, and apply it to two groups of 15 samples from Exponential(1) and Exponential(0.6). Compare its p-value with the z-test.
- (Peeking.) Simulate A/B tests with two identical versions (conversion rate 5%) and up to 10 000 users per group, computing the two-proportion z-test after every 200 users per group. What fraction of the experiments ever reaches $p < 0.05$? Compare with a single test at the end.

P15. Cross-validation. Use the polynomial regression data of the programming exercises of Lecture 2 (20 points $y = \sin(3x) + \varepsilon$).

- Implement k-fold cross-validation, and choose the polynomial degree from 1 to 12 with 5-fold cross-validation. Plot the cross-validation error and the training error against the degree.

- Repeat the cross-validation with 50 different random splits. How much does the selected degree, and the cross-validation error of a fixed degree, vary? Compare the spread with the naive standard error $\frac{s_e}{\sqrt{k}}$.
- Generate a large independent test set from the same distribution. Is the cross-validation error of the selected degree optimistic? Implement nested cross-validation and compare.

P16. Stochastic differential equations.

- Simulate 10 000 Brownian paths on $[0, 1]$ with the step $h = 0.001$. Check $\mathrm{Var}(B(t)) = t$ and $\mathrm{Cov}(B(0.3), B(0.8)) = 0.3$ empirically, and plot a few paths. Zoom into one path: plot it on $[0.5, 0.5 + \delta]$ for $\delta = 0.1, 0.01, 0.001$ (simulating the path with a step much smaller than δ). Why does it never look smooth?
- Simulate the Ornstein–Uhlenbeck process $dX = -\theta X\,dt + \sigma\,dB$ with Euler–Maruyama for $\theta = 1$, $\sigma = \sqrt{2}$, starting 10 000 paths at $x_0 = 3$. Plot the mean and the variance of $X(t)$ against t, together with $x_0 e^{-t}$ and $1 - e^{-2t}$. Compare the histogram at $t = 5$ with $\mathcal{N}(0, 1)$, and check v_h from Example 4.15.4 for $h = 0.5$.
- (Langevin sampling.) Use the unadjusted Langevin algorithm to sample from the mixture $p = \frac{1}{2}\mathcal{N}(-2, 0.5^2) + \frac{1}{2}\mathcal{N}(2, 0.5^2)$, i.e., $U = -\log p$ (compute ∇U by hand). Compare a histogram of one long chain with p, and count how often the chain moves between the two modes. Then sample the posterior of the logistic regression model of Lecture 3 on a small two-dimensional data set, with the prior $\mathcal{N}(\mathbf{0}, I)$, and plot the samples of $\mathbf{w}$ together with the MAP estimate.

P17. ROC curves. Generate 500 examples of each class with the scores $s \sim \mathcal{N}(0, 1)$ for the negatives and $s \sim \mathcal{N}(1, 1)$ for the positives.

- Compute the TPR and the FPR for all thresholds, plot the ROC curve, and compute the AUC as the area under the curve and as the fraction of correctly ordered pairs (Theorem 4.11.1). Compare with the exact value $\Phi(1/\sqrt{2}) \approx 0.76$ for these distributions.
- Plot the precision and the recall as functions of the threshold. Repeat with only 25 positives and 500 negatives: which curves change, and which do not?

P18. Gaussian mixtures and k-means. Generate 300 points in $\mathbb{R}^2$ from a mixture of three Gaussians with different means and covariance matrices.

- Implement Lloyd's algorithm for k-means with $K = 3$. Plot J after each step, and check that it never increases (Proposition 4.14.7). Run it from 10 random initializations; do they all give the same J?
- Implement EM for Gaussian mixtures (Theorem 4.14.5), initialized with the k-means result. Compute the responsibilities with logarithms and the log-sum-exp trick to avoid underflow. Plot the log-likelihood after each step, and check that it never decreases (Theorem 4.14.4).
- Plot the data colored by their most likely component, and the ellipses of the fitted covariance matrices. Where do EM and k-means disagree, and why?

P19. Naive Bayes. Generate 2000 examples with a label $y \in \{0, 1\}$ ($P(y = 1) = 0.3$) and 20 binary features that are conditionally independent given y, with $P(x_j = 1 \mid y)$ drawn uniformly from $[0.1, 0.9]$ for each j and y.

- Fit naive Bayes by counting, with Laplace smoothing, and logistic regression by gradient descent (Lecture 3), on 1000 training examples. Compare their test accuracies, and plot the predicted probabilities of one model against those of the other.

- Now copy the first feature 10 times, so that the features are no longer conditionally independent, and repeat. Which model's probabilities become overconfident, and why?

P20. Nonparametric estimation and probabilistic PCA.

- Draw $N = 100$ and $N = 1000$ samples from Exponential(1), plot the empirical CDF with the band $\hat{F}_N \pm \sqrt{\log(40)/(2N)}$ and the true CDF, and check in 1000 repetitions how often the true CDF leaves the band.
- Draw 500 samples from the mixture $0.5\,\mathcal{N}(-2,1) + 0.5\,\mathcal{N}(2, 0.5^2)$, and plot kernel density estimates with the Gaussian kernel and $h = 0.05$, 0.3 and 1, together with the true density. Choose h by maximizing the leave-one-out log-likelihood $\sum_i \log \hat{f}_{h,-i}(x_i)$, where $\hat{f}_{h,-i}$ omits the i-th point, and compare with the rule of thumb $h = 1.06\,\hat{\sigma} N^{-1/5}$.
- Generate $N = 1000$ points $\mathbf{x} = W\mathbf{z} + \boldsymbol{\varepsilon}$ in $\mathbb{R}^{10}$ with a random 10×2 matrix W, $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, I_2)$ and $\boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, 0.25I)$. Fit probabilistic PCA with Theorem 4.14.9, compare $\hat{\sigma}^2$ with 0.25, and compare the orthogonal projections onto the column spaces of $\hat{W}$ and W.

P21. Initialization of deep networks (Example 4.4.13).

- Pass a batch of 1000 standard normal inputs in $\mathbb{R}^{512}$ through 50 ReLU layers of width 512 with weights drawn from $\mathcal{N}(0, v)$, for $v = \frac{2}{512}$ (He), $\frac{1}{512}$ and $\frac{4}{512}$. Plot the mean of the squared activations against the layer on a logarithmic scale, and compare with the prediction $q_l = q_0(\frac{nv}{2})^l$.
- Repeat with tanh activations and $v = \frac{1}{512}$. Then backpropagate the gradient of the loss $\sum_i y_i$ of the last layer, and plot the mean squared gradient with respect to the activations of each layer. Which initialization keeps both the forward signal and the gradients at a constant size?

P22. Policy evaluation and value iteration (Section 4.8.2). A robot moves on a 4×4 grid with the actions up, down, left and right; it moves in the chosen direction with probability 0.8 and in one of the two perpendicular directions with probability 0.1 each, and stays in place if it would leave the grid. Entering the corner $(4, 4)$ gives the reward 1 and restarts the robot at $(1, 1)$, and every other step gives the reward -0.04. Let $\gamma = 0.95$.

- Compute the value of the uniformly random policy by solving the Bellman equation with `np.linalg.solve`, and by iterative policy evaluation. Plot the error $\|\mathbf{V}_k - \mathbf{V}\|_\infty$ against k on a logarithmic scale, together with $\gamma^k \|\mathbf{V}_0 - \mathbf{V}\|_\infty$.
- Find the optimal values with value iteration, and draw the greedy policy as arrows on the grid. Compare with policy iteration (Exercise 25): how many iterations does each method need?
- Implement Q-learning with ε-greedy exploration (a random action with probability $\varepsilon = 0.1$, otherwise the action with the largest Q), using simulated transitions only, and compare the learned greedy policy and $\max_a Q(s, a)$ with the results of value iteration.

P23. Time series (Section 4.8.3).

- Simulate AR(1) processes with $\phi = 0.5$, 0.9, 0.99 and 1 (a random walk) for $T = 1000$ steps with standard normal noise, starting at $X_0 = 0$, and plot them. Compute the sample autocorrelation $\hat{\rho}(h) = \frac{\sum_t (x_t - \bar{x})(x_{t+h} - \bar{x})}{\sum_t (x_t - \bar{x})^2}$ for $h = 0, \ldots, 30$, and compare with ϕ^h. What goes wrong for the random walk, and what happens after differencing?
- Estimate ϕ by least squares for 100 simulated series with $\phi = 0.9$ and $T = 50$, and for $T = 500$. Is $\hat{\phi}$ biased? Compare its standard deviation with $\sqrt{(1 - \phi^2)/T}$.

- Simulate the AR(2) processes of Section 4.8.3 and of Exercise 26, compute the eigenvalues of their companion matrices with `np.linalg.eigvals`, and relate them to the behavior of the simulated paths and of their autocorrelations.

P24. Survival analysis (Section 4.9.9). Generate $N = 200$ lifetimes from the Weibull distribution with the shape $k = 1.5$ and the scale $s = 10$ (by inverse transform sampling), and independent censoring times uniform on $[0, 15]$.

- Implement the Kaplan–Meier estimator, and plot it together with the true survival function $e^{-(t/s)^k}$ and with the empirical survival function of the observed times t_i that ignores the censoring. Which one is biased, and in which direction?
- Fit an exponential model with Proposition 4.9.28, and a Weibull model by maximizing the censored log-likelihood numerically over k and s (for example, with gradient ascent on $\log k$ and $\log s$). Plot both fitted survival functions together with the Kaplan–Meier estimate. Which model fits, and how do the fitted hazard rates differ?

P25. Inference for linear regression (Section 4.10.1).

- Generate $N = 20$ points $y = 1{+}2x{+}\varepsilon$ with x uniform on $[0, 10]$ and $\varepsilon \sim \mathcal{N}(0, 4)$. Compute $\hat{\mathbf{w}}$, s^2, the standard errors, the t-statistics and the 95% confidence intervals, and check them against `statsmodels` if you have it installed. In 10 000 repetitions, estimate the coverage probability of the interval for the slope.
- Repeat with heteroscedastic noise $\varepsilon \sim \mathcal{N}(0, (0.3x)^2)$. Compare the coverage of the usual interval with that of the interval based on the robust standard errors, and plot the residuals against x. How do the results change with $N = 200$?

Appendix A

Topology, Differential Topology and Functional Analysis

Introduction

In Lecture 3, we used several facts without proof: that a continuous function on a closed and bounded set attains its maximum, that a function with continuous partial derivatives is differentiable (Theorem 3.3.6), that integrals depending on a parameter are continuous (in the proof of Proposition 3.14.6), the implicit function theorem in the proof of the Lagrange multiplier theorem (Theorem 3.12.1), the necessity of the KKT conditions (Theorem 3.12.5), and the existence and uniqueness of solutions of differential equations (Theorem 3.15.2). We also met sets that are not open subsets of $\mathbb{R}^n$, but curved objects: spheres, the set of orthogonal matrices, and level sets of functions. And in Lectures 2 and 4, we worked with vector spaces of functions and of random variables, which are infinite-dimensional. This appendix provides the mathematical background.

- Sections A.1–A.4 are about topology, the mathematics of nearness and continuity. Its central notions are open sets, compactness and connectedness. Topology studies the properties of sets that do not change under continuous deformations.
- Sections A.5–A.10 are about differential topology, which studies manifolds, i.e., sets that look like $\mathbb{R}^k$ near each of their points, and differentiable maps between them. Its main tools are the inverse and implicit function theorems. The matrix groups of Lectures 1 and 2, such as the rotations, are manifolds as well, and they are Lie groups. Morse theory connects the critical points of a function, classified by the eigenvalues of its Hessian as in Lecture 3, with the shape of the space on which it is defined. Finally, differential forms, which are built from the tensors of Lecture 1, unify the integral theorems of Lecture 3 in one theorem of Stokes, and they detect holes in a space.
- Section A.11 collects some places where these ideas appear in machine learning.
- Sections A.12–A.15 are about functional analysis, i.e., linear algebra and analysis in infinite-dimensional vector spaces, such as spaces of functions. Its central notions are completeness, orthogonal projections and the spectral theorem for compact operators. They prove the existence theorem for differential equations of Lecture 3 and the Radon–Nikodym theorem of Appendix B, and they lead to Fourier series, to expansions of random functions, and to the theory of kernel methods. Section A.16 returns to machine learning.

We use the notation of Lecture 3: vectors in $\mathbb{R}^n$ are bold, and $\|\cdot\|$ is the Euclidean norm unless we say otherwise. Elements of general vector spaces, such as functions, are written in normal font.

A.1 Metric Spaces and Open Sets

Definition A.1.1 (Metric space)**.** A metric on a set X is a function $d : X \times X \to \mathbb{R}$ such that for all $x, y, z \in X$:

1. $d(x, y) \geq 0$, and $d(x, y) = 0$ if and only if $x = y$;
2. $d(x, y) = d(y, x)$;
3. $d(x, z) \leq d(x, y) + d(y, z)$ (triangle inequality).

A set with a metric is a metric space.

Example A.1.2.
1. Every norm on a vector space defines a metric $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x}-\mathbf{y}\|$ (Lecture 2). In particular, $\mathbb{R}^n$ with the ℓ_1, ℓ_2 or ℓ_∞ norm is a metric space.
2. On the space $C[a, b]$ of continuous functions on $[a, b]$, $d(f, g) = \max_{x\in[a,b]} |f(x) - g(x)|$ is a metric.
3. Every subset of a metric space is a metric space with the same metric. For example, the unit sphere $S^{n-1} = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\| = 1\}$ is a metric space with the Euclidean distance.
4. On any set, the discrete metric $d(x, y) = 1$ for $x \neq y$ (and $d(x, x) = 0$) is a metric.

Definition A.1.3 (Open and closed sets)**.** Let (X, d) be a metric space. The open ball with center x and radius $r > 0$ is $B(x, r) = \{y \in X : d(x, y) < r\}$. A set $U \subseteq X$ is open if for every $x \in U$ there is an $r > 0$ with $B(x, r) \subseteq U$. A set $C \subseteq X$ is closed if its complement $X \setminus C$ is open.

For $X = \mathbb{R}^n$, this is the definition of open sets in Lecture 3. Closed is not the opposite of open: $\mathbb{R}^n$ and $\emptyset$ are both open and closed, and the half-open interval $[0, 1)$ is neither open nor closed in $\mathbb{R}$.

Proposition A.1.4. *In a metric space X:*

1. *every open ball is open;*
2. *every union of open sets is open, and every intersection of finitely many open sets is open;*
3. *every intersection of closed sets is closed, and every union of finitely many closed sets is closed.*

Proof. (1) If $y \in B(x, r)$, let $s = r - d(x, y) > 0$. For $z \in B(y, s)$, the triangle inequality gives $d(x, z) \leq d(x, y) + d(y, z) < d(x, y) + s = r$, so $B(y, s) \subseteq B(x, r)$.

(2) If x lies in a union of open sets, it lies in one of them, U, and some ball $B(x, r) \subseteq U$ lies in the union. If $x \in U_1 \cap \cdots \cap U_m$ with $B(x, r_i) \subseteq U_i$, then $B(x, \min_i r_i)$ lies in the intersection.

(3) Take complements in (2). □

Infinite intersections of open sets need not be open: $\bigcap_{k\geq1}(-\frac{1}{k}, \frac{1}{k}) = \{0\}$. For a set $A \subseteq X$, the interior $\operatorname{int} A$ is the union of all open sets contained in A (the largest open subset of A), the closure $\overline{A}$ is the intersection of all closed sets containing A (the smallest closed set containing A), and the boundary is $\partial A = \overline{A} \setminus \operatorname{int} A$. For example, the closed ball $\{\mathbf{x} : \|\mathbf{x}\| \leq 1\}$ in $\mathbb{R}^n$ has the open ball as interior and the sphere S^{n-1} as boundary.

Definition A.1.5 (Convergence)**.** A sequence (x_k) in a metric space converges to x if $d(x_k, x) \to 0$ as $k \to \infty$. It is a Cauchy sequence if for every $\varepsilon > 0$ there is an N with $d(x_k, x_l) < \varepsilon$ for all $k, l \geq N$.

Closed sets are exactly the sets that cannot be left by taking limits.

Proposition A.1.6. *A subset C of a metric space X is closed if and only if the limit of every convergent sequence of points of C lies in C.*

Proof. Let C be closed, and let $x_k \in C$ converge to x. If $x \notin C$, then some ball $B(x, r)$ lies in the open set $X \setminus C$; but $x_k \in B(x, r)$ for large k, a contradiction. Conversely, suppose that C is not closed. Then $X \setminus C$ is not open, so there is an $x \notin C$ such that every ball $B(x, \frac{1}{k})$ contains a point $x_k \in C$. Then $x_k \to x$, and $x \notin C$. □

A.2 Continuity and Topological Spaces

Continuity can be described with open sets alone, without mentioning distances.

Proposition A.2.1. *Let $f : X \to Y$ be a map between metric spaces. Then f is continuous (at every point, in the ε-δ sense of Lecture 3) if and only if for every open set $V \subseteq Y$, the preimage $f^{-1}(V) = \{x \in X : f(x) \in V\}$ is open.*

Proof. Let f be continuous, $V \subseteq Y$ open, and $x \in f^{-1}(V)$. There is an $\varepsilon > 0$ with $B(f(x), \varepsilon) \subseteq V$, and by continuity a $\delta > 0$ such that $d(x, x') < \delta$ implies $d(f(x), f(x')) < \varepsilon$, i.e., $B(x, \delta) \subseteq f^{-1}(V)$. Conversely, if preimages of open sets are open, then for $x \in X$ and $\varepsilon > 0$, the set $f^{-1}(B(f(x), \varepsilon))$ is open and contains x, so it contains a ball $B(x, \delta)$; this is the ε-δ condition. □

Taking complements, f is continuous if and only if preimages of closed sets are closed.

Corollary A.2.2. *Let $f : \mathbb{R}^n \to \mathbb{R}$ be continuous and $c \in \mathbb{R}$. Then the set $\{f < c\}$ is open, and the sublevel set $\{f \leq c\}$ and the level set $\{f = c\}$ are closed. The same holds for intersections of finitely many such sets.*

Proof. These are the preimages of the open set $(-\infty, c)$ and the closed sets $(-\infty, c]$ and $\{c\}$. □

So the feasible sets of the constrained problems in Lecture 3, such as $\{\mathbf{x} : g_1(\mathbf{x}) \leq 0, \ldots, g_m(\mathbf{x}) \leq 0\}$ for continuous g_i, are closed. Similarly, the set $\mathrm{GL}(n, \mathbb{R})$ of invertible $n \times n$ matrices is open in $\mathbb{R}^{n \times n}$, as the preimage of the open set $\mathbb{R} \setminus \{0\}$ under the continuous function det (a polynomial in the entries).

Proposition A.2.1 suggests an abstraction: forget the metric, and only remember which sets are open.

Definition A.2.3 (Topological space)**.** A topology on a set X is a collection $\mathcal{T}$ of subsets of X, called open sets, such that $\emptyset, X \in \mathcal{T}$, every union of sets in $\mathcal{T}$ lies in $\mathcal{T}$, and every intersection of finitely many sets in $\mathcal{T}$ lies in $\mathcal{T}$. A map $f : X \to Y$ between topological spaces is continuous if preimages of open sets are open. A homeomorphism is a continuous bijection whose inverse is also continuous; if one exists, X and Y are homeomorphic.

By Proposition A.1.4, the open sets of a metric space form a topology. Two metrics can give the same topology: we will see in Theorem A.3.10 that all norms on $\mathbb{R}^n$ do. Homeomorphic spaces are the same from the point of view of topology: a homeomorphism matches up their open sets, convergent sequences, continuous functions, and so on.

Example A.2.4.

1. The sigmoid function is a homeomorphism from $\mathbb{R}$ to $(0, 1)$, with the continuous inverse $p \mapsto \log \frac{p}{1-p}$ (the logit). Similarly, tanh is a homeomorphism $\mathbb{R} \to (-1, 1)$. So bounded and unbounded sets can be homeomorphic: lengths and distances are not topological properties.
2. The open ball $B(\mathbf{0}, 1) \subseteq \mathbb{R}^n$ is homeomorphic to $\mathbb{R}^n$, via $\mathbf{x} \mapsto \frac{\mathbf{x}}{1-\|\mathbf{x}\|}$, with the inverse $\mathbf{y} \mapsto \frac{\mathbf{y}}{1+\|\mathbf{y}\|}$.
3. A square and a disk in $\mathbb{R}^2$ are homeomorphic (rescale each ray from the center), and so are a circle and the boundary of a square: corners are not a topological property. But a circle and an interval are not homeomorphic (Example A.4.3).

A.3 Compactness

The real numbers have no gaps. This is the completeness axiom, which we take as one of the defining properties of $\mathbb{R}$: every nonempty set of real numbers that is bounded above has a least upper bound (supremum). It implies that every bounded monotone sequence of real numbers converges (to the supremum or the infimum of its values).

Theorem A.3.1 (Bolzano–Weierstrass)**.** *Every bounded sequence in $\mathbb{R}^n$ has a convergent subsequence.*

Proof. First let $n = 1$. Call an index k a peak if $x_k \geq x_l$ for all $l > k$. If there are infinitely many peaks $k_1 < k_2 < \cdots$, then $x_{k_1} \geq x_{k_2} \geq \cdots$ is a monotone subsequence. Otherwise, there is an index k_1 after which there are no peaks; then k_1 is not a peak, so $x_{k_2} > x_{k_1}$ for some $k_2 > k_1$; k_2 is not a peak either, and so on. This gives an increasing subsequence. In both cases, the monotone subsequence is bounded, so it converges. For $n > 1$, choose a subsequence whose first coordinates converge, then a subsequence of it whose second coordinates converge, and so on; after n steps all coordinates converge. □

Corollary A.3.2. *$\mathbb{R}^n$ is complete: every Cauchy sequence in $\mathbb{R}^n$ converges.*

Proof. A Cauchy sequence $(\mathbf{x}_k)$ is bounded (all terms after some N lie within distance 1 of $\mathbf{x}_N$), so by Theorem A.3.1 a subsequence converges to some $\mathbf{x}$. For $\varepsilon > 0$, choose N such that $\|\mathbf{x}_k - \mathbf{x}_l\| < \varepsilon$ for $k, l \geq N$, and an element $\mathbf{x}_l$ of the subsequence with $l \geq N$ and $\|\mathbf{x}_l - \mathbf{x}\| < \varepsilon$. Then $\|\mathbf{x}_k - \mathbf{x}\| < 2\varepsilon$ for all $k \geq N$. □

Definition A.3.3 (Compact set)**.** A subset K of a metric space is compact if every sequence in K has a subsequence that converges to a point of K.

(In general topological spaces, one defines compactness with open covers: K is compact if every collection of open sets that covers K contains finitely many sets that already cover K. For metric spaces, the two definitions agree; we do not prove this.)

Theorem A.3.4 (Heine–Borel)**.** *A subset of $\mathbb{R}^n$ is compact if and only if it is closed and bounded.*

Proof. Let K be closed and bounded, and let $(\mathbf{x}_k)$ be a sequence in K. By Theorem A.3.1, a subsequence converges, and by Proposition A.1.6, its limit lies in K. Conversely, if K is unbounded, there are $\mathbf{x}_k \in K$ with $\|\mathbf{x}_k\| \geq k$, and no subsequence converges, since convergent sequences are bounded. If K is not closed, there is a sequence in K that converges to a point outside K (Proposition A.1.6), and all its subsequences converge to the same point, so none converges to a point of K. □

For example, spheres, closed balls (for any norm), and the set $\mathrm{O}(n)$ of orthogonal matrices are compact. $\mathrm{O}(n)$ is closed, as the preimage of $\{I\}$ under the continuous map $Q \mapsto Q^\top Q$, and bounded, since $\|Q\|_F^2 = \operatorname{tr}(Q^\top Q) = n$. The hyperbola $\{xy = 1\}$ is closed but unbounded, and the open ball is bounded but not closed; neither is compact.

Theorem A.3.5 (Extreme value theorem)**.** *Let K be a nonempty compact set, and let $f : K \to \mathbb{R}$ be continuous. Then f attains a minimum and a maximum on K.*

Proof. Let $m = \inf\{f(\mathbf{x}) : \mathbf{x} \in K\}$, where we allow $m = -\infty$. Choose $\mathbf{x}_k \in K$ with $f(\mathbf{x}_k) \to m$. A subsequence converges to some $\mathbf{x}^* \in K$, and by continuity, $f(\mathbf{x}^*) = m$. So m is finite, and it is the minimum. For the maximum, apply this to $-f$. □

This justifies the claims in Lecture 3 that $f(x, y) = x + y$ and the Rayleigh quotient $\mathbf{x}^\top A\mathbf{x}$ attain their maxima on the unit sphere (Examples 3.12.2 and 3.12.3). On $\mathbb{R}^n$, which is not compact, a minimum need not exist, but the following condition is often enough.

Proposition A.3.6 (Coercive functions)**.** *Let $f : \mathbb{R}^n \to \mathbb{R}$ be continuous and coercive, i.e., $f(\mathbf{x}) \to \infty$ as $\|\mathbf{x}\| \to \infty$. Then f has a global minimum.*

Proof. Fix any $\mathbf{x}_0$. The sublevel set $S = \{f \le f(\mathbf{x}_0)\}$ is closed (Corollary A.2.2) and bounded (by coercivity), so it is compact and nonempty, and f has a minimum $\mathbf{x}^*$ on S. Points outside S have values larger than $f(\mathbf{x}_0) \ge f(\mathbf{x}^*)$, so $\mathbf{x}^*$ is a global minimum. □

Example A.3.7. A loss $L(\mathbf{w}) \ge 0$ plus a ridge penalty, $L(\mathbf{w}) + \lambda\|\mathbf{w}\|^2$ with $\lambda > 0$, is coercive, so it has a global minimum whenever L is continuous. Without regularization, a minimum need not exist: for logistic regression with the single training example $x = 1$, $y = 1$, the loss $\log(1 + e^{-w})$ decreases towards 0 as $w \to \infty$, but never reaches it. The same happens for every linearly separable data set, where scaling up a separating weight vector always decreases the loss.

On a compact set, continuity is automatically uniform.

Proposition A.3.8 (Uniform continuity)**.** *Let $K \subseteq \mathbb{R}^n$ be compact, and let $f : K \to \mathbb{R}$ be continuous. Then f is uniformly continuous: for every $\varepsilon > 0$, there is a $\delta > 0$ such that $|f(\mathbf{x}) - f(\mathbf{y})| < \varepsilon$ for all $\mathbf{x}, \mathbf{y} \in K$ with $\|\mathbf{x} - \mathbf{y}\| < \delta$.*

Proof. Otherwise, there are an $\varepsilon > 0$ and points $\mathbf{x}_k, \mathbf{y}_k \in K$ with $\|\mathbf{x}_k - \mathbf{y}_k\| < \frac{1}{k}$ and $|f(\mathbf{x}_k) - f(\mathbf{y}_k)| \ge \varepsilon$ for every k. A subsequence $(\mathbf{x}_{k_l})$ converges to some $\mathbf{x} \in K$, and then $\mathbf{y}_{k_l} \to \mathbf{x}$ as well. By continuity, $f(\mathbf{x}_{k_l}) - f(\mathbf{y}_{k_l}) \to f(\mathbf{x}) - f(\mathbf{x}) = 0$, which contradicts $|f(\mathbf{x}_{k_l}) - f(\mathbf{y}_{k_l})| \ge \varepsilon$. □

The point is that δ depends only on ε, not on the points. On a set that is not compact, this can fail: $f(x) = x^2$ is continuous on $\mathbb{R}$, but $f(x + \delta) - f(x) = 2x\delta + \delta^2$ is large for large x, however small $\delta > 0$ is.

Corollary A.3.9 (Integrals depending on a parameter)**.** *Let $h(x, \theta)$ be continuous for $x \in [a, b]$ and θ in an open interval I. Then $H(\theta) = \int_a^b h(x, \theta)\, dx$ is continuous on I.*

Proof. Fix $\theta_0 \in I$, and choose $r > 0$ with $[\theta_0 - r, \theta_0 + r] \subseteq I$. The set $K = [a, b] \times [\theta_0 - r, \theta_0 + r]$ is closed and bounded, hence compact, so h is uniformly continuous on K. For $\varepsilon > 0$, choose $\delta \le r$ as in Proposition A.3.8. If $|\theta - \theta_0| < \delta$, then (x, θ) and (x, θ_0) have the distance $|\theta - \theta_0| < \delta$ for every x, so

$$|H(\theta) - H(\theta_0)| \le \int_a^b |h(x, \theta) - h(x, \theta_0)|\, dx \le (b - a)\varepsilon. \qquad \square$$

Applied to $h = \frac{\partial f}{\partial \theta}$, this fills the gap in the proof of the Leibniz rule (Proposition 3.14.6).

Compactness also explains why all norms on $\mathbb{R}^n$ lead to the same notions of convergence and continuity, as claimed in Lecture 3.

Theorem A.3.10 (Equivalence of norms)**.** *For every norm N on $\mathbb{R}^n$, there are constants $0 < c \le C$ with $c\|\mathbf{x}\| \le N(\mathbf{x}) \le C\|\mathbf{x}\|$ for all $\mathbf{x}$. Consequently, all norms on $\mathbb{R}^n$ define the same open sets.*

Proof. Writing $\mathbf{x} = \sum_i x_i \mathbf{e}_i$, the triangle inequality and the Cauchy–Schwarz inequality give

$$N(\mathbf{x}) \le \sum_i |x_i| N(\mathbf{e}_i) \le \Big(\sum_i N(\mathbf{e}_i)^2 \Big)^{1/2} \|\mathbf{x}\| = C\|\mathbf{x}\|.$$

In particular, $|N(\mathbf{x}) - N(\mathbf{y})| \le N(\mathbf{x} - \mathbf{y}) \le C\|\mathbf{x} - \mathbf{y}\|$, so N is continuous. By Theorem A.3.5, N attains a minimum c on the compact unit sphere S^{n-1}, and $c > 0$ since N is positive on S^{n-1}. For $\mathbf{x} \ne \mathbf{0}$, $N(\mathbf{x}) = \|\mathbf{x}\|\, N\big(\frac{\mathbf{x}}{\|\mathbf{x}\|}\big) \ge c\|\mathbf{x}\|$. Finally, the inequalities show that every N-ball around a point contains a Euclidean ball around it and vice versa, so the open sets are the same. □

Remark A.3.11 (Infinite dimensions). In infinite-dimensional spaces, closed and bounded sets need not be compact, and different norms can give different topologies. For example, in the space ℓ^2 of square-summable sequences (Section A.12), the unit sequences $e_1, e_2, \ldots$ lie in the closed unit ball, but $\|e_j - e_k\| = \sqrt{2}$ for $j \neq k$, so no subsequence converges. This is one reason why analysis in function spaces, such as the spaces of functions in Lecture 2, is harder than in $\mathbb{R}^n$ (Sections A.12–A.15).

A.4 Connectedness

Definition A.4.1 (Connected sets)**.** A subset X of a metric space is disconnected if $X \subseteq U \cup V$ for two open sets U and V such that $X \cap U$ and $X \cap V$ are nonempty and disjoint. Otherwise, X is connected. It is path-connected if any two points $x, y \in X$ can be joined by a path, i.e., a continuous map $\gamma : [0,1] \to X$ with $\gamma(0) = x$ and $\gamma(1) = y$.

Proposition A.4.2. *1. Every interval of $\mathbb{R}$ is connected.*

2. If X is connected and f is continuous on X, then $f(X)$ is connected.

3. Every path-connected set is connected.

Proof. (1) Suppose an interval I is covered by open sets U and V as in the definition, with $a \in I \cap U$ and $b \in I \cap V$; by symmetry, we may assume $a < b$. Then $[a,b] \subseteq I$. Let $s = \sup\{t \in [a,b] : [a,t] \subseteq U\}$; then $[a,t] \subseteq U$ for every $t < s$, and $s \in [a,b] \subseteq I$, so s lies in U or in V. If $s \in V$, then since V is open, $(s-\varepsilon, s] \subseteq V$ for some $\varepsilon > 0$; but points of $(s-\varepsilon, s)$ also lie in U, which contradicts $I \cap U \cap V = \emptyset$. If $s \in U$, then $s \neq b$ (since $b \in V$), so $s < b$, and since U is open, $[s, s+\varepsilon] \subseteq U$ for some small $\varepsilon > 0$ with $s + \varepsilon \leq b$. Then $[a, s+\varepsilon] \subseteq U$, which contradicts the definition of s.

(2) If $f(X)$ were covered by open sets U and V as in the definition, then the preimages $f^{-1}(U)$ and $f^{-1}(V)$ would disconnect X (they are open by continuity, in the sense that they are of the form $X \cap U'$ for open sets U').

(3) Suppose that open sets U, V disconnect a path-connected set X, with $x \in U$ and $y \in V$. Then the preimages of U and V under a path from x to y would disconnect the interval $[0,1]$, which contradicts (1). □

Together, (1) and (2) give the intermediate value theorem: a continuous function $f : [a,b] \to \mathbb{R}$ takes every value between $f(a)$ and $f(b)$, since its image is a connected subset of $\mathbb{R}$, i.e., an interval.

Example A.4.3.

1. $\mathbb{R}^n$, balls, and the sphere S^{n-1} for $n \geq 2$ are path-connected, hence connected.

2. The hyperbola $\{xy = 1\}$ is not connected: it lies in the union of the open half-planes $\{x > 0\}$ and $\{x < 0\}$, and meets both.

3. $\mathrm{GL}(n, \mathbb{R})$ is not connected: it is the union of the disjoint open sets $\{\det > 0\}$ and $\{\det < 0\}$. In particular, an invertible matrix with $\det < 0$ (a reflection) cannot be continuously deformed into the identity through invertible matrices. Similarly, $\mathrm{O}(n)$ consists of the two pieces $\det Q = 1$ (the rotations, Lecture 2) and $\det Q = -1$; one can show that the rotations form a connected set.

4. A circle and an interval are not homeomorphic. If we remove a point from a circle, the rest is still connected (it is an arc); if we remove an interior point from an interval, the rest is disconnected. A homeomorphism would map a point and its complement to a point and its complement, and it preserves connectedness by Proposition A.4.2(2).

A.5 The Inverse Function Theorem

From now on, we combine topology with derivatives. A map is smooth if all its partial derivatives of all orders exist and are continuous, and a diffeomorphism is a smooth bijection with a smooth inverse. We first fill a gap from Lecture 3, using a vector-valued version of the mean value theorem.

Lemma A.5.1 (Mean value inequality)**.** *Let $\mathbf{g} : U \to \mathbb{R}^m$ be continuously differentiable on an open set $U \subseteq \mathbb{R}^n$ that contains the segment between $\mathbf{x}$ and $\mathbf{y}$. Then*

$$\|\mathbf{g}(\mathbf{y}) - \mathbf{g}(\mathbf{x})\| \le \sup_{\mathbf{z}\in[\mathbf{x},\mathbf{y}]} \|D\mathbf{g}(\mathbf{z})\|_2 \, \|\mathbf{y} - \mathbf{x}\|,$$

where $\|\cdot\|_2$ is the spectral norm of Lecture 2.

Proof. Let $\mathbf{u} = \mathbf{g}(\mathbf{y}) - \mathbf{g}(\mathbf{x})$, and assume $\mathbf{u} \neq \mathbf{0}$. The function $\psi(t) = \mathbf{u}^\top \mathbf{g}(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))$ has $\psi(1) - \psi(0) = \|\mathbf{u}\|^2$. By the mean value theorem and the chain rule, $\psi(1) - \psi(0) = \psi'(\tau) = \mathbf{u}^\top D\mathbf{g}(\mathbf{z})(\mathbf{y} - \mathbf{x})$ with $\mathbf{z} = \mathbf{x} + \tau(\mathbf{y} - \mathbf{x})$. By the Cauchy–Schwarz inequality, $\|\mathbf{u}\|^2 \le \|\mathbf{u}\| \, \|D\mathbf{g}(\mathbf{z})\|_2 \, \|\mathbf{y} - \mathbf{x}\|$; divide by $\|\mathbf{u}\|$. □

Proposition A.5.2 (Theorem 3.3.6)**.** *If all partial derivatives of $f : U \to \mathbb{R}$ exist in a ball around $\mathbf{x}$ and are continuous at $\mathbf{x}$, then f is differentiable at $\mathbf{x}$.*

Proof. For small $\mathbf{h}$, let $\mathbf{z}_0 = \mathbf{x}$ and $\mathbf{z}_j = \mathbf{x} + h_1\mathbf{e}_1 + \cdots + h_j\mathbf{e}_j$, so that $\mathbf{z}_n = \mathbf{x} + \mathbf{h}$. The points $\mathbf{z}_{j-1}$ and $\mathbf{z}_j$ differ only in the j-th coordinate, so by the mean value theorem in one variable, $f(\mathbf{z}_j) - f(\mathbf{z}_{j-1}) = \frac{\partial f}{\partial x_j}(\boldsymbol{\xi}_j) h_j$ for a point $\boldsymbol{\xi}_j$ between them. Adding these equations,

$$\big|f(\mathbf{x} + \mathbf{h}) - f(\mathbf{x}) - \nabla f(\mathbf{x})^\top \mathbf{h}\big| = \Big|\sum_{j=1}^{n} \Big(\frac{\partial f}{\partial x_j}(\boldsymbol{\xi}_j) - \frac{\partial f}{\partial x_j}(\mathbf{x})\Big) h_j\Big| \le \sum_{j=1}^{n} \Big|\frac{\partial f}{\partial x_j}(\boldsymbol{\xi}_j) - \frac{\partial f}{\partial x_j}(\mathbf{x})\Big| \, \|\mathbf{h}\|.$$

As $\mathbf{h} \to \mathbf{0}$, all $\boldsymbol{\xi}_j \to \mathbf{x}$, so the sum tends to 0 by the continuity of the partial derivatives at $\mathbf{x}$. □

The key tool of this section is a simple but powerful theorem about solving equations by iteration.

Theorem A.5.3 (Banach fixed-point theorem)**.** *Let $C \subseteq \mathbb{R}^n$ be closed and nonempty, and let $T : C \to C$ be a contraction, i.e., $\|T(\mathbf{x}) - T(\mathbf{y})\| \le q\|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in C$, with a constant $q < 1$. Then T has exactly one fixed point $\mathbf{x}^* = T(\mathbf{x}^*)$. For every $\mathbf{x}_0 \in C$, the iterates $\mathbf{x}_{k+1} = T(\mathbf{x}_k)$ converge to $\mathbf{x}^*$, and*

$$\|\mathbf{x}_k - \mathbf{x}^*\| \le \frac{q^k}{1 - q}\|\mathbf{x}_1 - \mathbf{x}_0\|.$$

Proof. By induction, $\|\mathbf{x}_{j+1} - \mathbf{x}_j\| \le q^j \|\mathbf{x}_1 - \mathbf{x}_0\|$. For $l > k$, the triangle inequality and the geometric series give

$$\|\mathbf{x}_l - \mathbf{x}_k\| \le \sum_{j=k}^{l-1} q^j \|\mathbf{x}_1 - \mathbf{x}_0\| \le \frac{q^k}{1 - q}\|\mathbf{x}_1 - \mathbf{x}_0\|.$$

So $(\mathbf{x}_k)$ is a Cauchy sequence, which converges to some $\mathbf{x}^*$ by Corollary A.3.2, and $\mathbf{x}^* \in C$ since C is closed. Since T is continuous, $T(\mathbf{x}^*) = \lim T(\mathbf{x}_k) = \lim \mathbf{x}_{k+1} = \mathbf{x}^*$. Letting $l \to \infty$ in the inequality gives the error bound. If $\mathbf{y}^*$ is another fixed point, then $\|\mathbf{x}^* - \mathbf{y}^*\| = \|T(\mathbf{x}^*) - T(\mathbf{y}^*)\| \le q\|\mathbf{x}^* - \mathbf{y}^*\|$, so $\mathbf{x}^* = \mathbf{y}^*$. □

Example A.5.4. The function $T(x) = \cos x$ maps $[0, 1]$ into $[\cos 1, 1] \subseteq [0, 1]$, and $|T'(x)| = |\sin x| \le \sin 1 \approx 0.841$ there, so it is a contraction with $q = \sin 1$ (by the mean value theorem). So the equation $\cos x = x$ has exactly one solution in $[0, 1]$, $x^* = 0.739085\ldots$, and pressing the cosine key of a calculator repeatedly finds it. Starting at $x_0 = 1$, the error bound guarantees an accuracy of 10^{-6} after 87 steps; in fact, 32 steps suffice. Gradient descent on a quadratic function with $\rho < 1$ (Theorem 3.10.7) is another example of a contraction.

Corollary A.5.5 (Invertible residual layers)**.** *Let $\mathbf{g} : \mathbb{R}^n \to \mathbb{R}^n$ satisfy $\|\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{y})\| \le q\|\mathbf{x} - \mathbf{y}\|$ with $q < 1$. Then the residual layer $\mathbf{f}(\mathbf{x}) = \mathbf{x} + \mathbf{g}(\mathbf{x})$ is a homeomorphism of $\mathbb{R}^n$, and $\mathbf{f}^{-1}(\mathbf{y})$ is the limit of the iteration $\mathbf{x}_{k+1} = \mathbf{y} - \mathbf{g}(\mathbf{x}_k)$.*

Proof. For fixed $\mathbf{y}$, the map $T(\mathbf{x}) = \mathbf{y} - \mathbf{g}(\mathbf{x})$ is a contraction of $\mathbb{R}^n$, and $\mathbf{f}(\mathbf{x}) = \mathbf{y}$ if and only if $\mathbf{x} = T(\mathbf{x})$. By Theorem A.5.3, there is exactly one solution, so $\mathbf{f}$ is bijective. It is continuous, and so is its inverse: $\|\mathbf{x} - \mathbf{x}'\| \le \|\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{x}')\| + \|\mathbf{g}(\mathbf{x}) - \mathbf{g}(\mathbf{x}')\| \le \|\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{x}')\| + q\|\mathbf{x} - \mathbf{x}'\|$, so $\|\mathbf{f}^{-1}(\mathbf{y}) - \mathbf{f}^{-1}(\mathbf{y}')\| \le \frac{1}{1-q}\|\mathbf{y} - \mathbf{y}'\|$. □

Invertible neural networks used in normalizing flows (Remark 3.4.8) are built from such layers: constraining the Lipschitz constant of each residual branch (e.g., by normalizing the spectral norms of its weight matrices) makes the whole network invertible. A diffeomorphism must have invertible Jacobians everywhere, by the chain rule applied to $\mathbf{f}^{-1} \circ \mathbf{f} = \mathrm{id}$. The inverse function theorem is a local converse.

Theorem A.5.6 (Inverse function theorem)**.** *Let $\mathbf{f} : U \to \mathbb{R}^n$ be continuously differentiable on an open set $U \subseteq \mathbb{R}^n$, and let $D\mathbf{f}(\mathbf{a})$ be invertible. Then there are open sets $V \ni \mathbf{a}$ and $W \ni \mathbf{f}(\mathbf{a})$ such that $\mathbf{f}$ maps V bijectively onto W, and the inverse $\mathbf{f}^{-1} : W \to V$ is continuously differentiable, with*

$$D(\mathbf{f}^{-1})(\mathbf{f}(\mathbf{x})) = D\mathbf{f}(\mathbf{x})^{-1} \qquad \text{for } \mathbf{x} \in V.$$

If $\mathbf{f}$ is smooth, so is $\mathbf{f}^{-1}$.

Sketch of proof. Replacing $\mathbf{f}(\mathbf{x})$ by $D\mathbf{f}(\mathbf{a})^{-1}(\mathbf{f}(\mathbf{a} + \mathbf{x}) - \mathbf{f}(\mathbf{a}))$, we may assume $\mathbf{a} = \mathbf{0}$, $\mathbf{f}(\mathbf{0}) = \mathbf{0}$ and $D\mathbf{f}(\mathbf{0}) = I$. Then $\mathbf{g}(\mathbf{x}) = \mathbf{x} - \mathbf{f}(\mathbf{x})$ has $D\mathbf{g}(\mathbf{0}) = 0$, so by continuity there is an $r > 0$ such that the closed ball $\overline{B}_r = \{\|\mathbf{x}\| \le r\}$ lies in U and $\|D\mathbf{g}(\mathbf{x})\|_2 \le \frac{1}{2}$ on it. By Lemma A.5.1, $\mathbf{g}$ is a contraction on $\overline{B}_r$ with $q = \frac{1}{2}$, and $\|\mathbf{g}(\mathbf{x})\| \le \frac{1}{2}\|\mathbf{x}\|$. For $\|\mathbf{y}\| < \frac{r}{2}$, the map $T(\mathbf{x}) = \mathbf{y} + \mathbf{g}(\mathbf{x})$ maps $\overline{B}_r$ into itself, since $\|T(\mathbf{x})\| \le \frac{r}{2} + \frac{r}{2}$, and it is a contraction. Its unique fixed point is the unique $\mathbf{x} \in \overline{B}_r$ with $\mathbf{f}(\mathbf{x}) = \mathbf{y}$, and it satisfies $\|\mathbf{x}\| \le \|\mathbf{y}\| + \frac{1}{2}\|\mathbf{x}\|$, i.e., $\|\mathbf{x}\| \le 2\|\mathbf{y}\| < r$. So with $W = B(\mathbf{0}, \frac{r}{2})$ and $V = B(\mathbf{0}, r) \cap \mathbf{f}^{-1}(W)$, $\mathbf{f} : V \to W$ is bijective, and as in Corollary A.5.5, the inverse is continuous. One can show that it is also differentiable; then the chain rule applied to $\mathbf{f}^{-1}(\mathbf{f}(\mathbf{x})) = \mathbf{x}$ gives $D(\mathbf{f}^{-1})(\mathbf{f}(\mathbf{x}))\, D\mathbf{f}(\mathbf{x}) = I$. □

Example A.5.7.

1. Polar coordinates $\mathbf{f}(r, \varphi) = (r\cos\varphi, r\sin\varphi)$ have $\det D\mathbf{f} = r$ (Example 3.4.7). So near every point with $r > 0$, they have a smooth local inverse. But $\mathbf{f}$ is not injective, since $\mathbf{f}(r, \varphi + 2\pi) = \mathbf{f}(r, \varphi)$: the theorem is only local.
2. The map $F(x, y) = (x^2 - y^2, 2xy)$, i.e., $z \mapsto z^2$ for $z = x + iy$, has $\det DF = 4(x^2 + y^2)$. Near every point $z \neq 0$, it has a local inverse, a "branch of the square root". Globally, it is two-to-one, since $F(-z) = F(z)$, and at $z = 0$, it is not even locally injective.

A.6 The Implicit Function Theorem

When can an equation $F(\mathbf{x}, \mathbf{y}) = \mathbf{0}$ be solved for some of the variables? We write points of $\mathbb{R}^{k+m}$ as $(\mathbf{x}, \mathbf{y})$ with $\mathbf{x} \in \mathbb{R}^k$ and $\mathbf{y} \in \mathbb{R}^m$, and split the Jacobian of $F : \mathbb{R}^{k+m} \to \mathbb{R}^m$ into the $m \times k$ matrix $D_{\mathbf{x}}F$ and the square $m \times m$ matrix $D_{\mathbf{y}}F$ of partial derivatives with respect to $\mathbf{x}$ and $\mathbf{y}$.

Theorem A.6.1 (Implicit function theorem)**.** *Let $F : U \to \mathbb{R}^m$ be continuously differentiable on an open set $U \subseteq \mathbb{R}^{k+m}$, let $F(\mathbf{a}, \mathbf{b}) = \mathbf{0}$, and let $D_{\mathbf{y}}F(\mathbf{a}, \mathbf{b})$ be invertible. Then there are open sets $V \ni \mathbf{a}$ and $W \ni (\mathbf{a}, \mathbf{b})$ and a continuously differentiable function $\mathbf{g} : V \to \mathbb{R}^m$ with $\mathbf{g}(\mathbf{a}) = \mathbf{b}$, such that for $(\mathbf{x}, \mathbf{y}) \in W$,*

$$F(\mathbf{x}, \mathbf{y}) = \mathbf{0} \quad \Longleftrightarrow \quad \mathbf{x} \in V \text{ and } \mathbf{y} = \mathbf{g}(\mathbf{x}).$$

Moreover, $D\mathbf{g}(\mathbf{x}) = -D_{\mathbf{y}}F(\mathbf{x}, \mathbf{g}(\mathbf{x}))^{-1}\, D_{\mathbf{x}}F(\mathbf{x}, \mathbf{g}(\mathbf{x}))$.

Sketch of proof. Let $\Phi(\mathbf{x},\mathbf{y}) = (\mathbf{x}, F(\mathbf{x},\mathbf{y}))$. Its Jacobian $\begin{pmatrix} I & 0 \\ D_{\mathbf{x}}F & D_{\mathbf{y}}F \end{pmatrix}$ is invertible at $(\mathbf{a},\mathbf{b})$: the equations $\mathbf{h} = \mathbf{u}$, $D_{\mathbf{x}}F\,\mathbf{h} + D_{\mathbf{y}}F\,\mathbf{k} = \mathbf{v}$ have the unique solution $\mathbf{h} = \mathbf{u}$, $\mathbf{k} = D_{\mathbf{y}}F^{-1}(\mathbf{v} - D_{\mathbf{x}}F\,\mathbf{u})$. By Theorem A.5.6, Φ has a local inverse near $\Phi(\mathbf{a},\mathbf{b}) = (\mathbf{a},\mathbf{0})$. Since Φ does not change $\mathbf{x}$, neither does its inverse, which therefore has the form $(\mathbf{x},\mathbf{z}) \mapsto (\mathbf{x}, G(\mathbf{x},\mathbf{z}))$. Near $(\mathbf{a},\mathbf{b})$, $F(\mathbf{x},\mathbf{y}) = \mathbf{0}$ means $\Phi(\mathbf{x},\mathbf{y}) = (\mathbf{x},\mathbf{0})$, i.e., $\mathbf{y} = G(\mathbf{x},\mathbf{0}) =: \mathbf{g}(\mathbf{x})$. Differentiating $F(\mathbf{x},\mathbf{g}(\mathbf{x})) = \mathbf{0}$ with the chain rule gives $D_{\mathbf{x}}F + D_{\mathbf{y}}F\,D\mathbf{g} = 0$. □

Example A.6.2.

1. The unit circle $F(x,y) = x^2 + y^2 - 1 = 0$ has $\frac{\partial F}{\partial y} = 2y$. Near every point with $y \neq 0$, the circle is the graph of a function of x ($y = \pm\sqrt{1-x^2}$), with $\frac{dy}{dx} = -\frac{2x}{2y}$. Near $(\pm 1, 0)$, it is not, but it is the graph of a function of y, since $\frac{\partial F}{\partial x} = 2x \neq 0$ there.
2. For the folium of Descartes $x^3 + y^3 = 3xy$ (Example 3.9.4), $\frac{\partial F}{\partial y} = 3y^2 - 3x$ vanishes on the curve only at $(0,0)$, where the curve crosses itself, and at $(2^{2/3}, 2^{1/3})$, where its tangent is vertical. Near every other point of the curve, y is a smooth function of x.

Remark A.6.3 (Implicit differentiation in machine learning). Sometimes a quantity is defined as the solution of an equation, $G(\mathbf{z},\boldsymbol{\theta}) = \mathbf{0}$, and we need its derivative with respect to $\boldsymbol{\theta}$. Examples are the minimizer $\mathbf{z}^*(\boldsymbol{\theta})$ of an inner optimization problem with hyperparameters $\boldsymbol{\theta}$ (here $G = \nabla_{\mathbf{z}}$ of the inner objective), and the fixed point $\mathbf{z} = \mathbf{h}(\mathbf{z},\boldsymbol{\theta})$ of an "implicit layer" in deep equilibrium models. By Theorem A.6.1, $D\mathbf{z}^*(\boldsymbol{\theta}) = -D_{\mathbf{z}}G^{-1}D_{\boldsymbol{\theta}}G$, so we do not need to backpropagate through all the iterations of the solver that computed $\mathbf{z}^*$. For example, the ridge regression solution $\mathbf{w}^*(\lambda)$ solves $G(\mathbf{w},\lambda) = X^\top(X\mathbf{w} - \mathbf{y}) + \lambda\mathbf{w} = \mathbf{0}$, so

$$\frac{d\mathbf{w}^*}{d\lambda} = -(X^\top X + \lambda I)^{-1}\mathbf{w}^*(\lambda),$$

which can be used to tune λ by gradient descent on a validation loss.

A.7 Submanifolds and Tangent Spaces

A k-dimensional manifold is a set that looks like $\mathbb{R}^k$ near each of its points: a curve looks like a line, and a surface like a plane. We consider manifolds that lie in $\mathbb{R}^n$ and are described by equations.

Definition A.7.1 (Submanifold)**.** A set $M \subseteq \mathbb{R}^n$ is a k-dimensional submanifold of $\mathbb{R}^n$ if for every $\mathbf{p} \in M$, there are an open set $U \ni \mathbf{p}$ and a smooth map $F : U \to \mathbb{R}^{n-k}$ such that

$$M \cap U = \{\mathbf{x} \in U : F(\mathbf{x}) = \mathbf{0}\} \qquad \text{and} \qquad \operatorname{rank} DF(\mathbf{p}) = n - k.$$

(For $k = n$, there is no equation: M is an open subset of $\mathbb{R}^n$.)

So locally, M is cut out by $n-k$ independent equations. In particular, if $F : U \to \mathbb{R}^m$ is smooth and $\mathbf{c} \in \mathbb{R}^m$ is a regular value of F, i.e., $\operatorname{rank} DF(\mathbf{x}) = m$ for every $\mathbf{x}$ with $F(\mathbf{x}) = \mathbf{c}$, then the level set $F^{-1}(\mathbf{c})$ is an $(n-m)$-dimensional submanifold (use $F - \mathbf{c}$ in the definition). The implicit function theorem shows that submanifolds really look like $\mathbb{R}^k$.

Proposition A.7.2 (Submanifolds are local graphs)**.** *Let M be a k-dimensional submanifold and $\mathbf{p} \in M$. After renumbering the coordinates of $\mathbb{R}^n = \mathbb{R}^k \times \mathbb{R}^{n-k}$, there are open sets $V \subseteq \mathbb{R}^k$ and $W \ni \mathbf{p}$ and a smooth map $\mathbf{g} : V \to \mathbb{R}^{n-k}$ such that $M \cap W = \{(\mathbf{x}, \mathbf{g}(\mathbf{x})) : \mathbf{x} \in V\}$. In particular, $\mathbf{x} \mapsto (\mathbf{x}, \mathbf{g}(\mathbf{x}))$ is a homeomorphism from the open set $V \subseteq \mathbb{R}^k$ onto $M \cap W$.*

Proof. The $(n-k) \times n$ matrix $DF(\mathbf{p})$ has rank $n-k$, so it has $n-k$ linearly independent columns (Lecture 1). Renumber the coordinates so that these are the last $n-k$ columns, and write $\mathbf{x} \in \mathbb{R}^n$ as $(\mathbf{x}', \mathbf{y})$. Then $D_{\mathbf{y}}F(\mathbf{p})$ is invertible, and Theorem A.6.1 gives $\mathbf{g}$. The inverse of $\mathbf{x}' \mapsto (\mathbf{x}', \mathbf{g}(\mathbf{x}'))$ is the projection to the first k coordinates, which is continuous. □

Example A.7.3. 1. The sphere $S^{n-1} = \{\mathbf{x} : \|\mathbf{x}\|^2 = 1\}$ is an $(n-1)$-dimensional submanifold, since $F(\mathbf{x}) = \|\mathbf{x}\|^2 - 1$ has $DF(\mathbf{x}) = 2\mathbf{x}^\top \neq \mathbf{0}$ on the sphere.

2. The graph $\{(\mathbf{x}, \mathbf{g}(\mathbf{x}))\}$ of a smooth $\mathbf{g} : \mathbb{R}^k \to \mathbb{R}^{n-k}$ is a k-dimensional submanifold: $F(\mathbf{x}, \mathbf{y}) = \mathbf{y} - \mathbf{g}(\mathbf{x})$ has $DF = (-D\mathbf{g},\, I)$, of rank $n-k$.

3. $\mathrm{GL}(n, \mathbb{R})$ is open in $\mathbb{R}^{n\times n}$, hence an n^2-dimensional submanifold.

4. The special linear group $\mathrm{SL}(n, \mathbb{R}) = \{X : \det X = 1\}$ is a submanifold of dimension $n^2 - 1$. By Jacobi's formula (Proposition 3.6.4), $D\det(X)H = \det(X)\,\mathrm{tr}(X^{-1}H)$, which for $\det X = 1$ and $H = X$ equals $n \neq 0$. So $D\det(X) \neq 0$, and 1 is a regular value of det.

5. The orthogonal group $\mathrm{O}(n) = \{Q : Q^\top Q = I\}$ is a submanifold of dimension $\frac{n(n-1)}{2}$. Let $F(Q) = Q^\top Q$, a map from $\mathbb{R}^{n\times n}$ to the space Sym_n of symmetric matrices, which has dimension $\frac{n(n+1)}{2}$. Its derivative is $DF(Q)H = H^\top Q + Q^\top H$. For $Q \in \mathrm{O}(n)$ and any symmetric S, the matrix $H = \frac{1}{2}QS$ satisfies $DF(Q)H = \frac{1}{2}SQ^\top Q + \frac{1}{2}Q^\top QS = S$, so $DF(Q)$ is surjective, i.e., has full rank $\frac{n(n+1)}{2}$, and $\mathrm{O}(n)$ has dimension $n^2 - \frac{n(n+1)}{2} = \frac{n(n-1)}{2}$. For $n = 2$, this is 1 (one rotation angle), and for $n = 3$, it is 3 (three angles, as for the rotations R_1, R_2, R_3 of Lecture 2).

Not every level set is a submanifold. The level set $\{x^2 - y^2 = 0\}$ of Figure 3.1 in Lecture 3 consists of two lines that cross at the origin, where the gradient $(2x, -2y)$ vanishes. It is not a 1-dimensional submanifold near the origin: removing the origin from a small piece of it leaves four connected pieces, while removing a point from a small piece of a curve that looks like a line leaves two. Similarly, the double cone $\{x^2 + y^2 = z^2\}$ is not a submanifold at its tip. On the other hand, the rank condition is sufficient, but not necessary: $\{x^2 = 0\}$ is the y-axis, a perfectly good submanifold, although $D(x^2) = (2x, 0)$ vanishes on it.

Definition A.7.4 (Tangent space)**.** Let M be a submanifold and $\mathbf{p} \in M$. The tangent space $T_\mathbf{p}M$ is the set of velocities $\boldsymbol{\gamma}'(0)$ of all smooth curves $\boldsymbol{\gamma} : (-\varepsilon, \varepsilon) \to \mathbb{R}^n$ with $\boldsymbol{\gamma}(t) \in M$ for all t and $\boldsymbol{\gamma}(0) = \mathbf{p}$. Its orthogonal complement $N_\mathbf{p}M = (T_\mathbf{p}M)^\perp$ is the normal space.

Theorem A.7.5. *Let M be a k-dimensional submanifold, and let $M \cap U = F^{-1}(\mathbf{0})$ near $\mathbf{p}$ as in Definition A.7.1. Then*

$$T_\mathbf{p}M = \mathrm{null}\, DF(\mathbf{p}) \qquad \textit{and} \qquad N_\mathbf{p}M = \mathrm{row}\, DF(\mathbf{p}) = \mathrm{span}\{\nabla F_1(\mathbf{p}), \ldots, \nabla F_{n-k}(\mathbf{p})\}.$$

In particular, $T_\mathbf{p}M$ is a k-dimensional subspace.

Proof. If $\boldsymbol{\gamma}$ is a curve in M through $\mathbf{p}$, then $F(\boldsymbol{\gamma}(t)) = \mathbf{0}$ for small t, and the chain rule gives $DF(\mathbf{p})\boldsymbol{\gamma}'(0) = \mathbf{0}$. So $T_\mathbf{p}M \subseteq \mathrm{null}\, DF(\mathbf{p})$, which has dimension $n - (n-k) = k$ by the rank–nullity theorem (Lecture 1). Conversely, write M near $\mathbf{p} = (\mathbf{a}, \mathbf{g}(\mathbf{a}))$ as a graph as in Proposition A.7.2. For every $\mathbf{v} \in \mathbb{R}^k$, the curve $t \mapsto (\mathbf{a} + t\mathbf{v}, \mathbf{g}(\mathbf{a} + t\mathbf{v}))$ lies in M and has velocity $(\mathbf{v}, D\mathbf{g}(\mathbf{a})\mathbf{v})$ at $t = 0$. These velocities form a k-dimensional subspace (the image of an injective linear map), so $T_\mathbf{p}M$ contains a k-dimensional subspace of the k-dimensional space $\mathrm{null}\, DF(\mathbf{p})$, and the two are equal. Finally, $\mathrm{null}(A)^\perp = \mathrm{row}(A)$ for every matrix A (Lecture 2), and the rows of $DF(\mathbf{p})$ are the transposed gradients $\nabla F_i(\mathbf{p})^\top$. □

Example A.7.6. 1. $T_\mathbf{p}S^{n-1} = \mathrm{null}(2\mathbf{p}^\top) = \{\mathbf{v} : \mathbf{p}^\top\mathbf{v} = 0\} = \mathbf{p}^\perp$: the tangent vectors of a sphere are orthogonal to the radius (Figure A.1).

2. $T_I\mathrm{SL}(n, \mathbb{R}) = \{H : D\det(I)H = 0\} = \{H : \mathrm{tr}\, H = 0\}$, the traceless matrices: to first order, $\det(I + tH) = 1 + t\,\mathrm{tr}\, H$ (Lemma 3.6.3), so a matrix curve through I keeps determinant 1 only if its velocity has trace 0.




Figure A.1: The circle $M = \{F = 0\}$ with $F(x, y) = x^2 + y^2 - 1$. The tangent space $T_{\mathbf{p}}M = \operatorname{null} DF(\mathbf{p}) = \mathbf{p}^{\perp}$ is the tangent line (shifted to $\mathbf{p}$), and the normal space is spanned by the gradient $\nabla F(\mathbf{p})$.

3. $T_Q\mathrm{O}(n) = \{H : H^\top Q + Q^\top H = 0\}$. With $K = Q^\top H$, the condition says $K^\top + K = 0$, so $T_Q\mathrm{O}(n) = \{QK : K^\top = -K\}$, and $T_I\mathrm{O}(n)$ is the space of skew-symmetric matrices, of dimension $\frac{n(n-1)}{2}$. For example, the rotations $R(t)$ by the angle t in the plane have $R'(0) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

A.7.1 Lie groups

The examples $\mathrm{GL}(n, \mathbb{R})$, $\mathrm{SL}(n, \mathbb{R})$ and $\mathrm{O}(n)$ are both groups (Definition 1.6.70) and submanifolds. Such objects describe continuous symmetries, and they have a name.

Definition A.7.7 (Lie group)**.** A Lie group is a group that is also a manifold, such that the multiplication $(a, b) \mapsto ab$ and the inversion $a \mapsto a^{-1}$ are smooth. A matrix Lie group is a subgroup G of $\mathrm{GL}(n, \mathbb{R})$ that is a submanifold of $\mathbb{R}^{n\times n}$. Its tangent space $\mathfrak{g} = T_I G$ at the identity is the Lie algebra of G.

A matrix Lie group is a Lie group in the sense of the first sentence: its multiplication and inversion are restrictions of smooth maps on open sets, namely of $(A, B) \mapsto AB$, whose entries are polynomials in the entries of A and B, and of $A \mapsto A^{-1}$ on the open set $\mathrm{GL}(n, \mathbb{R})$, whose entries are rational functions of the entries of A by the cofactor formula (Proposition 1.6.96). The dimension of G is the number of independent parameters needed to describe its elements near I.

Proposition A.7.8 (The classical matrix groups)**.** *1.* $\mathrm{GL}(n, \mathbb{R})$, $\mathrm{SL}(n, \mathbb{R})$, $\mathrm{O}(n)$ *and* $\mathrm{SO}(n)$ *are matrix Lie groups of the dimensions* n^2, $n^2 - 1$, $\frac{n(n-1)}{2}$ *and* $\frac{n(n-1)}{2}$. *Their Lie algebras are all of* $\mathbb{R}^{n\times n}$, *the traceless matrices* $\mathfrak{sl}(n)$, *and (for both* $\mathrm{O}(n)$ *and* $\mathrm{SO}(n)$*) the skew-symmetric matrices* $\mathfrak{so}(n)$.

2. For every $H \in \mathbb{R}^{n\times n}$ *and* $s, t \in \mathbb{R}$, $e^{(s+t)H} = e^{sH}e^{tH}$, *so* $e^{tH} \in \mathrm{GL}(n, \mathbb{R})$ *with the inverse* e^{-tH}, *and* $\det e^{tH} = e^{t \operatorname{tr} H}$.

3. The exponential maps the Lie algebras into the groups: $e^{tH} \in \mathrm{SL}(n, \mathbb{R})$ *for* $H \in \mathfrak{sl}(n)$, *and* $e^{tK} \in \mathrm{SO}(n)$ *for* $K \in \mathfrak{so}(n)$.

4. $\mathfrak{sl}(n)$ *and* $\mathfrak{so}(n)$ *are closed under the commutator* $[A, B] = AB - BA$.

Proof. (1) These sets are groups by Lecture 1 (for $\mathrm{SO}(n)$, see Lecture 2), and $\mathrm{GL}(n, \mathbb{R})$, $\mathrm{SL}(n, \mathbb{R})$ and $\mathrm{O}(n)$ are submanifolds of the stated dimensions by Example A.7.3. Since $\det Q = \pm 1$ on $\mathrm{O}(n)$, $\mathrm{SO}(n)$ is the intersection of $\mathrm{O}(n)$ with the open set $\{\det > 0\}$, so it is a submanifold of the same

dimension (in Definition A.7.1, replace U by $U \cap \{\det > 0\}$), and it has the same tangent space at I. The Lie algebras are the tangent spaces of Example A.7.6; for the open set $\mathrm{GL}(n,\mathbb{R})$, every direction is tangent.

(2) The power series e^{sH} and e^{tH} are limits of polynomials in H, so they commute. As functions of t, both $e^{(s+t)H}$ and $e^{tH}e^{sH}$ solve the linear ODE $X' = HX$ (Lecture 3: $\frac{d}{dt}e^{tH} = He^{tH}$), column by column, with the same initial value e^{sH} at $t = 0$, so they agree by the uniqueness of solutions (Theorem A.12.4). With $s = -t$, $e^{-tH}e^{tH} = e^0 = I$. For $\phi(t) = \det e^{tH}$, Jacobi's formula (Proposition 3.6.4) and the chain rule give

$$\phi'(t) = \det(e^{tH}) \operatorname{tr}\left(e^{-tH}He^{tH}\right) = \phi(t) \operatorname{tr} H,$$

since $\operatorname{tr}(AB) = \operatorname{tr}(BA)$. So $\left(\phi(t)e^{-t\operatorname{tr}H}\right)' = 0$, and $\phi(t) = \phi(0)e^{t\operatorname{tr}H} = e^{t\operatorname{tr}H}$. (This extends Liouville's formula of Lecture 3 to all matrices.)

(3) If $\operatorname{tr} H = 0$, then $\det e^{tH} = 1$ by (2). If $K^\top = -K$, then $(e^{tK})^\top = e^{tK^\top} = e^{-tK}$ (transpose the power series term by term), so $(e^{tK})^\top e^{tK} = e^{-tK}e^{tK} = I$ by (2); moreover, $\operatorname{tr} K = 0$, so $\det e^{tK} = 1$.

(4) $\operatorname{tr}(AB - BA) = \operatorname{tr}(AB) - \operatorname{tr}(BA) = 0$. If $A^\top = -A$ and $B^\top = -B$, then $(AB - BA)^\top = B^\top A^\top - A^\top B^\top = BA - AB = -(AB - BA)$. □

By (2), $t \mapsto e^{tH}$ is a curve in the group with $e^{(s+t)H} = e^{sH}e^{tH}$, a one-parameter subgroup, whose velocity at $t = 0$ is H. So the Lie algebra consists of the infinitesimal generators of the group, and the commutator measures how much the group fails to be abelian. For example, $\mathfrak{so}(2)$ is spanned by $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, and e^{tJ} is the rotation by the angle t (Lecture 3). In $\mathfrak{so}(3)$, each skew-symmetric matrix K acts as $K\mathbf{v} = \boldsymbol{\omega} \times \mathbf{v}$ for a vector $\boldsymbol{\omega} \in \mathbb{R}^3$, and e^{tK} is the rotation about the axis $\boldsymbol{\omega}$ with the angular speed $\|\boldsymbol{\omega}\|$ (compare the axis of a rotation, Lecture 2).

Remark A.7.9 (Lie groups in machine learning). Lie groups describe the symmetries of data: rotations and translations of 3D molecules and point clouds (the rotations SO(3) and the rigid motions, which combine them with translations), and rotations and shifts of images. Equivariant neural networks build these symmetries into their layers, generalizing the translation equivariance of convolutions (Lecture 1). Lie groups also appear in optimization: a weight matrix that must stay orthogonal can be parametrized as e^K with a skew-symmetric K, which is unconstrained, or updated as $Q \mapsto Qe^{tK}$, which stays in $\mathrm{O}(n)$; this is an alternative to the retractions of Section A.8.

A.8 Lagrange Multipliers and Optimization on Manifolds

We can now complete the proof of the Lagrange multiplier theorem (Theorem 3.12.1).

Theorem A.8.1 (Lagrange multipliers)**.** *Let f and $g_1, \dots, g_m$ be smooth on an open set $U \subseteq \mathbb{R}^n$, let $\mathbf{x}^*$ be a local minimum of f on $M = \{\mathbf{x} \in U : g_1(\mathbf{x}) = \cdots = g_m(\mathbf{x}) = 0\}$, and let $\nabla g_1(\mathbf{x}^*), \dots, \nabla g_m(\mathbf{x}^*)$ be linearly independent. Then $\nabla f(\mathbf{x}^*) \in \operatorname{span}\{\nabla g_1(\mathbf{x}^*), \dots, \nabla g_m(\mathbf{x}^*)\}$, i.e., there are λ_i with $\nabla f(\mathbf{x}^*) + \sum_i \lambda_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.*

Proof. The linear independence says that $DG(\mathbf{x}^*)$ has rank m for $G = (g_1, \dots, g_m)$, so near $\mathbf{x}^*$, M is an $(n-m)$-dimensional submanifold (the rank stays m near $\mathbf{x}^*$, since a nonzero $m \times m$ minor stays nonzero by continuity). For every $\mathbf{v} \in T_{\mathbf{x}^*}M$, choose a curve $\boldsymbol{\gamma}$ in M with $\boldsymbol{\gamma}(0) = \mathbf{x}^*$ and $\boldsymbol{\gamma}'(0) = \mathbf{v}$. The function $t \mapsto f(\boldsymbol{\gamma}(t))$ has a local minimum at $t = 0$, so its derivative $\nabla f(\mathbf{x}^*)^\top \mathbf{v}$ is 0. Hence $\nabla f(\mathbf{x}^*) \in (T_{\mathbf{x}^*}M)^\perp = N_{\mathbf{x}^*}M = \operatorname{span}\{\nabla g_i(\mathbf{x}^*)\}$ by Theorem A.7.5. □

The same method proves the necessity of the KKT conditions for inequality constraints (Theorem 3.12.5). The new point is the sign of the multipliers: we move from $\mathbf{x}^*$ into the feasible region along a curve on which one active constraint becomes negative and the other active constraints stay 0.

Theorem A.8.2 (KKT conditions)**.** *Let $f, g_1, \dots, g_m$ be smooth on an open set $U \subseteq \mathbb{R}^n$, and let $\mathbf{x}^*$ be a local minimum of f on the feasible set $\{\mathbf{x} \in U : g_1(\mathbf{x}) \le 0, \dots, g_m(\mathbf{x}) \le 0\}$, such that the gradients $\nabla g_i(\mathbf{x}^*)$ of the active constraints, i.e., of those with $g_i(\mathbf{x}^*) = 0$, are linearly independent. Then there are numbers $\mu_1, \dots, \mu_m \ge 0$ such that $\nabla f(\mathbf{x}^*) + \sum_i \mu_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$, and $\mu_i = 0$ for every inactive constraint.*

Proof. Let A be the set of the indices of the active constraints. The inactive constraints satisfy $g_i(\mathbf{x}^*) < 0$, so by continuity, there is a ball $W \subseteq U$ around $\mathbf{x}^*$ on which $g_i < 0$ for all $i \notin A$, and we may choose it so small that $f(\mathbf{x}) \ge f(\mathbf{x}^*)$ for all feasible $\mathbf{x} \in W$. We set $\mu_i = 0$ for $i \notin A$. If A is empty, then all points of W are feasible, so $\mathbf{x}^*$ is a local minimum of f on an open set, and $\nabla f(\mathbf{x}^*) = \mathbf{0}$. Otherwise, all points of $M = \{\mathbf{x} \in W : g_i(\mathbf{x}) = 0 \text{ for all } i \in A\}$ are feasible, so $\mathbf{x}^*$ is a local minimum of f on M, and Theorem A.8.1 gives numbers μ_i, $i \in A$, with $\nabla f(\mathbf{x}^*) + \sum_{i \in A} \mu_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.

It remains to show that $\mu_j \ge 0$ for every $j \in A$. The matrix with the rows $\nabla g_i(\mathbf{x}^*)^\top$, $i \in A$, has linearly independent rows, so its column space is all of $\mathbb{R}^{|A|}$ (Lecture 1), and there is a vector $\mathbf{v}$ with

$$\nabla g_j(\mathbf{x}^*)^\top \mathbf{v} = -1 \qquad \text{and} \qquad \nabla g_i(\mathbf{x}^*)^\top \mathbf{v} = 0 \quad \text{for } i \in A,\ i \ne j.$$

If $A = \{j\}$, let $\boldsymbol{\gamma}(t) = \mathbf{x}^* + t\mathbf{v}$. Otherwise, the gradients of the constraints g_i, $i \in A \setminus \{j\}$, are linearly independent, so after shrinking W, $M_j = \{\mathbf{x} \in W : g_i(\mathbf{x}) = 0 \text{ for } i \in A \setminus \{j\}\}$ is a submanifold, as in the proof of Theorem A.8.1. By Theorem A.7.5, $\mathbf{v} \in T_{\mathbf{x}^*} M_j$, so there is a smooth curve $\boldsymbol{\gamma}$ in M_j with $\boldsymbol{\gamma}(0) = \mathbf{x}^*$ and $\boldsymbol{\gamma}'(0) = \mathbf{v}$. In both cases, the constraints g_i with $i \in A \setminus \{j\}$ are 0 along $\boldsymbol{\gamma}$, and the derivative of $g_j(\boldsymbol{\gamma}(t))$ at $t = 0$ is $\nabla g_j(\mathbf{x}^*)^\top \mathbf{v} = -1 < 0$, so $g_j(\boldsymbol{\gamma}(t)) < g_j(\mathbf{x}^*) = 0$ for small $t > 0$. Hence $\boldsymbol{\gamma}(t)$ is feasible for small $t > 0$, and $f(\boldsymbol{\gamma}(t)) \ge f(\mathbf{x}^*)$. So the difference quotients of $f(\boldsymbol{\gamma}(t))$ at $t = 0$ are nonnegative for $t > 0$, and so is its derivative:

$$0 \le \nabla f(\mathbf{x}^*)^\top \mathbf{v} = -\sum_{i \in A} \mu_i \nabla g_i(\mathbf{x}^*)^\top \mathbf{v} = \mu_j. \qquad \square$$

The proof suggests how to do calculus on a manifold: only the component of ∇f in the tangent space matters.

Definition A.8.3 (Riemannian gradient)**.** Let M be a submanifold of $\mathbb{R}^n$, $\mathbf{p} \in M$, and f smooth near M. The Riemannian gradient of f on M at $\mathbf{p}$ is the orthogonal projection (Lecture 2) of $\nabla f(\mathbf{p})$ onto $T_{\mathbf{p}} M$:

$$\operatorname{grad}_M f(\mathbf{p}) = P_{\mathbf{p}} \nabla f(\mathbf{p}).$$

Since $\nabla f(\mathbf{p}) - \operatorname{grad}_M f(\mathbf{p})$ is orthogonal to $T_{\mathbf{p}} M$, we have $\nabla f(\mathbf{p})^\top \mathbf{v} = \operatorname{grad}_M f(\mathbf{p})^\top \mathbf{v}$ for all tangent vectors $\mathbf{v}$: the Riemannian gradient gives the rates of change of f along M. As in Theorem 3.3.9, it points in the direction of steepest ascent among the tangent directions, and by Theorem A.8.1, $\operatorname{grad}_M f(\mathbf{p}) = \mathbf{0}$ exactly at the points that satisfy the Lagrange condition. These points are the critical points of f on M.

Riemannian gradient descent minimizes f on M by taking a step in the direction $-\operatorname{grad}_M f(\mathbf{p}_k)$, which leaves M slightly, and mapping the result back to M with a retraction R:

$$\mathbf{p}_{k+1} = R(\mathbf{p}_k - \eta \operatorname{grad}_M f(\mathbf{p}_k)).$$

On the sphere, $P_{\mathbf{p}} = I - \mathbf{p}\mathbf{p}^\top$, so $\operatorname{grad}_M f(\mathbf{p}) = \nabla f(\mathbf{p}) - (\mathbf{p}^\top \nabla f(\mathbf{p}))\mathbf{p}$, and the retraction can simply normalize: $R(\mathbf{x}) = \frac{\mathbf{x}}{\|\mathbf{x}\|}$. On $\mathrm{O}(n)$, one can retract with the Q factor of the QR decomposition (Lecture 2). Such methods are used for PCA, for orthogonality constraints on weight matrices, and for low-rank matrix problems.

Example A.8.4 (The Rayleigh quotient on the sphere)**.** Let A be symmetric with eigenvalues $\lambda_1 > \lambda_2 > \cdots > \lambda_n$ and unit eigenvectors $\mathbf{q}_1, \dots, \mathbf{q}_n$, and let $f(\mathbf{x}) = \mathbf{x}^\top A \mathbf{x}$ on S^{n-1}. Then

$\operatorname{grad}_M f(\mathbf{x}) = 2A\mathbf{x} - 2(\mathbf{x}^\top A\mathbf{x})\mathbf{x}$, which is $\mathbf{0}$ if and only if $\mathbf{x}$ is an eigenvector: the critical points are the $2n$ points $\pm\mathbf{q}_i$. To see their types, move from $\mathbf{q}_i$ towards $\mathbf{q}_j$ along the great circle $\boldsymbol{\gamma}(t) = \cos t\,\mathbf{q}_i + \sin t\,\mathbf{q}_j$:

$$f(\boldsymbol{\gamma}(t)) = \lambda_i \cos^2 t + \lambda_j \sin^2 t = \lambda_i + (\lambda_j - \lambda_i)\sin^2 t.$$

So f decreases in the $n-i$ tangent directions $\mathbf{q}_j$ with $\lambda_j < \lambda_i$, and increases in the $i-1$ directions with $\lambda_j > \lambda_i$. Hence $\pm\mathbf{q}_1$ are the maxima, $\pm\mathbf{q}_n$ the minima, and all other eigenvectors are saddle points. In particular, Riemannian gradient ascent for the first principal component (Lecture 2) has exactly one pair of local maxima, the correct answer; the other critical points are saddle points, which gradient ascent avoids from almost every starting point.

A.9 Critical Points and Morse Theory

A.9.1 Regular and critical values

Definition A.9.1 (Critical points and values of maps)**.** Let $F : U \to \mathbb{R}^m$ be smooth on an open set $U \subseteq \mathbb{R}^n$. A point $\mathbf{x}$ is a critical point of F if $\operatorname{rank} DF(\mathbf{x}) < m$, and a value $\mathbf{c}$ is a critical value if $\mathbf{c} = F(\mathbf{x})$ for a critical point $\mathbf{x}$. All other values in $\mathbb{R}^m$ are regular values.

For $m = 1$, the critical points are the points with $\nabla F(\mathbf{x}) = \mathbf{0}$, as in Lecture 3. By Section A.7, the level set of a regular value is a submanifold (or empty). How many critical values can there be? A set $Z \subseteq \mathbb{R}^m$ has measure zero if for every $\varepsilon > 0$ it can be covered by countably many boxes whose volumes add up to less than ε. For example, finite and countable sets, and lines in $\mathbb{R}^2$, have measure zero.

Theorem A.9.2 (Sard)**.** *The set of critical values of a smooth map $F : U \to \mathbb{R}^m$ has measure zero.*

We do not prove this theorem. It says that almost every value is regular, so almost every level set of a smooth function $f : U \to \mathbb{R}$ is an $(n-1)$-dimensional submanifold or empty. Note that the theorem is about values, not points: a constant function has only critical points, but only one critical value.

A.9.2 Nondegenerate critical points

From now on, let $f : U \to \mathbb{R}$ be smooth. Lecture 3 classified critical points by the eigenvalues of the Hessian (Theorem 3.9.3). The test was inconclusive when an eigenvalue is 0.

Definition A.9.3 (Nondegenerate critical points and the index)**.** A critical point $\mathbf{p}$ of f is nondegenerate if the Hessian $\nabla^2 f(\mathbf{p})$ is invertible, i.e., has no eigenvalue 0. Its index is the number of negative eigenvalues of $\nabla^2 f(\mathbf{p})$, counted with multiplicity. A function whose critical points are all nondegenerate is a Morse function.

So a nondegenerate critical point of index 0 is a strict local minimum, one of index n is a strict local maximum, and all others are saddle points. The index counts the independent directions in which f decreases. For example, $x^3 + y^3 - 3xy$ (Example 3.9.4) is a Morse function, with a critical point of index 1 at the origin and one of index 0 at $(1, 1)$. The "monkey saddle" $x^3 - 3xy^2$ has a degenerate critical point at the origin, where its Hessian is 0; near the origin, it goes up and down three times, unlike any nondegenerate saddle.

Proposition A.9.4. *Nondegenerate critical points are isolated: each one has a neighborhood that contains no other critical point.*

Proof. Let $H = \nabla^2 f(\mathbf{p})$, and let $\mu > 0$ be the smallest absolute value of an eigenvalue of H. Writing $\mathbf{h}$ in an orthonormal eigenbasis (Lecture 2) shows $\|H\mathbf{h}\| \geq \mu\|\mathbf{h}\|$. The gradient is differentiable with derivative H at $\mathbf{p}$, so $\nabla f(\mathbf{p}+\mathbf{h}) = H\mathbf{h} + \mathbf{r}(\mathbf{h})$ with $\|\mathbf{r}(\mathbf{h})\| \leq \frac{\mu}{2}\|\mathbf{h}\|$ for small $\mathbf{h}$. Then $\|\nabla f(\mathbf{p}+\mathbf{h})\| \geq \frac{\mu}{2}\|\mathbf{h}\| > 0$ for small $\mathbf{h} \neq \mathbf{0}$. □

In Lecture 3, we saw that near a critical point, f is approximately a sum of parabolas, $f(\mathbf{p}) + \frac{1}{2}\sum_i \lambda_i y_i^2$, in the coordinates given by the eigenvectors of the Hessian. For nondegenerate critical points, this becomes exact after a nonlinear change of coordinates.

Theorem A.9.5 (Morse lemma)**.** *Let $\mathbf{p}$ be a nondegenerate critical point of f with index k. Then there is a diffeomorphism $\boldsymbol{\varphi}$ from a neighborhood of $\mathbf{p}$ onto a neighborhood of $\mathbf{0}$, with $\boldsymbol{\varphi}(\mathbf{p}) = \mathbf{0}$, such that in the coordinates $\mathbf{y} = \boldsymbol{\varphi}(\mathbf{x})$,*

$$f = f(\mathbf{p}) - y_1^2 - \cdots - y_k^2 + y_{k+1}^2 + \cdots + y_n^2.$$

We do not prove the Morse lemma. It says that near a nondegenerate critical point, the higher-order terms of the Taylor expansion do not change the picture at all: up to a smooth deformation, the function is a bowl, a saddle with k downward directions, or an upside-down bowl. Degenerate critical points can be much more complicated, but they are rare.

Theorem A.9.6 (Morse functions are generic)**.** *Let $f : U \to \mathbb{R}$ be smooth. For almost every $\mathbf{a} \in \mathbb{R}^n$ (all except a set of measure zero), the function $f_{\mathbf{a}}(\mathbf{x}) = f(\mathbf{x}) - \mathbf{a}^\top\mathbf{x}$ is a Morse function.*

Proof. We have $\nabla f_{\mathbf{a}} = \nabla f - \mathbf{a}$ and $\nabla^2 f_{\mathbf{a}} = \nabla^2 f$. So $\mathbf{x}$ is a critical point of $f_{\mathbf{a}}$ if and only if $\nabla f(\mathbf{x}) = \mathbf{a}$, and it is degenerate if and only if $\nabla^2 f(\mathbf{x})$, the Jacobian of the map $\nabla f : U \to \mathbb{R}^n$, is singular. Hence $f_{\mathbf{a}}$ has a degenerate critical point if and only if $\mathbf{a}$ is a critical value of ∇f, and by Sard's theorem, these $\mathbf{a}$ form a set of measure zero. □

For example, $f(x) = x^3$ has a degenerate critical point at 0, but $x^3 - ax$ is a Morse function for every $a \neq 0$: it has two nondegenerate critical points if $a > 0$, and none if $a < 0$. An arbitrarily small, random linear perturbation removes all degenerate critical points.

A.9.3 Critical points and the shape of sublevel sets

Morse theory studies how the sublevel sets $M^c = \{\mathbf{x} : f(\mathbf{x}) \leq c\}$ change as c increases. We state its two main results informally, for a Morse function f whose sublevel sets are compact (for example, a Morse function on a compact manifold, or a coercive Morse function on $\mathbb{R}^n$).

- If there are no critical values in $[a, b]$, then M^a and M^b have the same shape: M^b can be continuously deformed into M^a, by flowing downhill along $-\nabla f$. Nothing interesting happens between critical values.
- When c passes the value of a critical point of index k, the sublevel set changes by attaching a k-dimensional "cell" (a point for $k = 0$, a segment for $k = 1$, a disk for $k = 2$, and so on). A minimum ($k = 0$) creates a new connected component. A saddle point of index 1 attaches a bridge, which either connects two components or closes a loop.

Example A.9.7 (A double well)**.** The function $f(x, y) = (x^2 - 1)^2 + y^2$ has $\nabla f = (4x(x^2 - 1), 2y)$, so its critical points are $(\pm 1, 0)$, with $f = 0$ and $\nabla^2 f = \mathrm{diag}(8, 2)$ (index 0), and $(0, 0)$, with $f = 1$ and $\nabla^2 f = \mathrm{diag}(-4, 2)$ (index 1). For $c < 0$, the sublevel set is empty; at $c = 0$, two components appear at the minima; at $c = 1$, the saddle point attaches a bridge that joins them, and for $c > 1$, the sublevel set is connected (Figure A.2). For gradient descent, this means that the two basins of the minima are separated by the saddle: an iteration that starts with $f(\mathbf{x}_0) < 1$ and decreases f in every step cannot move from one basin to the other.




Figure A.2: Sublevel sets $\{f \le c\}$ of the double well $f(x,y) = (x^2-1)^2 + y^2$, with the two minima (black, index 0, value 0) and the saddle point (red, index 1, value 1). Below the critical value 1, the sublevel set has two components; at $c = 1$, they touch at the saddle point; above it, they have merged into one.

On a compact manifold, these changes must add up. The Euler characteristic $\chi(M)$ is a number that only depends on the topology of M: for a surface divided into V vertices, E edges and F faces, it is $V - E + F$, which does not depend on the subdivision. For example, the sphere has $\chi = 2$ (a cube has $8 - 12 + 6 = 2$), and the torus (the surface of a doughnut) has $\chi = 0$.

Theorem A.9.8 (Morse relation)**.** *Let f be a Morse function on a compact manifold M, and let c_k be the number of its critical points of index k. Then*

$$\chi(M) = c_0 - c_1 + c_2 - \cdots + (-1)^n c_n.$$

We do not prove this theorem; it follows from the fact that attaching a k-cell changes the Euler characteristic by $(-1)^k$. It shows that the topology of M forces critical points to exist, whatever function we choose.

Example A.9.9.

1. The height function on the sphere S^2 has a minimum and a maximum: $\chi(S^2) = 1 - 0 + 1 = 2$.

2. The height function on an upright torus has a minimum, two saddle points and a maximum (Figure A.3): $\chi = 1 - 2 + 1 = 0$. In fact, every Morse function on the torus has at least four critical points, at least two of which are saddle points (this follows from the Morse inequalities of Section A.10).

3. For the Rayleigh quotient on S^{n-1} (Example A.8.4), the two critical points $\pm\mathbf{q}_i$ have index $n - i$, so $\sum_k (-1)^k c_k = 2\sum_{i=1}^{n} (-1)^{n-i}$, which is 2 for odd n and 0 for even n. This agrees with the Euler characteristic of the sphere S^{n-1}, which is 2 if $n-1$ is even and 0 if $n-1$ is odd. For $n = 3$ and $A = \operatorname{diag}(3,2,1)$: two maxima $\pm\mathbf{e}_1$, two saddle points $\pm\mathbf{e}_2$, two minima $\pm\mathbf{e}_3$, and $2 - 2 + 2 = 2$.

A.10 Differential Forms and Stokes' Theorem

Lecture 3 contains several theorems of the same shape: the fundamental theorem for line integrals, Green's theorem and the divergence theorem (Theorems 3.14.8, 3.14.14 and 3.14.12) all say that the integral of a derivative over a region equals an integral over its boundary. The derivatives involved, the gradient, the curl and the divergence, look different, and the curl was only defined in two and three dimensions. Differential forms unify all of this: they are the natural integrands of k-dimensional integrals in $\mathbb{R}^n$, they have one derivative d, and they satisfy one theorem of Stokes.




Figure A.3: The height function on an upright torus has four critical points: the bottom (a minimum), the bottom and the top of the hole (two saddle points), and the top (a maximum). As the height c increases, the sublevel set is first a disk around the bottom (after the minimum), then a bent tube, i.e., a cylinder (after the first saddle point, where the disk closes up around the tube), then a torus with a hole at its top (after the second saddle point, where the two ends of the tube meet and close a loop around the hole of the torus), and finally the whole torus (after the maximum).

They are built from alternating tensors, the linear algebra of oriented areas and volumes. We use the tensors of Lecture 1 (Section 1.11), and we write $\mathbf{e}_1, \dots, \mathbf{e}_n$ for the standard basis of $\mathbb{R}^n$. As in Lecture 1, elements of a general vector space V are written in normal font.

A.10.1 Alternating forms

Definition A.10.1 (Alternating form)**.** Let V be a finite-dimensional vector space and $k \geq 1$. A k-linear form ω on $V \times \cdots \times V$ (k factors) is alternating if $\omega(v_1, \dots, v_k) = 0$ whenever $v_i = v_j$ for some $i \neq j$. The alternating k-forms form a subspace $\Lambda^k V^*$ of the space of all k-linear forms on V, which is $V^* \otimes \cdots \otimes V^*$ by Theorem 1.11.11. We also set $\Lambda^0 V^* = \mathbb{R}$. Note that $\Lambda^1 V^* = V^*$.

Lemma A.10.2. *Let ω be an alternating k-form on V.*

1. *Swapping two arguments changes the sign of $\omega(v_1, \dots, v_k)$.*
2. *If $v_1, \dots, v_k$ are linearly dependent, then $\omega(v_1, \dots, v_k) = 0$. In particular, $\Lambda^k V^* = \{0\}$ for $k > \dim V$.*

Proof. (1) Put $v + w$ into both positions i and j, and expand by multilinearity:

$$0 = \omega(\dots, v + w, \dots, v + w, \dots) = \omega(\dots, v, \dots, w, \dots) + \omega(\dots, w, \dots, v, \dots),$$

since the terms with v in both positions and with w in both positions vanish. (2) Some v_j is a linear combination of the others; by (1), we may assume that $j = k$, so $v_k = \sum_{i<k} a_i v_i$. Then $\omega(v_1, \dots, v_k) = \sum_{i<k} a_i\, \omega(v_1, \dots, v_{k-1}, v_i) = 0$, since each term contains the vector v_i twice. More than $\dim V$ vectors are always linearly dependent. □

The determinant of an $n \times n$ matrix, as a function of its columns, is an alternating n-form on $\mathbb{R}^n$: it is multilinear (Example 1.11.2), and it vanishes when two columns are equal, since swapping them changes its sign (Theorem 1.6.84). More generally, determinants produce alternating forms from covectors.

Definition A.10.3 (Wedge product of covectors)**.** For $\varphi_1, \dots, \varphi_k \in V^*$, the wedge product $\varphi_1 \wedge \cdots \wedge \varphi_k \in \Lambda^k V^*$ is

$$(\varphi_1 \wedge \cdots \wedge \varphi_k)(v_1, \dots, v_k) = \det\left(\varphi_i(v_j)\right)_{i,j=1,\dots,k}.$$

This form is multilinear, since the j-th column of the matrix depends linearly on v_j and the determinant is linear in each column, and it is alternating, since two equal arguments give two equal columns. Swapping two covectors swaps two rows, so $\varphi \wedge \psi = -\psi \wedge \varphi$, and in particular $\varphi \wedge \varphi = 0$. For $k = 2$,

$$(\varphi \wedge \psi)(v, w) = \varphi(v)\psi(w) - \varphi(w)\psi(v), \qquad \text{i.e.,} \qquad \varphi \wedge \psi = \varphi \otimes \psi - \psi \otimes \varphi.$$

On $\mathbb{R}^n$, we write dx_i for the coordinate functional $\mathbf{x} \mapsto x_i$, i.e., for the dual basis vector e^i; the reason for this notation will appear soon. For an increasing sequence of indices $I = (i_1 < i_2 < \cdots < i_k)$, we write $dx_I = dx_{i_1} \wedge \cdots \wedge dx_{i_k}$. For example, $(dx_1 \wedge dx_2)(\mathbf{v}, \mathbf{w}) = v_1 w_2 - v_2 w_1$ is the signed area of the parallelogram spanned by the projections of $\mathbf{v}$ and $\mathbf{w}$ to the x_1x_2-plane, and $(dx_1 \wedge \cdots \wedge dx_n)(\mathbf{v}_1, \dots, \mathbf{v}_n) = \det(\mathbf{v}_1, \dots, \mathbf{v}_n)$ is the signed volume of the parallelepiped spanned by $\mathbf{v}_1, \dots, \mathbf{v}_n$.

Theorem A.10.4. *The forms dx_I, for all increasing sequences $I = (i_1 < \cdots < i_k)$ in $\{1, \dots, n\}$, form a basis of $\Lambda^k(\mathbb{R}^n)^*$. In particular, $\dim \Lambda^k(\mathbb{R}^n)^* = \binom{n}{k}$, and every alternating k-form on $\mathbb{R}^n$ is*

$$\omega = \sum_I \omega(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k})\, dx_I.$$

Proof. For increasing sequences I and J, $dx_I(\mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_k}) = \det(\delta_{i_a j_b})_{a,b}$, which is 1 if $I = J$, and 0 otherwise: if $I \neq J$, some i_a is not among the j_b, and the a-th row of the matrix is zero. So applying a relation $\sum_I c_I\, dx_I = 0$ to $(\mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_k})$ gives $c_J = 0$, and the forms dx_I are linearly independent. Now let ω be alternating, and let $\eta = \sum_I \omega(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k})\, dx_I$. The alternating forms ω and η agree on $(\mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_k})$ for every increasing J. By Lemma A.10.2, they also agree on all sequences of standard basis vectors with distinct indices, which are obtained from the increasing ones by swaps that change both values by the same sign, and both vanish on sequences with a repeated vector. A multilinear form is determined by its values on all sequences of basis vectors (Theorem 1.11.11), so $\omega = \eta$. □

For $k = n$, the theorem says that every alternating n-form on $\mathbb{R}^n$ is a multiple of the determinant $dx_1 \wedge \cdots \wedge dx_n$. The wedge product extends to forms of all degrees: for $\omega = \sum_I a_I\, dx_I \in \Lambda^k(\mathbb{R}^n)^*$ and $\eta = \sum_J b_J\, dx_J \in \Lambda^l(\mathbb{R}^n)^*$, let

$$\omega \wedge \eta = \sum_{I,J} a_I b_J\, dx_{i_1} \wedge \cdots \wedge dx_{i_k} \wedge dx_{j_1} \wedge \cdots \wedge dx_{j_l},$$

where each term is a wedge product of $k + l$ covectors, which is 0 if I and J have a common index, and $\pm dx_{I \cup J}$ otherwise. The wedge product is bilinear and associative, and since swapping two covectors changes the sign, moving each of the l covectors of dx_J past the k covectors of dx_I gives

$$\omega \wedge \eta = (-1)^{kl}\, \eta \wedge \omega.$$

For a number $a \in \Lambda^0$, we set $a \wedge \omega = a\omega$. So $dx_1 \wedge dx_2 = -dx_2 \wedge dx_1$ and $dx_1 \wedge dx_1 = 0$, but a 2-form can have a nonzero wedge product with itself (Exercise 17).

A.10.2 Differential forms and the exterior derivative

Definition A.10.5 (Differential form)**.** Let $U \subseteq \mathbb{R}^n$ be open. A differential k-form on U is a map ω that assigns to every point $\mathbf{x} \in U$ an alternating k-form $\omega_{\mathbf{x}} \in \Lambda^k(\mathbb{R}^n)^*$, such that the coefficients in

$$\omega = \sum_I f_I\, dx_I, \qquad \text{i.e.,} \qquad \omega_{\mathbf{x}} = \sum_I f_I(\mathbf{x})\, dx_I,$$

are smooth functions $f_I : U \to \mathbb{R}$. A 0-form is a smooth function. Forms are added, multiplied by functions and wedged pointwise.

Example A.10.6. 1. A 1-form $\omega = F_1\,dx_1 + \cdots + F_n\,dx_n$ satisfies $\omega_{\mathbf{x}}(\mathbf{v}) = F(\mathbf{x})^\top \mathbf{v}$, the work done by the vector field F along a small displacement $\mathbf{v}$ at $\mathbf{x}$. So 1-forms correspond to vector fields (Definition 3.13.1).

2. In $\mathbb{R}^3$, the 2-form $\omega = G_1\,dx_2 \wedge dx_3 + G_2\,dx_3 \wedge dx_1 + G_3\,dx_1 \wedge dx_2$ satisfies $\omega_{\mathbf{x}}(\mathbf{v}, \mathbf{w}) = G(\mathbf{x})^\top(\mathbf{v} \times \mathbf{w})$, since $(dx_2 \wedge dx_3)(\mathbf{v}, \mathbf{w}) = v_2 w_3 - v_3 w_2$ is the first entry of $\mathbf{v} \times \mathbf{w}$, and similarly for the others. This is the flux of G through the small parallelogram spanned by $\mathbf{v}$ and $\mathbf{w}$, so in $\mathbb{R}^3$, 2-forms also correspond to vector fields.

3. An n-form $f\,dx_1 \wedge \cdots \wedge dx_n$ is a function times the volume form $dx_1 \wedge \cdots \wedge dx_n$, like a density.

Definition A.10.7 (Exterior derivative)**.** The exterior derivative of a 0-form f is the 1-form

$$df = \frac{\partial f}{\partial x_1}\,dx_1 + \cdots + \frac{\partial f}{\partial x_n}\,dx_n,$$

so that $df_{\mathbf{x}}(\mathbf{v}) = Df(\mathbf{x})\mathbf{v}$ is the derivative of f in the direction $\mathbf{v}$. The exterior derivative of a k-form $\omega = \sum_I f_I\,dx_I$ is the $(k+1)$-form $d\omega = \sum_I df_I \wedge dx_I$.

For the coordinate function $f(\mathbf{x}) = x_i$, we get $df = dx_i$, which explains the notation. Since $dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ is $\pm dx_I$ or 0 for any (not necessarily increasing) indices, the formula $d(f\,dx_{i_1} \wedge \cdots \wedge dx_{i_k}) = df \wedge dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ holds for all indices, by linearity.

Proposition A.10.8. *1. d is linear.*

2. (Product rule.) $d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^k\,\omega \wedge d\eta$ for a k-form ω and any form η.

3. $d(d\omega) = 0$ for every form ω.

Proof. (1) is clear from the definition. (2) By linearity, it suffices to consider $\omega = f\,dx_I$ and $\eta = g\,dx_J$. Then $\omega \wedge \eta = fg\,dx_I \wedge dx_J$, and the product rule $d(fg) = g\,df + f\,dg$ gives

$$d(\omega \wedge \eta) = g\,df \wedge dx_I \wedge dx_J + f\,dg \wedge dx_I \wedge dx_J = d\omega \wedge \eta + (-1)^k\,\omega \wedge d\eta,$$

since moving the 1-form dg past the k covectors of dx_I gives the sign $(-1)^k$. (3) For $\omega = f\,dx_I$,

$$d(d\omega) = d\Big(\sum_j \frac{\partial f}{\partial x_j}\,dx_j \wedge dx_I\Big) = \sum_{i,j} \frac{\partial^2 f}{\partial x_i \partial x_j}\,dx_i \wedge dx_j \wedge dx_I = 0,$$

since the second derivatives are symmetric in i and j (Theorem 3.8.2), while $dx_i \wedge dx_j = -dx_j \wedge dx_i$: the terms for (i,j) and (j,i) cancel, and the terms with $i = j$ vanish. □

Example A.10.9 (Vector calculus)**.** In $\mathbb{R}^3$, let $\omega_F = F_1\,dx_1 + F_2\,dx_2 + F_3\,dx_3$ and $\eta_G = G_1\,dx_2 \wedge dx_3 + G_2\,dx_3 \wedge dx_1 + G_3\,dx_1 \wedge dx_2$ be the 1-form and the 2-form of vector fields F and G, as in Example A.10.6. A short computation gives

$$\begin{aligned}
df &= \tfrac{\partial f}{\partial x_1}\,dx_1 + \tfrac{\partial f}{\partial x_2}\,dx_2 + \tfrac{\partial f}{\partial x_3}\,dx_3,\\
d\omega_F &= \Big(\tfrac{\partial F_3}{\partial x_2} - \tfrac{\partial F_2}{\partial x_3}\Big)dx_2 \wedge dx_3 + \Big(\tfrac{\partial F_1}{\partial x_3} - \tfrac{\partial F_3}{\partial x_1}\Big)dx_3 \wedge dx_1\\
&\quad + \Big(\tfrac{\partial F_2}{\partial x_1} - \tfrac{\partial F_1}{\partial x_2}\Big)dx_1 \wedge dx_2,\\
d\eta_G &= \Big(\tfrac{\partial G_1}{\partial x_1} + \tfrac{\partial G_2}{\partial x_2} + \tfrac{\partial G_3}{\partial x_3}\Big)\,dx_1 \wedge dx_2 \wedge dx_3.
\end{aligned}$$

With the correspondences of Example A.10.6, d is the gradient on 0-forms, the curl on 1-forms, and the divergence on 2-forms, and $d \circ d = 0$ contains the identities $\operatorname{curl} \nabla f = 0$ and $\operatorname{div} \operatorname{curl} F = 0$ (Proposition 3.13.8). In the plane, $d(F_1\,dx_1 + F_2\,dx_2) = \operatorname{curl} F\,dx_1 \wedge dx_2$, and in $\mathbb{R}^n$, $d(\sum_j F_j\,dx_j) = \sum_{i<j} (\frac{\partial F_j}{\partial x_i} - \frac{\partial F_i}{\partial x_j})\,dx_i \wedge dx_j$, whose coefficients are those of the skew-symmetric part of the Jacobian of F: this is the curl in every dimension.

A.10.3 Pullback and integration

Definition A.10.10 (Pullback)**.** Let $U \subseteq \mathbb{R}^m$ and $U' \subseteq \mathbb{R}^n$ be open, let $\boldsymbol{\phi} : U \to U'$ be smooth, and let ω be a k-form on U'. The pullback of ω is the k-form $\boldsymbol{\phi}^*\omega$ on U with

$$(\boldsymbol{\phi}^*\omega)_{\mathbf{x}}(\mathbf{v}_1, \dots, \mathbf{v}_k) = \omega_{\boldsymbol{\phi}(\mathbf{x})}\big(D\boldsymbol{\phi}(\mathbf{x})\mathbf{v}_1, \dots, D\boldsymbol{\phi}(\mathbf{x})\mathbf{v}_k\big),$$

and $\boldsymbol{\phi}^* f = f \circ \boldsymbol{\phi}$ for a 0-form f.

In words: to evaluate $\boldsymbol{\phi}^*\omega$ on vectors at $\mathbf{x}$, push the vectors forward with the derivative of $\boldsymbol{\phi}$, and evaluate ω on them at $\boldsymbol{\phi}(\mathbf{x})$. Writing $y_1, \dots, y_n$ for the coordinates on U', we get

$$\boldsymbol{\phi}^*(dy_i) = d\phi_i = \sum_{j=1}^{m} \frac{\partial \phi_i}{\partial x_j}\, dx_j,$$

since $dy_i\big(D\boldsymbol{\phi}(\mathbf{x})\mathbf{v}\big) = D\phi_i(\mathbf{x})\mathbf{v}$. One checks that the pullback respects the wedge product, $\boldsymbol{\phi}^*(\omega \wedge \eta) = \boldsymbol{\phi}^*\omega \wedge \boldsymbol{\phi}^*\eta$, that $(\boldsymbol{\psi} \circ \boldsymbol{\phi})^* = \boldsymbol{\phi}^* \circ \boldsymbol{\psi}^*$ by the chain rule, and that the pullback commutes with the exterior derivative,

$$\boldsymbol{\phi}^*(d\omega) = d(\boldsymbol{\phi}^*\omega).$$

For 0-forms, this is the chain rule, $d(f \circ \boldsymbol{\phi}) = \boldsymbol{\phi}^*(df)$; for $\omega = f\, dy_I$, it follows with Proposition A.10.8, since $\boldsymbol{\phi}^*\omega = (f \circ \boldsymbol{\phi})\, d\phi_{i_1} \wedge \cdots \wedge d\phi_{i_k}$ and $d(d\phi_i) = 0$. So pullbacks are computed by substituting $y_i = \phi_i(\mathbf{x})$ and $dy_i = d\phi_i$. For forms of top degree, the Jacobian determinant appears.

Proposition A.10.11. *If $m = n$, then $\boldsymbol{\phi}^*(f\, dy_1 \wedge \cdots \wedge dy_n) = (f \circ \boldsymbol{\phi}) \det D\boldsymbol{\phi}\; dx_1 \wedge \cdots \wedge dx_n$.*

Proof. At each point, $\boldsymbol{\phi}^*(dy_1 \wedge \cdots \wedge dy_n)$ is an alternating n-form, so it is a multiple of $dx_1 \wedge \cdots \wedge dx_n$ (Theorem A.10.4), and the factor is its value on $(\mathbf{e}_1, \dots, \mathbf{e}_n)$:

$$(dy_1 \wedge \cdots \wedge dy_n)\big(D\boldsymbol{\phi}(\mathbf{x})\mathbf{e}_1, \dots, D\boldsymbol{\phi}(\mathbf{x})\mathbf{e}_n\big) = \det\Big(\frac{\partial \phi_i}{\partial x_j}(\mathbf{x})\Big)_{i,j} = \det D\boldsymbol{\phi}(\mathbf{x}),$$

since $dy_i\big(D\boldsymbol{\phi}(\mathbf{x})\mathbf{e}_j\big)$ is the (i,j) entry of $D\boldsymbol{\phi}(\mathbf{x})$. □

For example, for polar coordinates $\boldsymbol{\phi}(r, \theta) = (r\cos\theta, r\sin\theta)$, we get $\boldsymbol{\phi}^*(dx_1 \wedge dx_2) = r\, dr \wedge d\theta$, the factor r of Example 3.4.7.

Definition A.10.12 (Integral of a differential form)**.**

1. Let $\omega = f\, dx_1 \wedge \cdots \wedge dx_k$ be a k-form on an open set that contains a box $Q \subseteq \mathbb{R}^k$. Then $\int_Q \omega = \int_Q f(\mathbf{x})\, d\mathbf{x}$, the multiple integral of Lecture 3.

2. A k-cube in an open set $U \subseteq \mathbb{R}^n$ is a smooth map $\mathbf{c} : [0,1]^k \to U$ (more precisely, the restriction of a smooth map on an open set that contains $[0,1]^k$). For a k-form ω on U, we define $\int_{\mathbf{c}} \omega = \int_{[0,1]^k} \mathbf{c}^*\omega$. A 0-cube is a point $\mathbf{c} \in U$, and $\int_{\mathbf{c}} f = f(\mathbf{c})$.

A k-cube is a parametrized k-dimensional piece of U: a curve for $k = 1$, a surface patch for $k = 2$. Integrals of k-forms do not depend on the parametrization, except for the orientation.

Proposition A.10.13. *Let $\boldsymbol{\psi}$ be a smooth bijection of $[0,1]^k$ onto itself with a smooth inverse. If $\det D\boldsymbol{\psi} > 0$ everywhere, then $\int_{\mathbf{c}\circ\boldsymbol{\psi}} \omega = \int_{\mathbf{c}} \omega$. If $\det D\boldsymbol{\psi} < 0$ everywhere, then $\int_{\mathbf{c}\circ\boldsymbol{\psi}} \omega = -\int_{\mathbf{c}} \omega$.*

Proof. Write $\mathbf{c}^*\omega = g\, dx_1 \wedge \cdots \wedge dx_k$. By Proposition A.10.11, $(\mathbf{c} \circ \boldsymbol{\psi})^*\omega = \boldsymbol{\psi}^*(\mathbf{c}^*\omega) = (g \circ \boldsymbol{\psi}) \det D\boldsymbol{\psi}\; dx_1 \wedge \cdots \wedge dx_k$. Now the claim is the change of variables formula (Theorem 3.14.4), applied to the interior of the cube, in which $|\det D\boldsymbol{\psi}| = \pm \det D\boldsymbol{\psi}$. □

Example A.10.14. 1. (Line integrals.) For a curve $\mathbf{c} : [0,1] \to U$ and the 1-form $\omega = \sum_i F_i\, dx_i$, we get $\mathbf{c}^*\omega = \sum_i F_i(\mathbf{c}(t))\, c_i'(t)\, dt$, so

$$\int_{\mathbf{c}} \omega = \int_0^1 F(\mathbf{c}(t))^\top \mathbf{c}'(t)\, dt,$$

the line integral of the vector field F along $\mathbf{c}$ (Definition 3.14.7).

2. (Flux integrals.) For a surface patch $\mathbf{c} : [0,1]^2 \to \mathbb{R}^3$ and the 2-form ω of a vector field G as in Example A.10.6, the pullback is $\mathbf{c}^*\omega = G(\mathbf{c})^\top(\partial_1\mathbf{c} \times \partial_2\mathbf{c})\, du_1 \wedge du_2$, so $\int_{\mathbf{c}} \omega$ is the flux of G through the surface.

A.10.4 Manifolds with boundary and orientations

Stokes' theorem relates an integral over a k-dimensional region to an integral over its boundary, as for a ball and its boundary sphere, or for a surface and the curve around it. Such regions are manifolds with boundary: near their boundary points, they look like a half-space. Let

$$\mathbb{H}^k = \{\mathbf{u} \in \mathbb{R}^k : u_k \geq 0\}, \qquad \partial\mathbb{H}^k = \{\mathbf{u} \in \mathbb{R}^k : u_k = 0\}.$$

A set $V \subseteq \mathbb{H}^k$ is open in $\mathbb{H}^k$ if $V = V' \cap \mathbb{H}^k$ for an open set $V' \subseteq \mathbb{R}^k$, and a map on V is smooth if it is the restriction of a smooth map on such a set V'.

Definition A.10.15 (Submanifold with boundary)**.** Let $1 \leq k \leq n$. A set $M \subseteq \mathbb{R}^n$ is a k-dimensional submanifold with boundary if every $\mathbf{p} \in M$ has a local parametrization: a set V that is open in $\mathbb{R}^k$ or in $\mathbb{H}^k$, an open set $W \ni \mathbf{p}$, and a smooth map $\boldsymbol{\phi} : V \to \mathbb{R}^n$ that maps V homeomorphically onto $M \cap W$, with $\operatorname{rank} D\boldsymbol{\phi}(\mathbf{u}) = k$ for all $\mathbf{u} \in V$. The boundary ∂M is the set of the points $\boldsymbol{\phi}(\mathbf{u})$ with $\mathbf{u} \in \partial\mathbb{H}^k$, for the local parametrizations $\boldsymbol{\phi}$ on open subsets of $\mathbb{H}^k$.

With the inverse function theorem (Theorem A.5.6), one shows that the transition map $\boldsymbol{\phi}_2^{-1} \circ \boldsymbol{\phi}_1$ between two local parametrizations is smooth, with invertible derivatives, and that whether a point lies in ∂M does not depend on the parametrization. The maps $\mathbf{u}' \mapsto \boldsymbol{\phi}(\mathbf{u}', 0)$ are local parametrizations of ∂M by open subsets of $\mathbb{R}^{k-1}$, so ∂M is a $(k-1)$-dimensional submanifold with empty boundary. By Proposition A.7.2, the submanifolds of Definition A.7.1 are submanifolds with empty boundary.

Example A.10.16. 1. Let $g : \mathbb{R}^n \to \mathbb{R}$ be smooth, with $\nabla g(\mathbf{x}) \neq \mathbf{0}$ whenever $g(\mathbf{x}) = 0$. Then $M = \{\mathbf{x} : g(\mathbf{x}) \leq 0\}$ is an n-dimensional submanifold with boundary $\partial M = \{\mathbf{x} : g(\mathbf{x}) = 0\}$. Near a point $\mathbf{p}$ with $g(\mathbf{p}) < 0$, M contains an open set, on which the identity is a local parametrization. Near a point with $g(\mathbf{p}) = 0$, renumber the coordinates so that $\frac{\partial g}{\partial x_n}(\mathbf{p}) \neq 0$. Then $\boldsymbol{\Phi}(\mathbf{x}) = (x_1, \ldots, x_{n-1}, -g(\mathbf{x}))$ has an invertible derivative at $\mathbf{p}$, so it maps an open set $W \ni \mathbf{p}$ bijectively onto an open set V', with a smooth inverse. It maps $M \cap W$ onto $V' \cap \mathbb{H}^n$ and $\partial M \cap W$ onto $V' \cap \partial\mathbb{H}^n$, so the restriction of $\boldsymbol{\Phi}^{-1}$ to $V' \cap \mathbb{H}^n$ is a local parametrization. For $g(\mathbf{x}) = \|\mathbf{x}\|^2 - 1$, M is the closed unit ball B, with $\partial B = S^{n-1}$.

2. The closed upper hemisphere $S^2_+ = \{\mathbf{x} \in S^2 : x_3 \geq 0\}$ is a 2-dimensional submanifold with boundary, whose boundary is the equator $\{\mathbf{x} \in S^2 : x_3 = 0\}$. Near the equator, the longitude θ (in an open interval of length less than 2π) and the latitude $s \in [0, \frac{\pi}{2})$ give the local parametrization $\boldsymbol{\phi}(\theta, s) = (\cos s \cos\theta,\, \cos s \sin\theta,\, \sin s)$.

3. A compact interval $[a,b]$ is a 1-dimensional submanifold of $\mathbb{R}$ with boundary $\{a,b\}$, and the sphere S^{n-1} and the orthogonal group $\mathrm{O}(n)$ have empty boundary. The square $[0,1]^2$ is not a submanifold with boundary, because of its corners; Stokes' theorem for cubes below covers it.

By Proposition A.10.13, integrals of k-forms change their sign under reparametrizations with negative Jacobian determinants. So to integrate over M, we must choose its parametrizations consistently.

Definition A.10.17 (Orientation)**.** An orientation of a k-dimensional submanifold M with boundary is a collection of local parametrizations whose images cover M, such that the transition map between any two of them has a positive Jacobian determinant everywhere. These parametrizations, and all others whose transition maps with them have positive Jacobian determinants, are called positively oriented; those whose transition maps with them have negative Jacobian determinants are negatively oriented. M is orientable if it has an orientation, and oriented if one has been chosen.

Example A.10.18.

1. An n-dimensional submanifold $M \subseteq \mathbb{R}^n$ with boundary has the standard orientation, for which the positively oriented parametrizations are those with $\det D\boldsymbol{\phi} > 0$.

2. A 2-dimensional submanifold $S \subseteq \mathbb{R}^3$ with boundary (a surface) is oriented by a continuous unit normal field $\mathbf{n}$: the positively oriented parametrizations are those for which $\partial_1\boldsymbol{\phi} \times \partial_2\boldsymbol{\phi}$ is a positive multiple of $\mathbf{n}$. Conversely, every orientation of S comes from such a normal field. For example, S^2 and S^2_+ are oriented by the outward normal $\mathbf{n}(\mathbf{x}) = \mathbf{x}$, and the parametrization $\boldsymbol{\phi}(\theta, s)$ of Example A.10.16 is positively oriented, since $\partial_\theta\boldsymbol{\phi} \times \partial_s\boldsymbol{\phi} = \cos s \;\boldsymbol{\phi}(\theta, s)$. The Möbius strip is not orientable: a normal vector that is moved continuously once around the strip comes back as its negative.

Definition A.10.19 (Induced orientation)**.** Let M be an oriented k-dimensional submanifold with boundary, where $k \geq 2$. The induced orientation of ∂M is the one for which, for every positively oriented local parametrization $\boldsymbol{\phi}$ of M on an open subset of $\mathbb{H}^k$, the parametrization $\boldsymbol{\phi}_\partial(\mathbf{u}') = \boldsymbol{\phi}(\mathbf{u}', 0)$ of ∂M is positively oriented if k is even, and negatively oriented if k is odd. For $k = 1$ and a compact M, ∂M is a finite set of points, and the induced orientation gives a point $\boldsymbol{\phi}(0)$ the sign -1 if $\boldsymbol{\phi}$ is positively oriented, and $+1$ if it is negatively oriented; the integral of a function f over ∂M is the sum of the values $\pm f(\mathbf{p})$ over $\mathbf{p} \in \partial M$, with these signs.

(This is consistent: a transition map $\boldsymbol{\psi}$ between two parametrizations on open subsets of $\mathbb{H}^k$ maps points of $\mathbb{H}^k$ to $\mathbb{H}^k$ and points of $\partial\mathbb{H}^k$ to $\partial\mathbb{H}^k$, so at a point of $\partial\mathbb{H}^k$, the last row of $D\boldsymbol{\psi}$ is $(0, \ldots, 0, \frac{\partial\psi_k}{\partial u_k})$ with $\frac{\partial\psi_k}{\partial u_k} > 0$, and the restriction of $\boldsymbol{\psi}$ to $\partial\mathbb{H}^k$ also has a positive Jacobian determinant.) In words, a basis $\mathbf{v}_1, \ldots, \mathbf{v}_{k-1}$ of the boundary is positively oriented if $\mathbf{n}, \mathbf{v}_1, \ldots, \mathbf{v}_{k-1}$ is positively oriented for an outward vector $\mathbf{n}$: "outward normal first". For $\mathbb{H}^k$, the vector $-\mathbf{e}_k$ points outward, and $\det(-\mathbf{e}_k, \mathbf{e}_1, \ldots, \mathbf{e}_{k-1}) = (-1)^k$. For a region $M \subseteq \mathbb{R}^2$ with the standard orientation, ∂M is traversed with M on its left, i.e., counterclockwise for a disk, as in Green's theorem. For a region $M \subseteq \mathbb{R}^3$, the positively oriented parametrizations $\boldsymbol{\psi}$ of ∂M are those for which $\partial_1\boldsymbol{\psi} \times \partial_2\boldsymbol{\psi}$ points outward, as in the divergence theorem. For a surface S oriented by a normal field $\mathbf{n}$, ∂S is traversed with S on the left of a walker whose head points in the direction of $\mathbf{n}$. For example, the equator of S^2_+ with the outward normal is traversed counterclockwise, as seen from above: $\boldsymbol{\phi}_\partial(\theta) = (\cos\theta, \sin\theta, 0)$.

Definition A.10.20 (Integrals over manifolds)**.** Let M be a compact oriented k-dimensional submanifold with boundary, and let ω be a k-form on an open set that contains M.

1. Suppose that ω vanishes at all points of M outside $\boldsymbol{\phi}(K)$, for a positively oriented local parametrization $\boldsymbol{\phi} : V \to M \cap W$ and a compact set $K \subseteq V$. Then $\boldsymbol{\phi}^*\omega = g\, du_1 \wedge \cdots \wedge du_k$, where g vanishes on V outside K, and we define $\int_M \omega = \int_V g(\mathbf{u})\, d\mathbf{u}$, the integral of g, extended by 0, over a box that contains K.

2. In general, let $\rho_1, \ldots, \rho_m \geq 0$ be smooth functions on $\mathbb{R}^n$ with $\sum_i \rho_i = 1$ on an open set that contains M, such that each ρ_i vanishes at all points of M outside $\boldsymbol{\phi}_i(K_i)$, for some $\boldsymbol{\phi}_i$ and K_i as in 1 (a partition of unity). We define $\int_M \omega = \sum_{i=1}^m \int_M \rho_i\omega$.

In 1, the integral does not depend on $\boldsymbol{\phi}$: for a second positively oriented parametrization, the two integrals agree by Proposition A.10.11 and the change of variables formula, applied to the transition map, as in the proof of Proposition A.10.13. For a negatively oriented $\boldsymbol{\phi}$, the same argument gives $\int_M \omega = -\int_V \boldsymbol{\phi}^*\omega$. Partitions of unity exist because M is compact: M is covered by finitely many open balls whose closures lie in sets W with positively oriented parametrizations, and dividing smooth bump functions that are positive exactly on these balls (such as $e^{-1/(1-\|\mathbf{x}\|^2)}$ on the unit ball) by their sum gives the functions ρ_i; we omit the details. The definition in 2 does not depend on the partition of unity: for a second one, $\sigma_1, \dots, \sigma_l$, both sums are equal to $\sum_{i,j} \int_M \rho_i \sigma_j \omega$. Integrals over ∂M are defined in the same way, with the induced orientation.

For $k = n$ and the standard orientation, Proposition A.10.11 and the change of variables formula give $\int_M f\, dx_1 \wedge \cdots \wedge dx_n = \int_M f(\mathbf{x})\, d\mathbf{x}$, the multiple integral of Lecture 3. For a surface $S \subseteq \mathbb{R}^3$ oriented by $\mathbf{n}$ and the 2-form ω of a vector field G, $\int_S \omega$ is the flux of G through S in the direction of $\mathbf{n}$, by Example A.10.14.

A.10.5 Stokes' theorem

Theorem A.10.21 (Stokes)**.** *Let $M \subseteq \mathbb{R}^n$ be a compact oriented k-dimensional submanifold with boundary, where $k \geq 1$, let ∂M have the induced orientation, and let ω be a $(k-1)$-form on an open set that contains M. Then*

$$\int_M d\omega = \int_{\partial M} \omega.$$

In particular, $\int_M d\omega = 0$ if ∂M is empty.

We prove the theorem for cubes, which contains all the analysis, and then explain how the general case reduces to cubes. The boundary of the square $[0,1]^2$ consists of four edges, which we traverse counterclockwise. In general, the boundary of a k-cube $\mathbf{c}$ consists of its $2k$ faces, with signs. For $j = 1, \dots, k$ and $\alpha \in \{0,1\}$, the face $\mathbf{c}_{(j,\alpha)}$ is the $(k-1)$-cube obtained by fixing the j-th coordinate to α:

$$\mathbf{c}_{(j,\alpha)}(u_1, \dots, u_{k-1}) = \mathbf{c}(u_1, \dots, u_{j-1}, \alpha, u_j, \dots, u_{k-1}).$$

The boundary of $\mathbf{c}$ is the formal sum $\partial\mathbf{c} = \sum_{j=1}^{k} \sum_{\alpha=0,1} (-1)^{j+\alpha}\, \mathbf{c}_{(j,\alpha)}$, and for a $(k-1)$-form ω, we define

$$\int_{\partial\mathbf{c}} \omega = \sum_{j=1}^{k} \sum_{\alpha=0,1} (-1)^{j+\alpha} \int_{\mathbf{c}_{(j,\alpha)}} \omega.$$

For a curve ($k = 1$), $\int_{\partial\mathbf{c}} f = f(\mathbf{c}(1)) - f(\mathbf{c}(0))$: the boundary is the end point minus the starting point. For the square, the edges $(u,0)$, $(1,u)$, $(u,1)$ and $(0,u)$ get the signs $+$, $+$, $-$ and $-$, so they are traversed counterclockwise, as in Green's theorem.

Theorem A.10.22 (Stokes' theorem for cubes)**.** *Let ω be a $(k-1)$-form on an open set $U \subseteq \mathbb{R}^n$, and let $\mathbf{c}$ be a k-cube in U. Then*

$$\int_{\mathbf{c}} d\omega = \int_{\partial\mathbf{c}} \omega.$$

Proof. First, let $\mathbf{c}$ be the identity map of $[0,1]^k$, so that ω is a $(k-1)$-form on an open set containing $[0,1]^k \subseteq \mathbb{R}^k$. By linearity, we may assume that $\omega = f\, du_1 \wedge \cdots \wedge \widehat{du_j} \wedge \cdots \wedge du_k$, where the hat means that du_j is left out. Then

$$d\omega = \frac{\partial f}{\partial u_j}\, du_j \wedge du_1 \wedge \cdots \wedge \widehat{du_j} \wedge \cdots \wedge du_k = (-1)^{j-1} \frac{\partial f}{\partial u_j}\, du_1 \wedge \cdots \wedge du_k,$$

since the other terms of df contain some du_i twice, and moving du_j to its place takes $j-1$ swaps. By Fubini's theorem and the fundamental theorem of calculus, integrating over u_j first,

$$\int_{[0,1]^k} d\omega = (-1)^{j-1} \int_{[0,1]^{k-1}} \Big(f(\dots, \underset{j}{1}, \dots) - f(\dots, \underset{j}{0}, \dots) \Big)\, d\mathbf{u}',$$

where the j-th argument is 1 or 0, and $\mathbf{u}'$ denotes the other $k-1$ variables. On the other side, the i-th coordinate is constant on the face $\mathbf{c}_{(i,\alpha)}$, so the pullback of du_i to this face is 0. Hence only the two faces with $i = j$ contribute, and the pullback of ω to them is $f(\dots,\alpha,\dots)\,du_1 \wedge \cdots \wedge du_{k-1}$. So

$$\int_{\partial \mathbf{c}} \omega = (-1)^{j+1} \int_{[0,1]^{k-1}} f(\dots,\underset{j}{1},\dots)\,d\mathbf{u}' + (-1)^{j} \int_{[0,1]^{k-1}} f(\dots,\underset{j}{0},\dots)\,d\mathbf{u}',$$

which is the same. For a general k-cube $\mathbf{c}$, the faces of $\mathbf{c}$ are $\mathbf{c} \circ \mathbf{i}_{(j,\alpha)}$, where the $\mathbf{i}_{(j,\alpha)}$ are the faces of the identity cube $\mathbf{i}$ of $[0,1]^k$. Since the pullback commutes with d, the first case gives

$$\int_{\mathbf{c}} d\omega = \int_{[0,1]^k} \mathbf{c}^*(d\omega) = \int_{[0,1]^k} d(\mathbf{c}^*\omega) = \int_{\partial \mathbf{i}} \mathbf{c}^*\omega = \int_{\partial \mathbf{c}} \omega. \qquad \square$$

A k-cube need not be injective, so Theorem A.10.22 is not a special case of Theorem A.10.21; conversely, $[0,1]^k$ is not a submanifold with boundary for $k \geq 2$. But the analysis is the same, and Theorem A.10.21 reduces to Theorem A.10.22.

Idea of the proof of Theorem A.10.21. For $k = 1$, M consists of finitely many closed curves and arcs. Each of them can be parametrized, in the direction of its orientation, by a 1-cube $\mathbf{c}$ that is injective, except that $\mathbf{c}(0) = \mathbf{c}(1)$ for a closed curve, and Theorem A.10.22 gives $\int_{\mathbf{c}} df = f(\mathbf{c}(1)) - f(\mathbf{c}(0))$, the signed sum of f over the end points. Now let $k \geq 2$.

1. (Localization.) For a partition of unity $\rho_1, \dots, \rho_m$ as in Definition A.10.20, we have $\omega = \sum_i \rho_i \omega$ and $d\omega = \sum_i d(\rho_i \omega)$ near M, and both sides of Stokes' formula are linear in ω. So we may assume that ω and $d\omega$ vanish at all points of M outside $\boldsymbol{\phi}(K)$, for a positively oriented local parametrization $\boldsymbol{\phi} : V \to M \cap W$ and a compact set $K \subseteq V$. Shrinking and translating V if necessary, we may assume that V is open in $\mathbb{H}^k$.

2. (Pullback.) Since the pullback commutes with d, $\int_M d\omega = \int_V \boldsymbol{\phi}^*(d\omega) = \int_V d\eta$ for the $(k-1)$-form $\eta = \boldsymbol{\phi}^*\omega$, which vanishes outside K. On the boundary, the pullback of η to $\partial\mathbb{H}^k$ is $\boldsymbol{\phi}_\partial^*\omega$, so $\int_{\partial M} \omega$ is also an integral of η.

3. (A box.) Choose $a > 0$ such that K lies in the box $Q = [-a,a]^{k-1} \times [0,a]$, away from all its faces except the bottom face $u_k = 0$. The box Q is the image of the k-cube $\mathbf{A}(\mathbf{t}) = (2at_1 - a, \dots, 2at_{k-1} - a,\ at_k)$, an affine map with a positive Jacobian determinant. Extending η by 0, Proposition A.10.11, the change of variables formula and Theorem A.10.22 give

$$\int_M d\omega = \int_Q d\eta = \int_{\mathbf{A}} d\eta = \int_{\partial \mathbf{A}} \eta = (-1)^{k+0} \int_{\mathbf{A}_{(k,0)}} \eta,$$

since η vanishes on the other $2k-1$ faces of $\mathbf{A}$. The bottom face $\mathbf{A}_{(k,0)}(\mathbf{t}') = (2at_1 - a, \dots, 2at_{k-1} - a,\ 0)$ maps $[0,1]^{k-1}$ onto $[-a,a]^{k-1} \times \{0\}$ with a positive Jacobian determinant, so the right side is $(-1)^k \int_{[-a,a]^{k-1}} \boldsymbol{\phi}_\partial^*\omega$. This is $\int_{\partial M} \omega$, since $\boldsymbol{\phi}_\partial$ is positively oriented if k is even, and negatively oriented if k is odd (Definition A.10.19). If K does not meet $\partial\mathbb{H}^k$, both sides are 0.

This is where the sign $(-1)^k$ in the induced orientation comes from: it is the sign of the bottom face of a cube. Instead of a partition of unity, one can also cut M into pieces parametrized by cubes, as for the divergence theorem in Lecture 3. The integrals over the faces shared by two neighboring pieces cancel, since the two pieces induce opposite orientations on them, and only the integral over ∂M remains. $\square$

All the integral theorems of Lecture 3 are special cases of Theorem A.10.21.

- For $k = 1$ and a 0-form f: if M is an arc from $\mathbf{p}$ to $\mathbf{q}$, then $\int_M df = f(\mathbf{q}) - f(\mathbf{p})$, the fundamental theorem for line integrals. For $M = [a, b] \subseteq \mathbb{R}$, it is the fundamental theorem of calculus.
- For a compact region $M \subseteq \mathbb{R}^2$ with smooth boundary (a compact 2-dimensional submanifold with boundary, such as the sets $\{g \leq 0\}$ of Example A.10.16), with the standard orientation, and for $\omega = F_1\,dx_1 + F_2\,dx_2$: $d\omega = \operatorname{curl} F\,dx_1 \wedge dx_2$, and Stokes' theorem is Green's theorem, with ∂M traversed counterclockwise, i.e., with M on its left.
- For a compact region $M \subseteq \mathbb{R}^3$ with smooth boundary and the 2-form ω of a vector field G: $d\omega = \operatorname{div} G\,dx_1 \wedge dx_2 \wedge dx_3$, the integral $\int_{\partial M} \omega$ is the flux of G through ∂M with the outward normal, and Stokes' theorem is the divergence theorem.
- For a compact surface $S \subseteq \mathbb{R}^3$ with boundary, oriented by a unit normal field $\mathbf{n}$, and for $\omega = F_1\,dx_1 + F_2\,dx_2 + F_3\,dx_3$: $d\omega$ is the 2-form of the vector field $\operatorname{curl} F$, and Stokes' theorem is the classical Stokes theorem: the circulation of F around ∂S equals the flux of $\operatorname{curl} F$ through S in the direction of $\mathbf{n}$.

Example A.10.23. 1. The 2-form $\omega = x_1\,dx_2 \wedge dx_3 + x_2\,dx_3 \wedge dx_1 + x_3\,dx_1 \wedge dx_2$ of the vector field $G(\mathbf{x}) = \mathbf{x}$ has $d\omega = 3\,dx_1 \wedge dx_2 \wedge dx_3$. For the closed unit ball $B \subseteq \mathbb{R}^3$, whose boundary S^2 gets the outward normal, Stokes' theorem gives $\int_{S^2} \omega = 3 \cdot \frac{4}{3}\pi = 4\pi$, as in Example 3.14.13.

2. Let S^2_+ be the closed upper hemisphere with the outward normal, and let $\omega = -x_2\,dx_1 + x_1\,dx_2$ be the 1-form of $F(\mathbf{x}) = (-x_2, x_1, 0)$. Then $d\omega = 2\,dx_1 \wedge dx_2$, the 2-form of $\operatorname{curl} F = (0, 0, 2)$. The equator is traversed counterclockwise by $\boldsymbol{\phi}_\partial(\theta) = (\cos\theta, \sin\theta, 0)$, with $\boldsymbol{\phi}_\partial^*\omega = (\sin^2\theta + \cos^2\theta)\,d\theta$, so $\int_{S^2_+} d\omega = \int_0^{2\pi} d\theta = 2\pi$. The closed unit disk D in the plane $x_3 = 0$, oriented by the upward normal $\mathbf{e}_3$, has the same oriented boundary, so $\int_D d\omega = 2\pi$ as well, which is twice the area of D. In general, the flux of $\operatorname{curl} F$ through an oriented surface only depends on its oriented boundary.

A.10.6 Closed and exact forms

Definition A.10.24. A k-form ω is closed if $d\omega = 0$, and exact if $\omega = d\eta$ for a $(k-1)$-form η.

Every exact form is closed, since $d(d\eta) = 0$. For a 1-form $\omega = \sum_i F_i\,dx_i$, exact means that $F = \nabla f$ is a gradient field, and closed means that $\frac{\partial F_j}{\partial x_i} = \frac{\partial F_i}{\partial x_j}$ for all i and j, i.e., that the Jacobian of F is symmetric. The Poincaré lemma of Lecture 3 (Theorem 3.14.10) says that on $\mathbb{R}^n$, every closed 1-form is exact. More generally, on a star-shaped open set, every closed k-form with $k \geq 1$ is exact. On other sets, this can fail, and the failure detects holes.

Example A.10.25 (The angle form)**.** On $U = \mathbb{R}^2 \setminus \{\mathbf{0}\}$, the 1-form

$$\omega = \frac{-x_2\,dx_1 + x_1\,dx_2}{x_1^2 + x_2^2}$$

is closed: it belongs to the vortex field of Example 3.14.11, whose curl is 0. It is not exact: for the circle $\mathbf{c}(t) = (\cos 2\pi t, \sin 2\pi t)$, the pullback is $\mathbf{c}^*\omega = 2\pi\,dt$, so $\int_{\mathbf{c}} \omega = 2\pi$, while $\int_{\mathbf{c}} df = f(\mathbf{c}(1)) - f(\mathbf{c}(0)) = 0$ for every function f, by Theorem A.10.22, since $\mathbf{c}(1) = \mathbf{c}(0)$. Near every point of U, $\omega = d\theta$ for the polar angle θ, which explains its name; but the angle cannot be defined as a smooth function on all of U. For every closed curve $\mathbf{c}$ in U, the number $\frac{1}{2\pi}\int_{\mathbf{c}} \omega$ is an integer, the winding number of $\mathbf{c}$ around the origin.

Example A.10.26 (The solid angle form)**.** On $U = \mathbb{R}^3 \setminus \{\mathbf{0}\}$, let $\tau = \omega/\|\mathbf{x}\|^3$, where ω is the 2-form of Example A.10.23. It is the 2-form of the vector field $G(\mathbf{x}) = \mathbf{x}/\|\mathbf{x}\|^3$, the electric field of

a point charge (up to a constant), and it is closed, since $\operatorname{div} G = \sum_i \left(\|\mathbf{x}\|^{-3} - 3x_i^2\|\mathbf{x}\|^{-5}\right) = 0$. For a small parallelogram spanned by $\mathbf{v}$ and $\mathbf{w}$ at $\mathbf{x}$, $\tau_{\mathbf{x}}(\mathbf{v}, \mathbf{w})$ is approximately the solid angle under which it is seen from the origin, hence the name. Since $\|\mathbf{x}\| = 1$ on S^2, the pullbacks of τ and ω to S^2 agree, so $\int_{S^2} \tau = 4\pi$. But if $\tau = d\eta$ for a 1-form η on U, then Theorem A.10.21 for $S^2 \subseteq U$, which has empty boundary, would give $\int_{S^2} \tau = 0$. So τ is closed but not exact: it detects the hole of U with a sphere, as the angle form detects the hole of $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ with a circle.

The closed k-forms on U form a vector space, the kernel of d, and the exact k-forms form a subspace of it, the image of d. The quotient space (Section 1.9)

$$H^k(U) = \{\text{closed } k\text{-forms on } U\}/\{\text{exact } k\text{-forms on } U\}$$

is the k-th de Rham cohomology space of U. It measures how far closed forms are from being exact:

- $H^0(U)$ consists of the functions with $df = 0$, i.e., of the functions that are constant on each connected component of U (Section A.4). So $\dim H^0(U)$ is the number of connected components of U.
- By the Poincaré lemma, $H^k(\mathbb{R}^n) = \{0\}$ for $k \geq 1$.
- $H^1(\mathbb{R}^2 \setminus \{\mathbf{0}\})$ is one-dimensional, spanned by the class of the angle form: every closed 1-form η on $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ is $c\,\omega + df$ for some function f, with $c = \frac{1}{2\pi}\int_{\mathbf{c}} \eta$ for the unit circle $\mathbf{c}$.
- $H^2(\mathbb{R}^3 \setminus \{\mathbf{0}\}) \neq \{0\}$, by Example A.10.26. In fact, it is one-dimensional, spanned by the class of the solid angle form, while $H^1(\mathbb{R}^3 \setminus \{\mathbf{0}\}) = \{0\}$: closed curves cannot detect a point removed from $\mathbb{R}^3$.

De Rham's theorem says that these spaces only depend on the topology of U (more precisely, on its homotopy type), although they are defined with derivatives. They can also be defined for manifolds. For a compact manifold M, their dimensions $b_k = \dim H^k(M)$, the Betti numbers, refine the Euler characteristic: $\chi(M) = \sum_k (-1)^k b_k$. Together with the Morse relation (Theorem A.9.8), this leads to the Morse inequalities $c_k \geq b_k$: a Morse function on M has at least b_k critical points of index k. For example, the torus has $b_0 = 1$, $b_1 = 2$ and $b_2 = 1$, so every Morse function on it has at least four critical points, at least two of which are saddle points.

Remark A.10.27 (Differential forms in machine learning). Differential forms appear in machine learning in several places.

1. Probability densities transform like n-forms. If $\mathbf{y} = \boldsymbol{\phi}(\mathbf{x})$ for a diffeomorphism $\boldsymbol{\phi}$ with $\det D\boldsymbol{\phi} > 0$, Proposition A.10.11 gives $\boldsymbol{\phi}^*(p_Y\, dy_1 \wedge \cdots \wedge dy_n) = (p_Y \circ \boldsymbol{\phi}) \det D\boldsymbol{\phi}\, dx_1 \wedge \cdots \wedge dx_n$, which is the change of variables formula $p_X = (p_Y \circ \boldsymbol{\phi}) \det D\boldsymbol{\phi}$ behind normalizing flows (Remark 3.4.8).

2. Hamiltonian Monte Carlo samples from a density proportional to $e^{-U(\mathbf{q})}$ by simulating the motion of a particle with position $\mathbf{q}$ and momentum $\mathbf{p}$ in the potential U. The flow of these equations preserves the 2-form $\sum_i dq_i \wedge dp_i$ (it is symplectic), and hence volume (compare Liouville's formula, Proposition 3.16.1). The leapfrog integrator used in practice is symplectic as well, which keeps the error in the energy small over long trajectories, and the acceptance rate of the method high.

3. On a graph, functions on the nodes, on the edges and on the triangles are discrete analogues of 0-, 1- and 2-forms, the incidence matrices play the role of d, and $d \circ d = 0$ still holds. The resulting decomposition of flows on the edges into a gradient part, a curl part and a harmonic part (Hodge theory) is used, for example, to rank items from pairwise comparisons, and in neural networks on simplicial complexes.

A.11 Topology in Machine Learning

We end with a few places where the ideas of this appendix appear in machine learning. Some of them are theorems, others are well-supported heuristics and active research topics.

- **The manifold hypothesis.** High-dimensional data, such as images with millions of pixels, are believed to lie close to submanifolds of much smaller dimension: the images of a face, for example, vary with a small number of parameters such as pose, lighting and expression. PCA finds the best approximating linear subspace (Lecture 2); autoencoders and other nonlinear methods try to find curved, manifold-like structures.

- **Normalizing flows and topology.** A normalizing flow maps a simple distribution, such as a Gaussian on $\mathbb{R}^n$, to a complicated one by a diffeomorphism (Corollary A.5.5 and Remark 3.4.8). Diffeomorphisms preserve topology. So, by Proposition A.4.2, a flow cannot map $\mathbb{R}^n$, which is connected, exactly onto a distribution supported on two separate pieces, such as two clusters: it can only approximate it by squeezing a little probability mass through thin bridges between them.

- **Symmetries make critical points degenerate.** If a loss is invariant under a continuous family of transformations of the parameters, its minima are not isolated. For example, the ReLU satisfies $\max(0, ct) = c\max(0, t)$ for $c > 0$, so multiplying the incoming weights of a hidden unit by c and its outgoing weights by $\frac{1}{c}$ does not change the network. The set of minima then contains curves (in the simplest case, the hyperbola $w_1 w_2 = 1$ for the loss $(w_1 w_2 - 1)^2$), and by Proposition A.9.4, the Hessian at a minimum must have the eigenvalue 0, in the tangent directions of these curves. Loss functions of neural networks are therefore not Morse functions.

- **Saddle points in high dimensions.** At a critical point of a loss with millions of parameters, all eigenvalues of the Hessian must be nonnegative for a local minimum. A heuristic, supported by experiments and by random matrix theory, says that critical points with high loss usually have many negative eigenvalues, i.e., a large index: they are saddle points rather than bad local minima. Gradient descent with noise, as in SGD, tends to escape from saddle points, which is part of the explanation why training works.

- **Connected sublevel sets.** Experiments show that two minima found by training the same network twice can often be joined by a path along which the loss stays low ("mode connectivity"): the sublevel sets $\{L \leq c\}$ for small c seem to be connected, up to the symmetries of the network. Morse theory gives a language for such statements: a sublevel set can only become connected by passing saddle points of index 1.

A.12 Banach Spaces

In Lectures 2 and 4, we met vector spaces whose elements are functions, such as polynomials, continuous functions and random variables. These spaces are infinite-dimensional, and Remark A.3.11 showed that some results of the previous sections fail in them. Functional analysis studies such spaces. Its starting point is the property that makes $\mathbb{R}^n$ so convenient for analysis: completeness (Corollary A.3.2).

Definition A.12.1 (Normed and Banach spaces)**.** A norm on a real vector space V is a function $\|\cdot\| : V \to [0, \infty)$ such that $\|v\| = 0$ only for $v = 0$, $\|cv\| = |c|\|v\|$ and $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in V$ and $c \in \mathbb{R}$. A normed space is a metric space with the metric $d(u, v) = \|u - v\|$. It is a Banach space if it is complete, i.e., if every Cauchy sequence in it converges.

By Corollary A.3.2 and Theorem A.3.10, every finite-dimensional normed space is a Banach space, since a basis identifies it with $\mathbb{R}^n$.

Example A.12.2 (Banach spaces)**.**

1. The space $C[a,b]$ of continuous functions $f:[a,b]\to\mathbb{R}$ with the maximum norm $\|f\|_\infty = \max_{a\le x\le b}|f(x)|$ is a Banach space (Proposition A.12.3 below). Convergence in this norm is uniform convergence: $\|f_k - f\|_\infty \to 0$ means that $f_k(x)\to f(x)$ at the same speed for all x.

2. For $1\le p<\infty$, the sequences $x=(x_1,x_2,\ldots)$ with $\|x\|_p = (\sum_k |x_k|^p)^{1/p}<\infty$ form a Banach space ℓ^p, and the bounded sequences with $\|x\|_\infty = \sup_k |x_k|$ form a Banach space ℓ^∞.

3. Let $(\Omega,\mathcal{F},\mu)$ be a measure space and $1\le p<\infty$. The measurable functions with $\|f\|_p = (\int |f|^p\,d\mu)^{1/p}<\infty$, where functions that agree almost everywhere are identified, form a Banach space $L^p(\mu)$ (the Riesz–Fischer theorem, which we do not prove). The integral is the Lebesgue integral (Definition B.3.1). For a probability measure, these are the spaces L^p of random variables of Appendix B; for the Lebesgue measure on $[a,b]$, we write $L^p[a,b]$; and for the counting measure on $\{1,2,\ldots\}$, $L^p(\mu)$ is ℓ^p.

4. Not every normed space is complete. On $C[0,1]$, $\|f\|_1 = \int_0^1 |f(x)|\,dx$ is also a norm. The ramps f_k that are 0 on $[0,\frac12]$, 1 on $[\frac12+\frac1k,1]$ and linear in between satisfy $\|f_k - f_l\|_1 \le \frac{1}{2\min(k,l)}$, so they form a Cauchy sequence for this norm. But they converge in this norm to the step function $\mathbf{1}_{(1/2,1]}$, and no continuous function is their limit. So $(C[0,1],\|\cdot\|_1)$ is not complete. Adding the limits of all Cauchy sequences, as the real numbers complete the rational numbers, gives $L^1[0,1]$.

Proposition A.12.3. *$C[a,b]$ with the maximum norm is complete.*

Proof. Let (f_k) be a Cauchy sequence. For each x, $|f_k(x)-f_l(x)|\le\|f_k-f_l\|_\infty$, so $(f_k(x))$ is a Cauchy sequence of real numbers, with a limit $f(x)$. Given $\varepsilon>0$, choose N with $\|f_k-f_l\|_\infty\le\varepsilon$ for $k,l\ge N$. Letting $l\to\infty$ gives $|f_k(x)-f(x)|\le\varepsilon$ for all x and $k\ge N$, so $f_k\to f$ uniformly. It remains to show that f is continuous. Fix x, and choose $\delta>0$ with $|f_N(y)-f_N(x)|<\varepsilon$ for $|y-x|<\delta$. Then for $|y-x|<\delta$,

$$|f(y)-f(x)|\le|f(y)-f_N(y)|+|f_N(y)-f_N(x)|+|f_N(x)-f(x)|<3\varepsilon. \qquad \square$$

In particular, a uniform limit of continuous functions is continuous. A pointwise limit need not be: on $[0,1]$, x^k converges to the function that is 0 for $x<1$ and 1 for $x=1$.

Fixed points and differential equations. The proof of the Banach fixed-point theorem (Theorem A.5.3) only used the triangle inequality and completeness. So the theorem holds, with the same proof, for every contraction $T:C\to C$ of a closed nonempty subset C of a Banach space. This proves the existence theorem for differential equations of Lecture 3.

Theorem A.12.4 (Picard–Lindelöf)**.** *Let $F:\mathbb{R}^n\to\mathbb{R}^n$ be Lipschitz continuous, $\|F(\mathbf{x})-F(\mathbf{y})\|\le L\|\mathbf{x}-\mathbf{y}\|$, and let $T>0$. Then for every $\mathbf{x}_0$, the ODE $\mathbf{x}'=F(\mathbf{x})$ has exactly one solution on $[0,T]$ with $\mathbf{x}(0)=\mathbf{x}_0$. Consequently, it has exactly one solution defined for all $t\in\mathbb{R}$ (Theorem 3.15.2).*

Proof. By the fundamental theorem of calculus, a continuous function $\mathbf{x}:[0,T]\to\mathbb{R}^n$ is a solution if and only if it is a fixed point of the map

$$(\Phi\mathbf{x})(t) = \mathbf{x}_0 + \int_0^t F(\mathbf{x}(s))\,ds,$$

which maps the space $C([0,T],\mathbb{R}^n)$ of continuous functions into itself. We give this space the weighted norm $\|\mathbf{x}\|_w = \max_{0\le t\le T} e^{-2Lt}\|\mathbf{x}(t)\|$. Since $e^{-2LT}\le e^{-2Lt}\le 1$, it is equivalent to the

maximum norm, and the space is complete, as in Proposition A.12.3. For continuous $\mathbf{x}$ and $\mathbf{y}$, and $0 \le t \le T$,

$$\|(\Phi\mathbf{x})(t) - (\Phi\mathbf{y})(t)\| \le \int_0^t L\|\mathbf{x}(s) - \mathbf{y}(s)\|\,ds \le \|\mathbf{x} - \mathbf{y}\|_w \int_0^t Le^{2Ls}\,ds \le \frac{e^{2Lt}}{2}\|\mathbf{x} - \mathbf{y}\|_w.$$

Multiplying by e^{-2Lt} gives $\|\Phi\mathbf{x} - \Phi\mathbf{y}\|_w \le \frac{1}{2}\|\mathbf{x} - \mathbf{y}\|_w$. So Φ is a contraction, and it has exactly one fixed point. By uniqueness, the solutions on intervals $[0, T]$ of different lengths agree where they are both defined, so they form one solution on $[0, \infty)$. For negative times, apply the same argument to $\mathbf{y}(t) = \mathbf{x}(-t)$, which solves $\mathbf{y}' = -F(\mathbf{y})$. □

With the maximum norm, the same estimate only shows that Φ is a contraction if $LT < 1$; the weight e^{-2Lt} removes this restriction. The iteration $\mathbf{x}_{k+1} = \Phi\mathbf{x}_k$ is called Picard iteration. For $x' = x$ with $x(0) = 1$, starting from the constant $x_0(t) = 1$, it gives $x_1(t) = 1 + t$, $x_2(t) = 1 + t + \frac{t^2}{2}$, and in general the Taylor polynomials of the solution e^t.

Bounded operators. Linear maps between infinite-dimensional spaces need not be continuous. A linear map $A : V \to W$ between normed spaces is bounded if its operator norm

$$\|A\| = \sup_{\|v\| \le 1} \|Av\|$$

is finite, and then $\|Av\| \le \|A\|\|v\|$ for all v, as for the matrix norm of Lecture 2. A linear map is continuous if and only if it is bounded: if it is bounded, then $\|Au - Av\| \le \|A\|\|u - v\|$; if it is continuous at 0, there is a $\delta > 0$ with $\|Av\| \le 1$ for $\|v\| \le \delta$, so $\|A\| \le \frac{1}{\delta}$. In finite dimensions, every linear map is bounded.

Example A.12.5.

1. (Integration.) The Volterra operator $(Vf)(x) = \int_0^x f(t)\,dt$ on $C[0, 1]$ with the maximum norm satisfies $|(Vf)(x)| \le x\|f\|_\infty$, so $\|V\| = 1$, with equality for $f = 1$. By induction, $|(V^k f)(x)| \le \frac{x^k}{k!}\|f\|_\infty$, so $\|V^k\| = \frac{1}{k!}$: the powers of V decay much faster than $\|V\|^k = 1$.
2. (Differentiation.) On the polynomials on $[0, 1]$ with the maximum norm, the derivative D is linear but not bounded: the polynomials x^k have the norm 1, but their derivatives kx^{k-1} have the norm k. Differentiation amplifies small wiggles of high frequency, which is why numerical differentiation by finite differences (Proposition 3.2.9) is sensitive to noise.

Dual spaces. The bounded linear functionals $\varphi : V \to \mathbb{R}$ on a normed space V form its dual space V^*, a Banach space with the operator norm. On $\mathbb{R}^n$, every linear functional is $\mathbf{x} \mapsto \mathbf{a}^\top\mathbf{x}$. The Hahn–Banach theorem, which we do not prove, guarantees that every normed space has many bounded linear functionals: for every closed subspace $U \ne V$, there is a $\varphi \in V^*$ with $\varphi \ne 0$ and $\varphi(u) = 0$ for all $u \in U$. The dual space of $C[a, b]$ consists of the integrals $f \mapsto \int f\,d\mu$ with respect to signed measures μ, i.e., differences of two finite measures on the Borel sets (the Riesz–Markov theorem). For example, the evaluation $f \mapsto f(x_0) = \int f\,d\delta_{x_0}$ at a point is a bounded linear functional on $C[a, b]$ with the norm 1. On $L^2[a, b]$, evaluation at a point is not even defined, since the elements of L^2 are only defined up to null sets. The dual of a Hilbert space is the space itself, as we show in the next section.

A.13 Hilbert Spaces

Definition A.13.1 (Hilbert space)**.** An inner product space is a real vector space H with an inner product $\langle\cdot,\cdot\rangle$ (Lecture 2) and the norm $\|v\| = \sqrt{\langle v, v\rangle}$. It is a Hilbert space if it is complete with this norm.

The Cauchy–Schwarz inequality $|\langle u, v\rangle| \le \|u\|\|v\|$ and Pythagoras' theorem hold as in Lecture 2, since their proofs only use the properties of an inner product, and so does the parallelogram law $\|u+v\|^2 + \|u-v\|^2 = 2\|u\|^2 + 2\|v\|^2$ (expand both squares). By the Cauchy–Schwarz inequality, the inner product is continuous.

Example A.13.2 (Hilbert spaces)**.** 1. $\mathbb{R}^n$ with the dot product, and ℓ^2 with $\langle x, y\rangle = \sum_k x_k y_k$, which converges by the Cauchy–Schwarz inequality.

2. $L^2(\mu)$ with $\langle f, g\rangle = \int fg\, d\mu$, for example $L^2[a,b]$ with $\langle f, g\rangle = \int_a^b f(x)g(x)\, dx$. For a probability space, it is the space L^2 of random variables with finite variance and the inner product $\langle X, Y\rangle = \mathbb{E}[XY]$ of Lecture 4.

3. None of the other spaces ℓ^p is a Hilbert space, since their norms do not satisfy the parallelogram law; in ℓ^1, for example, the unit sequences $e_1 = (1, 0, 0, \ldots)$ and $e_2 = (0, 1, 0, \ldots)$ give $4 + 4 \ne 2 + 2$. And $C[0,1]$ with the inner product of $L^2[0,1]$ is an inner product space that is not complete, by the argument of Example A.12.2(4).

The most important property of Hilbert spaces is that we can project onto closed subspaces, as in Lecture 2.

Theorem A.13.3 (Projection theorem)**.** *Let C be a nonempty closed convex subset of a Hilbert space H. Then every $x \in H$ has exactly one closest point $P_C x$ in C. If $C = U$ is a closed subspace, then $P_U x$ is the only $p \in U$ with $x - p \perp U$, and every $x \in H$ can be written in exactly one way as $x = u + w$ with $u \in U$ and $w \perp U$.*

Proof. Let $d = \inf_{c\in C}\|x - c\|$, and choose $c_k \in C$ with $\|x - c_k\| \to d$. The parallelogram law for $x - c_k$ and $x - c_l$ gives

$$\|c_k - c_l\|^2 = 2\|x - c_k\|^2 + 2\|x - c_l\|^2 - 4\left\|x - \frac{c_k + c_l}{2}\right\|^2 \le 2\|x - c_k\|^2 + 2\|x - c_l\|^2 - 4d^2,$$

since $\frac{c_k+c_l}{2} \in C$ by convexity. The right side tends to 0 as $k, l \to \infty$, so (c_k) is a Cauchy sequence. Its limit c lies in C, since C is closed, and $\|x - c\| = d$ by the continuity of the norm. If c' is another closest point, the same inequality for c and c' gives $\|c - c'\|^2 \le 0$. Now let $C = U$ be a closed subspace, and $p = P_U x$. For $u \in U$, the function $t \mapsto \|x - p - tu\|^2 = \|x - p\|^2 - 2t\langle x - p, u\rangle + t^2\|u\|^2$ is minimal at $t = 0$, so $\langle x - p, u\rangle = 0$. Conversely, if $p' \in U$ and $x - p' \perp U$, then $\|x - u\|^2 = \|x - p'\|^2 + \|p' - u\|^2$ for all $u \in U$ by Pythagoras, so p' is the closest point. Finally, $x = P_U x + (x - P_U x)$, and this decomposition is unique since $U \cap U^\perp = \{0\}$. □

For a subspace U of $\mathbb{R}^n$, this is the least squares projection of Lecture 2. The subspace must be closed: the polynomials form a subspace of $L^2[0,1]$ that is not closed, and the function e^x has the distance 0 from it (by the Weierstrass approximation theorem, Section A.16), but no closest polynomial. One can show that the random variables in L^2 that are measurable with respect to a σ-algebra $\mathcal{G}$ form a closed subspace, and the projection onto it is the conditional expectation $\mathbb{E}[Y \mid \mathcal{G}]$ (Theorem B.7.2).

Theorem A.13.4 (Riesz representation)**.** *For every bounded linear functional φ on a Hilbert space H, there is exactly one $y \in H$ with $\varphi(x) = \langle x, y\rangle$ for all $x \in H$, and $\|\varphi\| = \|y\|$.*

Proof. If $\varphi = 0$, take $y = 0$. Otherwise, the kernel $U = \{x : \varphi(x) = 0\}$ is a closed subspace (the preimage of $\{0\}$ under a continuous map), and $U \ne H$. By the projection theorem, there is a $z \ne 0$ with $z \perp U$ (take $z = v - P_U v$ for some $v \notin U$), and we may scale it so that $\varphi(z) = 1$. For every x, the vector $x - \varphi(x)z$ lies in U, so it is orthogonal to z, i.e., $\langle x, z\rangle = \varphi(x)\|z\|^2$. So $y = \frac{z}{\|z\|^2}$ has the required property. If y' has it too, then $\langle x, y - y'\rangle = 0$ for all x, in particular for $x = y - y'$, so $y = y'$. Finally, $|\varphi(x)| \le \|x\|\|y\|$ by the Cauchy–Schwarz inequality, with equality for $x = y$. □

The representing vector depends on the inner product. This is why the gradient depends on the inner product (Remark 3.3.8): the derivative $Df(\mathbf{x})$ is a linear functional, and the gradient is its Riesz representer. The Riesz representation theorem also proves the Radon–Nikodym theorem of Appendix B, by an argument of von Neumann.

Theorem A.13.5 (Radon–Nikodym)**.** *Let μ and ν be finite measures on a measurable space $(\Omega, \mathcal{F})$ such that $\mu(A) = 0$ implies $\nu(A) = 0$. Then there is a measurable $p \geq 0$ with $\nu(A) = \int_A p\, d\mu$ for all $A \in \mathcal{F}$.*

Sketch of proof. Let $\rho = \mu + \nu$. For $f \in L^2(\rho)$, the Cauchy–Schwarz inequality gives $|\int f\, d\nu| \leq \int |f|\, d\rho \leq \rho(\Omega)^{1/2}\|f\|_{L^2(\rho)}$, so $f \mapsto \int f\, d\nu$ is a bounded linear functional on $L^2(\rho)$. By the Riesz representation theorem, there is a $g \in L^2(\rho)$ with

$$\int f\, d\nu = \int fg\, d\rho = \int fg\, d\mu + \int fg\, d\nu \qquad \text{for all } f \in L^2(\rho).$$

For $f = \mathbf{1}_A$, this gives $\nu(A) = \int_A g\, d\rho$, and since $0 \leq \nu(A) \leq \rho(A)$ for all A, one shows that $0 \leq g \leq 1$ almost everywhere. Rearranging, $\int f(1-g)\, d\nu = \int fg\, d\mu$. For $f = \mathbf{1}_N$ with $N = \{g = 1\}$, the left side is 0, so $\mu(N) = 0$, and hence also $\nu(N) = 0$. Now let $A \in \mathcal{F}$, and take $f = \mathbf{1}_A(1 + g + \cdots + g^k)$, so that $f(1-g) = \mathbf{1}_A(1 - g^{k+1})$:

$$\int_A (1 - g^{k+1})\, d\nu = \int_A (g + g^2 + \cdots + g^{k+1})\, d\mu.$$

Outside the null set N, $g^{k+1} \to 0$ as $k \to \infty$, and the monotone convergence theorem (Theorem B.4.1) on both sides gives $\nu(A) = \int_A \frac{g}{1-g}\, d\mu$. So $p = \frac{g}{1-g}$ is the density. □

This is Theorem B.5.2 for finite measures; the σ-finite case follows by splitting Ω into countably many pieces of finite measure.

Orthonormal bases. In infinite dimensions, a basis in the sense of Lecture 1, in which every vector is a finite linear combination of basis vectors, is not useful. Instead, we allow infinite sums.

Theorem A.13.6 (Orthonormal bases)**.** *Let $e_1, e_2, \ldots$ be orthonormal vectors in a Hilbert space H, and let $x \in H$.*

1. *(Bessel's inequality) $\sum_k \langle x, e_k\rangle^2 \leq \|x\|^2$.*
2. *The series $\sum_k \langle x, e_k\rangle e_k$ converges, and its sum is the orthogonal projection of x onto the closure U of $\operatorname{span}\{e_1, e_2, \ldots\}$.*
3. *If $U = H$, then $x = \sum_k \langle x, e_k\rangle e_k$ and $\|x\|^2 = \sum_k \langle x, e_k\rangle^2$ (Parseval's identity) for every $x \in H$. Then $e_1, e_2, \ldots$ is called an orthonormal basis of H.*

Proof. Let $c_k = \langle x, e_k\rangle$ and $s_n = \sum_{k \leq n} c_k e_k$. Then $x - s_n \perp e_1, \ldots, e_n$, so Pythagoras gives $\|x\|^2 = \|x - s_n\|^2 + \sum_{k \leq n} c_k^2$, which proves (1). For $m > n$, $\|s_m - s_n\|^2 = \sum_{k=n+1}^m c_k^2$, which tends to 0 as $n \to \infty$ by (1). So (s_n) is a Cauchy sequence, with a limit $s \in U$. By the continuity of the inner product, $\langle x - s, e_k\rangle = \lim_n \langle x - s_n, e_k\rangle = 0$ for all k, so $x - s$ is orthogonal to the span of the e_k, and, again by continuity, to its closure U. By Theorem A.13.3, $s = P_U x$. If $U = H$, then $s = x$, so $\|x - s_n\| \to 0$, and the identity $\|x\|^2 = \|x - s_n\|^2 + \sum_{k \leq n} c_k^2$ gives Parseval's identity in the limit. □

So in a Hilbert space with an orthonormal basis, a vector behaves like the sequence of its coefficients: the map $x \mapsto (\langle x, e_k\rangle)_k$ is a linear bijection from H onto ℓ^2 that preserves inner products. One can show that every Hilbert space with a countable dense subset, such as $L^2[a, b]$, has an orthonormal basis, by the Gram–Schmidt process of Lecture 2 applied to a dense sequence. So all

these spaces of infinite dimension are copies of ℓ^2. For example, the Gram–Schmidt process applied to $1, x, x^2, \dots$ in $L^2[-1,1]$ gives the normalized Legendre polynomials $\frac{1}{\sqrt{2}}$, $\sqrt{\frac{3}{2}}\,x$, $\sqrt{\frac{5}{8}}\,(3x^2-1)$, …, which form an orthonormal basis, since the polynomials are dense in $L^2[-1,1]$.

Example A.13.7 (Fourier series)**.** The functions $\frac{1}{\sqrt{2\pi}}$, $\frac{1}{\sqrt{\pi}}\cos kx$ and $\frac{1}{\sqrt{\pi}}\sin kx$ for $k = 1, 2, \dots$ are orthonormal in $L^2[-\pi,\pi]$, by the product formulas for sines and cosines, and one can show that they form an orthonormal basis. The expansion of a function f in this basis is its Fourier series,

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx), \qquad a_k = \frac{1}{\pi}\int_{-\pi}^{\pi} f(x)\cos kx\,dx, \quad b_k = \frac{1}{\pi}\int_{-\pi}^{\pi} f(x)\sin kx\,dx,$$

which converges in $L^2[-\pi,\pi]$. For $f(x) = x$, integration by parts gives $a_k = 0$ and $b_k = \frac{2(-1)^{k+1}}{k}$, so

$$x = 2\Big(\sin x - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \cdots\Big) \qquad \text{in } L^2[-\pi,\pi].$$

The coefficients with respect to the orthonormal functions $\frac{1}{\sqrt{\pi}}\sin kx$ are $\sqrt{\pi}\,b_k = \frac{2\sqrt{\pi}(-1)^{k+1}}{k}$, so Parseval's identity gives $\int_{-\pi}^{\pi} x^2\,dx = \frac{2\pi^3}{3} = \sum_k \frac{4\pi}{k^2}$, i.e.,

$$\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}.$$

The partial sums converge to f in L^2, but not uniformly: near the jumps of the periodic extension of f at $\pm\pi$, they overshoot by about 9% of the jump, however many terms we take (the Gibbs phenomenon). The Fourier coefficients of smooth periodic functions decay quickly, so that a few of them describe the function well. Compression methods for sound and images, such as MP3 and JPEG, are based on this idea, with related bases.

A.14 Compact Operators and the Spectral Theorem

The spectral theorem of Lecture 2 says that a symmetric matrix has an orthonormal basis of eigenvectors. In infinite dimensions, a bounded symmetric operator need not have any eigenvectors: the multiplication operator $(Mf)(x) = xf(x)$ on $L^2[0,1]$ satisfies $\langle Mf, g\rangle = \langle f, Mg\rangle$, but $Mf = \lambda f$ forces $f = 0$ outside the null set $\{\lambda\}$, i.e., $f = 0$ in L^2. The spectral theorem holds for operators that are close to matrices in the following sense.

Definition A.14.1 (Compact and self-adjoint operators)**.** A bounded linear operator $K : H \to H$ on a Hilbert space is compact if for every bounded sequence (x_k) in H, the sequence (Kx_k) has a convergent subsequence. It is self-adjoint if $\langle Kx, y\rangle = \langle x, Ky\rangle$ for all $x, y \in H$, and positive semidefinite if moreover $\langle Kx, x\rangle \geq 0$ for all x.

By the Bolzano–Weierstrass theorem (Theorem A.3.1), every bounded operator whose image is finite-dimensional is compact. The identity of an infinite-dimensional Hilbert space is not compact (Remark A.3.11).

Theorem A.14.2 (Spectral theorem for compact operators)**.** *Let K be a compact self-adjoint operator on a Hilbert space H. Then there are finitely or countably many orthonormal eigenvectors $e_1, e_2, \dots$ of K with nonzero eigenvalues $\lambda_1, \lambda_2, \dots$, ordered so that $|\lambda_1| \geq |\lambda_2| \geq \cdots$, such that*

$$Kx = \sum_k \lambda_k \langle x, e_k\rangle e_k \qquad \textit{for all } x \in H.$$

If there are infinitely many, then $\lambda_k \to 0$, and the eigenspace of each nonzero eigenvalue is finite-dimensional. If K is positive semidefinite, all λ_k are positive.

Idea of proof. One shows that $|\langle Kx, x\rangle|$ attains its maximum on the unit ball, by compactness (this replaces the compactness of the unit sphere in $\mathbb{R}^n$), and that a maximizer is an eigenvector with the eigenvalue $\pm\|K\|$, as for the Rayleigh quotient in Lecture 2. Then one repeats this on the orthogonal complement of the eigenvectors found so far, which K maps into itself. If there were infinitely many orthonormal eigenvectors with $|\lambda_k| \geq \varepsilon > 0$, then $\|Ke_k - Ke_l\|^2 = \lambda_k^2 + \lambda_l^2 \geq 2\varepsilon^2$ for $k \neq l$, so the sequence (Ke_k) would have no convergent subsequence, although (e_k) is bounded. This shows that $\lambda_k \to 0$, and that the eigenspaces of nonzero eigenvalues are finite-dimensional. □

Integral operators. Let $k : [a,b]^2 \to \mathbb{R}$ be continuous and symmetric, $k(x,y) = k(y,x)$. The integral operator

$$(Kf)(x) = \int_a^b k(x,y) f(y)\, dy$$

is the infinite-dimensional analogue of the product of a symmetric matrix with a vector, with $k(x,y)$ in the role of the entries. By Fubini's theorem, it is self-adjoint on $L^2[a,b]$, and one can show that it is compact. We call k positive semidefinite if the matrix $(k(x_i,x_j))_{i,j}$ is positive semidefinite for all points $x_1, \ldots, x_n \in [a,b]$, as for the kernels of Gaussian processes in Lecture 4; then K is positive semidefinite.

Theorem A.14.3 (Mercer)**.** *Let k be continuous, symmetric and positive semidefinite on $[a,b]^2$, and let λ_k and e_k be the positive eigenvalues and the orthonormal eigenfunctions of K. Then the e_k are continuous, and*

$$k(x,y) = \sum_k \lambda_k e_k(x) e_k(y),$$

with absolute and uniform convergence on $[a,b]^2$. In particular, $\sum_k \lambda_k = \int_a^b k(x,x)\, dx$.

We do not prove Mercer's theorem. It says that $k(x,y) = \langle \boldsymbol{\phi}(x), \boldsymbol{\phi}(y)\rangle$ is the inner product in ℓ^2 of the features $\boldsymbol{\phi}(x) = (\sqrt{\lambda_k}\, e_k(x))_k$: every continuous positive semidefinite kernel is an inner product of (possibly infinitely many) features, as the kernels of the kernel trick in Lecture 2.

Example A.14.4 (The Karhunen–Loève expansion of Brownian motion)**.** The covariance function $k(s,t) = \min(s,t)$ of Brownian motion on $[0,1]$ (Proposition 4.15.2) is continuous, symmetric and positive semidefinite, since covariance matrices are positive semidefinite. Its integral operator is

$$(Kf)(t) = \int_0^t s f(s)\, ds + t \int_t^1 f(s)\, ds.$$

Differentiating, $(Kf)'(t) = tf(t) + \int_t^1 f(s)\, ds - tf(t) = \int_t^1 f(s)\, ds$, so $(Kf)'' = -f$, $(Kf)(0) = 0$ and $(Kf)'(1) = 0$. Hence an eigenfunction with $Ke = \lambda e$ and $\lambda > 0$ satisfies $\lambda e'' = -e$ with $e(0) = 0$ and $e'(1) = 0$. So $e(t)$ is a multiple of $\sin(t/\sqrt{\lambda})$ with $\cos(1/\sqrt{\lambda}) = 0$, which gives the normalized eigenfunctions and the eigenvalues

$$e_k(t) = \sqrt{2}\sin\left((k - \tfrac{1}{2})\pi t\right), \qquad \lambda_k = \frac{1}{(k-\frac{1}{2})^2\pi^2}, \qquad k = 1, 2, \ldots$$

As Mercer's theorem predicts, $\sum_k \lambda_k = \frac{4}{\pi^2}\sum_k \frac{1}{(2k-1)^2} = \frac{4}{\pi^2}\cdot\frac{\pi^2}{8} = \frac{1}{2} = \int_0^1 \min(t,t)\, dt$, where the sum over the odd numbers is $\frac{\pi^2}{6} - \frac{1}{4}\cdot\frac{\pi^2}{6}$, by Example A.13.7. Now let $Z_1, Z_2, \ldots$ be i.i.d. $\mathcal{N}(0,1)$, and

$$B(t) = \sum_{k=1}^{\infty} \sqrt{\lambda_k}\, Z_k e_k(t).$$

By independence and Mercer's theorem, $\mathbb{E}[B(s)B(t)] = \sum_k \lambda_k e_k(s) e_k(t) = \min(s,t)$: B has the covariance function of Brownian motion, and one can show that the series converges and defines a Brownian motion. This is the Karhunen–Loève expansion. It writes a random function as a series

with independent coefficients, ordered by their variances: it is the principal component analysis (Lecture 2) of a random function. The first term carries the fraction $\frac{\lambda_1}{1/2} = \frac{8}{\pi^2} \approx 81\%$ of the total variance $\mathbb{E}\big[\int_0^1 B(t)^2\,dt\big] = \frac{1}{2}$, and the first three terms carry 93%.

A.15 Reproducing Kernel Hilbert Spaces

Kernel methods and Gaussian processes (Lecture 4) work with functions through a kernel. The natural Hilbert spaces for them are spaces of functions in which evaluating a function at a point is continuous, unlike in L^2.

Definition A.15.1 (Reproducing kernel Hilbert space)**.** Let X be a set. A Hilbert space H of functions $f : X \to \mathbb{R}$ is a reproducing kernel Hilbert space (RKHS) if for every $x \in X$, the evaluation $f \mapsto f(x)$ is a bounded linear functional on H.

By the Riesz representation theorem, for every $x \in X$ there is a function $k_x \in H$ with

$$f(x) = \langle f, k_x \rangle \qquad \text{for all } f \in H,$$

the reproducing property. The function $k(x, y) = \langle k_x, k_y \rangle = k_y(x)$ is called the reproducing kernel of H. It is symmetric, and it is positive semidefinite: for points $x_1, \dots, x_n$ and $\mathbf{a} \in \mathbb{R}^n$,

$$\sum_{i,j} a_i a_j k(x_i, x_j) = \Big\| \sum_i a_i k_{x_i} \Big\|^2 \geq 0.$$

Conversely, by the Moore–Aronszajn theorem, which we do not prove, every symmetric positive semidefinite kernel k on X is the reproducing kernel of exactly one RKHS H_k. It is the completion of the space of the finite combinations $\sum_i a_i k_{x_i}$ with the inner product $\langle \sum_i a_i k_{x_i}, \sum_j b_j k_{y_j} \rangle = \sum_{i,j} a_i b_j k(x_i, y_j)$. In particular, $\|\sum_i a_i k_{x_i}\|^2 = \mathbf{a}^\top K \mathbf{a}$, with the kernel matrix $K_{ij} = k(x_i, x_j)$.

Example A.15.2.

1. (Linear kernel.) For $k(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top \mathbf{y}$ on $\mathbb{R}^d$, H_k consists of the linear functions $f_{\mathbf{w}}(\mathbf{x}) = \mathbf{w}^\top \mathbf{x}$, with $\langle f_{\mathbf{v}}, f_{\mathbf{w}} \rangle = \mathbf{v}^\top \mathbf{w}$: indeed, $k_{\mathbf{x}} = f_{\mathbf{x}}$, and $\langle f_{\mathbf{w}}, k_{\mathbf{x}} \rangle = \mathbf{w}^\top \mathbf{x} = f_{\mathbf{w}}(\mathbf{x})$. So the RKHS norm $\|f_{\mathbf{w}}\| = \|\mathbf{w}\|$ of a linear function is the norm of its weight vector, which ridge regression penalizes.

2. (Feature maps.) Similarly, for $k(x, y) = \boldsymbol{\phi}(x)^\top \boldsymbol{\phi}(y)$ with a feature map $\boldsymbol{\phi} : X \to \mathbb{R}^m$, H_k consists of the functions $f_{\mathbf{w}}(x) = \mathbf{w}^\top \boldsymbol{\phi}(x)$, and $\|f\|$ is the smallest norm of a $\mathbf{w}$ with $f = f_{\mathbf{w}}$. For a kernel as in Mercer's theorem, with the features $\sqrt{\lambda_k}\, e_k(x)$, H_k consists of the functions $f = \sum_k c_k e_k$ with $\|f\|^2 = \sum_k \frac{c_k^2}{\lambda_k} < \infty$. Since $\lambda_k \to 0$, the coefficients of the functions in H_k decay quickly.

3. (Smoothness.) For the kernel $\min(s, t)$ on $[0, 1]$, H_k consists of the functions $f(t) = \int_0^t g(s)\,ds$ with $g \in L^2[0, 1]$, with $\langle f_1, f_2 \rangle = \int_0^1 f_1'(s) f_2'(s)\,ds$. Indeed, $k_s(t) = \min(s, t)$ has the derivative $\mathbf{1}_{[0,s]}$, so $\langle f, k_s \rangle = \int_0^s f'(u)\,du = f(s)$. The RKHS norm $\|f\| = \|f'\|_{L^2}$ measures how much f varies. Similarly, one can show that all functions in the RKHS of the squared exponential kernel are infinitely differentiable.

The RKHS norm is a natural regularizer for learning a function from data, and the following theorem makes such learning problems finite-dimensional.

Theorem A.15.3 (Representer theorem)**.** *Let H be an RKHS on X with the kernel k, and let $\lambda > 0$. For any points $x_1, \dots, x_N$ in X and any function $L : \mathbb{R}^N \to \mathbb{R}$, every minimizer f over H of*

$$L(f(x_1), \dots, f(x_N)) + \lambda \|f\|^2$$

has the form $f = \sum_{i=1}^N \alpha_i k_{x_i}$ with $\alpha_1, \dots, \alpha_N \in \mathbb{R}$.

Proof. The span U of $k_{x_1}, \ldots, k_{x_N}$ is finite-dimensional, hence closed. Write $f = u + w$ with $u \in U$ and $w \perp U$ (Theorem A.13.3). Then $f(x_i) = \langle f, k_{x_i} \rangle = \langle u, k_{x_i} \rangle = u(x_i)$, so the first term is the same for f and u, while $\|f\|^2 = \|u\|^2 + \|w\|^2$. If $w \neq 0$, then u has a smaller objective than f, so f is not a minimizer. □

So an optimization problem over an infinite-dimensional space of functions reduces to the N coefficients α_i.

Example A.15.4 (Kernel ridge regression)**.** For the squared loss, we minimize $\sum_{i=1}^N (y_i - f(x_i))^2 + \lambda \|f\|^2$. By the representer theorem, $f = \sum_j \alpha_j k_{x_j}$, so $f(x_i) = (K\boldsymbol{\alpha})_i$ and $\|f\|^2 = \boldsymbol{\alpha}^\top K \boldsymbol{\alpha}$, and we have to minimize $\|\mathbf{y} - K\boldsymbol{\alpha}\|^2 + \lambda \boldsymbol{\alpha}^\top K \boldsymbol{\alpha}$ over $\boldsymbol{\alpha} \in \mathbb{R}^N$. This convex function has the gradient $2K\big((K + \lambda I)\boldsymbol{\alpha} - \mathbf{y}\big)$, which vanishes for $\boldsymbol{\alpha} = (K + \lambda I)^{-1}\mathbf{y}$. So the prediction at a new point x_* is

$$f(x_*) = \mathbf{k}_*^\top (K + \lambda I)^{-1} \mathbf{y}, \qquad \mathbf{k}_* = (k(x_1, x_*), \ldots, k(x_N, x_*))^\top,$$

the posterior mean of Gaussian process regression with the noise variance $\sigma^2 = \lambda$ (Proposition 4.12.11). For the data of Example 4.12.12, with $\lambda = \sigma^2 = 0.1$, the coefficients are $\boldsymbol{\alpha} = (1.306, -0.720)$, the prediction at $x_* = 0.5$ is $f(0.5) = 0.517$, and the RKHS norm of the regression function is given by $\|f\|^2 = \boldsymbol{\alpha}^\top K \boldsymbol{\alpha} \approx 1.084$.

A.16 Functional Analysis in Machine Learning

- **Universal approximation.** By the Weierstrass approximation theorem, which we do not prove, the polynomials are dense in $C[a, b]$: every continuous function is a uniform limit of polynomials. Since $\|f\|_{L^2} \leq \sqrt{b-a}\, \|f\|_\infty$, and one can show that the continuous functions are dense in $L^2[a, b]$, the polynomials are also dense in $L^2[a, b]$. Neural networks have the same property. Cybenko's theorem (1989) says that for a continuous function σ with $\sigma(t) \to 0$ as $t \to -\infty$ and $\sigma(t) \to 1$ as $t \to \infty$, such as the logistic sigmoid, the networks $\sum_{i=1}^m c_i \sigma(\mathbf{w}_i^\top \mathbf{x} + b_i)$ with one hidden layer are dense in $C(Q)$, the continuous functions on a box $Q \subseteq \mathbb{R}^n$ with the maximum norm. The proof is functional analysis. If the closure U of the subspace of these networks were not all of $C(Q)$, the Hahn–Banach theorem would give a nonzero bounded linear functional that vanishes on U, and by the Riesz–Markov theorem, it would be a signed measure $\mu \neq 0$ with $\int \sigma(\mathbf{w}^\top \mathbf{x} + b)\, d\mu(\mathbf{x}) = 0$ for all $\mathbf{w}$ and b. One then shows that this forces $\mu = 0$. The theorem says nothing about how many neurons are needed, or whether training finds good weights.

- **Kernel methods.** Support vector machines, kernel ridge regression and Gaussian processes work in the RKHS of a kernel, which is often infinite-dimensional. By the representer theorem, all computations only involve the $N \times N$ kernel matrix: this is the kernel trick of Lecture 2. The RKHS norm is the regularizer, and for kernels such as the squared exponential kernel, it penalizes functions that are not smooth.

- **Gaussian processes and their RKHS.** The posterior mean of a Gaussian process lies in the RKHS of its kernel, but the paths of the process are rougher than the functions in the RKHS. For Brownian motion, the RKHS consists of the functions with square-integrable derivatives (Example A.15.2), while the paths are nowhere differentiable (Section 4.15). In terms of the Karhunen–Loève expansion $B = \sum_k \sqrt{\lambda_k} Z_k e_k$, the RKHS norm of a path would be $\sum_k Z_k^2$, which is infinite almost surely, by the strong law of large numbers (Theorem B.9.6).

- **The neural tangent kernel.** For a network $f(\mathbf{x}; \boldsymbol{\theta})$, the kernel

$$\Theta(\mathbf{x}, \mathbf{x}') = \nabla_{\boldsymbol{\theta}} f(\mathbf{x}; \boldsymbol{\theta})^\top \nabla_{\boldsymbol{\theta}} f(\mathbf{x}'; \boldsymbol{\theta})$$

describes how a gradient step on one example changes the prediction at another input. One can show that for infinitely wide networks with a suitably scaled random initialization, this kernel stays constant during training, and that training by gradient descent on the squared loss then amounts to kernel regression with this kernel. This connects deep learning with the kernel methods of this appendix, at least in this limit.

- **Functional gradients and boosting.** Gradient boosting performs gradient descent in a space of functions: at each step, it fits a simple model, such as a small decision tree, to the negative gradient of the loss with respect to the predictions $f(\mathbf{x}_i)$, and adds a small multiple of it to the current model.
- **Neural operators.** Some learning problems map functions to functions, for example the initial condition of a differential equation to the solution at a later time. Neural operators learn such maps between infinite-dimensional spaces, for example by acting on the Fourier coefficients of their inputs (Example A.13.7).

Key Takeaways

- Open sets, closed sets, convergence and continuity only depend on the notion of nearness. A map is continuous if and only if preimages of open sets are open; so sublevel sets $\{f \le c\}$ and level sets of continuous functions are closed. All norms on $\mathbb{R}^n$ define the same open sets.
- A subset of $\mathbb{R}^n$ is compact if and only if it is closed and bounded (Heine–Borel). A continuous function on a nonempty compact set attains its minimum and maximum, and a continuous coercive function on $\mathbb{R}^n$ has a global minimum; regularization makes losses coercive.
- Continuous maps preserve connectedness; this gives the intermediate value theorem, and shows, e.g., that $\mathrm{GL}(n,\mathbb{R})$ has two pieces and that a circle is not homeomorphic to an interval.
- A contraction of a closed set has a unique fixed point, found by iteration (Banach). This proves the inverse function theorem: a map with an invertible Jacobian is locally invertible, with $D(\mathbf{f}^{-1}) = (D\mathbf{f})^{-1}$. Residual layers $\mathbf{x} + \mathbf{g}(\mathbf{x})$ with a contractive $\mathbf{g}$ are invertible.
- The implicit function theorem solves $F(\mathbf{x},\mathbf{y}) = \mathbf{0}$ locally for $\mathbf{y} = \mathbf{g}(\mathbf{x})$ when $D_{\mathbf{y}}F$ is invertible, with $D\mathbf{g} = -D_{\mathbf{y}}F^{-1}D_{\mathbf{x}}F$; this is implicit differentiation, also used for implicit layers and hyperparameter optimization.
- The level set of a regular value of $F : \mathbb{R}^n \to \mathbb{R}^m$ is an $(n-m)$-dimensional submanifold, which is locally a graph. Its tangent space is $\operatorname{null} DF(\mathbf{p})$, and its normal space is spanned by the gradients $\nabla F_i(\mathbf{p})$. Examples: spheres, $\mathrm{SL}(n)$ (dimension n^2-1, tangent space at I: traceless matrices), $\mathrm{O}(n)$ (dimension $\frac{n(n-1)}{2}$, tangent space at I: skew-symmetric matrices).
- A matrix Lie group is a subgroup of $\mathrm{GL}(n,\mathbb{R})$ that is a submanifold, such as $\mathrm{GL}(n,\mathbb{R})$, $\mathrm{SL}(n,\mathbb{R})$, $\mathrm{O}(n)$ and $\mathrm{SO}(n)$; multiplication and inversion are smooth. Its Lie algebra T_IG is closed under the commutator $[A,B] = AB - BA$, and $t \mapsto e^{tH}$ is a one-parameter subgroup; $\det e^{tH} = e^{t\operatorname{tr} H}$, and e^{tK} is a rotation for every skew-symmetric K.
- Lagrange multipliers say that ∇f is normal to the constraint manifold at a constrained extremum. The Riemannian gradient is the projection of ∇f onto the tangent space; Riemannian gradient descent alternates such steps with retractions.
- By Sard's theorem, almost all values of a smooth map are regular. A critical point is nondegenerate if its Hessian is invertible, and its index is the number of negative eigenvalues. Nondegenerate critical points are isolated, and by the Morse lemma f is exactly a sum of $\pm y_i^2$ near them. Almost every linear perturbation of a function is a Morse function.

- Sublevel sets change their topology only at critical values: a critical point of index k attaches a k-cell. On a compact manifold, $\chi(M) = \sum_k (-1)^k c_k$.

- Alternating k-forms on $\mathbb{R}^n$ have the basis $dx_I = dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ with $i_1 < \cdots < i_k$, so $\dim \Lambda^k(\mathbb{R}^n)^* = \binom{n}{k}$, and $(\varphi_1 \wedge \cdots \wedge \varphi_k)(v_1, \ldots, v_k) = \det(\varphi_i(v_j))$. The exterior derivative of $\sum_I f_I\, dx_I$ is $\sum_I df_I \wedge dx_I$, with $d \circ d = 0$; in $\mathbb{R}^3$, it is the gradient, the curl and the divergence. Pullbacks of top-degree forms produce Jacobian determinants, and integrals of forms over k-cubes do not depend on the parametrization (up to orientation). Forms are integrated over compact oriented manifolds with boundary with local parametrizations and partitions of unity. Stokes' theorem $\int_M d\omega = \int_{\partial M} \omega$, with the orientation "outward normal first" on ∂M, contains all the integral theorems of Lecture 3; it is proved for cubes with Fubini's theorem and the fundamental theorem of calculus, and the general case reduces to cubes by pullbacks. Closed forms that are not exact, such as the angle form on $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ and the solid angle form on $\mathbb{R}^3 \setminus \{\mathbf{0}\}$, detect holes; the de Rham cohomology spaces H^k count them.

- A Banach space is a complete normed space, such as $C[a, b]$ with the maximum norm, ℓ^p and L^p. The Banach fixed-point theorem holds in it; with a weighted maximum norm, it proves the existence and uniqueness of solutions of ODEs (Picard–Lindelöf). A linear map is continuous if and only if it is bounded; integration is bounded, differentiation is not.

- In a Hilbert space, every closed convex set has a unique closest point (projection theorem), every bounded linear functional is an inner product with a fixed vector (Riesz), and an orthonormal basis gives the expansion $x = \sum_k \langle x, e_k \rangle e_k$ with Parseval's identity, such as Fourier series. Conditional expectations are projections in L^2, and the Riesz representation theorem proves the Radon–Nikodym theorem.

- A compact self-adjoint operator has orthonormal eigenvectors with eigenvalues that tend to 0, and it is diagonal in them. For a continuous positive semidefinite kernel, Mercer's theorem gives $k(x, y) = \sum_k \lambda_k e_k(x) e_k(y)$, and the Karhunen–Loève expansion of a random function, such as Brownian motion, is its principal component analysis.

- In an RKHS, evaluation is continuous, $f(x) = \langle f, k_x \rangle$, and every positive semidefinite kernel has one. By the representer theorem, regularized learning in an RKHS reduces to N coefficients; kernel ridge regression gives $\boldsymbol{\alpha} = (K + \lambda I)^{-1}\mathbf{y}$, the posterior mean of a Gaussian process. Networks with one hidden layer are dense in the continuous functions (Cybenko).

Exercises

1. Which of the following subsets of $\mathbb{R}^2$ are open, closed, both or neither? Find their interior, closure and boundary: (a) $\{x > 0, y > 0\}$; (b) $\{x^2 + y^2 \leq 1\}$; (c) $\{(x, 0) : 0 \leq x < 1\}$; (d) $\mathbb{Q}^2$, the points with rational coordinates.

2. Show that the union of infinitely many closed sets need not be closed. Show that a set $U \subseteq X$ is open if and only if it is a union of open balls.

3. (a) Show that the set of $m \times n$ matrices of rank at most r is closed in $\mathbb{R}^{m \times n}$. (Hint: the rank is at most r if and only if all $(r + 1) \times (r + 1)$ minors are 0.)
   (b) Show that every square matrix A is a limit of invertible matrices. (Hint: $\det(A + tI)$ is a polynomial in t.) So $\mathrm{GL}(n, \mathbb{R})$ is open and dense in $\mathbb{R}^{n \times n}$.

4. Which of the following sets are compact? (a) The closed unit ball of the ℓ_1 norm in $\mathbb{R}^n$; (b) $\{(x, y) : xy = 1\}$; (c) $\mathrm{SO}(n)$; (d) $\mathrm{GL}(n, \mathbb{R})$; (e) the ellipsoid $\{\mathbf{x} : \mathbf{x}^\top A \mathbf{x} \leq 1\}$ for a symmetric positive definite A; (f) the same set for $A = \mathrm{diag}(1, -1)$.

5. Show that the regularized logistic regression loss $\frac{1}{N}\sum_i \log(1 + e^{-y_i \mathbf{w}^\top \mathbf{x}_i}) + \lambda\|\mathbf{w}\|^2$ with $\lambda > 0$ has a global minimum, and that it is unique. (Use Proposition A.3.6 and Lecture 3.)

6. Let $A \in \mathbb{R}^{n\times n}$. Show that $N(\mathbf{x}) = \|A\mathbf{x}\|$ is a norm if and only if A is invertible, and that the best constants in $c\|\mathbf{x}\| \le N(\mathbf{x}) \le C\|\mathbf{x}\|$ are the smallest and the largest singular value of A (Lecture 2).

7. (a) Show that the unit circle S^1 is path-connected, and that the hyperbola $\{x^2 - y^2 = 1\}$ is not connected.
   (b) Show that $\mathrm{SO}(2)$ is path-connected, and that $\mathrm{O}(2)$ is not connected.
   (c) Show that no continuous injective map from S^1 to $\mathbb{R}$ exists. (Hint: intermediate value theorem.)

8. Use the iteration $x_{k+1} = \cos x_k$ with $x_0 = 1$ to compute the solution of $\cos x = x$ to six decimal places. How many steps does the a priori bound of Theorem A.5.3 require, and how many do you actually need?

9. Show that $f(x) = x + \frac{1}{2}\tanh(x)$ is a homeomorphism of $\mathbb{R}$, and compute $f^{-1}(1)$ with the iteration of Corollary A.5.5, starting at $x_0 = 0$. Is f a diffeomorphism?

10. At which points of the folium $x^3 + y^3 = 3xy$ can the curve not be written locally as a graph $y = g(x)$? Compute $g'(x)$ at the point $(\frac{3}{2}, \frac{3}{2})$.

11. Let $F(x, y, z) = x^2 + y^2 - z^2$. Find all critical points and critical values of F. For which c is $F^{-1}(c)$ a submanifold? Describe the level sets for $c > 0$, $c = 0$ and $c < 0$; which of them are connected?

12. (a) Find the tangent space of the sphere $x^2+y^2+z^2 = 9$ at $(1, 2, 2)$ as null DF, and the tangent plane $(1, 2, 2) + T_{(1,2,2)}M$ as an equation.
    (b) Show that $T_Q\mathrm{O}(n) = \{QK : K^\top = -K\}$ and $T_I\mathrm{SL}(n, \mathbb{R}) = \{H : \operatorname{tr} H = 0\}$ have the dimensions predicted by Example A.7.3.

13. (Lie groups.)
    (a) Show that the upper triangular $n \times n$ matrices with positive diagonal entries form a matrix Lie group of dimension $\frac{n(n+1)}{2}$, whose Lie algebra consists of the upper triangular matrices. (This group appears in the QR and Cholesky decompositions of Lecture 2.)
    (b) For $K_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$, compute K_1^2 and K_1^3, and show that e^{tK_1} is the rotation by the angle t about the x_1-axis.
    (c) With K_2 and K_3 the analogous generators of the rotations about the x_2- and x_3-axes, check that $[K_1, K_2] = K_3$. What does this say about rotations about different axes?

14. (Orthogonal Procrustes problem.) Given $A \in \mathbb{R}^{n\times n}$ with SVD $A = U\Sigma V^\top$, show that $\operatorname{tr}(Q^\top A)$ is maximal over $Q \in \mathrm{O}(n)$ for $Q = UV^\top$. (Hint: $\operatorname{tr}(Q^\top A) = \operatorname{tr}(\Sigma Z)$ with the orthogonal matrix $Z = V^\top Q^\top U$, whose entries are at most 1 in absolute value.) This is how one aligns two point clouds by a rotation or reflection.

15. (a) Find all critical points of the monkey saddle $f(x, y) = x^3 - 3xy^2$. Show that the origin is a degenerate critical point. In polar coordinates, $f = r^3 \cos 3\varphi$; describe the regions where $f > 0$ and $f < 0$.
    (b) For which $a \in \mathbb{R}$ is $x^3 - ax$ a Morse function? Find its critical points and their indices.

(c) Find the critical points and their indices of $f(x,y) = x^4 - 2x^2 + y^2 + \varepsilon x$ for a small $\varepsilon > 0$, and describe how the sublevel sets change as c increases.

16. Let $A = \operatorname{diag}(3,2,1)$ and $f(\mathbf{x}) = \mathbf{x}^\top A\mathbf{x}$ on the sphere S^2. Find all critical points of f and their indices, and check the Morse relation $\chi(S^2) = 2$. Repeat for $A = \operatorname{diag}(4,3,2,1)$ on S^3, whose Euler characteristic is 0.

17. (Alternating forms.)

(a) For $\mathbf{u} = (1,0,0)$, $\mathbf{v} = (1,2,3)$ and $\mathbf{w} = (0,1,4)$, compute $(dx_1 \wedge dx_2)(\mathbf{v},\mathbf{w})$, $(dx_1 \wedge dx_3)(\mathbf{v},\mathbf{w})$ and $(dx_1 \wedge dx_2 \wedge dx_3)(\mathbf{u},\mathbf{v},\mathbf{w})$, and interpret the results as areas and a volume.

(b) Write down a basis of $\Lambda^2(\mathbb{R}^4)^*$.

(c) Show that $\varphi \wedge \varphi = 0$ for every 1-form φ, but that $\omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_4$ satisfies $\omega \wedge \omega = 2\, dx_1 \wedge dx_2 \wedge dx_3 \wedge dx_4 \neq 0$.

18. Compute $d\omega$ for (a) $\omega = x_1 x_2\, dx_1 + x_1^2\, dx_2$ on $\mathbb{R}^2$; (b) $\omega = x_2 x_3\, dx_1 + x_1 x_3\, dx_2 + x_1 x_2\, dx_3$ on $\mathbb{R}^3$; (c) $\omega = x_1\, dx_2 \wedge dx_3 + x_2\, dx_3 \wedge dx_1 + x_3\, dx_1 \wedge dx_2$ on $\mathbb{R}^3$. Which of these forms are closed, and which are exact? Relate the result of (c) to the divergence of the vector field $F(\mathbf{x}) = \mathbf{x}$.

19. (Pullbacks.)

(a) For the spherical coordinates $\boldsymbol{\phi}(r,\theta,\varphi) = (r\sin\theta\cos\varphi,\, r\sin\theta\sin\varphi,\, r\cos\theta)$, show that $\boldsymbol{\phi}^*(dx_1 \wedge dx_2 \wedge dx_3) = r^2 \sin\theta\, dr \wedge d\theta \wedge d\varphi$.

(b) For the curve $\mathbf{c}(t) = (a\cos 2\pi t,\, b\sin 2\pi t)$ with $a, b > 0$, compute $\mathbf{c}^*(x_1\, dx_2 - x_2\, dx_1)$. Use Stokes' theorem for the 2-cube $(r,t) \mapsto (ar\cos 2\pi t,\, br\sin 2\pi t)$ to explain why $\frac{1}{2}\int_{\mathbf{c}}(x_1\, dx_2 - x_2\, dx_1)$ is the area πab of the ellipse. (Which faces of the 2-cube contribute to the boundary integral?)

20. Let ω be the angle form of Example A.10.25.

(a) Show that $d\omega = 0$, and that $\eta = \frac{x_1\, dx_1 + x_2\, dx_2}{x_1^2 + x_2^2}$ is exact on $\mathbb{R}^2 \setminus \{\mathbf{0}\}$.

(b) Compute $\int_{\mathbf{c}} \omega$ for the circle $\mathbf{c}(t) = (3 + \cos 2\pi t,\, \sin 2\pi t)$, which does not go around the origin. (Hint: on the half-plane $x_1 > 0$, $\omega = d\arctan(x_2/x_1)$.)

(c) Show that the integral of ω over the boundary of every 2-cube in $\mathbb{R}^2 \setminus \{\mathbf{0}\}$ is 0. Why does this not contradict $\int_{\mathbf{c}} \omega = 2\pi$ for the unit circle?

21. Verify Stokes' theorem for the 1-form $\omega = -x_2^2\, dx_1 + x_1^2\, dx_2$ and the identity 2-cube of $[0,1]^2$ by computing both sides.

22. (Stokes' theorem on manifolds.)

(a) Let $M = \{\mathbf{x} \in \mathbb{R}^2 : 1 \le \|\mathbf{x}\| \le 2\}$ be the closed annulus, with the standard orientation. Show that M is a submanifold with boundary, describe ∂M with the induced orientation, and verify Theorem A.10.21 for $\omega = x_1\, dx_2$ by computing both sides.

(b) Let ω be a 1-form on $\mathbb{R}^3$. Show in two ways that $\int_{S^2} d\omega = 0$: with Theorem A.10.21 for S^2, and with Theorem A.10.21 for the closed upper and lower hemispheres, whose induced orientations traverse the equator in opposite directions.

23. (a) Show that $f_n(x) = \sqrt{x^2 + \frac{1}{n}}$ converges to $|x|$ uniformly on $[-1,1]$. Conclude that the continuously differentiable functions do not form a closed subset of $C[-1,1]$.

(b) Show that $f_n(x) = x^n$ is a Cauchy sequence in $L^2[0,1]$, but not in $C[0,1]$ with the maximum norm.

24. (A Neumann series.) Let V be the Volterra operator of Example A.12.5. Show that the equation $f = 1 + Vf$, i.e., $f(x) = 1 + \int_0^x f(t)\,dt$, has exactly one solution in $C[0,1]$ (Hint: such an f solves a differential equation), and that the iteration $f_0 = 0$, $f_{k+1} = 1 + Vf_k$ gives the partial sums of the series $\sum_{j\geq 0} V^j 1$, whose terms have the norms $\frac{1}{j!}$. Which function is the solution?

25. (The best quadratic approximation.) Use the normalized Legendre polynomials of Section A.13 to find the polynomial p of degree at most 2 that minimizes $\int_{-1}^{1}(|x| - p(x))^2\,dx$, and compute the minimal value.

26. Compute the Fourier series of $f(x) = x^2$ on $[-\pi, \pi]$, and use Parseval's identity to show that $\sum_{k=1}^{\infty} \frac{1}{k^4} = \frac{\pi^4}{90}$. Why do the partial sums converge uniformly here, unlike for $f(x) = x$? (Hint: compare the decay of the coefficients.)

27. (Riesz representers.)

    (a) Let A be symmetric positive definite, and give $\mathbb{R}^n$ the inner product $\langle \mathbf{x}, \mathbf{y}\rangle_A = \mathbf{x}^\top A\mathbf{y}$. Find the vector that represents the functional $\mathbf{x} \mapsto \mathbf{c}^\top\mathbf{x}$.

    (b) Find the function in $L^2[0,1]$ that represents $f \mapsto \int_0^{1/2} f(x)\,dx$.

    (c) Show that $f \mapsto f(\frac{1}{2})$ is not bounded on $C[0,1]$ with the norm of $L^2[0,1]$. (Hint: narrow tents.)

28. Check by integrating that $e_1(t) = \sqrt{2}\sin\frac{\pi t}{2}$ is an eigenfunction of the integral operator of Example A.14.4 with the eigenvalue $\frac{4}{\pi^2}$, and that $\|e_1\|_{L^2} = 1$.

29. (Minimal-norm interpolation.) Let $0 < t_1 < \cdots < t_N \leq 1$ and $y_1, \ldots, y_N \in \mathbb{R}$. In the RKHS of the kernel $\min(s,t)$ on $[0,1]$ (Example A.15.2), find the function f with the smallest norm such that $f(t_i) = y_i$ for all i. (Hint: as in the proof of the representer theorem, $f = \sum_i \alpha_i k_{t_i}$.) Show that f is the piecewise linear function through the points $(0,0), (t_1, y_1), \ldots, (t_N, y_N)$, constant on $[t_N, 1]$.

30. For the linear kernel $k(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top\mathbf{y}$, show that the prediction $\mathbf{k}_*^\top(K + \lambda I)^{-1}\mathbf{y}$ of kernel ridge regression is $\mathbf{x}_*^\top\hat{\mathbf{w}}$ with the ridge regression solution $\hat{\mathbf{w}} = (X^\top X + \lambda I)^{-1}X^\top\mathbf{y}$, where X has the rows $\mathbf{x}_i^\top$. (Compare with the exercise on Gaussian processes and Bayesian linear regression in Lecture 4.)

Programming Exercises

As in Lecture 3, use NumPy (and Matplotlib for plots), and start each file with

```
import numpy as np
import matplotlib.pyplot as plt
rng = np.random.default_rng(0)
```

P1. Inverting a residual layer. Let `W = rng.normal(size=(5, 5))` and $\mathbf{g}(\mathbf{x}) = c\tanh(W\mathbf{x})$ with $c = 0.9/\|W\|_2$.

- Show that $\mathbf{g}$ is Lipschitz continuous with constant 0.9 (use $|\tanh'| \leq 1$ and Lemma A.5.1).
- For a random $\mathbf{x}$, compute $\mathbf{y} = \mathbf{x} + \mathbf{g}(\mathbf{x})$, and recover $\mathbf{x}$ with the iteration of Corollary A.5.5. Plot the error against the number of iterations on a logarithmic scale, together with the bound of Theorem A.5.3.
- Repeat with $c = 1.5/\|W\|_2$. Does the iteration still converge? Is $\mathbf{x} \mapsto \mathbf{x} + \mathbf{g}(\mathbf{x})$ still injective? (Try to find two points with the same image.)

P2. Riemannian gradient ascent on the sphere. For a random symmetric $A = B + B^\top$ with `B = rng.normal(size=(6, 6))`, maximize $f(\mathbf{x}) = \mathbf{x}^\top A\mathbf{x}$ on S^5 with the Riemannian gradient and the normalization retraction (Section A.8).

- Start from 100 random points, and check that you always reach $\pm\mathbf{q}_1$ (compare with `np.linalg.eigh`). Plot $f(\mathbf{p}_k)$ for a few runs.
- Start exactly at an eigenvector $\mathbf{q}_i$ with $i > 1$. What happens? Now add a perturbation of size 10^{-8}. How many steps does it take to escape from the saddle point, and how does this depend on $\lambda_1 - \lambda_i$?

P3. Orthogonal Procrustes. Generate a point cloud `X = rng.normal(size=(50, 3))`, a random orthogonal matrix Q_0 (from `np.linalg.qr`), and $Y = XQ_0^\top + 0.01\cdot\text{noise}$. Recover Q_0 by maximizing $\operatorname{tr}(Q^\top Y^\top X)$ over $\mathrm{O}(3)$ with the SVD formula of Exercise 14. Then solve the same problem with Riemannian gradient ascent on $\mathrm{O}(3)$: project $G = \nabla_Q$ onto $T_Q\mathrm{O}(3)$ by $Q\,\frac{1}{2}(Q^\top G - G^\top Q)$, take a step, and retract with the Q factor of a QR decomposition (fixing the signs so that the diagonal of R is positive). Compare the results.

P4. Counting components of sublevel sets. For $f(x, y) = x^4 - 2x^2 + y^2 + 0.3x$, evaluate f on a fine grid over $[-2, 2]^2$, and for 50 values of c, count the connected components of the grid points with $f \le c$ (write a breadth-first search over neighboring grid points). Plot the number of components against c, and compare the values of c where it changes with the critical values of f, which you can find with Newton's method applied to $\nabla f = \mathbf{0}$.

P5. Integrating differential forms.

- Write a function that computes $\int_{\mathbf{c}} \omega$ for a 1-form $\omega = \sum_i F_i\,dx_i$ on $\mathbb{R}^n$ and a curve $\mathbf{c} : [0, 1] \to \mathbb{R}^n$, using the pullback $\int_0^1 F(\mathbf{c}(t))^\top \mathbf{c}'(t)\,dt$ and the midpoint rule with N points. Pass F, $\mathbf{c}$ and $\mathbf{c}'$ as Python functions.
- For the angle form of Example A.10.25, compute $\frac{1}{2\pi}\int_{\mathbf{c}} \omega$ for the unit circle, for the circle of radius 1 around $(3, 0)$, for the unit circle traversed twice, $\mathbf{c}(t) = (\cos 4\pi t, \sin 4\pi t)$, and for the curve $\mathbf{c}(t) = \big((2 + \cos 10\pi t)\cos 2\pi t,\ (2 + \cos 10\pi t)\sin 2\pi t\big)$. Check that the results are close to integers, and explain them.
- Verify Stokes' theorem numerically for the 2-cube $\mathbf{c}(u, v) = (u, v, u^2 - v^2)$ in $\mathbb{R}^3$ and the 1-form $\omega = x_3\,dx_1 + x_1\,dx_2$: compute $\int_{\mathbf{c}} d\omega$ from the pullback of $d\omega = dx_3 \wedge dx_1 + dx_1 \wedge dx_2$, and $\int_{\partial\mathbf{c}} \omega$ as the signed sum of four line integrals. Both should be 2.

P6. Picard iteration. Consider the logistic equation $x' = x(1 - x)$ with $x(0) = 0.2$, whose solution is $x(t) = \frac{1}{1+4e^{-t}}$.

- On a grid of 1001 points in $[0, 5]$, implement the Picard iteration $x_{k+1}(t) = 0.2 + \int_0^t x_k(s)(1 - x_k(s))\,ds$ of Theorem A.12.4, starting from $x_0(t) = 0.2$ and computing the integrals with the cumulative trapezoidal rule. Plot the iterates $x_0, \dots, x_8$ together with the solution.
- Plot the maximum error of x_k against k on a logarithmic scale, on $[0, 1]$ and on $[0, 5]$. Where does the iteration converge first, and why?

P7. Fourier series and the Gibbs phenomenon.

- Plot the partial sums of the Fourier series of $f(x) = x$ on $[-\pi, \pi]$ with 5, 20 and 100 terms (Example A.13.7), and compute their errors in the L^2 norm and in the maximum norm. Which error tends to 0?

- For $N = 50, 100, 200, 400, 800$ terms, compute the maximum of the partial sum on a fine grid near π, and the overshoot above π as a fraction of the jump 2π. Compare with the limit ≈ 0.0895.
- Check Parseval's identity numerically, and compare the partial sums of $\sum_k \frac{1}{k^2}$ with $\frac{\pi^2}{6}$. How fast do they converge?

P8. The Karhunen–Loève expansion. Let $t_i = \frac{i}{n}$ for $i = 1, \dots, n$ with $n = 500$.

- The matrix $\frac{1}{n}\big(\min(t_i, t_j)\big)_{i,j}$ approximates the integral operator of Example A.14.4 (why?). Compute its largest eigenvalues with `np.linalg.eigh`, and compare them with $\frac{1}{(k-1/2)^2\pi^2}$. Compare the eigenvectors, scaled by $\sqrt{n}$, with $\sqrt{2}\sin\big((k-\frac{1}{2})\pi t\big)$.
- Sample Brownian paths from the expansion truncated after 5, 20 and 200 terms, and compare them with paths simulated as random walks (Lecture 4). Check $\mathbb{E}[B(0.5)^2] = 0.5$ for the truncated expansions with 10^4 samples.

P9. Kernel ridge regression.

- Implement kernel ridge regression with the squared exponential kernel (Example A.15.4), and check that it gives the posterior mean of your Gaussian process regression in the programming exercises of Lecture 4, with $\lambda = \sigma^2$. Plot the fits for $\lambda = 10^{-4}$, 10^{-2} and 1.
- For the polynomial kernel $k(x, y) = (1 + xy)^2$ on $\mathbb{R}$, show that $k(x, y) = \boldsymbol{\phi}(x)^\top \boldsymbol{\phi}(y)$ with $\boldsymbol{\phi}(x) = (1, \sqrt{2}x, x^2)$, and check numerically that kernel ridge regression gives the same predictions as ridge regression with these features. Compare the running times of the two methods for $N = 10^2, 10^3, 10^4$ data points.

Appendix B

Measure-Theoretic Probability

Introduction

In Lecture 4, we assigned probabilities to events and computed expectations as sums and integrals, without asking which sets of outcomes are events (Remark 4.1.2). This is enough for discrete random variables and for random variables with densities. But some distributions are neither discrete nor continuous, such as the output of a ReLU unit with a Gaussian input, and some questions are about infinitely many random variables at once, such as whether the sample means converge along a sequence of observations. Measure theory treats all of them in a single framework: events form a σ-algebra, probabilities are measures, random variables are measurable functions, and expectations are Lebesgue integrals, with powerful theorems for exchanging limits and expectations. This appendix builds probability theory on this foundation.

- Sections B.1–B.3 introduce σ-algebras and measures, random variables as measurable functions, and the Lebesgue integral.
- Sections B.4–B.6 prove the convergence theorems for integrals, and discuss densities with the Radon–Nikodym theorem, and product measures with the Fubini–Tonelli theorem.
- Sections B.7 and B.8 define the conditional expectation given a σ-algebra, as an orthogonal projection, and study martingales, the mathematical model of fair games.
- Section B.9 compares the different ways in which random variables converge, and proves the strong law of large numbers.

We use the notation of Lecture 4. Some proofs use Hilbert spaces, which are treated in Appendix A.

B.1 Measures and Probability Spaces

A random experiment has a set Ω of possible outcomes, the sample space, and we want to assign probabilities to sets of outcomes, the events. If there are only finitely or countably many outcomes, every set of outcomes can be an event. For experiments with a continuum of outcomes, such as a uniformly random point of $[0, 1]$, this is impossible (Remark B.1.5), and we have to specify which sets are events. Measure theory provides the framework: the events form a σ-algebra, and the probability is a measure on it.

Definition B.1.1 (σ-algebra)**.** A σ-algebra on a set Ω is a collection $\mathcal{F}$ of subsets of Ω such that

1. $\Omega \in \mathcal{F}$;
2. if $A \in \mathcal{F}$, then $A^c = \Omega \setminus A \in \mathcal{F}$;

3. if $A_1, A_2, \ldots \in \mathcal{F}$, then $A_1 \cup A_2 \cup \cdots \in \mathcal{F}$.

The pair $(\Omega, \mathcal{F})$ is called a measurable space, and the sets in $\mathcal{F}$ are called measurable sets, or events.

A σ-algebra contains $\emptyset = \Omega^c$, and since $\bigcap_k A_k = \left(\bigcup_k A_k^c\right)^c$ (De Morgan's law), it is also closed under countable intersections, and under finite unions, intersections and differences $A \setminus B = A \cap B^c$. So the usual set operations, applied countably many times, lead from events to events. The "σ" stands for countable unions; closedness under countable, not only finite, operations is what allows us to take limits.

Example B.1.2 (σ-algebras)**.** 1. The power set 2^Ω, the set of all subsets of Ω, is a σ-algebra; we use it whenever Ω is countable. The smallest σ-algebra on Ω is $\{\emptyset, \Omega\}$.

2. (Information.) Let Ω be partitioned into disjoint sets $B_1, \ldots, B_n$. The unions of some of the B_i form a σ-algebra $\mathcal{G}$ with 2^n elements. It describes the information of an observer who only learns which B_i contains the outcome: an event belongs to $\mathcal{G}$ if and only if this observer can decide whether it has happened. For two dice, an observer who only learns the sum knows whether the event "the sum is even" has happened, but not whether "the first die shows 6" has. We use σ-algebras in this way to describe information in Sections B.7 and B.8.

3. (Generated σ-algebras.) An intersection of σ-algebras on Ω is again a σ-algebra. So for every collection $\mathcal{C}$ of subsets of Ω, there is a smallest σ-algebra $\sigma(\mathcal{C})$ that contains $\mathcal{C}$, namely the intersection of all σ-algebras that contain $\mathcal{C}$. It is called the σ-algebra generated by $\mathcal{C}$.

4. (Borel sets.) The Borel σ-algebra $\mathcal{B}(\mathbb{R}^n)$ is the σ-algebra generated by the open subsets of $\mathbb{R}^n$, and its elements are called Borel sets. They include all open and closed sets, all boxes and all countable sets, and everything that can be built from them by countably many set operations. The Borel σ-algebra is also generated by the boxes $(a_1, b_1] \times \cdots \times (a_n, b_n]$, and for $n = 1$ by the intervals $(-\infty, x]$ with $x \in \mathbb{R}$.

Definition B.1.3 (Measure and probability space)**.** A measure on a measurable space $(\Omega, \mathcal{F})$ is a function $\mu : \mathcal{F} \to [0, \infty]$ with $\mu(\emptyset) = 0$ that is countably additive: if $A_1, A_2, \ldots \in \mathcal{F}$ are pairwise disjoint, then

$$\mu(A_1 \cup A_2 \cup \cdots) = \mu(A_1) + \mu(A_2) + \cdots .$$

The triple $(\Omega, \mathcal{F}, \mu)$ is called a measure space. A probability measure is a measure P with $P(\Omega) = 1$, and a probability space is a measure space $(\Omega, \mathcal{F}, P)$ with a probability measure P.

A probability measure satisfies the axioms of Definition 4.1.1, except that only the sets in $\mathcal{F}$ are events. These are Kolmogorov's axioms of probability (1933). A measure is finite if $\mu(\Omega) < \infty$, and σ-finite if Ω is a countable union of sets of finite measure.

Example B.1.4 (Measures)**.** 1. (Discrete probability.) Let $\Omega = \{\omega_1, \omega_2, \ldots\}$ be countable, and let $p_1, p_2, \ldots \geq 0$ with $\sum_i p_i = 1$. Then $P(A) = \sum_{i:\omega_i \in A} p_i$ defines a probability measure on 2^Ω; it is countably additive, since series of nonnegative terms can be rearranged. If Ω is finite and all outcomes are equally likely, then $P(A) = \frac{|A|}{|\Omega|}$.

2. (Counting and Dirac measures.) The counting measure $\mu(A) = |A|$, the number of elements of A (or ∞), is a σ-finite measure on $2^\mathbb{Z}$. The Dirac measure $\delta_x(A) = \mathbf{1}_A(x)$ is the probability measure of an outcome that is x with certainty.

3. (Lebesgue measure.) There is exactly one measure λ on $\mathcal{B}(\mathbb{R}^n)$ that assigns to every box its volume. We do not prove this theorem; it follows from Carathéodory's extension theorem, which extends a function on boxes to the σ-algebra that they generate. Since $\lambda(\{x\}) = 0$, countable additivity gives $\lambda(A) = 0$ for every countable set A, for example $\lambda(\mathbb{Q}) = 0$. The uniform distribution on $[0, 1]$ is the probability measure $P(A) = \lambda(A)$ on the Borel subsets of $[0, 1]$.

Remark B.1.5 (Why not all subsets?). There is no measure on all subsets of $[0,1)$ that gives every interval its length and is invariant under translations modulo 1. Here is the idea of Vitali's argument (1905). Call $x, y \in [0,1)$ equivalent if $x - y$ is rational, and choose one point from each equivalence class; let V be the set of chosen points (this needs the axiom of choice). The translates $V_q = \{v + q \bmod 1 : v \in V\}$ for the countably many rationals $q \in [0,1)$ are pairwise disjoint and cover $[0,1)$. If V had a measure m, then so would every V_q, and countable additivity would give $1 = m + m + m + \cdots$, which is impossible both for $m = 0$ and for $m > 0$. Sets like V never arise from explicit constructions; all sets that we meet in practice are Borel sets.

The rules of Proposition 4.1.3 hold for the events of every probability space, with the same proofs, and two of them, monotonicity and the union bound, hold for every measure. Measures have one more property, which allows us to take limits.

Proposition B.1.6 (Continuity)**.** *Let $(\Omega, \mathcal{F}, \mu)$ be a measure space, and let $A_1, A_2, \ldots \in \mathcal{F}$.*

1. *If $A_1 \subseteq A_2 \subseteq \cdots$, then $\mu(\bigcup_k A_k) = \lim_{k\to\infty} \mu(A_k)$.*
2. *If $A_1 \supseteq A_2 \supseteq \cdots$ and $\mu(A_1) < \infty$, then $\mu(\bigcap_k A_k) = \lim_{k\to\infty} \mu(A_k)$.*

Proof. (1) Let $A_0 = \emptyset$ and $B_k = A_k \setminus A_{k-1}$. The B_k are disjoint, $A_n = B_1 \cup \cdots \cup B_n$, and the B_k have the same union as the A_k, so $\mu(\bigcup_k A_k) = \sum_k \mu(B_k) = \lim_{n\to\infty} \sum_{k\le n} \mu(B_k) = \lim_{n\to\infty} \mu(A_n)$. (2) The sets $A_1 \setminus A_k$ increase to $A_1 \setminus \bigcap_k A_k$, and $\mu(A_1 \setminus A_k) = \mu(A_1) - \mu(A_k)$, since $\mu(A_1) < \infty$. Now apply (1). □

Without the condition $\mu(A_1) < \infty$, part (2) can fail: the intervals $[k, \infty)$ decrease to $\emptyset$, but all have infinite Lebesgue measure.

An event with probability 1 happens almost surely (a.s.), and an event with probability 0 is a null event. Null events need not be impossible: a uniformly random point of $[0,1]$ is irrational almost surely, since $\lambda(\mathbb{Q} \cap [0,1]) = 0$, although every rational number in $[0,1]$ is a possible outcome.

Definition B.1.7 (Independence)**.** Infinitely many events are independent if any finitely many of them are, in the sense of Definition 4.1.8. More generally, σ-algebras $\mathcal{G}_1, \ldots, \mathcal{G}_n \subseteq \mathcal{F}$ are independent if $P(G_1 \cap \cdots \cap G_n) = P(G_1) \cdots P(G_n)$ for all $G_1 \in \mathcal{G}_1, \ldots, G_n \in \mathcal{G}_n$: they describe pieces of information that tell us nothing about each other.

Example B.1.8 (Infinitely many coin flips)**.** Many questions about random sequences need infinitely many independent events on one probability space. Let $\Omega = [0,1)$ with the uniform distribution, and let $D_1(\omega), D_2(\omega), \ldots \in \{0,1\}$ be the digits of the binary expansion $\omega = \sum_k D_k(\omega) 2^{-k}$ (choosing the expansion that does not end with infinitely many 1s). For every pattern $d_1, \ldots, d_n$, the set of ω whose first n digits are $d_1, \ldots, d_n$ is an interval of length 2^{-n}, so $P(D_1 = d_1, \ldots, D_n = d_n) = 2^{-n}$. Summing over the digits that we do not care about, it follows that the events $\{D_k = 1\}$ are independent, with probability $\frac{1}{2}$ each: the digits behave exactly like independent fair coin flips, with 1 for heads. The events "the first n flips are tails" have the probabilities 2^{-n} and decrease to the event "all flips are tails", so by continuity, its probability is $\lim_n 2^{-n} = 0$: almost surely, heads eventually appears. (The event "all flips are tails" is $\{0\}$, a possible but null event.) In the same way, one can construct a probability space that carries an infinite sequence of independent random variables (Definition B.6.2) with any given distributions; this is a theorem of Kolmogorov, which we do not prove.

B.2 Random Variables and Their Distributions

A random variable is a numerical quantity that depends on the outcome of the experiment. To ask for the probability that it lies in a set B, the set of outcomes for which it does must be an event.

Definition B.2.1 (Random variable and distribution)**.** Let $(\Omega, \mathcal{F}, P)$ be a probability space. A random variable is a measurable function $X : \Omega \to \mathbb{R}$, i.e., a function with

$$\{X \in B\} = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{F} \qquad \text{for every Borel set } B \subseteq \mathbb{R}.$$

Its distribution (or law) is the probability measure $P_X(B) = P(X \in B)$ on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$, and its cumulative distribution function (CDF) is $F(x) = P(X \le x)$.

The distribution P_X is a probability measure, since taking preimages commutes with complements and countable unions; it is called the image (or pushforward) of P under X. To check that X is measurable, it suffices to check $\{X \le x\} \in \mathcal{F}$ for all x: the Borel sets B with $\{X \in B\} \in \mathcal{F}$ form a σ-algebra, and if it contains the intervals $(-\infty, x]$, it contains all Borel sets. One can show that continuous functions of random variables, and sums, products, maxima and limits of sequences of random variables, are again random variables. We use such facts without comment: in practice, every function that we write down is measurable.

Theorem B.2.2 (The CDF determines the distribution)**.** *The CDF F of a random variable X is nondecreasing and right-continuous, with $\lim_{x\to-\infty} F(x) = 0$ and $\lim_{x\to\infty} F(x) = 1$, and $P(a < X \le b) = F(b) - F(a)$ for $a < b$. Random variables with the same CDF have the same distribution. Conversely, every function F with these four properties is the CDF of a random variable.*

Proof of some parts. F is nondecreasing by part (2) of Proposition 4.1.3. If $x_k \downarrow x$, the events $\{X \le x_k\}$ decrease to $\{X \le x\}$, so $F(x_k) \to F(x)$ by continuity (Proposition B.1.6); the limits at $\pm\infty$ follow in the same way from $\bigcap_k \{X \le -k\} = \emptyset$ and $\bigcup_k \{X \le k\} = \Omega$. That the CDF determines P_X follows from a uniqueness theorem of measure theory, which we do not prove: two probability measures that agree on a collection of sets that is closed under intersections, such as the intervals $(-\infty, x]$, agree on the σ-algebra that it generates. The converse follows from inverse transform sampling (Theorem 4.2.7): on the probability space $\big((0,1), \mathcal{B}((0,1)), \lambda\big)$, the random variable $U(\omega) = \omega$ is uniform, and $F^{-1}(U)$ has the CDF F. □

So in practice, we rarely look at the sample space: a random variable is described by its distribution, i.e., by its CDF. The jumps of the CDF are the values that have positive probability: since $\{X < x\} = \bigcup_k \{X \le x - \frac{1}{k}\}$, continuity gives $P(X < x) = \lim_{y\uparrow x} F(y)$, so $P(X = x) = F(x) - \lim_{y\uparrow x} F(y)$.

The discrete and continuous random variables of Definition 4.2.2 are the two most common kinds. The density p of a continuous random variable has to be measurable, and then $P(X \in B) = \int_B p(x)\,dx$ for every Borel set B, since both sides are probability measures that agree on intervals. Here, and from now on, integrals are Lebesgue integrals (Section B.3), which agree with the integrals of Lecture 3 whenever these converge absolutely. In Section B.5, we will see that pmfs and pdfs are both densities with respect to a measure, and that some distributions have neither.

B.3 The Lebesgue Integral and Expectation

The expectation of a random variable is its average value, weighted by probability. For a random variable with finitely many values $x_1, \dots, x_k$, it is the sum $\sum_i x_i P(X = x_i)$. In general, it is an integral with respect to the probability measure P, the Lebesgue integral. The construction works for every measure, and for the Lebesgue measure on $\mathbb{R}^n$, it extends the integral of Lecture 3.

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. A function $f : \Omega \to \mathbb{R}$ or $f : \Omega \to [0, \infty]$ is measurable if $\{f \le x\} \in \mathcal{F}$ for all $x \in \mathbb{R}$; on a probability space, the measurable real-valued functions are the random variables.

Definition B.3.1 (Lebesgue integral)**.** 1. A simple function is a finite sum $s = \sum_{i=1}^k a_i \mathbf{1}_{A_i}$ with $a_i \ge 0$ and $A_i \in \mathcal{F}$, and its integral is $\int s\,d\mu = \sum_{i=1}^k a_i \mu(A_i)$, with $0 \cdot \infty = 0$.

2. For a measurable $f : \Omega \to [0, \infty]$,
$$\int f \, d\mu = \sup \Big\{ \int s \, d\mu : s \text{ simple}, \, 0 \le s \le f \Big\} \in [0, \infty].$$
3. A measurable $f : \Omega \to \mathbb{R}$ is integrable if $\int |f| \, d\mu < \infty$, and then $\int f \, d\mu = \int f^+ \, d\mu - \int f^- \, d\mu$, where $f^+ = \max(f, 0)$ and $f^- = \max(-f, 0)$, so that $f = f^+ - f^-$ and $|f| = f^+ + f^-$.

For $A \in \mathcal{F}$, we write $\int_A f \, d\mu = \int f \mathbf{1}_A \, d\mu$. Other notations are $\int f(\omega) \, \mu(d\omega)$, and for the Lebesgue measure, $\int f(\mathbf{x}) \, d\mathbf{x}$.

(One checks that the integral of a simple function does not depend on how it is written as a sum.) The Riemann integral of Lecture 3 divides the domain into small boxes. The Lebesgue integral divides the range instead: the simple functions
$$s_n = \min \big(n, \, 2^{-n} \lfloor 2^n f \rfloor\big),$$
which round f down to a multiple of 2^{-n} and cut it off at n, increase to f, and $\int s_n \, d\mu$ adds up the values $j2^{-n}$, weighted by the measures of the sets $\{j2^{-n} \le f < (j+1)2^{-n}\}$. One can show that the integral is linear and monotone ($f \le g$ implies $\int f \, d\mu \le \int g \, d\mu$), that $\int |f| \, d\mu = 0$ if and only if $f = 0$ almost everywhere, i.e., outside a set of measure 0, and that functions which agree almost everywhere have the same integral. For bounded Riemann integrable functions on boxes, and for absolutely convergent improper Riemann integrals, the Lebesgue integral agrees with the Riemann integral. But it integrates many more functions. For example, the indicator function $\mathbf{1}_{\mathbb{Q}}$ of the rationals on $[0, 1]$ is not Riemann integrable, since every Riemann sum can be made 0 or 1 by choosing irrational or rational points, but its Lebesgue integral is $\lambda(\mathbb{Q} \cap [0, 1]) = 0$. For the counting measure on $\mathbb{N}$, the integral is a series: $\int f \, d\mu = \sum_k f(k)$.

Definition B.3.2 (Expectation)**.** The expectation (expected value, or mean) of a random variable X on a probability space $(\Omega, \mathcal{F}, P)$ is
$$\mathbb{E}[X] = \int X \, dP,$$
defined if $X \ge 0$ (then possibly $\mathbb{E}[X] = \infty$) or if X is integrable, i.e., $\mathbb{E}[|X|] < \infty$.

For discrete random variables and random variables with a density, this is the expectation of Definition 4.3.1, by Theorem B.3.3 below. The expectation is linear, $\mathbb{E}[aX + bY + c] = a\mathbb{E}[X] + b\mathbb{E}[Y] + c$, and monotone, since the integral is, and $\mathbb{E}[\mathbf{1}_A] = P(A)$ for every event A. So linearity holds for all integrable random variables; Lecture 4 only proved it for discrete ones. To compute expectations, we do not need the sample space, only the distribution.

Theorem B.3.3 (Expectations from distributions)**.** *Let X be a random variable, and let $g : \mathbb{R} \to \mathbb{R}$ be measurable with $g \ge 0$ or $\mathbb{E}[|g(X)|] < \infty$. Then $\mathbb{E}[g(X)] = \int_{\mathbb{R}} g \, dP_X$. In particular,*
$$\mathbb{E}[g(X)] = \sum_i g(x_i) \, p(x_i) \qquad \text{or} \qquad \mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x) \, p(x) \, dx$$
for discrete X with the pmf p and continuous X with the density p, respectively.

Proof. For $g = \mathbf{1}_B$, both sides are $P(X \in B)$, by the definition of P_X. By linearity, the formula holds for simple functions g, and by the monotone convergence theorem (Theorem B.4.1), applied to simple functions that increase to g, for all $g \ge 0$. For integrable $g(X)$, apply it to g^+ and g^-. The two formulas follow in the same way from Definition B.5.1, with the counting measure and the Lebesgue measure. □

L^p spaces. For $p \geq 1$, the random variables with $\mathbb{E}[|X|^p] < \infty$ form a vector space L^p, and $\|X\|_p = \mathbb{E}[|X|^p]^{1/p}$ is a norm on it, if we do not distinguish random variables that are equal almost surely (the triangle inequality is Minkowski's inequality, which we do not prove). By Jensen's inequality (Theorem 4.3.7) for the convex function $y \mapsto y^{q/p}$ on $[0,\infty)$, $\|X\|_p \leq \|X\|_q$ for $p \leq q$, so $L^q \subseteq L^p$. The space L^2, with the inner product $\mathbb{E}[XY]$ of Lecture 4, is complete, which makes it a Hilbert space (see Appendix A); we use this for the conditional expectation in Section B.7.

B.4 Convergence Theorems

When can we exchange a limit and an expectation? This question comes up constantly: for limits of random variables, for infinite series, and for derivatives of expectations with respect to parameters. The answer is the main advantage of the Lebesgue integral.

Theorem B.4.1 (Monotone convergence)**.** *Let $0 \leq f_1 \leq f_2 \leq \cdots$ be measurable functions with $f_n \to f$ pointwise. Then $\int f_n \, d\mu \to \int f \, d\mu$.*

Proof. By monotonicity, the integrals $\int f_n \, d\mu$ increase to a limit $L \leq \int f \, d\mu$. Conversely, let $s = \sum_i a_i \mathbf{1}_{A_i}$ be simple with $0 \leq s \leq f$, and let $0 < c < 1$. The sets $E_n = \{f_n \geq cs\}$ increase to Ω: where $s > 0$, we have $f_n \to f \geq s > cs$, and where $s = 0$, trivially $f_n \geq cs$. Hence

$$\int f_n \, d\mu \geq \int cs\mathbf{1}_{E_n} \, d\mu = c \sum_i a_i \mu(A_i \cap E_n) \to c \int s \, d\mu,$$

by the continuity of measures (Proposition B.1.6). So $L \geq c \int s \, d\mu$ for all $c < 1$ and all such s, i.e., $L \geq \int f \, d\mu$. □

Applied to partial sums, monotone convergence shows that the expectation of a series of nonnegative random variables is the series of their expectations: $\mathbb{E}\big[\sum_k X_k\big] = \sum_k \mathbb{E}[X_k]$ for $X_k \geq 0$. For example, a random variable X with values in $\{0, 1, 2, \ldots\}$ is the series $\sum_{k \geq 0} \mathbf{1}_{\{X > k\}}$, which gives the tail sum formula $\mathbb{E}[X] = \sum_{k \geq 0} P(X > k)$ of Example 4.3.2.

Theorem B.4.2 (Fatou's lemma and dominated convergence)**.**

1. *(Fatou) Let $f_n \geq 0$ be measurable. Then $\int \liminf_n f_n \, d\mu \leq \liminf_n \int f_n \, d\mu$.*
2. *(Dominated convergence) Let $f_n \to f$ pointwise (or almost everywhere), and let $|f_n| \leq g$ for all n, with an integrable function g. Then f is integrable, and $\int f_n \, d\mu \to \int f \, d\mu$.*

Proof. (1) The functions $g_k = \inf_{n \geq k} f_n$ increase to $\liminf_n f_n$, and $\int g_k \, d\mu \leq \int f_n \, d\mu$ for all $n \geq k$. By monotone convergence, $\int \liminf_n f_n \, d\mu = \lim_k \int g_k \, d\mu \leq \liminf_n \int f_n \, d\mu$. (2) Since $|f| \leq g$, f is integrable. Fatou's lemma, applied to the nonnegative functions $g + f_n$ and $g - f_n$, gives

$$\int g \, d\mu + \int f \, d\mu \leq \int g \, d\mu + \liminf_n \int f_n \, d\mu \qquad \text{and} \qquad \int g \, d\mu - \int f \, d\mu \leq \int g \, d\mu - \limsup_n \int f_n \, d\mu.$$

Hence $\limsup_n \int f_n \, d\mu \leq \int f \, d\mu \leq \liminf_n \int f_n \, d\mu$. □

Example B.4.3. 1. (Mass can escape.) On $[0,1]$ with the Lebesgue measure, let $f_n = n\mathbf{1}_{(0,1/n)}$. Then $f_n(x) \to 0$ for every x, but $\int f_n \, d\lambda = 1$ for all n: the limit of the integrals is not the integral of the limit. In the language of probability, $X_n = n\mathbf{1}_{\{U < 1/n\}}$ for a uniform U is a lottery ticket that wins n euros with probability $\frac{1}{n}$; it is 0 for all large n, but $\mathbb{E}[X_n] = 1$. Dominated convergence does not apply, since $\sup_n f_n(x) \geq \frac{1}{2x}$ for $0 < x \leq \frac{1}{2}$, which is not integrable.

2. (Differentiating under the integral sign.) Let $f(x,\theta)$ be integrable in x and differentiable in θ, with $|\frac{\partial f}{\partial \theta}(x,\theta)| \le g(x)$ for all θ in an open interval, where $\int g\,d\mu < \infty$. Then

$$\frac{d}{d\theta}\int f(x,\theta)\,d\mu(x) = \int \frac{\partial f}{\partial \theta}(x,\theta)\,d\mu(x).$$

Indeed, for $h_n \to 0$, the difference quotients $\frac{f(x,\theta+h_n)-f(x,\theta)}{h_n}$ converge to $\frac{\partial f}{\partial \theta}(x,\theta)$, and by the mean value theorem, they are bounded by $g(x)$; now apply dominated convergence. This proves the Leibniz rule (Proposition 3.14.6) for unbounded regions, under the condition stated there, and for general measures. It also shows that the gradient of an expected loss is the expected gradient, $\nabla_{\boldsymbol{\theta}}\mathbb{E}[\ell(\boldsymbol{\theta};\mathbf{X})] = \mathbb{E}[\nabla_{\boldsymbol{\theta}}\ell(\boldsymbol{\theta};\mathbf{X})]$, if the partial derivatives are dominated in this way. This is why the gradient on a random training example is an unbiased estimate of the gradient of the expected loss.

B.5 Densities and the Radon–Nikodym Theorem

Discrete and continuous distributions look different, but they are two cases of one notion.

Definition B.5.1 (Density with respect to a measure)**.** Let μ be a measure on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$. A random vector $\mathbf{X}$, or its distribution, has the density $p \ge 0$ with respect to μ if

$$P(\mathbf{X} \in B) = \int_B p\,d\mu \qquad \text{for all Borel sets } B.$$

Then $\mathbb{E}[g(\mathbf{X})] = \int g\,p\,d\mu$ for every measurable $g \ge 0$, by the argument of Theorem B.3.3.

For the Lebesgue measure, p is the density of Definition 4.2.2. For the counting measure on a countable set $\{x_1, x_2, \ldots\}$ of values, $\int_B p\,d\mu = \sum_{x_i \in B} p(x_i)$, so p is the pmf. So sums with pmfs and integrals with pdfs are integrals with respect to different reference measures, and everything that we prove for integrals holds for both. A necessary condition for a density is that $\mu(B) = 0$ implies $P(\mathbf{X} \in B) = 0$, since an integral over a null set is 0. The distribution is then called absolutely continuous with respect to μ, and this condition is also sufficient.

Theorem B.5.2 (Radon–Nikodym)**.** *Let μ and ν be σ-finite measures on $(\Omega, \mathcal{F})$ such that $\mu(A) = 0$ implies $\nu(A) = 0$ (written $\nu \ll \mu$). Then there is a measurable $p \ge 0$ with $\nu(A) = \int_A p\,d\mu$ for all $A \in \mathcal{F}$, and p is unique up to changes on a set of μ-measure 0. It is called the Radon–Nikodym derivative $\frac{d\nu}{d\mu}$.*

We do not prove this theorem here; Theorem A.13.5 proves it with the Riesz representation theorem for Hilbert spaces. Not every distribution has a density with respect to the Lebesgue measure or a counting measure.

Example B.5.3 (The ReLU of a Gaussian)**.** Let $Z \sim \mathcal{N}(0,1)$, and let $Y = \max(0, Z)$ be the activation of a ReLU unit with a Gaussian input. Then $P(Y = 0) = P(Z \le 0) = \frac{1}{2}$, and $P(a \le Y \le b) = \int_a^b \varphi(y)\,dy$ for $0 < a < b$. So Y is neither discrete nor continuous: it has an atom of mass $\frac{1}{2}$ at 0, which the Lebesgue measure does not see. But Y has a density with respect to $\mu = \delta_0 + \lambda$, namely $p(0) = \frac{1}{2}$ and $p(y) = \varphi(y)\mathbf{1}_{\{y>0\}}$ for $y \ne 0$. Hence

$$\mathbb{E}[Y] = 0 \cdot \tfrac{1}{2} + \int_0^\infty y\varphi(y)\,dy = \frac{1}{\sqrt{2\pi}} \approx 0.399, \qquad \mathbb{E}[Y^2] = \int_0^\infty y^2\varphi(y)\,dy = \frac{1}{2},$$

so $\mathrm{Var}(Y) = \frac{1}{2} - \frac{1}{2\pi} \approx 0.341$. Mixed distributions of this kind also describe zero-inflated data, such as daily rainfall, or the amount that a visitor of a web shop spends.

Remark B.5.4 (Singular distributions and likelihood ratios). 1. Some distributions have neither atoms nor a density with respect to the Lebesgue measure. An example is the Cantor distribution, the distribution of $X = \sum_{k=1}^{\infty} 2D_k 3^{-k}$ with independent fair coin flips $D_k \in \{0, 1\}$, a random number whose ternary digits are all 0 or 2. Its CDF, the Cantor function, is continuous, so X has no atoms. But X lies in the Cantor set, which has Lebesgue measure 0, so it has no density. One can show that every distribution on $\mathbb{R}$ is a mixture of a discrete, an absolutely continuous, and such a singular part (the Lebesgue decomposition).

2. If $P \ll Q$, the Radon–Nikodym derivative $\frac{dP}{dQ}$ is the likelihood ratio, which is $\frac{p}{q}$ if P and Q have the densities p and q. The weights of importance sampling (Section 4.7.2) are likelihood ratios, $\mathbb{E}_P[f] = \mathbb{E}_Q[f\frac{dP}{dQ}]$, and the KL divergence of Section 4.13 is $\mathrm{KL}(P \,\|\, Q) = \mathbb{E}_P[\log \frac{dP}{dQ}]$ for $P \ll Q$, and ∞ otherwise. So these notions make sense for arbitrary distributions.

B.6 Product Measures and Independence

A random vector $\mathbf{X} = (X_1, \ldots, X_n)$ is a measurable map from Ω to $\mathbb{R}^n$, i.e., $\{\mathbf{X} \in B\} \in \mathcal{F}$ for all Borel sets $B \subseteq \mathbb{R}^n$; one can show that this holds if and only if every X_i is a random variable. Its joint distribution is the probability measure $P_{\mathbf{X}}(B) = P(\mathbf{X} \in B)$ on $\mathbb{R}^n$, and the distributions of the X_i are its marginals. Joint pmfs and joint densities (Definition 4.4.1) are densities of the joint distribution with respect to the counting measure and the Lebesgue measure, and the "reasonable" sets in that definition are the Borel sets. On a product of two measure spaces, there is a natural measure.

Theorem B.6.1 (Product measures and Fubini–Tonelli)**.** *Let μ and ν be σ-finite measures on $(\Omega_1, \mathcal{F}_1)$ and $(\Omega_2, \mathcal{F}_2)$, and let $\mathcal{F}_1 \otimes \mathcal{F}_2$ be the σ-algebra on $\Omega_1 \times \Omega_2$ generated by the products $A \times B$ with $A \in \mathcal{F}_1$ and $B \in \mathcal{F}_2$. There is exactly one measure $\mu \otimes \nu$ on it with $(\mu \otimes \nu)(A \times B) = \mu(A)\nu(B)$, the product measure. For every measurable $f \geq 0$ (Tonelli), and for every f that is integrable with respect to $\mu \otimes \nu$ (Fubini),*

$$\int f \, d(\mu \otimes \nu) = \int \Big(\int f(x, y) \, d\nu(y) \Big) d\mu(x) = \int \Big(\int f(x, y) \, d\mu(x) \Big) d\nu(y).$$

We do not prove this theorem. The Lebesgue measure on $\mathbb{R}^2$ is $\lambda \otimes \lambda$, and the theorem contains Fubini's theorem of Lecture 3, including its condition: for nonnegative functions, the order of integration never matters, and for functions of both signs, it does not matter if $\int |f| < \infty$ (compare with Example 3.14.3, in which the two orders give $\frac{\pi}{4}$ and $-\frac{\pi}{4}$).

Definition B.6.2 (Independent random variables)**.** Random variables $X_1, \ldots, X_n$ are independent if

$$P(X_1 \in B_1, \ldots, X_n \in B_n) = P(X_1 \in B_1) \cdots P(X_n \in B_n) \qquad \text{for all Borel sets } B_1, \ldots, B_n,$$

i.e., if their joint distribution is the product $P_{X_1} \otimes \cdots \otimes P_{X_n}$ of their distributions. Infinitely many random variables are independent if any finitely many of them are.

For discrete random variables, independence means $p(x, y) = p_X(x)\, p_Y(y)$ for all values x and y, and for random variables with a joint density, it means that the joint density factorizes as $p(x, y) = p_X(x)\, p_Y(y)$ (outside a null set), as in Definition 4.4.2. In the language of σ-algebras, the information contained in X is the σ-algebra $\sigma(X) = \{\{X \in B\} : B \text{ Borel}\}$ generated by X, and X and Y are independent if and only if $\sigma(X)$ and $\sigma(Y)$ are independent (Definition B.1.7). Since $\sigma(g(X)) \subseteq \sigma(X)$ for measurable g, functions $g(X)$ and $h(Y)$ of independent random variables are again independent.

Proposition B.6.3. *If X and Y are independent, and $g(X)$ and $h(Y)$ are integrable (or nonnegative), then $\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\,\mathbb{E}[h(Y)]$. In particular, $\mathbb{E}[XY] = \mathbb{E}[X]\,\mathbb{E}[Y]$ for integrable X and Y.*

Proof. The joint distribution is $P_X \otimes P_Y$, so by Theorem B.3.3 and Tonelli's theorem (for nonnegative functions) or Fubini's theorem,

$$\mathbb{E}[g(X)h(Y)] = \iint g(x)h(y)\,dP_X(x)\,dP_Y(y) = \int g\,dP_X \int h\,dP_Y = \mathbb{E}[g(X)]\,\mathbb{E}[h(Y)]. \qquad \square$$

This extends Proposition 4.4.4 to all random variables. Tonelli's theorem also proves the formulas of Examples 4.4.5 and 4.4.7 in general. For example, every random variable $X \geq 0$ satisfies $X = \int_0^\infty \mathbf{1}_{\{t<X\}}\,dt$, and Tonelli's theorem for $P \otimes \lambda$ gives

$$\mathbb{E}[X] = \int_0^\infty P(X > t)\,dt,$$

also if X is neither discrete nor continuous.

B.7 Conditional Expectation

The conditional expectation $\mathbb{E}[Y \mid X]$ of Definition 4.4.14 only depends on the information contained in X: it is a function of X, and by the proof of Theorem 4.4.15, with $h = \mathbf{1}_B$, $\mathbb{E}\big[(Y - \mathbb{E}[Y \mid X])\mathbf{1}_G\big] = 0$ for every event $G = \{X \in B\}$ in $\sigma(X)$. These two properties define the conditional expectation in general: also when $P(X = x) = 0$ for all x and there is no joint density, and for conditioning on infinitely many random variables.

Definition B.7.1 (Conditional expectation given a σ-algebra)**.** Let Y be integrable, and let $\mathcal{G} \subseteq \mathcal{F}$ be a σ-algebra. A conditional expectation of Y given $\mathcal{G}$ is a $\mathcal{G}$-measurable integrable random variable $\mathbb{E}[Y \mid \mathcal{G}]$ with

$$\mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{G}]\,\mathbf{1}_G\big] = \mathbb{E}[Y\mathbf{1}_G] \qquad \text{for all } G \in \mathcal{G}.$$

For a random variable or random vector X, $\mathbb{E}[Y \mid X] = \mathbb{E}[Y \mid \sigma(X)]$.

Theorem B.7.2. *The conditional expectation exists, and it is unique up to changes on a null set. If $\mathbb{E}[Y^2] < \infty$, then $\mathbb{E}[Y \mid \mathcal{G}]$ is the orthogonal projection of Y onto the subspace $L^2(\mathcal{G})$ of $\mathcal{G}$-measurable random variables in L^2: it minimizes $\mathbb{E}[(Y - Z)^2]$ over all $Z \in L^2(\mathcal{G})$. Moreover, for integrable Y:*

1. *(Tower property) $\mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{G}]\big] = \mathbb{E}[Y]$, and $\mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{G}] \mid \mathcal{H}\big] = \mathbb{E}[Y \mid \mathcal{H}]$ for σ-algebras $\mathcal{H} \subseteq \mathcal{G}$;*
2. *(Taking out what is known) $\mathbb{E}[ZY \mid \mathcal{G}] = Z\,\mathbb{E}[Y \mid \mathcal{G}]$ for every bounded $\mathcal{G}$-measurable random variable Z;*
3. *(Independence) if $\sigma(Y)$ and $\mathcal{G}$ are independent, then $\mathbb{E}[Y \mid \mathcal{G}] = \mathbb{E}[Y]$.*

Proof of some parts. Existence for $Y \in L^2$: one can show that $L^2(\mathcal{G})$ is a closed subspace of L^2, so by the projection theorem for Hilbert spaces (see Appendix A), Y has an orthogonal projection $\hat{Y}$ onto it. The error $Y - \hat{Y}$ is orthogonal to every element of $L^2(\mathcal{G})$, in particular to $\mathbf{1}_G$ for $G \in \mathcal{G}$, which is the defining property. For integrable $Y \geq 0$, one applies the Radon–Nikodym theorem to the measure $G \mapsto \mathbb{E}[Y\mathbf{1}_G]$ on $\mathcal{G}$. Uniqueness: if Z_1 and Z_2 both have the defining property, then $G = \{Z_1 > Z_2\} \in \mathcal{G}$ gives $\mathbb{E}[(Z_1 - Z_2)\mathbf{1}_G] = 0$, so $Z_1 \leq Z_2$ almost surely, and by symmetry, $Z_1 = Z_2$ almost surely. (1) The first statement is the defining property for $G = \Omega$. For the second, $\mathbb{E}[Y \mid \mathcal{H}]$ is $\mathcal{H}$-measurable, and for $H \in \mathcal{H} \subseteq \mathcal{G}$, $\mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{H}]\mathbf{1}_H\big] = \mathbb{E}[Y\mathbf{1}_H] = \mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{G}]\mathbf{1}_H\big]$, which is the defining property of $\mathbb{E}\big[\mathbb{E}[Y \mid \mathcal{G}] \mid \mathcal{H}\big]$. (3) The constant $\mathbb{E}[Y]$ is $\mathcal{G}$-measurable, and $\mathbb{E}[Y\mathbf{1}_G] = \mathbb{E}[Y]\,P(G) = \mathbb{E}\big[\mathbb{E}[Y]\mathbf{1}_G\big]$ by independence (Proposition B.6.3). We omit the proof of (2). $\square$

By the Doob–Dynkin lemma, which we do not prove, every $\sigma(X)$-measurable random variable is a measurable function of X. So $\mathbb{E}[Y \mid X] = m(X)$ for a measurable function m, and for discrete X or a joint density, $m(x)$ is the conditional mean $\mathbb{E}[Y \mid X = x]$ of Definition 4.4.14; Theorem 4.4.15 is the case $\mathcal{G} = \sigma(X)$ of Theorem B.7.2.

Example B.7.3. 1. (Finitely many pieces of information.) Let $\mathcal{G}$ be generated by a partition $B_1, \dots, B_n$ of Ω with $P(B_i) > 0$ (Example B.1.2). A $\mathcal{G}$-measurable random variable is constant on each B_i, and the defining property with $G = B_i$ gives

$$\mathbb{E}[Y \mid \mathcal{G}] = \sum_{i=1}^{n} \frac{\mathbb{E}[Y \mathbf{1}_{B_i}]}{P(B_i)} \mathbf{1}_{B_i} :$$

on each piece, $\mathbb{E}[Y \mid \mathcal{G}]$ is the average of Y over that piece.

2. (Law of total variance.) For $Y \in L^2$, the decomposition $Y - \mathbb{E}[Y] = (Y - \mathbb{E}[Y \mid \mathcal{G}]) + (\mathbb{E}[Y \mid \mathcal{G}] - \mathbb{E}[Y])$ is orthogonal, and Pythagoras gives

$$\operatorname{Var}(Y) = \mathbb{E}\big[\operatorname{Var}(Y \mid \mathcal{G})\big] + \operatorname{Var}\big(\mathbb{E}[Y \mid \mathcal{G}]\big), \qquad \text{where } \operatorname{Var}(Y \mid \mathcal{G}) = \mathbb{E}\big[(Y - \mathbb{E}[Y \mid \mathcal{G}])^2 \mid \mathcal{G}\big].$$

For $\mathcal{G} = \sigma(X)$, this is the law of total variance of Example 4.4.16.

B.8 Martingales

Conditional expectations describe what we expect, given the information that is available at a certain time. A filtration is an increasing sequence of σ-algebras $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \cdots \subseteq \mathcal{F}$, where $\mathcal{F}_n$ is the information available at time n; for example, $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$ for a sequence of observations $X_1, X_2, \dots$, and $\mathcal{F}_0 = \{\emptyset, \Omega\}$.

Definition B.8.1 (Martingale)**.** A sequence of integrable random variables $M_0, M_1, \dots$ is a martingale with respect to a filtration $(\mathcal{F}_n)$ if every M_n is $\mathcal{F}_n$-measurable and $\mathbb{E}[M_{n+1} \mid \mathcal{F}_n] = M_n$ for all n.

A martingale is a fair game: given the past, the expected next value is the current value. By the tower property, $\mathbb{E}[M_n] = \mathbb{E}[M_0]$ for all n.

Example B.8.2. 1. (Random walks.) Let $X_1, X_2, \dots$ be independent with mean 0, and let $S_n = X_1 + \cdots + X_n$, with $S_0 = 0$. Taking out what is known and independence give $\mathbb{E}[S_{n+1} \mid \mathcal{F}_n] = S_n + \mathbb{E}[X_{n+1}] = S_n$, so (S_n) is a martingale. If the X_i also have the variance σ^2, then $S_n^2 - n\sigma^2$ is a martingale, since $\mathbb{E}[S_{n+1}^2 \mid \mathcal{F}_n] = S_n^2 + 2S_n\mathbb{E}[X_{n+1}] + \mathbb{E}[X_{n+1}^2] = S_n^2 + \sigma^2$.

2. (Accumulating information.) For every integrable Y, $M_n = \mathbb{E}[Y \mid \mathcal{F}_n]$ is a martingale, by the tower property: it is the best prediction of Y, updated as information arrives. For example, the posterior mean $\mathbb{E}[\theta \mid X_1, \dots, X_n]$ of Bayesian inference (Section 4.12) is a martingale.

3. (Likelihood ratios.) Let $X_1, X_2, \dots$ be i.i.d. with the density q, and let p be another density with $p(x) = 0$ whenever $q(x) = 0$. Then the likelihood ratio $L_n = \prod_{i=1}^{n} \frac{p(X_i)}{q(X_i)}$ of the data under p versus q is a martingale with $\mathbb{E}[L_n] = 1$, since its factors are independent with the mean $\int \frac{p}{q}\, q\, dx = 1$. By Ville's inequality, which we do not prove, $P\big(L_n \geq \frac{1}{\alpha} \text{ for some } n\big) \leq \alpha$ for every nonnegative martingale with $L_0 = 1$. So we may compute L_n after every observation and reject the hypothesis that the data have the density q as soon as $L_n \geq \frac{1}{\alpha}$: unlike repeated z-tests (Section 4.11), this sequential test keeps its level α, however often we look at the data. If the data actually have the density p, the strong law of large numbers (Theorem B.9.6) gives $\frac{1}{n} \log L_n \to \mathbb{E}_p\big[\log \frac{p(X_1)}{q(X_1)}\big] = \mathrm{KL}(p \,\|\, q)$ (Section 4.13), so L_n grows exponentially fast, and the test eventually rejects.

A stopping time is a random time τ with values in $\{0, 1, 2, \ldots\} \cup \{\infty\}$ such that $\{\tau = n\} \in \mathcal{F}_n$ for all n: whether we stop at time n only depends on the information available at time n, not on the future.

Theorem B.8.3 (Optional stopping)**.** *Let (M_n) be a martingale and τ a stopping time with $\tau < \infty$ almost surely, and let $|M_{\min(n,\tau)}| \le C$ for all n, with a constant C. Then $\mathbb{E}[M_\tau] = \mathbb{E}[M_0]$.*

Proof. The stopped process $N_n = M_{\min(n,\tau)}$ is again a martingale: $N_{n+1} - N_n = \mathbf{1}_{\{\tau > n\}}(M_{n+1} - M_n)$, and since $\{\tau > n\} \in \mathcal{F}_n$, taking out what is known gives $\mathbb{E}[N_{n+1} - N_n \mid \mathcal{F}_n] = \mathbf{1}_{\{\tau>n\}}\mathbb{E}[M_{n+1} - M_n \mid \mathcal{F}_n] = 0$. So $\mathbb{E}[N_n] = \mathbb{E}[M_0]$ for all n. As $n \to \infty$, $N_n \to M_\tau$ almost surely, and $|N_n| \le C$, so dominated convergence gives $\mathbb{E}[M_\tau] = \mathbb{E}[M_0]$. □

So no stopping rule turns a fair game into a favorable one, as long as the stakes stay bounded. Without this condition, the theorem fails: a gambler who doubles the stake after every loss in a fair game wins 1 euro almost surely, but needs an unbounded fortune to do so.

Example B.8.4 (Gambler's ruin)**.** A gambler starts with a euros and bets 1 euro on fair coin flips, until the fortune reaches $a + b$ euros or 0. The fortune is $a + S_n$ with the simple random walk S_n (with steps ± 1 of probability $\frac{1}{2}$ each), and the game ends at the first time τ with $S_\tau = b$ or $S_\tau = -a$. One can show that $\tau < \infty$ almost surely, and $|S_{\min(n,\tau)}| \le \max(a, b)$. By optional stopping,

$$0 = \mathbb{E}[S_\tau] = b\,P(S_\tau = b) - a\,P(S_\tau = -a), \qquad \text{so} \qquad P(S_\tau = b) = \frac{a}{a+b}.$$

Applying optional stopping to the martingale $S_n^2 - n$ (with more care, since it is not bounded) gives $\mathbb{E}[\tau] = \mathbb{E}[S_\tau^2] = b^2\frac{a}{a+b} + a^2\frac{b}{a+b} = ab$. For $a = 10$ and $b = 90$, the gambler reaches 100 euros with probability 0.1, and the game lasts 900 rounds on average.

Martingales also exist in continuous time. With the filtration $\mathcal{F}_s = \sigma(B(u) : u \le s)$, the independence of the increments of Brownian motion (Definition 4.15.1) gives $\mathbb{E}[B(t) \mid \mathcal{F}_s] = B(s) + \mathbb{E}[B(t) - B(s)] = B(s)$ for $s \le t$: Brownian motion is a martingale, the continuous analogue of a fair random walk.

B.9 Modes of Convergence and the Strong Law

The weak law of large numbers (Theorem 4.7.2) is a statement about probabilities, for each N. Measure theory allows statements about the whole random sequence: for example, that $\bar{X}_N(\omega) \to \mu$ for almost every outcome ω, i.e., along almost every infinite sequence of observations. Random variables can converge in several different senses.

Definition B.9.1 (Modes of convergence)**.** Let $X, X_1, X_2, \ldots$ be random variables on the same probability space. We say that $X_n \to X$

1. almost surely if $P(X_n \to X) = 1$, i.e., $X_n(\omega) \to X(\omega)$ for all ω outside a null set;
2. in probability if $P(|X_n - X| > \varepsilon) \to 0$ for every $\varepsilon > 0$;
3. in L^p, for $p \ge 1$, if $\mathbb{E}[|X_n - X|^p] \to 0$; for $p = 2$, this is convergence in mean square.

Random variables X_n and X, possibly on different probability spaces, converge in distribution if $F_{X_n}(x) \to F_X(x)$ at every point x at which F_X is continuous.

Convergence in probability and in distribution are the notions of Definition 4.7.8: the weak law of large numbers is convergence in probability, and the central limit theorem (Theorem 4.7.3) is convergence in distribution of the standardized means to $\mathcal{N}(0, 1)$.

Proposition B.9.2. *Almost sure convergence and convergence in L^p each imply convergence in probability, which implies convergence in distribution. None of the converse implications holds in general.*

Proof. If $X_n \to X$ almost surely, the events $A_n = \{|X_m - X| > \varepsilon \text{ for some } m \geq n\}$ decrease to an event that is contained in the null event $\{X_n \not\to X\}$, so $P(|X_n - X| > \varepsilon) \leq P(A_n) \to 0$ by continuity. Convergence in L^p implies convergence in probability by Markov's inequality (Proposition 4.3.6): $P(|X_n - X| > \varepsilon) \leq \frac{\mathbb{E}[|X_n - X|^p]}{\varepsilon^p}$. If $X_n \to X$ in probability, then $X_n = X + (X_n - X)$ with $X_n - X \to 0$ in probability, so $X_n \to X$ in distribution by Slutsky's theorem (Theorem 4.7.10), applied to the constant sequence X. Example B.9.3 disproves the converses. □

Example B.9.3. Let U be uniform on $[0, 1)$.

1. The lottery tickets $X_n = n\mathbf{1}_{\{U<1/n\}}$ of Example B.4.3 converge to 0 almost surely, since $X_n = 0$ as soon as $n > \frac{1}{U}$, but not in L^1, since $\mathbb{E}[X_n] = 1$.
2. (A sliding block.) Write $n = 2^k + j$ with $0 \leq j < 2^k$, and let $X_n = \mathbf{1}_{\{j2^{-k} \leq U < (j+1)2^{-k}\}}$: the indicator of a block of length 2^{-k}, which slides across $[0, 1)$, and then starts again with half the length. Then $\mathbb{E}[|X_n|^p] = 2^{-k} \to 0$, so $X_n \to 0$ in every L^p and in probability. But for every value of U, one block of every generation k covers it, so $X_n = 1$ for infinitely many n, and the sequence $X_n(\omega)$ converges for no outcome ω.
3. If $X \sim \mathcal{N}(0, 1)$ and $X_n = -X$ for all n, then $X_n \to X$ in distribution, since $-X$ has the same distribution as X, but $|X_n - X| = 2|X|$ does not converge to 0 in probability.

The main tool for proving almost sure convergence is the following lemma.

Lemma B.9.4 (Borel–Cantelli)**.** *Let $A_1, A_2, \ldots$ be events, and let $\{A_n \text{ i.o.}\}$ be the event that infinitely many of them happen ("infinitely often").*

1. *If $\sum_n P(A_n) < \infty$, then $P(A_n \text{ i.o.}) = 0$.*
2. *If the A_n are independent and $\sum_n P(A_n) = \infty$, then $P(A_n \text{ i.o.}) = 1$.*

Proof. (1) For every k, $\{A_n \text{ i.o.}\} \subseteq \bigcup_{n \geq k} A_n$, which has probability at most $\sum_{n \geq k} P(A_n)$ by the union bound, and this tends to 0 as $k \to \infty$. (2) The complement of $\{A_n \text{ i.o.}\}$ is the union over k of the events $\bigcap_{n \geq k} A_n^c$. By independence and $1 - x \leq e^{-x}$, for all $m \geq k$,

$$P\Big(\bigcap_{n=k}^{m} A_n^c\Big) = \prod_{n=k}^{m} (1 - P(A_n)) \leq \exp\Big(-\sum_{n=k}^{m} P(A_n)\Big),$$

which tends to 0 as $m \to \infty$. So every $\bigcap_{n \geq k} A_n^c$ is a null event, and so is their countable union. □

Example B.9.5. Consider infinitely many fair coin flips (Example B.1.8). The events A_k that the flips $2k - 1$ and $2k$ are both heads are independent and have the probability $\frac{1}{4}$, so by part (2), almost surely, two heads in a row appear infinitely often; the same argument works for every finite pattern, for example for the text of this appendix, encoded in bits. On the other hand, let B_k be the event that the $\lceil 2 \log_2 k \rceil$ flips starting with flip k are all heads. Then $P(B_k) \leq k^{-2}$, which is summable, so by part (1), almost surely, only finitely many B_k happen: the longest run of heads among the first n flips is eventually shorter than about $2 \log_2 n$.

Theorem B.9.6 (Strong law of large numbers)**.** *Let $X_1, X_2, \ldots$ be i.i.d. and integrable with mean μ. Then $\bar{X}_N \to \mu$ almost surely.*

Proof if X_1 has a finite fourth moment. Let $Y_i = X_i - \mu$ and $S_N = Y_1 + \cdots + Y_N$. When we expand $S_N^4 = \sum_{i,j,k,l} Y_i Y_j Y_k Y_l$, every term in which some Y_i appears exactly once has expectation 0, by independence. What remains are the N terms Y_i^4 and the $3N(N-1)$ terms $Y_i^2 Y_j^2$ with $i \neq j$, so $\mathbb{E}[S_N^4] = N\mathbb{E}[Y_1^4] + 3N(N-1)\sigma^4 \leq CN^2$ for a constant C. By Markov's inequality,

$$P(|\bar{X}_N - \mu| \geq \varepsilon) = P(S_N^4 \geq N^4\varepsilon^4) \leq \frac{C}{N^2\varepsilon^4},$$

which is summable. By the Borel–Cantelli lemma, almost surely, $|\bar{X}_N - \mu| \geq \varepsilon$ only for finitely many N. Applying this to $\varepsilon = \frac{1}{m}$ for $m = 1, 2, \ldots$ (countably many null events) gives $\bar{X}_N \to \mu$ almost surely. □

Example B.9.7 (Normal numbers)**.** The binary digits of a uniformly random $U \in [0, 1)$ are i.i.d. fair coin flips (Example B.1.8). By the strong law, the fraction of 1s among the first N digits converges to $\frac{1}{2}$ almost surely: almost every number in $[0, 1)$ has asymptotically equally many 0s and 1s in its binary expansion (Borel's normal number theorem, 1909). Yet for natural constants such as $\sqrt{2} - 1$, it is not known whether they have this property. This is typical of probabilistic arguments: they show that almost all objects have a property, without exhibiting one.

By the strong law, applied to the X_i and the X_i^2, the sample standard deviation s_N also converges to σ almost surely. Similarly, every estimator that is a continuous function of sample means, such as the maximum likelihood estimator $\frac{1}{\bar{X}_N}$ of the rate of an exponential distribution, converges to the true parameter almost surely: it is strongly consistent, which is stronger than consistency (Definition 4.9.11).

Key Takeaways

- Events form a σ-algebra, which is closed under complements and countable unions, and a probability is a countably additive measure with $P(\Omega) = 1$. Not every subset of $[0, 1]$ can have a length (Vitali), but every Borel set has one (the Lebesgue measure). Measures are continuous along increasing and decreasing sequences of sets, and null events, such as a rational uniform point, can happen. σ-algebras describe information, and independent σ-algebras describe pieces of information that tell us nothing about each other.
- Random variables are measurable functions, and their distributions are the image measures P_X. The CDF determines the distribution, and every nondecreasing, right-continuous function with the limits 0 and 1 is a CDF.
- The Lebesgue integral divides the range instead of the domain; it extends the integral of Lecture 3 and contains series as integrals with respect to the counting measure. The expectation is $\mathbb{E}[X] = \int X\, dP$, computed from the distribution as $\mathbb{E}[g(X)] = \int g\, dP_X$. Monotone and dominated convergence allow us to exchange limits and expectations, and to differentiate under the integral sign; without a dominating function, mass can escape.
- Pmfs and pdfs are densities with respect to the counting and the Lebesgue measure. A distribution has a density with respect to μ if and only if it is absolutely continuous with respect to μ (Radon–Nikodym). Mixed distributions such as the ReLU of a Gaussian have densities with respect to other measures, singular distributions such as the Cantor distribution have none, and likelihood ratios are Radon–Nikodym derivatives.
- Independence means that the joint distribution is the product measure of the marginals. By Fubini–Tonelli, nonnegative or integrable functions can be integrated in either order, so expectations of products of independent random variables factorize, and $\mathbb{E}[X] = \int_0^\infty P(X > t)\, dt$ for every $X \geq 0$.

- The conditional expectation $\mathbb{E}[Y \mid \mathcal{G}]$ is the $\mathcal{G}$-measurable random variable with the same integrals as Y over the events of $\mathcal{G}$; for $Y \in L^2$, it is the orthogonal projection of Y onto $L^2(\mathcal{G})$, the best mean-squared-error prediction of Y from the information $\mathcal{G}$. It satisfies the tower property and the law of total variance, and it takes out what is known.
- A martingale is a fair game, $\mathbb{E}[M_{n+1} \mid \mathcal{F}_n] = M_n$; random walks, posterior means, likelihood ratios and Brownian motion are martingales. Stopping at a stopping time does not change the expectation if the stopped martingale stays bounded (optional stopping), which solves the gambler's ruin, and Ville's inequality gives likelihood ratio tests that stay valid when we look at the data repeatedly.
- Almost sure convergence and convergence in L^p imply convergence in probability, which implies convergence in distribution, but not conversely. The Borel–Cantelli lemma proves almost sure statements, and the strong law of large numbers says that $\bar{X}_N \to \mu$ almost surely; for example, almost every number is normal.

Exercises

1. (σ-algebras.)
   (a) Show that the intersection of two σ-algebras on Ω is a σ-algebra, but that their union need not be one. (Hint: $\Omega = \{1, 2, 3\}$.)
   (b) Let $\mathcal{G}$ be the σ-algebra on $\Omega = \{1, \dots, 6\}^2$ generated by the events $\{\text{the sum is } s\}$ for two dice. How many elements does $\mathcal{G}$ have? Does it contain the event "both dice show the same number"?
2. Let X be a random variable with the CDF F. Use the continuity of probability (Proposition B.1.6) to show that $P(X > t) \to 0$ as $t \to \infty$, and that $P(X = x) = F(x) - \lim_{y \uparrow x} F(y)$. Conclude that F has at most countably many jumps. (Hint: there are at most n jumps of size larger than $\frac{1}{n}$.)
3. (Limits and expectations.)
   (a) Use monotone convergence to show that $\lim_{n\to\infty} \int_0^n \left(1 - \frac{x}{n}\right)^n dx = 1$. (Hint: $(1 - \frac{x}{n})^n \mathbf{1}_{[0,n]}(x)$ increases to e^{-x}.)
   (b) Let $X \geq 0$ with $\mathbb{E}[X] < \infty$. Show that $\mathbb{E}[X \mathbf{1}_{\{X > t\}}] \to 0$ as $t \to \infty$, and conclude that $nP(X > n) \to 0$.
   (c) Find random variables $X_n \geq 0$ with $X_n \to 0$ almost surely and $\mathbb{E}[X_n] \to \infty$.
4. A waiting time $Z \sim$ Exponential(1) is cut off after 2 minutes: $Y = \min(Z, 2)$. Find the CDF of Y and a density of Y with respect to $\lambda + \delta_2$, and compute $\mathbb{E}[Y]$ in two ways: with this density, and with the tail integral $\mathbb{E}[Y] = \int_0^\infty P(Y > t)\, dt$ of Section B.6.
5. Let X_1 and X_2 be independent and Uniform$(0, 1)$, and $M = \max(X_1, X_2)$.
   (a) Show that $\mathbb{E}[X_1 \mathbf{1}_{\{M \leq t\}}] = \mathbb{E}\left[\frac{3}{4} M \mathbf{1}_{\{M \leq t\}}\right]$ for $0 \leq t \leq 1$. (One can show that this implies $\mathbb{E}[X_1 \mid M] = \frac{3}{4} M$, since the events $\{M \leq t\}$ generate $\sigma(M)$ and are closed under intersections.)
   (b) Explain the result: given $M = m$, $X_1 = m$ with probability $\frac{1}{2}$, and otherwise X_1 is uniform on $[0, m]$. Check the tower property.
6. (Pólya's urn.) An urn contains one red and one blue ball. Repeatedly, a ball is drawn at random and put back together with another ball of the same color. Let R_n be the fraction of red balls after n draws. Show that (R_n) is a martingale, and find $\mathbb{E}[R_n]$.

7. Let X_n be independent with $P(X_n = 1) = p_n$ and $P(X_n = 0) = 1 - p_n$.
   (a) For $p_n = \frac{1}{n}$, show that $X_n \to 0$ in probability and in L^1, but not almost surely.
   (b) For $p_n = \frac{1}{n^2}$, show that $X_n \to 0$ almost surely.

Programming Exercises

As in Lecture 4, implement the methods yourself with NumPy, and use library routines (such as `rng.normal` or `scipy.stats`) only to check your results. Start each file with

```
import numpy as np
import matplotlib.pyplot as plt
rng = np.random.default_rng(0)
```

P1. Distributions without densities.

- Sample 10^5 values of $\max(0, Z)$ for $Z \sim \mathcal{N}(0,1)$, and check $P(Y = 0) = \frac{1}{2}$, $\mathbb{E}[Y] = \frac{1}{\sqrt{2\pi}}$ and $\mathrm{Var}(Y) = \frac{1}{2} - \frac{1}{2\pi}$ (Example B.5.3).
- Sample from the Cantor distribution (Remark B.5.4) with 30 random ternary digits 0 or 2, plot the empirical CDF (the Cantor function), and check the mean $\frac{1}{2}$ and the variance $\frac{1}{8}$. What fraction of the samples lies in $(\frac{1}{3}, \frac{2}{3})$, and in $(\frac{1}{9}, \frac{2}{9})$?

P2. Modes of convergence.

- For the sequences (1) and (2) of Example B.9.3, plot the paths $n \mapsto X_n$ for $n \le 1000$ for a few values of U, and estimate $\mathbb{E}[X_n]$ and $P(X_n \neq 0)$ from 10^4 values of U. Which sequence converges almost surely, which in probability, and which in L^1?
- Simulate independent events A_n with $P(A_n) = \frac{1}{n}$ and with $P(A_n) = \frac{1}{n^2}$, for $n \le 10^6$. In 20 runs, count how many of the events happen, and record when the last one happens. Compare with the Borel–Cantelli lemma; for $P(A_n) = \frac{1}{n}$, compare the count with $\sum_{n \le N} \frac{1}{n} \approx \log N$.
- (The strong law.) Plot $\bar{X}_N$ against $N \le 10^4$ for 20 independent sequences of Exponential(1) variables, together with the curves $1 \pm 2/\sqrt{N}$. How does the spread of the paths shrink, and does every path converge?

P3. Martingales.

- Simulate the gambler's ruin of Example B.8.4 with $a = 10$ and $b = 90$, 10 000 times, and check the winning probability 0.1 and the mean duration 900. Then let the coin show heads with probability 0.49. How do the two numbers change?
- (A test that allows peeking.) For coin flips with H_0: $\theta = \frac{1}{2}$ and the alternative $\theta = 0.6$, compute the likelihood ratio $L_n = \prod_{i \le n} \frac{0.6^{x_i} 0.4^{1-x_i}}{0.5}$ after every flip, and reject H_0 as soon as $L_n \ge 20$ (Example B.8.2). In 10 000 simulations of a fair coin with up to 1000 flips, how often is H_0 rejected? How many flips are needed on average to reject H_0 when $\theta = 0.6$? Compare with the peeking z-test in the programming exercises of Lecture 4.