

Gain and Loss as Singular Limits of Gaussian Transfer Matrices

Asadullah Bhuiyan
(Dated: October 7, 2026)

Fermion creation and annihilation preserve Gaussian conditional states but are nilpotent, so they have no invertible conjugation action on fermion operators. We resolve this apparent mismatch by representing each jump as a projectively rescaled limit of an invertible odd Gaussian operator. Its polar decomposition is a weak occupation measurement followed by a Majorana flip; its action on Majoranas lies in the disconnected component of $O(2N, \mathbb{C})$. Outcome-dependent feedback gives finite-strength instruments whose limits are precisely the fill and deplete steps of a fresh-ancilla adaptive circuit. Ordered products have a well-defined normalized state limit for every fixed finite, nonzero branch, even when their transfer matrices diverge. We derive the exact Slater-projector updates at the endpoint and explain why regulator singularities, physical trajectory singular values, and tangent Lyapunov rates must be kept distinct.

CONTENTS

I. The physical question and conventions	1
II. From a nilpotent jump to an orthogonal transformation	2
A. An invertible regulator and its polar decomposition	2
B. The group and the meaning of its boundary	2
III. A physical measurement and the adaptive feedback choices	3
IV. Ordered products and a controlled singular limit	4
A. Composition before the limit	4
B. What converges, and under which assumptions	4
V. The exact endpoint without a regulator	5
VI. Consequences for trajectory spectra	6
References	6

I. THE PHYSICAL QUESTION AND CONVENTIONS

An adaptive occupation measurement may have to add or remove a fermion after detecting the wrong occupation. Such an operation is Gaussian in its action on states, but cannot be written as an invertible exponential of a quadratic operator: a creation or annihilation operator squares to zero. The useful statement is not that the jump itself belongs to a classical matrix group. Rather, it is a singular limit of operators whose Majorana conjugation actions belong to that group. The scalar used to take the limit and the state on which the operator acts are indispensable.

We use N complex modes $\hat{\psi}_i$ with $\{\hat{\psi}_i, \hat{\psi}_j^\dagger\} = \delta_{ij}$ and $\{\hat{\psi}_i, \hat{\psi}_j\} = 0$. A normalized column $\chi \in \mathbb{C}^N$ defines

$$\hat{\chi}^\dagger = \sum_i \chi_i \hat{\psi}_i^\dagger, \quad \hat{\chi} = \sum_i \chi_i^* \hat{\psi}_i, \quad \hat{n} = \hat{\chi}^\dagger \hat{\chi}, \quad \hat{n}^2 = \hat{n}. \quad (1)$$

Many-body operators carry hats. Following Bhuiyan, Pan, and Jian (BPJ) [1], the correlation matrix is $G_{ij} = \text{Tr}(\hat{\rho} \hat{\psi}_i^\dagger \hat{\psi}_j)$. For occupied columns in the convention (1), it is convenient to use

$$C = G^\text{T}, \quad C_{ij} = \text{Tr}(\hat{\rho} \hat{\psi}_j^\dagger \hat{\psi}_i). \quad (2)$$

Thus a Slater determinant with orthonormal occupied columns F has $C = FF^\dagger$; no centered covariance is denoted by G here.

The Majoranas

$$\hat{\gamma}_{2i-1} = \hat{\psi}_i + \hat{\psi}_i^\dagger, \quad \hat{\gamma}_{2i} = -i(\hat{\psi}_i - \hat{\psi}_i^\dagger), \quad \{\hat{\gamma}_a, \hat{\gamma}_b\} = 2\delta_{ab} \quad (3)$$

span a $2N$ -dimensional space. We allow both even Gaussian operators and their products with Majorana operators, which are odd. This extends the invertible quadratic-exponential description by its odd Clifford sector; the state and measurement formalism is consistent with fermionic linear optics [2]. A physical implementation of an odd system branch exchanges parity with an ancillary fermion, rather than violating total fermion-parity conservation.

II. FROM A NILPOTENT JUMP TO AN ORTHOGONAL TRANSFORMATION

A. An invertible regulator and its polar decomposition

For $0 < \varepsilon \leq 1$, define

$$\begin{aligned} \hat{j}_+(\varepsilon) &= \hat{\chi}^\dagger + \varepsilon \hat{\chi}, & \hat{j}_-(\varepsilon) &= \hat{\chi} + \varepsilon \hat{\chi}^\dagger, \\ \hat{v}_\pm(\varepsilon) &= \varepsilon^{-1/2} \hat{j}_\pm(\varepsilon). \end{aligned} \quad (4)$$

The canonical anticommutation relations give

$$\hat{j}_\pm^2 = \varepsilon \hat{\mathbf{1}}, \quad \hat{v}_\pm^2 = \hat{\mathbf{1}}, \quad \sqrt{\varepsilon} \hat{v}_+ \longrightarrow \hat{\chi}^\dagger, \quad \sqrt{\varepsilon} \hat{v}_- \longrightarrow \hat{\chi}. \quad (5)$$

Every nonzero regulator therefore supplies an inverse. Exact gain and loss do not: $\hat{\chi}^2 = (\hat{\chi}^\dagger)^2 = 0$.

To identify the physical content, set

$$\hat{\Gamma} = \hat{\chi} + \hat{\chi}^\dagger, \quad \hat{\Gamma}^\dagger = \hat{\Gamma}, \quad \hat{\Gamma}^2 = \hat{\mathbf{1}}, \quad \lambda = -\frac{1}{2} \log \varepsilon. \quad (6)$$

Direct multiplication gives

$$\begin{aligned} \hat{v}_+^\dagger \hat{v}_+ &= \varepsilon^{-1}(\hat{\mathbf{1}} - \hat{n}) + \varepsilon \hat{n} = e^{-2\lambda(2\hat{n}-\hat{\mathbf{1}})}, \\ \hat{v}_-^\dagger \hat{v}_- &= \varepsilon(\hat{\mathbf{1}} - \hat{n}) + \varepsilon^{-1} \hat{n} = e^{+2\lambda(2\hat{n}-\hat{\mathbf{1}})}. \end{aligned} \quad (7)$$

Since $\hat{\Gamma} \hat{n} = \hat{\chi}$ and $\hat{\Gamma}(\hat{\mathbf{1}} - \hat{n}) = \hat{\chi}^\dagger$, their polar decompositions are

$$\hat{v}_\pm = \hat{\Gamma} e^{\mp \lambda(2\hat{n}-\hat{\mathbf{1}})}. \quad (8)$$

The positive factor filters occupation; the odd unitary factor flips it. The filter has the form of the weak single-mode measurement used in BPJ Appendix B.2 [1]. A scalar normalization and a complete set of outcomes are still required to make a physical instrument.

B. The group and the meaning of its boundary

For any complex column z , let $\hat{\mu}(z) = \sum_a z_a \hat{\gamma}_a$. The relevant quadratic form is bilinear, not Hermitian:

$$\hat{\mu}(z)^2 = (z^\top z) \hat{\mathbf{1}}. \quad (9)$$

When $z^\top z \neq 0$, anticommuting $\hat{\gamma}_a$ through $\hat{\mu}(z)$ gives

$$\hat{\mu}(z) \hat{\gamma}_a \hat{\mu}(z)^{-1} = \sum_b t[z]_{ab} \hat{\gamma}_b, \quad t[z] = -\mathbf{1} + \frac{2zz^\top}{z^\top z}. \quad (10)$$

The rank-one factor in this expression is an idempotent, hence

$$t[z]^\top t[z] = \mathbf{1}, \quad t[z]^2 = \mathbf{1}, \quad \det t[z] = (-1)^{2N-1} = -1. \quad (11)$$

Thus an invertible odd Clifford operator induces a single-particle transfer matrix (sTM) in the disconnected component of $O(2N, \mathbb{C})$. On the two Majoranas of the measured mode,

$$z_+(\varepsilon) = \frac{1}{2} \begin{pmatrix} \varepsilon^{-1/2} + \varepsilon^{1/2} \\ i(\varepsilon^{1/2} - \varepsilon^{-1/2}) \end{pmatrix}, \quad z_+^\top z_+ = 1, \quad (12)$$

represents $\hat{v}_+$. The creation operator instead corresponds to $z_0 = (1, -i)^\top/2$, with $z_0^\top z_0 = 0$. The jump is a null Clifford direction, so Eq. (10) cannot be evaluated there by ordinary matrix substitution.

The same degeneration is transparent in the local Nambu basis:

$$\hat{v}_+ \begin{pmatrix} \hat{\chi} \\ \hat{\chi}^\dagger \end{pmatrix} \hat{v}_+^{-1} = \begin{pmatrix} 0 & \varepsilon^{-1} \\ \varepsilon & 0 \end{pmatrix} \begin{pmatrix} \hat{\chi} \\ \hat{\chi}^\dagger \end{pmatrix}. \quad (13)$$

Orthogonal spectator modes acquire a minus sign under this odd conjugation. In an orthonormal mode basis the full sTM has singular values $\{\varepsilon^{-1}, 1, \dots, 1, \varepsilon\}$, although all its eigenvalues are ± 1 . The divergent singular values are not divergent eigenvalues.

Here “boundary” means a boundary of the projective operator or normalized-state description: $[\hat{v}_+] \rightarrow [\hat{\chi}^\dagger]$, with the rescaling specified in Eq. (5). It is *not* a finite singular matrix in the ordinary closure of $O(2N, \mathbb{C})$. A finite limit of matrices satisfying $t^\top t = \mathbf{1}$ still satisfies that identity and is invertible. In the present limit the conjugation matrix escapes to infinity, while its rescaled many-body representative has a finite nonzero limit. Normalized Choi states supply a concrete way to retain that limit, as developed below.

III. A PHYSICAL MEASUREMENT AND THE ADAPTIVE FEEDBACK CHOICES

Define two positive weak-measurement operators,

$$\hat{M}_0 = \frac{\hat{\mathbf{1}} - \hat{n} + \varepsilon \hat{n}}{\sqrt{1 + \varepsilon^2}}, \quad \hat{M}_1 = \frac{\hat{n} + \varepsilon(\hat{\mathbf{1}} - \hat{n})}{\sqrt{1 + \varepsilon^2}}. \quad (14)$$

The subscripts label outcomes that become “empty” and “occupied” in the strong limit. Their effects obey

$$\hat{E}_0 = \frac{\hat{\mathbf{1}} - \hat{n} + \varepsilon^2 \hat{n}}{1 + \varepsilon^2}, \quad \hat{E}_1 = \frac{\hat{n} + \varepsilon^2(\hat{\mathbf{1}} - \hat{n})}{1 + \varepsilon^2}, \quad \hat{E}_0 + \hat{E}_1 = \hat{\mathbf{1}}. \quad (15)$$

This is a positive-operator-valued measure (POVM). In the weighted convention $\sum_m w_m \hat{K}_m^\dagger \hat{K}_m = \hat{\mathbf{1}}$, our displayed Kraus operators absorb $\sqrt{w_m}$, so all remaining weights equal one. For example, $p_0 = [1 - \langle \hat{n} \rangle + \varepsilon^2 \langle \hat{n} \rangle]/(1 + \varepsilon^2)$.

Outcome-dependent unitary feedback leaves these effects unchanged. The following three choices have distinct physical purposes:

Feedback	$\hat{K}_0$	$\hat{K}_1$	$\varepsilon \rightarrow 0$ endpoint
Fill	$\hat{\Gamma} \hat{M}_0$	$\hat{M}_1$	$\{\hat{\chi}^\dagger, \hat{n}\}$
Deplete	$\hat{M}_0$	$\hat{\Gamma} \hat{M}_1$	$\{\hat{\mathbf{1}} - \hat{n}, \hat{\chi}\}$
Flip both	$\hat{\Gamma} \hat{M}_0$	$\hat{\Gamma} \hat{M}_1$	$\{\hat{\chi}^\dagger, \hat{\chi}\}$

In particular,

$$\hat{\Gamma} \hat{M}_0 = \frac{\hat{j}_+(\varepsilon)}{\sqrt{1 + \varepsilon^2}}, \quad \hat{\Gamma} \hat{M}_1 = \frac{\hat{j}_-(\varepsilon)}{\sqrt{1 + \varepsilon^2}}. \quad (16)$$

Every listed finite- ε branch is invertible and Gaussian. The first two instruments approach exactly the target-fill and target-empty measurement-and-correction steps, up to irrelevant branch phases. They are the fresh-ancilla idealization of the occupation-measurement and fermionic-swap (fSWAP) construction in BPJ Sec. III [1], not its finite-bath dynamics. The third instrument flips either result and must not be substituted for targeted feedback. Pairing gain with loss is therefore one completion, not a unique requirement. At $\varepsilon = 1$ no occupation information is gained; these instruments reduce to random unitary evolution.

Equivalently, the flip-both pair uses the unnormalized $\hat{v}_\pm$ with equal weights

$$w_+ = w_- = \frac{1}{\varepsilon + \varepsilon^{-1}} = \frac{1}{2 \cosh(2\lambda)}. \quad (17)$$

This recovers the scalar that conjugation matrices discard. Multiplying operators along a selected record is legitimate, but does not specify a Born-sampled ensemble until the instrument and its branch probabilities are fixed.

There are two symmetry distinctions. First, finite- ε jump branches superpose charge raising and lowering and generally break $U(1)$ covariance. At $\varepsilon = 0$ every branch has a definite charge change, so $e^{i\theta \hat{N}_F} \hat{\chi}^\dagger e^{-i\theta \hat{N}_F} = e^{i\theta} \hat{\chi}^\dagger$ and

its conjugate make the branch map $\hat{\rho} \mapsto \hat{K}\hat{\rho}\hat{K}^\dagger$ $U(1)$ covariant. The regulator is an auxiliary deformation, not a claim that the original protocol leaves its symmetry class. Equality of the measurement effects with a BPJ construction alone does not classify all feedback-dressed many-body evolution operators (mEOs).

Second, an odd Kraus operator does not create a parity-odd density operator. If $\hat{P} = (-1)^{\hat{N}_F}$, $\hat{P}\hat{K}\hat{P} = -\hat{K}$, and $\hat{P}\hat{\rho}\hat{P} = \hat{\rho}$, then

$$\hat{P}(\hat{K}\hat{\rho}\hat{K}^\dagger)\hat{P} = \hat{K}\hat{\rho}\hat{K}^\dagger. \quad (18)$$

A definite-parity state can switch sectors without acquiring coherence between them. A globally parity-preserving dilation supplies the exchanged parity through an ancilla. No extension of symmetry-class POVM no-go theorems is needed for these elementary statements.

IV. ORDERED PRODUCTS AND A CONTROLLED SINGULAR LIMIT

A. Composition before the limit

Fix a finite record $\mathbf{m}$ and write its chronological operator as

$$\hat{V}_{\mathbf{m}} = \hat{O}_q \cdots \hat{O}_1, \quad (19)$$

where the rightmost factor acts first. The factors can include gain, loss, occupation projectors, and invertible charge-conserving Gaussian operations. Any feedback schedule is held fixed by the record. Replace each jump by $\hat{j}_\pm(\varepsilon_k)$ and each projector by $\hat{\mathbf{1}} - \hat{n} + \varepsilon_k \hat{n}$ or $\hat{n} + \varepsilon_k(\hat{\mathbf{1}} - \hat{n})$ as appropriate. At every nonzero regulator the product $\hat{W}(\varepsilon)$ is invertible and Gaussian.

With BPJ's row-index convention,

$$\hat{V}\hat{\gamma}_a\hat{V}^{-1} = \sum_b t[\hat{V}]_{ab}\hat{\gamma}_b, \quad t[\hat{V}_2\hat{V}_1] = t[\hat{V}_1]t[\hat{V}_2]. \quad (20)$$

The order reversal follows by applying $\hat{V}_2$ to each operator in $\hat{V}_1\hat{\gamma}_a\hat{V}_1^{-1}$. Thus $t[\hat{W}] = t[\hat{O}_1] \cdots t[\hat{O}_q]$ with the regulated factors understood; transposed sTMs instead multiply in the same order as Eq. (19). Scalar branch weights do not appear in either convention. An even quadratic factor has determinant +1; a product with r odd factors has determinant $(-1)^r$.

The product must be formed with its chronological data intact. Already on one mode,

$$\hat{\chi}\hat{\chi}^\dagger = \hat{\mathbf{1}} - \hat{n}, \quad \hat{\chi}^\dagger\hat{\chi} = \hat{n}, \quad (21)$$

whereas opposite diagonal “zero/pole” factors can multiply to the identity and erase this distinction. At finite regulator the full Nambu matrix, including its anomalous blocks, is needed. If its blocks are $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$, the upper-left block of a product contains $A_1A_2 + B_1C_2$, not just A_1A_2 .

B. What converges, and under which assumptions

For a fixed finite number of factors and fixed mode wavefunctions, polynomial continuity gives, in any finite-dimensional operator norm,

$$\hat{W}(\varepsilon) = \hat{V}_{\mathbf{m}} + O(\eta), \quad \eta = \max_k |\varepsilon_k|. \quad (22)$$

The same statement holds with the physical normalizations in Eqs. (14) and (16). For an input with $\|\hat{V}_{\mathbf{m}}|\psi\rangle\| > 0$, normalization is continuous, so the normalized branch converges independently of regulator ratios. The constants depend on the word and its nonzero success probability; this is not a uniform statement at large depth or near a forbidden outcome.

For an operator-level version choose a normalized, maximally entangled Gaussian reference with N reference modes and N physical modes,

$$|\text{REF}\rangle = 2^{-N/2} \prod_{i=1}^N (\hat{\psi}_{i,R}^\dagger + \hat{\psi}_{i,P}^\dagger)|0\rangle, \quad \|\hat{V}_P|\text{REF}\rangle\|^2 = 2^{-N} \text{Tr}(\hat{V}^\dagger\hat{V}). \quad (23)$$

A fixed fermionic mode ordering is implicit. If $\hat{V} \neq 0$, its normalized Choi vector and full covariance are therefore limits of those of $\hat{W}$, irrespective of how the regulators approach zero. This is a concrete state-level meaning of the boundary, with no assertion that a chosen homogeneous transfer chart remains finite or invertible.

There is a useful sharper rate for *jump-only regularization*, with charge-conserving invertible factors between jumps and a fixed-charge input (including the reference above). Flipping one regulated jump from gain to loss or vice versa changes the output charge by two. Thus

$$|\Psi(\varepsilon)\rangle = |\Psi_0\rangle + \sum_k \varepsilon_k |\Psi_k\rangle + O(\eta^2) \quad (24)$$

has no first-order cross term in its norm or in any charge-neutral bilinear expectation. For a nonzero limiting branch,

$$G(\varepsilon) - G(0) = O(\eta^2), \quad \langle \hat{\psi}_i \hat{\psi}_j \rangle_\varepsilon = O(\eta). \quad (25)$$

The *normal* correlation block converges quadratically; the full Majorana covariance generally converges only linearly. If projectors are also regularized, their first-order corrections need not change charge, so only the general $O(\eta)$ bound is asserted.

The nonzero-branch hypothesis cannot be removed. Two same-mode gains give zero exactly, but their regulated product is

$$(\hat{\chi}^\dagger + \varepsilon_2 \hat{\chi})(\hat{\chi}^\dagger + \varepsilon_1 \hat{\chi}) = \varepsilon_1 \hat{n} + \varepsilon_2 (\hat{\mathbf{1}} - \hat{n}). \quad (26)$$

Normalizing this vanishing operator can retain the arbitrary ratio $\varepsilon_1/\varepsilon_2$. There is no exact physical branch to which that normalized state should converge.

V. THE EXACT ENDPOINT WITHOUT A REGULATOR

The regularization explains the geometry, but exact state propagation need not use ill-conditioned matrices. Let a normalized pure Slater state have occupied projector $C = FF^\dagger$, with $F^\dagger F = \mathbf{1}$. For a gain, decompose its column as $\chi = C\chi + (\mathbf{1} - C)\chi$. Creation in the occupied component vanishes by Pauli exclusion. The surviving vector is orthogonal to the old occupied space, so it adds one normalized column. A loss removes the occupied component $C\chi$. Consequently,

$$\begin{aligned} p_+ &= \chi^\dagger (\mathbf{1} - C) \chi, & C_+ &= C + \frac{(\mathbf{1} - C) \chi \chi^\dagger (\mathbf{1} - C)}{p_+}, \\ p_- &= \chi^\dagger C \chi, & C_- &= C - \frac{C \chi \chi^\dagger C}{p_-}. \end{aligned} \quad (27)$$

These expressions require $p_\pm > 0$; a zero denominator signals a vanishing branch, not a prescription for a pseudoinverse. They give $C_\pm^2 = C_\pm$, $\text{tr } C_\pm = \text{tr } C \pm 1$, $C_+ \chi = \chi$, and $C_- \chi = 0$. They are the pure-state specialization of Gaussian measurement propagation [2].

The two projective branches needed alongside the jumps follow from $\hat{n} = \hat{\chi}^\dagger \hat{\chi}$ and $\hat{\mathbf{1}} - \hat{n} = \hat{\chi} \hat{\chi}^\dagger$. After loss the mode is empty, so refilling it adds $P_\chi = \chi \chi^\dagger$; after gain the mode is filled, so emptying it subtracts the same projector. Therefore

$$C_{\hat{n}} = C_- + P_\chi, \quad p_{\hat{n}} = p_-, \quad C_{\hat{\mathbf{1}} - \hat{n}} = C_+ - P_\chi, \quad p_{\hat{\mathbf{1}} - \hat{n}} = p_+. \quad (28)$$

Equations (27) and (28) handle all four outcomes of the exact fill/deplete instruments using only one-particle matrices. The second step of either projector factorization has conditional probability one, explaining the displayed probabilities.

For an intervening even operator $\hat{V} = \exp[-\sum_{ij} M_{ij} \hat{\psi}_i^\dagger \hat{\psi}_j]$, define $A = e^{-M} = t^{-1}$. It sends the occupied columns to AF , giving

$$C' = AF(F^\dagger A^\dagger AF)^{-1} F^\dagger A^\dagger, \quad \|\hat{V}|F\rangle\|^2 = \det(F^\dagger A^\dagger AF). \quad (29)$$

An additional scalar multiplying $\hat{V}$ contributes its squared modulus to the norm. These formulas follow from the Gram matrix of the transformed Slater columns, and do not assume A is unitary. Using Eq. (27) after each even step automatically retains chronology and Pauli blocking. The product of successive conditional norm factors is the norm squared of the complete word; those factors are Born probabilities only when the operations belong to the specified instrument.

VI. CONSEQUENCES FOR TRAJECTORY SPECTRA

The group description does not by itself identify a physical Lyapunov spectrum. A single regulated jump has logarithmic sTM singular values $\pm \log(1/\varepsilon)$, but these are regulator scales, not long-time rates. For a particularly simple counterexample, repeating the same normalized odd element twice gives $\hat{v}_+^2 = \hat{\mathbf{1}}$ and $t[\hat{v}_+]^2 = \mathbf{1}$. At fixed $\varepsilon > 0$ an arbitrarily long repetition has zero asymptotic sTM growth rates even though one step becomes arbitrarily ill-conditioned as $\varepsilon \rightarrow 0$. For the rescaled operator, $\hat{j}_+^2 = \varepsilon \hat{\mathbf{1}}$, so the many-body scalar has a different fate. The regulator limit and the long-record limit cannot be interchanged on the strength of Eq. (22).

The physical endpoint can instead be characterized directly. For a nonzero exact Gaussian record with a definite charge shift, propagate a maximally mixed input and set

$$\hat{\rho}_{\mathbf{m}} = \frac{\hat{V}_{\mathbf{m}} \hat{V}_{\mathbf{m}}^\dagger}{Z_{\mathbf{m}}}, \quad Z_{\mathbf{m}} = \text{Tr}(\hat{V}_{\mathbf{m}} \hat{V}_{\mathbf{m}}^\dagger) = 2^N p_{\mathbf{m}}. \quad (30)$$

The endpoint is number-conserving Gaussian. If its occupation eigenvalues are ν_a , natural-orbital factorization gives its Fock-space eigenvalues and hence the squared singular values of the mEO:

$$\sigma_{\mathbf{n}}(\hat{V}_{\mathbf{m}})^2 = Z_{\mathbf{m}} \prod_a \nu_a^{n_a} (1 - \nu_a)^{1-n_a}, \quad n_a \in \{0, 1\}. \quad (31)$$

The exact occupations 0 and 1 encode genuine zero singular values; the preferred occupation factor equals one, including at an endpoint. There is no need to turn them into small nonzero numbers by clipping. Covariance eigenvalues determine relative many-body singular values, while the record probability restores their absolute scale. This is an operator spectrum, distinct from the derivative of normalized conditional state evolution with respect to its initial state.

The conclusion is that gain/loss feedback admits a controlled Gaussian boundary representation, and exact covariance propagation remains well defined on each successful branch. This observation is not a proof of criticality, a determination of central charge, or an identification of every regulator-stable one-particle rate with a many-body or tangent Lyapunov exponent.

-
- [1] A. Bhuiyan, H. Pan, and C.-M. Jian, [Physical Review Research](#) **8**, 023147 (2026), [arXiv:2507.13437 \[quant-ph\]](#).
 - [2] S. Bravyi, [Quantum Information and Computation](#) **5**, 216 (2005), [arXiv:quant-ph/0404180](#).