

Optimization 1 — Tutorial 6

November 26, 2020

Convex Function

A function $f: C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ for a convex set C is called convex if

$$f(\lambda \mathbf{x} + (1 - \lambda) \mathbf{y}) \leq \lambda f(\mathbf{x}) + (1 - \lambda) f(\mathbf{y}),$$

for any $\mathbf{x}, \mathbf{y} \in C$ and $\lambda \in [0, 1]$. We say that f is strictly convex if the above is satisfied with a strict inequality and for every $\lambda \in (0, 1)$. We say that f is concave if $-f$ is convex.

Level Set

The level set of $f: S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ with level $\alpha \in \mathbb{R}$ is

$$\text{Lev}(f, \alpha) = \{\mathbf{x} \in S: f(\mathbf{x}) \leq \alpha\} \subseteq \mathbb{R}^n.$$

Epigraph

The epigraph of a function $f: C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ for a convex set C is

$$\text{epi}(f) = \{(\mathbf{x}, t) \in C \times \mathbb{R}: f(\mathbf{x}) \leq t\} \subseteq \mathbb{R}^{n+1}.$$

Quasi-Convex

A function $f: C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ for a convex set C is called quasi-convex if $\text{Lev}(f, \alpha)$ is a convex set for every $\alpha \in \mathbb{R}$.

Remark: a convex function is quasi-convex, but a quasi-convex function is not necessarily convex.

Characterization of Convex Functions

Let $f: C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be a function defined over a convex set C .

1. Gradient Inequality: suppose that f is continuously differentiable over C . Then, f is convex if and only if

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}), \quad \forall \mathbf{x}, \mathbf{y} \in C.$$

2. Second-order Characterization: suppose that f is twice continuously differentiable over C and that C is open. Then f is convex if and only if $\nabla^2 f(\mathbf{x}) \succeq 0$ for any $\mathbf{x} \in \mathbb{R}$. Additionally, if $\nabla^2 f(\mathbf{x}) \succ 0$ for any $\mathbf{x} \in \mathbb{R}$ then f is strictly convex.

Jensen's Inequality

Let $f: C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function defined over a convex set C . Then, for every $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in C$ and $\boldsymbol{\lambda} \in \Delta_k$ we have

$$f\left(\sum_{i=1}^k \lambda_i \mathbf{x}_i\right) \leq \sum_{i=1}^k \lambda_i f(\mathbf{x}_i).$$

Problem 1

For each of the following functions, determine whether it is convex, concave, quasi-convex or quasi-concave.

(a) $f(x) = e^x - 1$ over \mathbb{R} .

(b) $f(x, y) = xy$ over \mathbb{R}_{++}^2 .

(c) $f(x, y) = \frac{1}{xy}$ over \mathbb{R}_{++}^2 .

(d) $f(x, y) = \frac{x}{y}$ over \mathbb{R}_{++}^2 .

(e) $f(\mathbf{x}) = \sqrt{\|\mathbf{x}\|^2 + 1}$ over \mathbb{R}^n .

Problem 2

Arithmetic-Geometric Mean (AGM) Inequality. Prove that for any $x_1, x_2, \dots, x_n \geq 0$, the following inequality holds:

$$\frac{1}{n} \sum_{i=1}^n x_i \geq \sqrt[n]{\prod_{i=1}^n x_i}.$$

In general, for any $\boldsymbol{\lambda} \in \Delta_n$ we have

$$\sum_{i=1}^n \lambda_i x_i \geq \prod_{i=1}^n x_i^{\lambda_i}.$$

Problem 3

Young's Inequality. Prove that for any $s, t \geq 0$ and $p, q \geq 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$st \leq \frac{s^p}{p} + \frac{t^q}{q}.$$

Problem 4

Hölder's Inequality. Prove that for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $p, q \geq 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$|\mathbf{x}^T \mathbf{y}| \leq \|\mathbf{x}\|_p \|\mathbf{y}\|_q.$$