



# 2D Geometrical Transformations

# 2D Translation



# 2D Translation Equation




$$\mathbf{P} = (x, y)$$

$$\mathbf{t} = (t_x, t_y)$$

$$\mathbf{P}' = \mathbf{P} + \mathbf{t} = (x + t_x, y + t_y)$$

# 2D Translation using Matrices



$$\mathbf{P} = (x, y)$$

$$\mathbf{t} = (t_x, t_y)$$

$$\mathbf{P}' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates

- Multiply the coordinates by a non-zero scalar and add an extra coordinate equal to that scalar. For example,

$$(x, y) \rightarrow (x \cdot z, y \cdot z, z) \quad z \neq 0$$

$$(x, y, z) \rightarrow (x \cdot w, y \cdot w, z \cdot w, w) \quad w \neq 0$$

# Back to Cartesian Coordinates:


- Divide by the last coordinate and eliminate it. For example,

$$(x, y, z) \quad z \neq 0 \rightarrow (x / z, y / z)$$

$$(x, y, z, w) \quad w \neq 0 \rightarrow (x / w, y / w, z / w)$$

- NOTE: in our example the scalar was 1

# 2D Translation using Homogeneous Coordinates




$$\mathbf{P} = (x, y) \rightarrow (x, y, 1)$$

$$\mathbf{t} = (t_x, t_y) \rightarrow (t_x, t_y, 1)$$

$$\mathbf{P}' \rightarrow \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$


$= \begin{bmatrix} \mathbf{I} & \mathbf{t} \\ 0 & 1 \end{bmatrix} \cdot \mathbf{P} = \mathbf{T} \cdot \mathbf{P}$

# Scaling





# Scaling Equation




$$\mathbf{P} = (x, y) \rightarrow \mathbf{P}' = (s_x x, s_y y)$$

$$\mathbf{P} = (x, y) \rightarrow (x, y, 1)$$

$$\mathbf{P}' = (s_x x, s_y y) \rightarrow (s_x x, s_y y, 1)$$

$$\mathbf{P}' \rightarrow \begin{bmatrix} s_x x \\ s_y y \\ 1 \end{bmatrix} = \underbrace{\begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix}}_{\mathbf{S}} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} \mathbf{S}' & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{bmatrix} \cdot \mathbf{P} = \mathbf{S} \cdot \mathbf{P}$$

# Scaling & Translating



$$P' = S \cdot P$$

$$P'' = T \cdot P'$$

$$P'' = T \cdot P' = T \cdot (S \cdot P) = (T \cdot S) \cdot P = A \cdot P$$

# Scaling & Translating


$$\begin{aligned}\mathbf{P}'' &= \mathbf{T} \cdot \mathbf{S} \cdot \mathbf{P} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \\ &= \underbrace{\begin{bmatrix} s_x & 0 & t_x \\ 0 & s_y & t_y \\ 0 & 0 & 1 \end{bmatrix}}_A \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} S & t \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + t_x \\ s_y y + t_y \\ 1 \end{bmatrix}\end{aligned}$$

# Translating & Scaling = Scaling & Translating ?

$$\mathbf{P}''' = \mathbf{T} \cdot \mathbf{S} \cdot \mathbf{P} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & t_x \\ 0 & s_y & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + t_x \\ s_y y + t_y \\ 1 \end{bmatrix}$$


$$\begin{aligned} \mathbf{P}''' = \mathbf{S} \cdot \mathbf{T} \cdot \mathbf{P} &= \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \\ &= \begin{bmatrix} s_x & 0 & s_x t_x \\ 0 & s_y & s_y t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} s_x x + s_x t_x \\ s_y y + s_y t_y \\ 1 \end{bmatrix} \end{aligned}$$

# Rotation



# Rotation Equations

Counter-clockwise rotation by an angle  $\theta$



$$x' = \cos \theta \, x - \sin \theta \, y$$

$$y' = \cos \theta \, y + \sin \theta \, x$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\mathbf{P'} = \mathbf{R} \, \mathbf{P}$$

# Degrees of Freedom

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

R is 2x2  4 elements

Note: R belongs to the category of normal matrices and satisfies many interesting properties:

$$\mathbf{R} \cdot \mathbf{R}^T = \mathbf{R}^T \cdot \mathbf{R} = \mathbf{I}$$

$$\det(\mathbf{R}) = 1$$

# Rotation+ Scaling +Translation

$$\mathbf{P}' = (\mathbf{T} \mathbf{R} \mathbf{S}) \mathbf{P}$$

$$\mathbf{P}' = \mathbf{T} \cdot \mathbf{R} \cdot \mathbf{S} \cdot \mathbf{P} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} =$$

$$= \begin{bmatrix} \cos \theta & -\sin \theta & t_x \\ \sin \theta & \cos \theta & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} =$$

$$= \begin{bmatrix} \mathbf{R}' & \mathbf{t} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \mathbf{S} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \boxed{\begin{bmatrix} \mathbf{R}' \mathbf{S} & \mathbf{t} \\ 0 & 1 \end{bmatrix}} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

If  $s_x = s_y$ , this is a  
similarity  
transformation!