

Evaluation: document similarities

Integral formula of Minkowski type and new characterization of the Wulff shape

Yijun He *

Haizhong Li [†]

Abstract

Given a positive function F on S^n which satisfies a convexity condition, we introduce the r -th anisotropic mean curvature M_r for hypersurfaces in \mathbb{R}^{n+1} which is a generalization of the usual r -th mean curvature H_r . We get integral formulas of Minkowski type for compact hypersurfaces in \mathbb{R}^{n+1} . We give some new characterizations of the Wulff shape by use of our integral formulas of Minkowski type, in case $F = 1$ which reduces to some well-known results.

2000 Mathematics Subject Classification: Primary 53C42, 53A30; Secondary 53B25.

Key words and phrases: Wulff shape, F -Weingarten operator, anisotropic principal curvature, r -th anisotropic mean curvature, integral formula of Minkowski type.

[xiv.org/pdf/1408.0189](https://arxiv.org/pdf/1408.0189)

[xiv.org/pdf/math/0504051](https://arxiv.org/pdf/math/0504051)

[xiv.org/pdf/1112.3014](https://arxiv.org/pdf/1112.3014)

[xiv.org/pdf/1109.1922](https://arxiv.org/pdf/1109.1922)

[xiv.org/pdf/1408.4595](https://arxiv.org/pdf/1408.4595)

[xiv.org/pdf/0902.0616](https://arxiv.org/pdf/0902.0616)

[xiv.org/pdf/astro-ph/0508060](https://arxiv.org/pdf/astro-ph/0508060)