Forward · Inverse · Example's own settings, where a change helped

ODEs
- Integro-differential ODE
- Integro-differential ODE – Volterra kernel
- Linear ODE system †
- Lotka-Volterra ODE †
- Second-order ODE †
- Lorenz ODE – unknown C_1, C_2, C_3
- Lorenz ODE – unknown C_1, C_2, C_3, exogenous input

1D, steady
- Euler beam 1D †
- Fractional Poisson 1D †
- Poisson 1D – Dirichlet
- Poisson 1D – Dirichlet/Neumann
- Poisson 1D – Dirichlet/Periodic †
- Poisson 1D – Dirichlet/point-set operator
- Poisson 1D – Dirichlet/Robin
- Poisson 1D – hard BCs
- Poisson 1D – multi-scale Fourier features
- Brinkman-Forchheimer 1D – unknown ν_e, K †
- Diffusion-reaction 1D – unknown rate field † ‡
- Fractional Poisson 1D – unknown order α
- Poisson 1D – unknown forcing field †
- Poisson 1D – unknown forcing field (batches) †

1D, time-dependent
- Allen-Cahn 1D †
- Allen-Cahn 1D – Wight & Zhao (time-adaptive)
- Allen-Cahn 1D – Wight & Zhao (whole interval)
- Cahn-Hilliard 1D – Wight & Zhao
- Cahn-Hilliard 1D – Mattey & Ghosh † ‡
- Black-Scholes 1D – American put
- Black-Scholes 1D – European call
- Burgers 1D – Raissi et al. †
- Burgers 1D – RAR
- Diffusion 1D †
- Diffusion 1D – hard IC and BCs
- Diffusion 1D – resampling †
- Diffusion-reaction 1D †
- Fractional diffusion 1D
- Heat 1D – Dirichlet †
- Heat 1D – resampling †
- Klein-Gordon 1D †
- Schrödinger 1D †
- Wave 1D – space-time multi-scale Fourier features †
- Diffusion 1D – unknown coefficient †
- Diffusion-reaction 1D – unknown k_f, D

2D, steady
- Fractional Poisson 2D – disk
- Helmholtz 2D – Dirichlet
- Helmholtz 2D – hyperparameter search
- Helmholtz 2D – sound-hard scattering †
- Helmholtz 2D – square with hole †
- Laplace 2D – disk
- Linear elasticity 2D – plate †
- Linear elasticity 2D – plate, soft BCs
- Navier-Stokes 2D – Kovasznay flow †
- Poisson 2D – disk †
- Poisson 2D – L-shape †
- Fractional Poisson 2D – unknown order α

2D, time-dependent
- Allen-Cahn 2D – Wight & Zhao †
- Allen-Cahn 2D – Mattey & Ghosh, best setup †
- Cahn-Hilliard 2D – Wight & Zhao †
- Cahn-Hilliard 2D – Mattey & Ghosh
- Burgers 2D – Mathias et al. †
- Heat 2D †

3D
- Fractional Poisson 3D – sphere
- Poisson 3D – sphere
- Allen-Cahn 3D – Wight & Zhao †
- Heat 3D †
- Navier-Stokes 3D – Beltrami flow †

ω · d, a · a, b · C_1, C_3, C_2 · C_3, C_2, C_1 · EI · K, ν_e · α · c, d · γ_2, γ_1 · λ_2, λ_1 · D · a · c · C · D, k_f · Re · f · α · $\lambda\varepsilon^2, \lambda$ · c_2, c_1^2 · ε^2 · κ, α · ν · α · $\lambda\varepsilon^2, \lambda$ · α · ν

10^{-6} 10^{-5} 10^{-4} 10^{-3} 10^{-2} 10^{-1} 10^{0}

Relative L_2 error of the main output at the last reference time

0.001 0.1 10

Worst parameter error (%)

† in this study (filled: the chosen run); rows without † show the README's result. ‡ the run with the lowest training loss (shown) is more than 2× less accurate than another run.