Forward · Inverse · Example's own settings, where a change helped

ODEs
- Integro-differential ODE †
- Integro-differential ODE – Volterra kernel †
- Linear ODE system † — ω
- Lotka-Volterra ODE † — a, d
- Second-order ODE † — a, b
- Lorenz ODE – unknown C_1, C_2, C_3 † — C1, C3, C2
- Lorenz ODE – unknown C_1, C_2, C_3, exogenous input † — C3, C2, C1

1D, steady
- Euler beam 1D † — EI
- Fractional Poisson 1D †
- Poisson 1D – Dirichlet †
- Poisson 1D – Dirichlet/Neumann †
- Poisson 1D – Dirichlet/Periodic †
- Poisson 1D – Dirichlet/point-set operator †
- Poisson 1D – Dirichlet/Robin †
- Poisson 1D – hard BCs †
- Poisson 1D – multi-scale Fourier features †
- Brinkman-Forchheimer 1D – unknown ν_e, K † — K, ν_e
- Diffusion-reaction 1D – unknown rate field † ‡
- Fractional Poisson 1D – unknown order α † — α
- Poisson 1D – unknown forcing field †
- Poisson 1D – unknown forcing field (batches) †

1D, time-dependent
- Allen-Cahn 1D – Wight & Zhao † — gamma_2, gamma_1
- Cahn-Hilliard 1D – Wight & Zhao †
- Cahn-Hilliard 1D – Mattey & Ghosh † ‡
- Black-Scholes 1D – American put
- Black-Scholes 1D – European call †
- Burgers 1D – Raissi et al. † — lambda_1, lambda_2
- Burgers 1D – RAR †
- Diffusion 1D †
- Diffusion 1D – hard IC and BCs †
- Diffusion 1D – resampling †
- Diffusion-reaction 1D † — D
- Fractional diffusion 1D †
- Heat 1D – Dirichlet † — a
- Heat 1D – resampling †
- Klein-Gordon 1D † — γ
- Schrödinger 1D † — c
- Wave 1D – space-time multi-scale Fourier features † — C
- Diffusion 1D – unknown coefficient † — C
- Diffusion-reaction 1D – unknown k_f, D † — D, kf

2D, steady
- Fractional Poisson 2D – disk †
- Helmholtz 2D – Dirichlet † ‡ — k_0
- Helmholtz 2D – hyperparameter search
- Helmholtz 2D – sound-hard scattering †
- Helmholtz 2D – square with hole †
- Laplace 2D – disk †
- Linear elasticity 2D – plate † — μ, λ
- Linear elasticity 2D – plate, soft BCs †
- Mixed convection 2D – Soomro et al.
- Navier-Stokes 2D – Kovasznay flow † — Re
- Poisson 2D – disk † — f
- Poisson 2D – L-shape †
- Fractional Poisson 2D – unknown order α † — α

2D, time-dependent
- Allen-Cahn 2D – Wight & Zhao † — $\lambda\varepsilon^2$, λ
- Allen-Cahn 2D – Mattey & Ghosh † — c_2, c_1^2
- Cahn-Hilliard 2D – Wight & Zhao † ‡ — ε^2
- Cahn-Hilliard 2D – Mattey & Ghosh † ‡ — kappa, α
- Burgers 2D – Mathias et al. † — ν
- Heat 2D † ‡ — α

3D
- Fractional Poisson 3D – sphere †
- Poisson 3D – cube with spherical cut-out †
- Poisson 3D – sphere † — f
- Allen-Cahn 3D – Wight & Zhao † — $\lambda\varepsilon^2$, λ
- Heat 3D † — α
- Navier-Stokes 3D – Beltrami flow † — ν

10^{-6} 10^{-5} 10^{-4} 10^{-3} 10^{-2} 10^{-1} 10^{0}

Relative L_2 error of the main output at the last reference time

0.001 0.1 10

Worst parameter error (%)

† in this study (filled: the chosen run); rows without † show the README's result. ‡ the run with the lowest training loss (shown) is more than 2× less accurate than another run.