
Tuned settings
Starting settings
Starting settings kept
Inverse, each parameter
Forward problems
Inverse problems
ODEs
Lotka–Volterra ODE
Lorenz ODE
Integro-differential ODE
a, d
C3, C2, C1
1D, steady
Euler beam 1D
Poisson 1D – Dirichlet/Periodic
Poisson 1D – unknown forcing
Fractional Poisson 1D
EI
q(x)
α
1D, time-dependent
Burgers 1D
Allen–Cahn 1D
Cahn–Hilliard 1D
Schrödinger 1D
Klein–Gordon 1D
Wave 1D – MsFFN
Diffusion-reaction 1D
Black–Scholes 1D
λ1, λ2
γ2, γ1
c
γ
C
D
2D, steady
Helmholtz 2D – Dirichlet
Linear elasticity 2D
Navier–Stokes 2D – Kovasznay
Poisson 2D – disk
k0
μ, λ
Re
f
2D, time-dependent
Allen–Cahn 2D
Cahn–Hilliard 2D
Burgers 2D
Heat 2D
c2, c1²
ε²
ν
α
3D
Allen–Cahn 3D
Navier–Stokes 3D – Beltrami
λε², λ
ν
10⁻⁶ 10⁻⁵ 10⁻⁴ 10⁻³ 10⁻² 10⁻¹ 10⁰
Relative L2 error
at the final reference time
10⁻³ 10⁻¹ 10¹
Parameter or field
error (%)