





DEKA



INTRADERMAL VACCINE PATCH



COVID-19 VACCINE

MANUFACTURER: PFIZER-BIONTECH

INTRADERMAL PATCH APPLICATOR

ACTIVE INGREDIENTS: mRNA, lipids ((4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate), 2 [(polyethylene glycol)-2000]-N,N-ditetradecylacetamide, 1,2-Distearoyl-sn-glycero-3-phosphocholine, and cholesterol), potassium chloride, monobasic potassium phosphate, sodium chloride, dibasic sodium phosphate dihydrate, and sucrose.

USES: The Pfizer-BioNTech COVID-19 Vaccine is an unapproved vaccine that may prevent COVID-19. There is no FDA-approved vaccine to prevent COVID-19. The FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19 in individuals 16 years of age and older under an Emergency Use Authorization (EUA).

PRECAUTIONS: Tell the vaccination provider if you: • have any allergies • have a fever • have a bleeding disorder or are on a blood thinner • are immunocompromised or are on a medicine that affects your immune system • are pregnant or plan to become pregnant • are breastfeeding • have received another COVID-19 vaccine or another vaccine in the last 14 days

CONTRAINDICATIONS: You should not get the Pfizer-BioNTech COVID-19 Vaccine if you: • had a severe allergic reaction after a

previous dose of this vaccine • had a severe allergic reaction to any ingredient of this vaccine.

RISKS: There is a remote chance the Pfizer-BioNTech COVID-19 Vaccine could cause a severe allergic reaction. A severe allergic reaction would usually occur within a few minutes to one hour after. Your vaccination provider may ask you to stay for monitoring after vaccination.

Signs of a severe allergic reaction can include: • Difficulty breathing • Severe swelling of your face and throat • A fast heartbeat • A bad rash all over your body • Dizziness and weakness

Commonly reported side effects include: • rash, itching, hives, or mild swelling of the face • injection site pain • tiredness • headache • muscle pain • chills • joint pain • fever • injection site swelling • injection site redness • nausea • feeling unwell • swollen lymph nodes (lymphadenopathy) • diarrhea • vomiting • arm pain

How is the PFIZER-BIONTECH COVID-19 vaccine given? The Pfizer-BioNTech COVID-19 Vaccine will be given as an intradermal injection in 2 doses given 3 weeks apart.

INJECTION SITE SELECTION: Palm side of non-dominant forearm or either outer thigh. Do not use arm on the side of a prior mastectomy or a paralyzed limb, if possible. Delivery site should be free of pain, lumps, swelling, warmth, wounds, bruising, redness, infections, implanted ports or other devices. May be given through a tattoo.

DEKA
SELF ADMINSTRABLE VACCINE PATCH

PEEL

HOW TO USE THE PATCH APPLICATOR:

1. Cleanse hands (If handwashing is unavailable, use hand sanitizer with at least 60% alcohol to thoroughly sanitize hands).
2. Choose delivery site and clean 2 inch area with sterile alcohol prep pad.
3. Remove applicator patch from packaging.
4. Hold top of device in dominant hand and peel off removable liner from the flat side of the applicator with non-dominant hand.
5. Grasp the device firmly from the top and place the applicator securely onto skin at the pre-selected cleansed delivery site.
6. Push the center of the applicator firmly into skin to invert device petals and deliver vaccine (expect skin to spread taut beneath the applicator).
7. Leave the applicator in place for XX seconds.
8. To remove, quickly peel the applicator from skin.
9. Hold pressure to site as necessary.
10. Cleanse hands or use 60% alcohol hand sanitizer.
11. Return applicator to pharmacist and complete required observation period (15 or 30 minutes).
Report any adverse reactions.



Scan for detailed instructions