

Assignment 4

Team 7

June 23, 2021

Contents

1	Introduction	1
2	Archish S me20b032	1
3	Prabhat ME20B132	3
3.1	Terms used in the equation	3

List of Figures

1	An MDP models the synchronous interaction between agent and world	2
2	Propogation of Light with respect to magnetic and electric fields	3

1 Introduction

This is the submission file for Assignment 4.

The Authors:

- ME20B020 AKSHAT RAKESH GARHWAL
- ME20B032 ARCHISH S
- ME20B132 PRABHAT BEDIDA
- MM20B005 ALBIN GEORGE
- MM20B049 PRITHVIRAJ PRATAP BHOSLE

2 Archish S me20b032

A Markov decision process [2] can be described as a tuple $\langle S, A, T, R \rangle$, where

- S is a finite set of states of the world;
- A is a finite set of actions;
- $T : S \times A \rightarrow \Pi(S)$ is the *state-transition function*, giving for each world state and agent action, a probability distribution over world states (we write $T(s, a, s')$ for the probability of ending in state s' , gievn that the agent starts in state s and takes action a);

- $R : S \times A \rightarrow \mathbb{R}$ is the reward function, giving the expected immediate reward gained by the agent for taking each action in each state (we write $R(s, a)$ for the expected reward for taking action a in state s);
- A stationary policy, $\pi : S \rightarrow A$, is a situation-action mapping that specifies, for each state, an action to be taken.
- $V_\pi(s)$ is the expected discounted sum of future reward for starting in state s and executing policy π .




Figure 1: An MDP models the synchronous interaction between agent and world

In this model, as described by figure 1, the next state and the expected reward depend only on the previous state and the action taken; even if we were to condition on additional previous states, the transition probabilities and the expected rewards would remain the same. This is known as the Markov property.

$$V_\pi(s) = R(s, \pi(s)) + \gamma \sum_{s' \in S} T(s, \pi(s), s') V_\pi(s') \quad (1)$$

Given the Value Function 1 a greedy policy with respect to that value function, π_V , is defined as

$$\pi_V(s) = \operatorname{argmax}_a \left[R(s, a) + \gamma \sum_{s' \in S} T(s, a, s') V(s') \right] \quad (2)$$

3 Prabhat ME20B132

$$\nabla \times \vec{B} = \mu_0 \vec{J} + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t} \quad (3)$$

Eqn. 3 is commonly known as Maxwell's fourth electromagnetic equation. It is an expansion on Ampère's Circuital Law and is hence also known as the Ampère-Maxwell Law. Currently a more generalized form is used, which takes into account the behaviour and interference of material substances in electric and magnetic fields [1].

3.1 Terms used in the equation

- $\nabla \times$ - Curl Operator
- \vec{B} - Magnetic Field Vector
- \vec{J} - Current Density Vector
- \vec{E} - Electric Field Vector
- μ_0 (Constant) - Permeability of Free Space
- ϵ_0 (Constant) - Permittivity of Free Space

It is an important equation as it links two phenomena, which were previously thought to be separate, magnetism and electricity. It also finally proved that light is an electromagnetic wave, (as seen in Fig. 2 taken from a paper by J. Gratus, M.W.McCall and P.Kinsler [1]) leading to a whole new area and direction of research.




Figure 2: Propagation of Light with respect to magnetic and electric fields

References

- [1] J. Gratus, M.W. McCall, and P. Kinsler. Electromagnetism, axions, and topology: A first-order operator approach to constitutive responses provides greater freedom. *Physical Review A*, 101(4), 2020. cited By 2.
- [2] L.P. Kaelbling, M.L. Littman, and A.R. Cassandra. Planning and acting in partially observable stochastic domains. *Artificial Intelligence*, 101(1-2):99–134, 1998.