

Exam Notes

Made by Leon :) *Note: Any reference numbers are to the lecture notes*

1 Revisiting FPM

Definition 1.1: Nested Sequences

A sequence $(I_n)_{n \in \mathbb{N}}$ of sets is said to be **nested** if

$$I_1 \subset I_2 \subset I_3 \subset \dots$$

Theorem 1.1: Nested Interval Property

If (I_n) is a nested sequence of nonempty closed bounded intervals then

$$E = \bigcap_{n \in \mathbb{N}} I_n = \{x \in \mathbb{R} : x \in I_n, \forall n \in \mathbb{N}\}$$

is nonempty (i.e. it contains at least one number). Moreover if $\lambda(I_n) \rightarrow 0$, where $\lambda(I_n)$ denotes the length of interval I_n , then E contains exactly one number

Theorem 1.2: Covers

Let E be a subset of \mathbb{R}^n

- A **cover** of E is a collection of sets $\{I_\alpha\}_{\alpha \in A}$ such that

$$E \subseteq \bigcup_{\alpha \in A} I_\alpha$$

- An **open covering** of E is a cover such that each I_α is open, i.e. (a, b) compared to $[a, b]$
- A **finite subcover** of E is a collection of sets $(I_\alpha)_{\alpha \in A_0}$ where there exists a subset $A_0 = \{\alpha_1, \alpha_2, \dots, \alpha_N\}$ of A such that $(I_\alpha)_{\alpha \in A_0}$ is a finite subset of $(I_\alpha)_{\alpha \in A}$ that is also a cover
- The set E is said to be **compact** iff every open covering of E has a **finite subcovering**; that is

$$E \subseteq \bigcup_{j=1}^N I_{a_j} \quad \text{or} \quad E \subseteq I_{\alpha_1} \cup I_{\alpha_2} \cup \dots \cup I_{\alpha_N}$$

Definition 1.2: Epsilon-N Convergence of Sequence

A sequence of real numbers (x_n) is said to **converge** to a real number $a \in \mathbb{R}$ iff for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$n \geq N \text{ implies } |x_n - a| < \epsilon$$

If (x_n) converges to a , we will write $\lim_{n \rightarrow \infty} x_n = a$, or $x_n \rightarrow a$. The number a is called the limit of the sequence (x_n) . A sequence that does not converge to some real number is said to *diverge

Definition 1.3: Cauchy Sequence

A sequence (x_n) of numbers $x_n \in \mathbb{R}$ is said to be **Cauchy** if for every $\epsilon > 0$ there is $N \in \mathbb{N}$ such that

$$|x_n - x_m| < \epsilon \quad \forall n, m \geq N$$

Theorem 1.3: Convergent Sequences are Cauchy

Let (x_n) be a sequence of real numbers. Then (x_n) is a Cauchy sequence if and only if (x_n) is a convergent sequence.
Note: This works both ways $((x_n)$ is a convergent seq \implies Cauchy)

Definition 1.4: Subsequences

Suppose $(x_n)_{n \in \mathbb{N}}$ is a sequence. A subsequence of this sequence is a sequence of the form $(x_{n_k})_{k \in \mathbb{N}}$ where for each k there is a positive integer n_k such that

$$n_1 < n_2 < \dots < n_k < n_{k+1} < \dots$$

Thus, $(x_n)_{n \in \mathbb{N}}$ is just a selection of some (possibly all) of the x_n 's taken in order

Theorem 1.5: Bolzano-Weierstrass

Every bounded sequence of real numbers has a convergent subsequence

Definition 1.5: Limit Superior and Inferior

If (x_n) is a bounded sequence of real numbers we denote by

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} x_k \right), \quad \liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\inf_{k \geq n} x_k \right)$$

Note: These are only defined for bounded sequences

- If (x_n) is not bounded from above then we write $\limsup_{n \rightarrow \infty} x_n = +\infty$
- If (x_n) is not bounded from below then we write $\liminf_{n \rightarrow \infty} x_n = -\infty$

Theorem 1.6: Convergence from Limsup and Liminf

A sequence (x_n) of real numbers is convergent if and only if $\limsup_{n \rightarrow \infty} x_n$ and $\liminf_{n \rightarrow \infty} x_n$ are real numbers and

$$\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n$$

Definition 1.6: Convergent Infinite Series

Let $S = \sum_{k=1}^{\infty} a_k$ be an infinite series a_k . For each $n \in \mathbb{N}$, the partial sum of S of order n is defined by

$$s_n = \sum_{k=1}^n a_k$$

S is said to **converge** iff its sequence of partial sums (s_n) converges to some $s \in \mathbb{R}$ as $n \rightarrow \infty$; that is, iff for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that for all $n \geq N$ we have $|s_n - s| < \epsilon$. In this case we shall write

$$\sum_{k=1}^{\infty} a_k = s$$

and call s the **sum** or **value** of the series $\sum_{k=1}^{\infty} a_k$

A series $S = \sum_{k=1}^{\infty} a_k$ is said to be **absolutely convergent** if the series $\sum_{k=1}^{\infty} |a_k|$ is convergent. A series is called **conditionally convergent** if it is convergent but not absolutely convergent.

Theorem 1.7: Cauchy Criterion

Let $S = \sum_{k=1}^{\infty} a_k$ be a series. Then the series S is convergent iff for any $\epsilon > 0$ there exists N such that for all $m \geq n \geq N$ we have that

$$\left| \sum_{k=n+1}^m a_k \right| < \epsilon$$

Theorem 1.8: Rearrangements of Abs. Convergent Series

Let $S = \sum_{k=1}^{\infty} a_k$ be an absolutely convergent series. Then

- The series S is convergent
- Let $z : \mathbb{N} \rightarrow \mathbb{N}$ be a bijection. Then the series $\sum_{k=1}^{\infty} a_{z(k)}$ is convergent and

$$\sum_{k=1}^{\infty} a_k = \sum_{k=1}^{\infty} a_{z(k)}$$

The series $\sum_{k=1}^{\infty} a_{z(k)}$ is called a **rearrangement** of the series $\sum_{k=1}^{\infty} a_k$. What we do here is add the terms of the sum in a different order to the original one, for example

$$a_3 + a_7 + a_1 + a_{100} + a_2 + \dots$$

Since $z : \mathbb{N} \rightarrow \mathbb{N}$ is a bijection, we will miss no terms.

Theorem 1.9: Rearrangements of Cond. Convergent Series

Let $S = \sum_{k=1}^{\infty} a_k$ be any conditionally convergent series. Then there exists rearrangements $z : \mathbb{N} \rightarrow \mathbb{N}$ (where z is a bijection) such that

- For any $r \in \mathbb{R}$ the series $\sum_{k=1}^{\infty} a_{z(k)}$ is conditionally convergent and its sum is r
- The series $\sum_{k=1}^{\infty} a_{z(k)}$ diverges to $+\infty$
- The series $\sum_{k=1}^{\infty} a_{z(k)}$ diverges to $-\infty$
- The partial sums of the series $\sum_{k=1}^{\infty} a_{z(k)}$ oscillate between any two real numbers

Definition 1.7: Continuity

Let f be a function $f : \text{dom}(f) \rightarrow \mathbb{R}$ where $\text{dom}(f) \subset \mathbb{R}$. We say that f is continuous at some $a \in \text{dom}(f)$ if for any sequence (x_n) whose terms lie in $\text{dom}(f)$ and which converges to a , we have $\lim_{n \rightarrow \infty} f(x_n) = f(a)$. If f is continuous at each $a \in S \subset \text{dom}(f)$ then we say f is continuous on S . If f is continuous on $\text{dom}(f)$ then we say f is continuous

Theorem 1.10: Properties of Continuity

Let $f, g : D \rightarrow \mathbb{R}$ be continuous on D , and let $\alpha \in \mathbb{R}$. Then the following functions are continuous on D :

1. αf
2. $f + g$
3. $f g$

Definition 1.8: Composition

Let $A, B \subseteq \mathbb{R}$ be nonempty, let $f : A \rightarrow \mathbb{R}$, $g : B \rightarrow \mathbb{R}$ and $f(A) \subseteq B$. The composition of g with f is the function $g \circ f : A \rightarrow \mathbb{R}$ defined by

$$(g \circ f)(x) = g(f(x)), \quad \text{for all } x \in A$$

Theorem 1.11: Continuity of Composition

If f is continuous at $a \in \mathbb{R}$ and g is continuous at $f(a)$ then the composition $g \circ f$ is continuous at a

Theorem 1.12: $\epsilon - \delta$ definition of continuity

Let f be a function $f : \text{dom}(f) \rightarrow \mathbb{R}$ where $\text{dom}(f) \subset \mathbb{R}$. Then f is continuous at $a \in \text{dom}(f)$ iff for any $\epsilon > 0$ there exists $\delta > 0$ s.t. whenever $x \in \text{dom}(f)$ and $|x - a| < \delta$ we have $|f(x) - f(a)| < \epsilon$

Definition 1.13: Intermediate Value Theorem

Let $a < b$ real numbers and $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$. If $f(a)f(b) < 0$ then there exists at least one $c \in (a, b)$ s.t. $f(c) = 0$

Definition 1.14: Extreme Value Theorem

Let $a < b$ real numbers and $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$. Then there exists points $c, d \in [a, b]$ s.t.

$$f(c) = \inf\{f(x) : x \in [a, b]\}, \quad f(d) = \sup\{f(x) : x \in [a, b]\}$$

That is, the function f on the interval $[a, b]$ is bounded and attains its minimal value at some point $c \in [a, b]$. Similarly, the maximal value of f is also attained at some point $d \in [a, b]$

2 Uniform convergence

Definition 2.1: Pointwise Convergence

Let E be a nonempty subset of \mathbb{R} . A sequence of functions $f_n : E \rightarrow \mathbb{R}$ is said to **converge pointwise** on E , written $f_n \rightarrow f$ pointwise on E as $n \rightarrow \infty$, iff $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ exists for each $x \in E$

Let E be a nonempty subset of \mathbb{R} . Then a sequence of functions f_n converges pointwise on E , as $n \rightarrow \infty$, iff for every $\epsilon > 0$ and $x \in E$ there is an $N \in \mathbb{N}$ (which may depend on x as well as ϵ) such that

$$n \geq N \quad \text{implies} \quad |f_n(x) - f(x)| < \epsilon$$

Remarks:

- The pointwise limit of continuous (respectively, differentiable) functions is not necessarily continuous (respectively, differentiable).
- The pointwise limit of integrable functions is not necessarily integrable.
- There exist continuous functions f_n and f such that $f_n \rightarrow f$ pointwise on $[0, 1]$ but

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx \neq \int_0^1 \left(\lim_{n \rightarrow \infty} f_n(x) \right) dx$$

Definition 2.2: Uniform Convergence

Let E be a nonempty subset of \mathbb{R} . A sequence of functions $f_n : E \rightarrow \mathbb{R}$ is said to **converge uniformly** on E to a function f (notation: $f_n \rightarrow f$ uniformly on E as $n \rightarrow \infty$) if and only if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that for all $x \in E$

$$n \geq N \quad \text{implies} \quad |f_n(x) - f(x)| < \epsilon$$

Remark 2.2: Differences between Pointwise and Uniform

Let E be a nonempty subset of \mathbb{R} .

- A sequence of functions f_n **converges pointwise** on E , as $n \rightarrow \infty$, if and only if for every $\epsilon > 0$ and $x \in E$ there is an $N \in \mathbb{N}$ (which may depend on x as well as ϵ) such that

$$n \geq N \quad \text{implies} \quad |f_n(x) - f(x)| < \epsilon$$

- A sequence of functions $f_n : E \rightarrow \mathbb{R}$ **converges uniformly** on E iff for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that for all $x \in E$

$$n \geq N \quad \text{implies} \quad |f_n(x) - f(x)| < \epsilon$$

For a sequence of functions to be pointwise convergent, it is enough to have an N_n for every x_n , but for it to be uniformly convergent, it has to have **the same** N for every x in the sequence

Theorem 2.1: Equivalence of Uniform Convergence

The following are equivalent concerning a sequence of functions $f_n : E \rightarrow \mathbb{R}$ and $f : E \rightarrow \mathbb{R}$:

- $f_n \rightarrow f$ uniformly on E
- $\sup_{x \in E} |f_n(x) - f(x)| \rightarrow 0$ as $n \rightarrow \infty$
- there exists a sequence $a_n \rightarrow 0$ s.t. $|f_n(x) - f(x)| \leq a_n, \forall x \in E$

Theorem 2.1

Let E be a nonempty subset of \mathbb{R} and suppose that $f_n \rightarrow f$ uniformly on E as $n \rightarrow \infty$. If each f_n is continuous at some $x_0 \in E$, then f is continuous at x_0

Definition 2.2: Uniformly Bounded Sequences

A sequence of functions f_n is said to be **uniformly bounded** on a set E if there is a $M > 0$ such that $|f_n(x)| \leq M$ for all $x \in E$ and all $n \in \mathbb{N}$

Theorem 2.2

Suppose that $f_n \rightarrow f$ uniformly on a closed interval $[a, b]$. If each f_n is integrable on $[a, b]$, then so is f and

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b \left(\lim_{n \rightarrow \infty} f_n(x) \right) dx$$

Theorem 2.3

Let (a, b) be a bounded interval and suppose that f_n is a sequence of functions which converges at some $x_0 \in (a, b)$. If each f_n is differentiable on (a, b) , and f'_n converges uniformly on (a, b) as $n \rightarrow \infty$, then f_n converges uniformly on (a, b) and

$$\lim_{n \rightarrow \infty} f'_n(x) = \left(\lim_{n \rightarrow \infty} f_n(x) \right)'$$

Definition 2.3: Convergence of series

Let f_k be a sequence of a real functions defined on some set E and set

$$s_n(x) = \sum_{k=1}^n f_k(x), \quad x \in E, n \in \mathbb{N}$$

- The series $\sum_{k=1}^{\infty} f_k$ is said to **converge pointwise** on E if and only if the sequence $s_n(x)$ converges pointwise on E as $n \rightarrow \infty$
- The series $\sum_{k=1}^{\infty} f_k$ is said to **converge uniformly** on E if and only if the sequence $s_n(x)$ converges uniformly on E as $n \rightarrow \infty$
- The series $\sum_{k=1}^{\infty} f_k$ is said to **converge absolutely** (pointwise) on E if and only if $\sum_{k=1}^{\infty} |f_k(x)|$ converges for each $x \in E$

Theorem 2.5: Weierstrass M-test

Let E be a nonempty subset of \mathbb{R} , let $f_k : E \rightarrow \mathbb{R}$, $k \in \mathbb{N}$, and suppose that $M_k > 0$ satisfies $\sum_{k=1}^{\infty} M_k < \infty$. If $|f_k(x)| \leq M_k$ for $k \in \mathbb{N}$ and $x \in E$, then $f = \sum_{k=1}^{\infty} f_k$ converges absolutely and uniformly on E .

Definition 3.0: Power Series

Let (a_n) be a sequence of real numbers, and $c \in \mathbb{R}$. A **power series** is a series of the form

$$\sum_{n=0}^{\infty} a_n(x-c)^n$$

The numbers a_n are called the **coefficients** of the power series, and c is its **centre**. In many cases it suffices to set $c = 0$. Note that the series will always converge at the point $x = c$ as all terms beyond the first are 0.

Theorem 3.3: Differentiation of Power Series

Suppose the radius of convergence of a power series is R . Then the function

$$f(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$$

is infinitely differentiable on $|x-c| < R$, and for such x ,

$$f'(x) = \sum_{n=0}^{\infty} n a_n(x-c)^{n-1}$$

and the series converges absolutely, and also uniformly on $[c-r, c+r]$ for any $r < R$. Moreover,

$$a_n = \frac{f^{(n)}(c)}{n!}$$

Theorem 2.4: Results of Convergent Series

Let E be a nonempty subset of \mathbb{R} and let (f_k) be a sequence of real functions defined on E .

- Suppose that $x_0 \in E$ and that each f_k is continuous at $x_0 \in E$. If $f = \sum_{k=1}^{\infty} f_k$ converges uniformly on E , then f is continuous at $x_0 \in E$.
- Term-by-term integration: Suppose that $E = [a, b]$ and that each f_k is integrable on $[a, b]$. If $f = \sum_{k=1}^{\infty} f_k$ converges uniformly on $[a, b]$, then f is integrable on $[a, b]$ and

$$\int_a^b \sum_{k=1}^{\infty} f_k(x) dx = \sum_{k=1}^{\infty} \int_a^b f_k(x) dx$$

- Term-by-term differentiation: Suppose that E is a bounded, open interval and that each f_k is differentiable on E . If $\sum_{k=1}^{\infty} f_k(x_0)$ converges at some $x_0 \in E$, and $g = \sum_{k=1}^{\infty} f'_k(k)$ converges uniformly on E , then $f = \sum_{k=1}^{\infty} f_k$ converges uniformly on E , is differentiable on E , and

$$f'(x) = \left(\sum_{k=1}^{\infty} f_k(x) \right)' = \sum_{k=1}^{\infty} f'_k(x) = g(x)$$

for $x \in E$

Definition 3.1: Radius of Convergence

The **radius of convergence** R of the power series

$$\sum_{n=0}^{\infty} a_n(x-c)^n \quad (*)$$

is defined by

$$R = \sup\{r \geq 0 : (a_n r^n) \text{ is bounded}\}$$

unless $(a_n r^n)$ is bounded for all $r \geq 0$, in which case we say that $R = \infty$

Thm 3.1: Suppose the radius of convergence R of $*$ satisfies $0 < R < \infty$. If $|x-c| < R$, the power series $*$ converges absolutely. If $|x-c| > R$, the power series $*$ diverges

Theorem 3.2: Continuity of Power Series

Assume that $R > 0$. Suppose that $0 < r < R$. Then a power series converges uniformly and absolutely on $|x-c| \leq r$ to a continuous function f . Hence

$$f(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$$

defines a continuous function $f : (c-R, c+R) \rightarrow \mathbb{R}$

Lemma 3.1: The two power series

$$\sum_{n=1}^{\infty} a_n(x-c)^n \text{ and } \sum_{n=1}^{\infty} n a_n(x-c)^{n-1}$$

have the same radius of convergence

3 Lebesgue Integration

Definition 4.0: Characteristic Function

Let E be a subset of \mathbb{R} . We define its **characteristic function** $\chi_E : \mathbb{R} \rightarrow \mathbb{R}$ by $\chi_E(x) = 1$ if $x \in E$ and $\chi_E(x) = 0$ if $x \notin E$. In other words,

$$\chi_E = \begin{cases} 1 & x \in E \\ 0 & x \notin E \end{cases}$$

In other words, this is a function that is 1 at all points of a bounded interval, and 0 elsewhere

Let I be a bounded interval with endpoints a, b and $a \leq b$. We call the number $b - a$ the **length of the interval** I and we denote it by $\lambda(I)$. This might also be referred to as $|I|$. That is,

$$\lambda((a, b)) = \lambda([a, b]) = \lambda((a, b]) = \lambda([a, b)) = b - a$$

From our definition of a characteristic function and the length of an interval, we have that the area of the characteristic function is a rectangle with width $\lambda(I)$ and height 1, therefore

$$\int \chi_I = 1 \cdot \lambda(I) = \lambda(I)$$

Definition 4.1: Step function

We say that $\phi : \mathbb{R} \rightarrow \mathbb{R}$ is a **step function** if there exist real numbers $x_0 < x_1 < x_2 < \dots < x_n$ (for some $n \in \mathbb{N}$) such that

1. $\phi(x) = 0$ for $x < x_0$ and $x > x_n$
2. ϕ is constant on (x_{j-1}, x_j) for $1 \leq j \leq n$

We shall use the phrase " ϕ is a step function with respect to $\{x_0, x_1, \dots, x_n\}$ " to describe this situation

Properties of Step Functions

1. The class of step functions is a vector space - i.e. if ϕ and ψ are step functions and α and β are real numbers, then $\alpha\phi + \beta\psi$ is a step function, and that if ϕ and ψ are step functions, then $\max\{\phi, \psi\}$, $\min\{\phi, \psi\}$, $|\phi|$ and $\phi\psi$ are also step functions
2. If ϕ and ψ are step functions, then $\phi + \psi$ is a step function
3. ϕ is a step function if and only if it is of the form

$$\phi = \sum_{j=1}^n c_j \chi_{J_j}$$

for some n , c_j , and bounded intervals J_j

Def 4.2: Integral of a Step Function

If ϕ is a step function with respect to $\{x_0, x_1, \dots, x_n\}$ which takes the value c_j on (x_{j-1}, x_j) , then

$$\int \phi := \sum_{j=1}^n c_j (x_j - x_{j-1})$$

Therefore, using the characteristic definition of a step function, the integral is

$$\int \phi = \int \sum_{j=1}^n c_j \chi_{J_j} = \sum_{j=1}^n c_j \int \chi_{IJ_j} = \sum_{j=1}^n c_j \lambda(J_j)$$

Definition 4.3: Lebesgue Integrals

A function $f : I \rightarrow \mathbb{R}$ is said to be **integrable** or more precisely **Lebesgue integrable** on an interval I if there exist numbers c_j and bounded intervals $J_j \subset I$, $j = 1, 2, 3, \dots$ such that

$$\sum_{j=1}^{\infty} |c_j| \lambda(J_j) < \infty$$

and the equality

$$f(x) = \sum_{j=1}^{\infty} c_j \chi_{J_j}(x)$$

holds for all $x \in I$ at which

$$\sum_{j=1}^{\infty} |c_j| \chi_{J_j}(x) < \infty$$

We denote by $\int_I f$ the number

$$\int_I f = \sum_{j=1}^{\infty} c_j \lambda(J_j)$$

and call it the integral of f over the interval I . If the function f is not integrable on the interval I then we say that the integral of f on I does not exist. Hence if we say that the integral of f on I exists it just means that f is (Lebesgue) integrable on I .

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus

quis tortor vitae risus porta vehicula. Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur. Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio. Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis sem-

per, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui. Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit

ipsum dolor sed augue. Nulla nec lacus. Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi. Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien. Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit. Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor. Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.