

# CS7301: Advanced Topics in Optimization for Machine Learning

## Lecture 5.1: Second Order Methods Cont., Barzelia Borwein GD and Conjugate Gradient Method

Rishabh Iyer

Department of Computer Science  
University of Texas, Dallas

<https://github.com/rishabhk108/AdvancedOptML>

February 26, 2021



# Summary of Algorithms so Far

## First Order Methods (Unconstrained)

- Gradient Descent
- Proximal Gradient Descent (if non-smooth function is simple – i.e. proximal operator of that function can be computed efficiently.
- Accelerated Gradient versions in the unconstrained case, projected case and proximal case.

## First Order Methods (Constrained)

- Projected Gradient Descent (whenever the Projection step is easy)
- Conditional Gradient Descent



## Second Order Methods

- Newtons Method
- General Quasi Newton Template
- Symmetric Rank One Update
- BFGS Update
- DFP Update



# Outline of Lecture 5.1

- LBFGS (Limited Memory BFGS)
- Conjugate Gradient Descent
- Barzelia Borowein Gradient Descent
- Coordinate Descent Algorithms



# Recall: BFGS Update

- BFGS update is a rank two update:

$$B_k = B_{k-1} + a u u^T + b v v^T$$

- Recall BFGS update:

$$B_k = B_{k-1} - \frac{B_{k-1} s_{k-1} s_{k-1}^T B_{k-1}}{s_{k-1}^T B_{k-1} s_{k-1}} + \frac{y_{k-1} y_{k-1}^T}{y_{k-1}^T s_{k-1}}$$

- BFGS Inverse Update:

$$B_{k+1}^{-1} = \left( I - \frac{s_{k-1} y_{k-1}^T}{s_{k-1}^T y_{k-1}} \right) B_k^{-1} \left( I - \frac{y_{k-1} s_{k-1}^T}{y_{k-1}^T s_{k-1}} \right) + \frac{s_{k-1} s_{k-1}^T}{y_{k-1}^T s_{k-1}}$$



# Limited Memory BFGS

- The memory and run-time cost of  $O(d^2)$  is itself not scalable for large number of features!



# Limited Memory BFGS

- The memory and run-time cost of  $O(d^2)$  is itself not scalable for large number of features!
- We need the improved (e.g. super-linear) convergence of BFGS/Newton but with linear or sublinear per iteration complexity!



# Limited Memory BFGS

- The memory and run-time cost of  $O(d^2)$  is itself not scalable for large number of features!
- We need the improved (e.g. super-linear) convergence of BFGS/Newton but with linear or sublinear per iteration complexity!
- Idea of LBFGS: Do not store  $B_k^{-1}$  explicitly.





# Limited Memory BFGS

- The memory and run-time cost of  $O(d^2)$  is itself not scalable for large number of features!
- We need the improved (e.g. super-linear) convergence of BFGS/Newton but with linear or sublinear per iteration complexity!
- Idea of LBFGS: Do not store  $B_k^{-1}$  explicitly.
- Instead store upto  $l$  (e.g.  $l = 30$ ) values of  $s_k$  and  $y_k$  and compute  $B_k^{-1}$  using the recursive BFGS equation and assuming  $B_{k-l} = I$ :

$$B_{k+1}^{-1} = \left(I - \frac{s_{k-1}y_{k-1}^T}{s_{k-1}^T y_{k-1}}\right) B_k^{-1} \left(I - \frac{y_{k-1}s_{k-1}^T}{y_{k-1}^T s_{k-1}}\right) + \frac{s_{k-1}s_{k-1}^T}{y_{k-1}^T s_{k-1}}$$



# Limited Memory BFGS

- The memory and run-time cost of  $O(d^2)$  is itself not scalable for large number of features!
- We need the improved (e.g. super-linear) convergence of BFGS/Newton but with linear or sublinear per iteration complexity!
- Idea of LBFGS: Do not store  $B_k^{-1}$  explicitly.
- Instead store upto  $l$  (e.g.  $l = 30$ ) values of  $s_k$  and  $y_k$  and compute  $B_k^{-1}$  using the recursive BFGS equation and assuming  $B_{k-l} = I$ :

$$B_{k+1}^{-1} = \left(I - \frac{s_{k-1}y_{k-1}^T}{s_{k-1}^T y_{k-1}}\right) B_k^{-1} \left(I - \frac{y_{k-1}s_{k-1}^T}{y_{k-1}^T s_{k-1}}\right) + \frac{s_{k-1}s_{k-1}^T}{y_{k-1}^T s_{k-1}}$$

- This makes the cost per iteration of  $O(dl)$  and storage also as  $O(dl)$  instead of  $O(d^2)$ .



# Summary so Far...

- Newton's method: Faster than any other algorithm seen so far.  
However costly  $O(d^3)$  to Invert Hessian



# Summary so Far...

- Newton's method: Faster than any other algorithm seen so far. However costly  $O(d^3)$  to Invert Hessian
- SR1, BFGS and DFP are simpler updates (using the Secant condition) which are Newton like (hence Quasi-Newton) but do this in  $O(d^2)$  complexity!



# Summary so Far...

- Newton's method: Faster than any other algorithm seen so far. However costly  $O(d^3)$  to Invert Hessian
- SR1, BFGS and DFP are simpler updates (using the Secant condition) which are Newton like (hence Quasi-Newton) but do this in  $O(d^2)$  complexity!
- LBFGS is an approximation to BFGS to make the algorithm linear!



# Summary so Far...

- Newton's method: Faster than any other algorithm seen so far. However costly  $O(d^3)$  to Invert Hessian
- SR1, BFGS and DFP are simpler updates (using the Secant condition) which are Newton like (hence Quasi-Newton) but do this in  $O(d^2)$  complexity!
- LBFGS is an approximation to BFGS to make the algorithm linear!
- Next: Can we achieve Newton like behaviour with Gradient Descent like algorithms?



# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization



# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization
- Conjugate gradient solves the problem  $Ax = b$  (which can be seen as minimizing  $\frac{1}{2}x^T Ax - b^T x$ )





# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization
- Conjugate gradient solves the problem  $Ax = b$  (which can be seen as minimizing  $\frac{1}{2}x^T Ax - b^T x$ )
- Non-Linear Conjugate Gradient is an extension to general convex functions!



# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization
- Conjugate gradient solves the problem  $Ax = b$  (which can be seen as minimizing  $\frac{1}{2}x^T Ax - b^T x$ )
- Non-Linear Conjugate Gradient is an extension to general convex functions!
- Several variants of conjugate gradient descent. Basic idea: Instead of using  $-\nabla f(x_k)$  as the update direction (as in gradient descent), do the following:



# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization
- Conjugate gradient solves the problem  $Ax = b$  (which can be seen as minimizing  $\frac{1}{2}x^T Ax - b^T x$ )
- Non-Linear Conjugate Gradient is an extension to general convex functions!
- Several variants of conjugate gradient descent. Basic idea: Instead of using  $-\nabla f(x_k)$  as the update direction (as in gradient descent), do the following:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$



# Conjugate Gradient Descent

- Linear Conjugate Gradient is an important Algorithm for quadratic optimization
- Conjugate gradient solves the problem  $Ax = b$  (which can be seen as minimizing  $\frac{1}{2}x^T Ax - b^T x$ )
- Non-Linear Conjugate Gradient is an extension to general convex functions!
- Several variants of conjugate gradient descent. Basic idea: Instead of using  $-\nabla f(x_k)$  as the update direction (as in gradient descent), do the following:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$
  - Set  $d_{k+1} = -g_{k+1} + \beta_k d_k$  where  $\beta_k$  is appropriately set!




# Conjugate Gradient Descent for Quadratic Functions

A comparison of

- \* gradient descent with optimal step size (in green) and
- \* conjugate vector (in red)

for minimizing a quadratic function.

Conjugate gradient converges in at most  $n$  steps (here  $n=2$ ).



# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:



# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$



# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:

- Set  $x_{k+1} = x_k + \alpha_k d_k$
- Set  $d_{k+1} = -g_{k+1} + \beta_k d_k$  where  $\beta_k$  is appropriately set!





# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$
  - Set  $d_{k+1} = -g_{k+1} + \beta_k d_k$  where  $\beta_k$  is appropriately set!
- Fletcher–Reeves CG algorithm:

$$\beta_k = \frac{\|\nabla f(x_k)\|^2}{\|\nabla f(x_{k-1})\|^2}$$



# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$
  - Set  $d_{k+1} = -g_{k+1} + \beta_k d_k$  where  $\beta_k$  is appropriately set!
- Fletcher–Reeves CG algorithm:

$$\beta_k = \frac{\|\nabla f(x_k)\|^2}{\|\nabla f(x_{k-1})\|^2}$$

- Polak–Ribière:

$$\beta_k = \frac{\nabla f(x_k)^T (\nabla f(x_k) - \nabla f(x_{k-1}))}{\|\nabla f(x_{k-1})\|^2}$$



# Variants of Conjugate gradient Descent

- Conjugate Gradient Framework:
  - Set  $x_{k+1} = x_k + \alpha_k d_k$
  - Set  $d_{k+1} = -g_{k+1} + \beta_k d_k$  where  $\beta_k$  is appropriately set!
- Fletcher–Reeves CG algorithm:

$$\beta_k = \frac{\|\nabla f(x_k)\|^2}{\|\nabla f(x_{k-1})\|^2}$$

- Polak–Ribière:

$$\beta_k = \frac{\nabla f(x_k)^T (\nabla f(x_k) - \nabla f(x_{k-1}))}{\|\nabla f(x_{k-1})\|^2}$$

- Hestenes–Stiefel:

$$\beta_k = \frac{\nabla f(x_k)^T (\nabla f(x_k) - \nabla f(x_{k-1}))}{d_k^T (\nabla f(x_k) - \nabla f(x_{k-1}))}$$



# Conjugate Gradient and Quasi Newton

- Hestenes–Stiefel conjugate gradient update is equivalent to LBFGS with  $\ell = 1$ !



# Conjugate Gradient and Quasi Newton

- Hestenes–Stiefel conjugate gradient update is equivalent to LBFGS with  $l = 1$ !
- Moreover, similar to Newton and BFGS, one can also show superlinear local convergence and global convergence results for CG!



# Conjugate Gradient and Quasi Newton

- Hestenes–Stiefel conjugate gradient update is equivalent to LBFGS with  $l = 1$ !
- Moreover, similar to Newton and BFGS, one can also show superlinear local convergence and global convergence results for CG!
- Advantage of CG is that the complexity is exactly identical to Gradient Descent and yet it has some of the properties of Quasi Newton algorithms!



# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$



# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$

- Compute  $x_{t+1}$  by optimizing the quadratic function above using the (linear) conjugate gradient!





# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$

- Compute  $x_{t+1}$  by optimizing the quadratic function above using the (linear) conjugate gradient!
- Repeat these steps till converged



# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$

- Compute  $x_{t+1}$  by optimizing the quadratic function above using the (linear) conjugate gradient!
- Repeat these steps till converged
- Moreover, we can do this Hessian Free(!) because we just need to compute  $\nabla^2 f(x_t)v$ , i.e. we don't need to compute  $\nabla^2 f$  for this!



# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$

- Compute  $x_{t+1}$  by optimizing the quadratic function above using the (linear) conjugate gradient!
- Repeat these steps till converged
- Moreover, we can do this Hessian Free(!) because we just need to compute  $\nabla^2 f(x_t)v$ , i.e. we don't need to compute  $\nabla^2 f$  for this!
- Note that  $(\nabla^2 f(x)v)_i = \sum_{j=1}^d \frac{\partial^2 f}{\partial x_i \partial x_j}(x) \cdot v_j = \nabla \frac{\partial f}{\partial x_i}(x) \cdot v$



# Hessian Free Optimization (James Martin, ICML 2010)

- Recall the quadratic expansion:

$$f(x) \approx f(x_t) + \nabla f(x_t)^T (x - x_t) + \frac{1}{2} (x - x_t)^T \nabla^2 f(x_t) (x - x_t)$$

- Compute  $x_{t+1}$  by optimizing the quadratic function above using the (linear) conjugate gradient!
- Repeat these steps till converged
- Moreover, we can do this Hessian Free(!) because we just need to compute  $\nabla^2 f(x_t)v$ , i.e. we don't need to compute  $\nabla^2 f$  for this!
- Note that  $(\nabla^2 f(x)v)_i = \sum_{j=1}^d \frac{\partial^2 f}{\partial x_i \partial x_j}(x) \cdot v_j = \nabla \frac{\partial f}{\partial x_i}(x) \cdot v$
- This is the directional derivative of  $\frac{\partial f}{\partial x_i}$  in the direction of  $v$ ! and can be computed numerically!



# Barzilai Borwein (BB) Gradient Descent

- The BB Method was introduced in an 8-page paper in 1988 (J. Barzilai and J. Borwein. Two point step size gradient method, IMA J. Numerical Analysis, 1988)



# Barzilai Borwein (BB) Gradient Descent

- The BB Method was introduced in an 8-page paper in 1988 (J. Barzilai and J. Borwein. Two point step size gradient method, IMA J. Numerical Analysis, 1988)
- Its a Gradient method with special step sizes.



# Barzelai Borwein (BB) Gradient Descent

- The BB Method was introduced in an 8-page paper in 1988 (J. Barzilao and J. Borwein. Two point step size gradient method, IMA J. Numerical Analysis, 1988)
- Its a Gradient method with special step sizes.
- Motivated by Newtons method but does not compute the Hessian!



# Barzilai Borwein (BB) Gradient Descent

- The BB Method was introduced in an 8-page paper in 1988 (J. Barzilai and J. Borwein. Two point step size gradient method, IMA J. Numerical Analysis, 1988)
- Its a Gradient method with special step sizes.
- Motivated by Newtons method but does not compute the Hessian!
- At no extra cost over the standard gradient descent, this method can significantly outperform GD!





# Barzilai Borwein (BB) Gradient Descent

- The BB Method was introduced in an 8-page paper in 1988 (J. Barzilai and J. Borwein. Two point step size gradient method, IMA J. Numerical Analysis, 1988)
- Its a Gradient method with special step sizes.
- Motivated by Newtons method but does not compute the Hessian!
- At no extra cost over the standard gradient descent, this method can significantly outperform GD!
- Typically used with non-monotone line search as convergence safeguard against non-quadratic problems (we won't cover this in class).



# Barzilai Borwein (BB) Gradient Descent

- Recall Gradient Descent:

$$x_{t+1} = x_t - \alpha_t \nabla f(x_t)$$

. Pros: Simple! Cons: No second order information at all!



# Barzilai Borwein (BB) Gradient Descent

- Recall Gradient Descent:

$$x_{t+1} = x_t - \alpha_t \nabla f(x_t)$$

. Pros: Simple! Cons: No second order information at all!

- Newton's method (similarly BFGS):

$$x_{t+1} = x_t - \alpha_t H_t^{-1} \nabla f(x_t)$$

. Pros: Very Fast Convergence! Cons: Creating or updating  $H_k^{-1}$  superlinear in cost!



# Barzilai Borwein (BB) Gradient Descent

- Recall Gradient Descent:

$$x_{t+1} = x_t - \alpha_t \nabla f(x_t)$$

. Pros: Simple! Cons: No second order information at all!

- Newton's method (similarly BFGS):

$$x_{t+1} = x_t - \alpha_t H_t^{-1} \nabla f(x_t)$$

. Pros: Very Fast Convergence! Cons: Creating or updating  $H_k^{-1}$  superlinear in cost!

- BB Gradient Descent: Choose  $\alpha_t \nabla f(x_t)$  such that it approximates  $H_t^{-1} \nabla f(x_t)$



# Barzilai Borwein (BB) Gradient Descent

- Recall Secant Condition:

$$H_k s_{k-1} = y_{k-1}$$

where  $s_{k-1} = x_k - x_{k-1}$  and  $y_{k-1} = \nabla f(x_k) - \nabla f(x_{k-1})$



# Barzilai Borwein (BB) Gradient Descent

- Recall Secant Condition:

$$H_k s_{k-1} = y_{k-1}$$

where  $s_{k-1} = x_k - x_{k-1}$  and  $y_{k-1} = \nabla f(x_k) - \nabla f(x_{k-1})$

- Another way to see the secant condition is the fact that this is the expression of the Hessian for quadratic functions!



# Barzilai Borwein (BB) Gradient Descent

- Recall Secant Condition:

$$H_k s_{k-1} = y_{k-1}$$

where  $s_{k-1} = x_k - x_{k-1}$  and  $y_{k-1} = \nabla f(x_k) - \nabla f(x_{k-1})$

- Another way to see the secant condition is the fact that this is the expression of the Hessian for quadratic functions!
- Now, the BB update is very simple: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1} I) s_{k-1} \approx y_{k-1}$$



# Barzilai Borwein (BB) Gradient Descent

- Recall Secant Condition:

$$H_k s_{k-1} = y_{k-1}$$

where  $s_{k-1} = x_k - x_{k-1}$  and  $y_{k-1} = \nabla f(x_k) - \nabla f(x_{k-1})$

- Another way to see the secant condition is the fact that this is the expression of the Hessian for quadratic functions!
- Now, the BB update is very simple: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1} I) s_{k-1} \approx y_{k-1}$$

- Solve this using least squares!





# Barzilai Borwein (BB) Gradient Descent

- Goal: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1} I) s_{k-1} \approx y_{k-1}$$



# Barzilai Borwein (BB) Gradient Descent

- Goal: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1} I) s_{k-1} \approx y_{k-1}$$

- BB Method:



# Barzilai Borwein (BB) Gradient Descent

- Goal: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1}I)s_{k-1} \approx y_{k-1}$$

- BB Method:
  - Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$



# Barzilai Borwein (BB) Gradient Descent

- Goal: Select  $\alpha_k$  such that:

$$(\alpha_k^{-1}I)s_{k-1} \approx y_{k-1}$$

- BB Method:

- Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$

- Approach 2: Least squares in  $\alpha$  directly:

$$\alpha_k = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1} - y_{k-1}\alpha\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T y_{k-1}}{y_{k-1}^T y_{k-1}}$$



# BB Gradient Descent in Practice

- BB Method:



# BB Gradient Descent in Practice

- BB Method:
  - Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$



# BB Gradient Descent in Practice

- BB Method:

- Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$

- Approach 2: Least squares in  $\alpha$  directly:

$$\alpha_k = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1} - y_{k-1}\alpha\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T y_{k-1}}{y_{k-1}^T y_{k-1}}$$



# BB Gradient Descent in Practice

- BB Method:

- Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$

- Approach 2: Least squares in  $\alpha$  directly:

$$\alpha_k = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1} - y_{k-1}\alpha\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T y_{k-1}}{y_{k-1}^T y_{k-1}}$$

- In practice, use either of these two or pick the better among the two or even alternate by fixing one for a few steps.





# BB Gradient Descent in Practice

- BB Method:
  - Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$

- Approach 2: Least squares in  $\alpha$  directly:

$$\alpha_k = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1} - y_{k-1}\alpha\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T y_{k-1}}{y_{k-1}^T y_{k-1}}$$

- In practice, use either of these two or pick the better among the two or even alternate by fixing one for a few steps.
- However,  $f(x_k)$  and  $\nabla f(x_k)$  are not monotonic!



# BB Gradient Descent in Practice

- BB Method:

- Approach 1: Let  $\beta = \alpha^{-1}$ . Then:

$$\alpha_k^{-1} = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1}\beta - y_{k-1}\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T y_{k-1}}$$

- Approach 2: Least squares in  $\alpha$  directly:

$$\alpha_k = \operatorname{argmin}_{\alpha} \frac{1}{2} \|s_{k-1} - y_{k-1}\alpha\|^2 \Rightarrow \alpha_k = \frac{s_{k-1}^T y_{k-1}}{y_{k-1}^T y_{k-1}}$$

- In practice, use either of these two or pick the better among the two or even alternate by fixing one for a few steps.
- However,  $f(x_k)$  and  $\nabla f(x_k)$  are not monotonic!
- Non Monotonic line search required for convergence analysis



# Summary So Far...

- Gradient Descent is easy to implement, simple with low per iteration complexity. However, at best linear convergence! For most standard cases, sublinear convergence.



# Summary So Far...

- Gradient Descent is easy to implement, simple with low per iteration complexity. However, at best linear convergence! For most standard cases, sublinear convergence.
- Accelerated Gradient Descent is an improvement. Slightly more involved but similar per iteration complexity. Slightly faster but still similar to GD!



# Summary So Far...

- Gradient Descent is easy to implement, simple with low per iteration complexity. However, at best linear convergence! For most standard cases, sublinear convergence.
- Accelerated Gradient Descent is an improvement. Slightly more involved but similar per iteration complexity. Slightly faster but still similar to GD!
- Newton Method is superlinear convergence! However very slow per iterations:  $O(d^3)$  complexity.



# Summary So Far...

- Gradient Descent is easy to implement, simple with low per iteration complexity. However, at best linear convergence! For most standard cases, sublinear convergence.
- Accelerated Gradient Descent is an improvement. Slightly more involved but similar per iteration complexity. Slightly faster but still similar to GD!
- Newton Method is superlinear convergence! However very slow per iterations:  $O(d^3)$  complexity.
- BFGS improves it to  $O(d^2)$  which is further improved to  $O(d)$  by LBFGS!



# Summary So Far...

- Gradient Descent is easy to implement, simple with low per iteration complexity. However, at best linear convergence! For most standard cases, sublinear convergence.
- Accelerated Gradient Descent is an improvement. Slightly more involved but similar per iteration complexity. Slightly faster but still similar to GD!
- Newton Method is superlinear convergence! However very slow per iterations:  $O(d^3)$  complexity.
- BFGS improves it to  $O(d^2)$  which is further improved to  $O(d)$  by LBFGS!
- CG and BB are variants of Gradient Descent but try to mimic Newton's methods and in practice are much faster than simple GD while maintaining the simplicity!

