

Math Prep Course Day Four

Ashhad Shahzad

September 2025

1 Function Composition

Problem 1.1. Find The Domain of

$$f(x) = \frac{7 - \sqrt{x^2 - 9}}{\sqrt{25 - x^2}}$$

Solution. We can decompose $f(x)$ into its numerator and denominator functions and take the intersection of their domain. That can be done as follows

$$\begin{aligned}x^2 - 9 &\geq 0 \\x^2 &\geq 9 \\x &\geq |3|\end{aligned}$$

$$\begin{aligned}25 - x^2 &\geq 0 \\x^2 &\leq 25 \\x &\leq |5|\end{aligned}$$

And so the intersection of the domains is simply $(-5, -3] \cup [3, 5)$.

Problem 1.2. Let $f(x) = \sqrt{x}$, $g(x) = \frac{4}{5-x}$, $h(x) = x^2$. Find

$$(h(h \circ g \circ f - f))(4)$$

And it's domain

Solution.

$$\begin{aligned}
 h(h \circ g \circ f - f) &= \left(\left(\frac{4}{5 - \sqrt{x}} \right)^2 - \sqrt{x} \right)^2 \\
 (h(h \circ g \circ f - f))(4) &= \left(\left(\frac{4}{5 - \sqrt{4}} \right)^2 - \sqrt{4} \right)^2 \\
 &= \left(\left(\frac{4}{5 - 2} \right)^2 - 2 \right)^2 \\
 &= \left(\left(\frac{4}{3} \right)^2 - 2 \right)^2 \\
 &= \frac{4}{81}
 \end{aligned}$$

The domain is clearly $[0, \infty)/\{25\}$

2 Polynomials

I've solved some of these using long division instead of the remainder theorem for my own sake. The remainder theorem is however far more efficient.

Problem 2.1. Given that $(x - 3)$ is a factor of

$$f(x) = x^3 - 2x^2 + kx + 6$$

Show that $k = -5$.

Solution.

$$\begin{array}{r|l}
 & x - 3 \\
 \hline
 x^2 & x^3 - 2x^2 + kx + 6 \\
 & -x^3 + 3x^2 \\
 \hline
 x & x^2 + kx + 6 \\
 & -x^2 + 3x \\
 \hline
 k + 3 & (k + 3)x + 6 \\
 & -(k + 3)x + 3k + 39 \\
 \hline
 & 3k + 15
 \end{array}$$

Since $3k + 15 = 0$, we obtain $k = -5$

Problem 2.2. Evaluate

$$(x^5 - 4x^4 + 2x^3 - 3x^2 + 4x + 1) \div (x^2 + x + 1)$$

Solution.

$$\begin{array}{r|l}
 & (x^2 + x + 1) \\
 \hline
 x^3 & x^5 - 4x^4 + 2x^3 - 3x^2 + 4x + 1 \\
 & -x^5 - x^4 - x^3 \\
 \hline
 -5x^2 & -5x^4 + x^3 - 3x^3 + 4x + 1 \\
 & 5x^4 + 5x^3 + 5x^2 \\
 \hline
 6x & 6x^3 + 2x^2 + 4x + 1 \\
 & -6x^3 - 6x^2 - 6x \\
 \hline
 -4 & -4x^2 - 2x + 1 \\
 & 4x^2 + 4x + 4 \\
 \hline
 & 2x + 5
 \end{array}$$

Problem 2.3. Find the remainder

$$(4x^3 - 2x^2 + x + 1) \div (x - 1)$$

Solution.

$$\begin{array}{r|l}
 & x - 1 \\
 \hline
 4x^2 & 4x^3 - 2x^2 + x + 1 \\
 & -4x^3 + 4x^2 \\
 \hline
 2x & 2x^2 + x + 1 \\
 & -2x^2 + 2x \\
 \hline
 3 & 3x + 1 \\
 & -3x + 3 \\
 \hline
 & 4
 \end{array}$$

Problem 2.4.

$$f(x) = ax^3 - 7x^2 + 1$$

Has the factor $(x - 1)$. Find a .

Solution.

$$\begin{array}{r|l}
 & x - 1 \\
 \hline
 ax^2 & ax^3 - 7x^2 + 1 \\
 & -ax^3 + ax^2 \\
 \hline
 (a - 7)x & (a - 7)x^2 + 1 \\
 & (7 - a)x^2 + (a - 7)x \\
 \hline
 (7 - a) & (a - 7)x + 1 \\
 & (7 - a)x + (a - 7) \\
 \hline
 & a - 6
 \end{array}$$

Since the remainder must be zero, we obtain $a = 6$

Problem 2.5.

$$5x^6 - 3x^5 - x^4 + 1 = (x - 1)(x - 2) - x^2 + 3x - 1$$

Solution.

$$\begin{aligned}5x^6 - 3x^5 - x^4 + 1 &= x^2 - 3x + 2 - x^2 + 3x - 1 \\5x^6 - 3x^5 - x^4 &= 0 \\x^4(5x^2 - 3x - 1) &= 0\end{aligned}$$

Solving for the quadratic, we obtain the solutions

$$x = \left\{ 0, \frac{3 \pm \sqrt{29}}{10} \right\}$$

3 Logarithmic and Exponential Functions

Problem 3.1.

$$\log_2 x = \log_2 x^2 - 4$$

Solution.

$$\begin{aligned}\log_2 x &= 2 \log_2 x - 4 \\\log_2 x &= 4 \\x &= 2^4 \\x &= 16\end{aligned}$$

Problem 3.2.

$$2^x = 64$$

Solution.

$$\begin{aligned}2^x &= 64 \\x &= \log_2(2^6) \\x &= 6\end{aligned}$$

Problem 3.3.

$$\ln(x-1) - \ln(x^2-1) = e$$

Solution.

$$\begin{aligned}\ln\left(\frac{x-1}{x^2-1}\right) &= e \\\ln(x+1) &= -e \\x+1 &= e^{-e} \\x &= e^{-e} - 1\end{aligned}$$

However, $e^{-e} - 1 < 1$, which makes the expression $\ln(x-1)$ invalid, therefore we have an extraneous solution and the equation is unsatisfiable in \mathbb{R}

Problem 3.4. Find the domain and range of $x = \ln(e^x)$.

Solution. Observe that e^x is defined over all of \mathbb{R} and has range $(0, \infty)$. Note that this is precisely the domain of the \ln function, which has the range $(-\infty, \infty)$. Therefore the domain and range of the function are both \mathbb{R} .

In fact its just the constant function x since its the composition of a function with its inverse lmao.

Problem 3.5.

$$\log_2 x + \log_x 2 + 1 = 0$$

Solution.

$$\begin{aligned} \log_2 x + \frac{\log_2 2}{\log_2 x} + 1 &= 0 & \because \log_a b &= \frac{\log_c b}{\log_c a} \\ \log_2 x + \frac{1}{\log_2 x} + 1 &= 0 \\ (\log_2 x)^2 + \log_2 x + 1 &= 0 \\ y^2 + y + 1 &= 0 & \because \text{let } y &= \log_2 x \end{aligned}$$

Since the solution is complex, I'm not going to evaluate it further.

Problem 3.6.

$$\log_5 x + \log_{10} 8 = 1$$

Solution.

$$\begin{aligned} \log_5 x &= 1 - \log_{10} 8 \\ x &= 5^{(1 - \log_{10} 8)} \end{aligned}$$

Problem 3.7.

$$(e^x)^2 + \ln e^{e^x} + e^{\ln_e 5} = 0$$

Solution.

$$\begin{aligned} (e^x)^2 + e^x + 5 &= 0 \\ y^2 + y + 5 &= 0 & \because y &= e^x \end{aligned}$$

Again, the solution is complex.

Problem 3.8.

$$x^2 = e^{\ln(\frac{1}{5}x) + \ln(5x)} + \log_{10} e^{10}$$

Solution.

$$\begin{aligned}x^2 &= \left(\frac{x}{5}\right)(5x) + 10 \log_{10} e \\0 &= 10 \log_{10} e\end{aligned}$$

Thus the equation has no solutions.