Critical Phenomenon in Axisymmetric Vacuum Spacetimes: Role of Twist

Phase 1 Presentation

Ashique T Nizar

Indian Institute of Science Education and Research, Trivandrum

Under the guidance of Dr. Soumen Basak and Dr.David Hilditch

November 27, 2025

GR into a time evolution problem

"Matter tells spacetime how to curve, and spacetime tells matter how to move."

— John A. Wheeler

Often in time evolution PDE problems, the relevent equations can be formulated of the form.

$$\partial_t u = f(u, \partial_{x^i} u)$$

where t is the time and x^i is the spatial coordinates

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi T_{\mu\nu}$$

BUT, One of the most important equation of GR, the Einstein's Field equations, which are system of non-linear coupled differential equations, does'nt have that form explicitly

3+1 Decomposition

- **Spatial Metric and extrinsic curvature:**

$$\gamma_{ab} = g_{ab} + n_a n_b, \qquad K_{ab} = -\gamma^c{}_a \nabla_c n_b.$$



Figure: Slicing into spacelike hypersurfaces (Hilditch 2024)

3+1 Decomposition

- **Spatial Metric and extrinsic curvature:**

$$\gamma_{ab} = g_{ab} + n_a n_b, \qquad K_{ab} = -\gamma^c{}_a \nabla_c n_b.$$

- **Constraints:**

$$C_{Hamiltonian} : R + K^2 - K_{ab}K^{ab} - 16\pi\rho = 0,$$

$$C^a_{Momentum} : D_b K^{ab} - D^a K - 8\pi j^a = 0.$$



Figure: Slicing into spacelike hypersurfaces (Hilditch 2024)

3+1 Decomposition

- **Spatial Metric and extrinsic curvature:**

$$\gamma_{ab} = g_{ab} + n_a n_b, \qquad K_{ab} = -\gamma^c{}_a \nabla_c n_b.$$

- **Constraints:**

$$C_{Hamiltonian} : R + K^2 - K_{ab}K^{ab} - 16\pi\rho = 0,$$

$$C^a_{Momentum} : D_b K^{ab} - D^a K - 8\pi j^a = 0.$$

- **Evolution:**

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij},$$

$$\partial_t K_{ij} = -D_i D_j \alpha + \alpha(R_{ij} - 2K_{ik}K^k{}_j + KK_{ij}) + 4\pi\alpha[\gamma_{ij}(s-\rho) - 2s_{ij}] + \mathcal{L}_\beta K_{ij}.$$



Figure: Slicing into spacelike hypersurfaces (Hilditch 2024)

3+1 Decomposition

- **Spatial Metric and extrinsic curvature:**

$$\gamma_{ab} = g_{ab} + n_a n_b, \qquad K_{ab} = -\gamma^c{}_a \nabla_c n_b.$$

- **Constraints:**

$$C_{Hamiltonian} : R + K^2 - K_{ab}K^{ab} - 16\pi\rho = 0,$$

$$C^a_{Momentum} : D_b K^{ab} - D^a K - 8\pi j^a = 0.$$

- **Evolution:**

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij},$$

$$\partial_t K_{ij} = -D_i D_j \alpha + \alpha(R_{ij} - 2K_{ik}K^k{}_j + KK_{ij}) + 4\pi\alpha[\gamma_{ij}(s-\rho) - 2s_{ij}] + \mathcal{L}_\beta K_{ij}.$$



Figure: Slicing into spacelike hypersurfaces (Hilditch 2024)

3+1 Decomposition

- **Spatial Metric and extrinsic curvature:**

$$\gamma_{ab} = g_{ab} + n_a n_b, \qquad K_{ab} = -\gamma^c{}_a \nabla_c n_b.$$

- **Constraints:**

$$C_{Hamiltonian} : R + K^2 - K_{ab}K^{ab} - 16\pi\rho = 0,$$

$$C^a_{Momentum} : D_b K^{ab} - D^a K - 8\pi j^a = 0.$$

- **Evolution:**

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij},$$

$$\partial_t K_{ij} = -D_i D_j \alpha + \alpha(R_{ij} - 2K_{ik}K^k{}_j + KK_{ij}) + 4\pi\alpha[\gamma_{ij}(s-\rho) - 2s_{ij}] + \mathcal{L}_\beta K_{ij}.$$



Figure: Slicing into spacelike hypersurfaces (Hilditch 2024)

If we know these two quantities (as well as matter variables) at the initial hypersurface, we will be able to evolve it in time satisfying Einstein's equations even using regular computational methods

Critical Phenomenon in Gravitational Collapse

If we have some constraint satisfying initial data[1] (γ_{ab}, K_{ab},Matter fields)

- Sufficiently "weak" Initial data :- Mass energy will get dispersed
- Sufficiently "strong" Initial data :- Collapses to a singularity

Choptuik in his 1992 paper,

- Took a spherically symmetric scalar field as matter source for T_{ab}
- He took several one-parameter families of initial data (by parameterizing the scalar field)
- In each family, he saw that at parameter p $>$ some "Critical" parameter p*, it collapses (supercritical data) and p$<$p* the scalar field disperses (subcritical data)

[1] Christodoulou 1990-2000

Choptuik's findings

But the groundbreaking thing he found was that, if we try to tune to this critical parameter p* and see the evolution, three remarkable features emerges

1. Power-law mass scaling

   The mass of the black hole that forms when $p > p^*$ has no lower bound and follows the scaling law

   $$M_{\mathrm{BH}} \propto (p - p^*)^{\gamma}.$$

   When $p < p^*$,the maximum of curvature invariants follow the scaling law

   $$I_{\max} \propto (-p + p^*)^{-2\gamma}.$$



Figure: Choptuik 1992

Choptuik's findings

But the groundbreaking thing he found was that, if we try to tune to this critical parameter p* and see the evolution, three remarkable features emerges

2. Universality of Critical exponent

   The critical exponent γ is universal across all families of initial data that share the same matter model and symmetry.



Figure: Universality of critical exponent(Choptuik 1992)

Choptuik's findings

But the groundbreaking thing he found was that, if we try to tune to this critical parameter p* and see the evolution, three remarkable features emerges

3. Self similiarity and universality of Critical solution

   Critical solutions from all the families of initial data show self similarity in spatiotemporal scales

   $Z^*(a, b) \approx Z^*(a - \Delta, b - \Delta)$ where Δ is called scale echoing constant (Same irrespective of initial data family)

Deviation from spherical symmetry

- After this people did a lot of simulations with various matter models in spherical symmetry and was able to find similiar critical behaviour
- As we deviate from spherical symmetry, we began to see a "loss" in critical phenomenon. In the similiar massless scalar field test done in axisymmetry, they found

ϵ^2	γ
0	$0.382 \pm 2\%$
1/3	$0.380 \pm 2\%$
1/2	$0.375 \pm 3\%$
2/3	$0.346 \pm 3\%$
$3/4(a)$	$0.313 \pm 4\%$
$3/4(b)$	$0.40 \pm 10\%$
$5/6(a)$	$0.28 \pm 10\%$
$5/6(b)$	$0.41 \pm 10\%$
AS	$0.383 \pm 2\%$

Here ϵ marks the inital-data's deviation from spherical symmetry. Here as we can see that eventhough everyone of the families are axisymmetric,the *universality* of the critical exponent is gone now

Figure: Deviations observed (Choptuik et al, 2003)

Introduction of angular momentum

But on the introduction of angular momentum into the initial data, we begin to recover perfect critical phenomenon as before



Figure: Choptuik massless scalar field with angular momentum (Choptuik et al 2004)

$\Delta = 0.42 \pm 4\%, \qquad \gamma = 0.11 \pm 10\%$

- But what about vacuum ?
- Earlier spherical symmetry prevents pure gravitational wave and its collapse but axisymmetry does not ![a]

[a] Christodoulou 1990-2000

Vacuum and angular momentum



Figure: No proper critical phenomenon (Isabel Suarez Fernandez et al 2022)

- Even in axisymmetric vacuum gravitational wave collapse, we see the same
- But unlike in the case of matter, its not possible to generate angular momentum from regular initial data

- Axisymmetry is characterised by the presence of a spacelike Killing vector field and twist measures its vorticity
- Angular momentum implies twist but twist does not necessarily imply the presence of angular momentum

Motivation



(a) Matter without Angular momentum (Choptuik et al, 2003)



(b) Twist free vacuum (Isabel Suarez Fernandez et al 2022)

Motivation

- Until now all the vacuum tests that were done was in twist free conditions and it showed the same bad critical phenomenon as in angular momentum-free matter
- We are going to see how twist affects the critical phenomenon in this scenario and if it has some improvement angular momentum provides to matter

Twisted Brill Initial Data construction

- Axisymmetric vacuum gravitational waves with $K_{ij} = 0$ initial data is known as Brill waves. (This choice makes the momentum constraint be satisfied trivially)
- The usual twist excluded Brill wave ansatz

$$\gamma_{ij} = \psi^4 \tilde{\gamma}_{ij}(\rho, z) = \psi^4 \begin{pmatrix} e^{2q(\rho,z)} & 0 & 0 \\ 0 & e^{2q(\rho,z)} & 0 \\ 0 & 0 & \rho^2 \end{pmatrix},$$

when twist is zero, $\gamma_{i\phi} = 0$. Hence for including twist we add two more degrees of freedom at $\gamma_{i\phi}$

- We choose the twist included Brill wave ansatz

$$\gamma_{ij} = \psi^4 \tilde{\gamma}_{ij}(\rho, z) = \psi^4 \begin{pmatrix} e^{2q(\rho,z)} + f^2(\rho, z) & f(\rho, z)h(\rho, z) & \rho f(\rho, z) \\ f(\rho, z)h(\rho, z) & e^{2q(\rho,z)} + h(\rho, z)^2 & \rho h(\rho, z) \\ \rho f(\rho, z) & \rho h(\rho, z) & \rho^2 \end{pmatrix}$$

Twisted Brill Initial Data construction

- Since the initial data have to be constraint satisfying we use Conformal-York decomposition method which converts the Hamiltonian contraint into an elliptic equation of the form

$$8\tilde{D}^2\psi = \left(-\frac{e^{-2q}(\rho(\frac{\partial f}{\partial z} - \frac{\partial h}{\partial \rho})) + h)^2}{2\rho^2} - 2\left(\frac{\partial^2 q}{\partial \rho^2} + \frac{\partial^2 q}{\partial z^2}\right)\right)\psi$$

- We choose the the following ansatz for the f and h

$$f(\rho, z) = \mathtt{tp1}\,\rho^2 z\, e^{-(\rho^2+z^2)} \quad h(\rho, z) = \mathtt{tp2}\,\rho^3 e^{-(\rho^2+z^2)}.$$

- As for the first family, we choose pure twisting (q=0) with `tp1`=`tp2` =p
- We solved the differential equation using the Psuedospectral method and the residuals are in the order of 10^{-30} and was checked for convergence with increasing resolution.
  Hence, the initial data is ready to evolve now ✔

Results

- Currently, we have tuned the critical parameter to about 3 decimal places for p positive and negative $p^* = 4.110(\pm 0.003)$



How do we tune?

- Eventhough we see the evidence of a power-law, we don't currently report the critical exponent at face value since we haven't tuned enough
- It could be that we simply would not have reached the critical regime yet

How do we tune and why is it hard?

To observe subcriticality, we look at the dispersal of the Kretschmann Curvature scalar $K = R_{abcd}R^{abcd}$

How do we tune and why is it hard?

To observe supercriticality, we probe for the formation of **Apparent horizons** (AHs) in the slice. Singularity theorems guarantee a singular spacetime if AHs are formed in the spacetime.

How do we tune and why is it hard?

Often while tuning, we might not be able to classify the spacetimes due to some reasons which we hypothesised as

1. Coordinate artifacts and gauge/slicing effects Solution:- In NR, they are mitigated by utilising the gauge freedom and changing the gauges to overcome the artifact. But this process requires trial and error, which slows down the tuning process.

How do we tune and why is it hard?

2 Limitations of Apparent Horizon finder

Apparent horizons are defined as outermost MOTS (Marginally Outer Trapped Surface) which are closed 2-surfaces that obeys the condition that the expansion of outgoing null geodesics emanating from it vanishes i.e

$$\Theta = D_i s^i + K_{ij} s^i s^j - K = 0$$

where D is the induced 3-metric, K_{ij} the extrinsic curvature, s^i the normal vector to the surface. We find surfaces that satisfies this condition.



Figure: Apparent horizons that we usually find close to criticality in our experiments

Horizon finding limitations

- We at first used the finder `AHLoc3D` which uses a spherical Strahlkorper / star-shaped parameterization where assume that the AH is star-shaped / a ray from a choosen center only touches the surface once. ($r = F(\theta)$ where r is the radius from a chosen center and θ the polar angle.)
- Since we were not able to go beyond 4.115 (towards 4.100), we rethought on this assumption and recently accounted for twist into a cartesian based AH finder which previously was made for non-twisting axisymmetry.



Figure: An example for a non-ray body horizon[2]

Cartesian based finder

Previous studies in vacuum axisymmetry didnt necessitate the usage of a different parameterization since it very much simplifies the finding problem, and they were able to tune high values of precision.

Thornburg 2006

Horizon finding limitations



- Using the new AH finder, we were able to find disjoint AHs around z axis values where the Kretschmann scalar is the maximum in the slice
- We can see the limitation that `AHLoc3D` brings into the picture, the parameterization breaks at some points (shown in red) if we were to use the star shaped assumption
- The downside is that the new AH finder relies on ODE shooting method, which required substantial **manual intervention** unlike the automated `AHLoc3D`

Forward

- Hence we expect that breakdown of the parameterisation was the real cause behind why we couldn't classify
- With this new AH finder we will be able to tune and reach the critical regime in a much faster rate from now on

Conclusion and future directions

- We currently set up the framework to create and evolve twisting brill wave initial data for arbitrary families and to characterise the critical phenomenon
- With the initial family, we have tuned upto 3 decimal places and solved some of the issues that we face while tuning towards the threshold. Further tuning is being performed with improvements we made to the tuning process.
- After sufficient tuning, we will begin our analysis of the critical threshold of twisting vacuum spacetimes. (Critical exponents, self similarity and echoing constants.)
- Construct new families (Eg.Impure twisting) and do the similiar exercise and compare the phenomenon between families and check for universality
- Devise strategies to make the new AH finding process automative

Acknowledgement

- My sincere gratitude to my guide, Dr. Soumen Basak, for his continuous guidance and support.
- Thank you to Dr.David Hilditch (CENTRA, Lisbon), Dr.Ananya Adhikari(CENTRA, Lisbon), Dr.Prayush Kumar (ICTS,Bangalore) for their helpful discussion regarding Numerical Relativity, setting up the code base and computational time
- Heartfelt appreciation to my family and my friends for their encouragement throughout this work.

Deviation from spherical symmetry

- Hence a perfect notion of universal critical exponent and echoing period is lost which now varies with family
- Center of collapse bifurcates into two
- Complicated notion of self similarity of critical solution



Figure: Complicated self similarity(Choptuik et al, 2003)

Twist and angular momentum

- Axisymmetry is defined by the existence of a killing vector field $\xi^a = (\partial_\phi)^a$ and twist (ω_a) measures its failure to be hypersurface orthogonal.
- From the idea of Komar angular momentum and its conservation, J $= J_{BH} + J_{Matter}$ ie Black holes formed from vacuum axisymmetric collapse are non rotating (no angular momentum)

- But when twist is 0, there won't be angular momentum as well

$$J = \frac{1}{32\pi^2\lambda} \oint_{\partial S_t} \epsilon_{abcd} \xi^c \omega^d dS^{ab}$$

Hence in vacuum there can be twist even without angular momentum (since J_{BH} and J_{Matter} vanishes)

Construction of one-parameter families of initial data

- A general axisymmetric, regular metric can be written as: (Rinne 2006)

$$\gamma_{ij} = \begin{bmatrix} H + \rho^2 J & \rho E & \rho^3 K \\ \rho E & C & \rho^2 G \\ \rho^3 K & \rho^2 G & \rho^2 (H - \rho^2 J) \end{bmatrix}$$

  Here the functions H, E, C, G, K, J are even in ρ (i.e. they depend on ρ^2) and depend on z.

Choptuik's findings

But the groundbreaking thing he found was that, if we try to tune to this critical parameter p* and see the evolution, three remarkable features emerges
When $p < p^*$,the maximum of curvature invariants follow the scaling law

$$I_{\max} \propto (-p + p^*)^{-2\gamma}.$$

1. Power-law mass scaling

   When $p < p^*$,the maximum of curvature invariants follow the scaling law

   $$I_{\max} \propto (-p + p^*)^{-2\gamma}.$$



FIG. 1. $\ln R_{\max}$ is plotted vs. $\ln(p*-p)$. The result is a line with slope -2γ and a periodic wiggle with period $\Delta/(2\gamma)$

Figure: Scaling of curvature scalar (maximum)

Construction

- Additionally to check whether the constraints are satisified (If we have solved the correct differential equation) we use the `bamps` framework which we use for time evolutions of our initial data



(a) x direction



(b) z direction

Hence the initial data is ready to evolve now ✔