Assignment-11.14.7

EE:1205-Signals and Systems Indian Institute of Technology, Hyderabad

Md Ayaan Ashraf EE23BTECH11041

Question

The motion of a particle executing simple harmonic motion is described by the displacement function, $x(t) = A \cos(\omega t + \phi)$. If the initial (t = 0) position of the particle is 1cm and its initial velocity is $\omega cm/s$, what are its amplitude and initial phase angle? The angular frequency of the particle is πs^{-1} . If instead of the cosine function, we choose the sine function to describe the SHM: $x = B \sin(\omega t + \alpha)$, what are the amplitude and initial phase of the particle with the above initial conditions.

Solution

Parameter	Description	Value
<i>x</i> (0)	Initial position of particle	1cm
f	Frequency of particle	$\frac{\omega}{2\pi}$
x'(0)	Initial velocity of particle	ω
φ	Initial Phase Angle	?
α	New Phase Angle	?
A	Initial Amplitude	?
В	New Amplitude	?

TABLE 1: Parameter Table 11.14.7

1) In Fig. ??, the equation is

$$x(t) = A\cos(2\pi f t + \phi) \tag{1}$$

Given:

$$x(0) = A\cos(\phi) = 1 \text{ cm}$$
 (2)

$$x'(0) = -A2\pi f \sin(\phi) = 2\pi f \text{ cm/s}$$
 (3)

Solving for ϕ and A:

$$\tan(\phi) = -1 \tag{4}$$

$$\implies \phi = -\frac{\pi}{4} \tag{5}$$

$$\implies A = \sqrt{2}cm$$
 (6)




Fig. 1: $x(t) = \sqrt{2}\cos(\pi t - \frac{\pi}{4})$

2) In Fig. ??, the equation is

$$x(t) = B\sin(2\pi f t + \alpha) \tag{7}$$

1

Given:

$$B\sin(\alpha) = 1$$
 cm (8)

$$B2\pi f \cos(\alpha) = 2\pi f \text{ cm/s} \tag{9}$$

Solving for α and B:

$$tan(\alpha) = 1 \tag{10}$$

$$\implies \alpha = \frac{\pi}{4} \tag{11}$$

$$\implies B = \sqrt{2}cm$$
 (12)




Fig. 2: $x(t) = \sqrt{2}\sin(\pi t + \frac{\pi}{4})$