

Crop rotation refers to the practice of growing different types of crops over the same area in different seasons. This is because the same type of crop planted repeatedly over the same area keeps draining the same nutrients from the land needed for that plant's growth. Crop rotation not only enriches soil fertility but also helps in pest control.

9. With reference to soil conservation, consider the following practices :

1. Crop rotation 2. Sand fences
3. Terracing 4. Windbreaks

Which of the above are considered appropriate methods for soil conservation in India?

- (a) 1, 2 and 3 (b) 2 and 4
- (c) 1, 3 and 4 (d) 1, 2, 3 and 4

I.A.S. (Pre) 2010

Ans. (c)

The appropriate methods for soil conservation are done by, crop rotation, terracing and windbreaks.

10. In India, the problem of soil erosion is associated with which of the following?

1. Terrace cultivation
2. Deforestation
3. Tropical climate

Select the correct answer using the code given below.

- (a) 1 and 2 only (b) 2 only
- (c) 1 and 3 only (d) 1, 2 and 3

I.A.S. (Pre) 2014

Ans. (b)

The destruction of soil cover is described as soil erosion. The soil forming processes and the erosional processes of running water and wind go on simultaneously. But generally, there is a balance between these two processes. Sometimes, such a balance is disturbed by natural or human factors, leading to a greater rate of removal of soil. Deforestation is one of the major causes of soil erosion. Plants keep soils bound in locks of roots and thus prevent erosion. Terrace farming is a method of soil conservation. The tropical climate is not responsible for soil erosion.

11. Soil erosion can be checked by

- (a) Excess grazing (b) Removal of plants
- (c) Afforestation (d) Increasing number of birds

40th B.P.S.C. (Pre) 1995

Ans. (c)

Soil conservation includes all the methods to check soil erosion and maintain soil fertility. Some of the effective ways to prevent soil erosion are – afforestation, contour tillage, banning shifting cultivation, etc.

12. Soil erosion can be prevented by –

- (a) Heavy rains (b) Deforestation
- (c) Afforestation (d) Overgrazing

U.P.R.O./A.R.O. (Pre) 2014

Ans. (c)

See the explanation of the above question.

13. Which of the following practices can help in water conservation in agriculture?

1. Reduced on zero tillage of the land
2. Applying gypsum before irrigating the field
3. Allowing crop residue to remain in the field

Select the correct answer using the code given below:

- (a) 1 and 2 only (b) 3 only
- (c) 1 and 3 only (d) 1, 2 and 3

I.A.S. (Pre) 2017

Ans. (d)

Reduced or zero tillage of the land in farming is a way of growing crops or pasture year to year without disturbing the soil through tillage; thus, it helps in soil moisture conservation. Gypsum helps in dissolution of the salt clay pan found in soils that obstructs the percolation of water. therefore, it increases the efficiency of irrigation. Allowing crop residue to remain in the field reduces evaporation of water from the soil. Thus all the statements are correct.

Natural Vegetation

*Natural Vegetation refers to the flora that grows naturally without human aid and they change their physical stability, shape and size with respect to the environment. *The Type of vegetation found in an area depends on the geographical factors such as **humidity, temperature, soil, rainfall, altitude**, etc. On this basis, there are several types of vegetation found in India.

1. Tropical Evergreen Forest-


*The tropical evergreen and semi-evergreen forest is usually found in warm and humid areas receiving more than **200 cm** of rainfall and a mean annual temperature above **22° C**. The relative humidity of this region is more than 70%. *Due to high humidity and temperature, trees are **very tall** and

intensive (dense). They are also called as “**Selvas**”.

*There is a different time for leaf fall for different species of trees, due to which the entire forest is evergreen. The tropical evergreen forest in India is found in **Assam, Kerala, West Bengal, The Andaman-Nicobar Islands, Meghalaya, Tripura**, Mizoram and Western coastal plain.

The major species found in the tropical evergreen forest include **Cinchona, Rubber, Coconut, Bamboo, Mahogany, Ebony**, etc.

2. Tropical Deciduous or Monsoon Forests: *Tropical deciduous forests are also called **monsoon forests**. Trees shed their leaves in the dry season to conserve water. The average rainfall in this region is **70-200 cm**. On the basis of availability of water, these forests can be categorized as Tropical Moist Deciduous Forest and Tropical Dry Deciduous Forest. In India, Tropical Moist Deciduous Forests are found in the eastern slope of the Western Ghats, the northeastern region of the Indian peninsula and the Bhabar and Tarai (Terai) region. *Trees found in these forests include **Sal, Teak, Neem, Shisham and Mango**. Hardwood trees are extremely helpful for making furniture.



3. Tropical Dry Deciduous Forest:

*These forests are found in the region where there is annual rainfall ranging between **70 and 100 cm**. These forests are found mainly in **Uttar Pradesh, Andhra Pradesh, Maharashtra, Karnataka and Tamil Nadu**. *The main trees are

Rosewood, Mango, Sandalwood, and Mahua. These forests are economically important and valuable.

4. Arid Forest:

*Arid forests are also referred to as ‘**Tropical Thorn Forests**’. They are found in the areas receiving **less than 70 cm rainfall**. *They include semi-arid areas of **south-west Punjab, South-west Haryana, Rajasthan and Gujarat**.

*These forests consist of a variety of grasses and shrubs. In these forests, plants remain leafless for most of the year. Important species found are **babool, ber, wild date palm, khair, neem, khejri, palas**, etc.

5. Delta Forest :-

*Deltaic forest is also known as mangrove, swampy or **littoral forest**. Mangroves grow along the coast in the salt marshes, tidal creeks, mudflats and estuaries. Deltaic forest develops in the deltaic regions of the Ganga, Godavari, Brahmaputra, Krishna rivers, etc. The Sundarban forest is the best example of a deltaic forests.

6. Montane Forests:-

In mountainous areas, the decrease in temperature with increasing altitude leads to a corresponding change in natural vegetation. Mountain forests can be classified into two types: the northern mountain forest and the southern mountain forest.

*The Himalayan range shows a succession of vegetation from tropical to **tundra**.

Deciduous types of forests are found in the foothills of the Himalayas. It is succeeded by the wet temperate type of forests between an altitude of about **1000-2000m**. Evergreen broad-leaf trees such as **oak** and **chest nut** are predominantly found here. At about 2000-3000m altitude on the southern slopes of the Himalayas, temperate forests containing coniferous trees like **Pine, Deodar, Silver Fir, Spruce** and **Cedar** are found. These forests are also found in places with high altitudes in southern and north-east India.

*Alpine forests grow all along the Himalayas at altitudes ranging between about **3000-4900 meters**. **Deodar**, a highly valued endemic species grows mainly in the western part of the Himalayan range at an altitude of about **1500-3000 meter**.

*The southern mountain forests include the forests found in three distinct areas of the peninsula e.g. the Western Ghats, the Vindhyas and the Nilgiris. Vegetation in these forests is temperate in higher regions and sub tropical in lower regions. The temperate forests are called **sholas** in the Nilgiris, Anaimalai and Palani Hills.

***Butea monosperma** is called the “**flame of the forest**”. It is also known as Dhak or **Palash**. Palash has been declared as the **state flower of Uttar Pradesh**.

1. The birch tree is found in –

- (a) Aravallis (b) Himalayas
(b) Nilgiris (d) Vindhya

Uttarakhand P.C.S. (Pre) 2006

Ans. (b)

The birch tree is found in the Himalayas. The white paper-like bark of the tree was used in ancient times for writing scriptures and texts.

2. Which of the following woods is used for manufacturing Katha?

- (a) Sal (b) Khair
(c) Babool (d) Saaja

M.P.P.C.S. (Pre) 1990

Ans. (b)

The Katha is obtained from the wood of Khair. Katha is a white substance found in Khair wood. It grows throughout India from the Himalayas to the south of India, especially in arid regions.

3. In India, which type of forest among the following occupies the largest area?

- (a) Montane Wet Temperate Forest
(b) Sub-tropical Dry Evergreen Forest
(c) Tropical Moist Deciduous Forest
(d) Tropical Wet Evergreen Forest

I.A.S. (Pre) 2010

Ans. (c)

Tropical (Dry) deciduous forest occupies the largest area in India. The tropical moist deciduous forest is the second largest forest area in India. Teak and Sal are significant products of it. According to ISFR, 2023, montane wet temperate forest is found on 2.72 percent area, subtropical dry evergreen forest on 0.02% area, tropical wet evergreen forest on 3.34% area and tropical dry deciduous forest on 39.22% area.

4. Teak and Sal are the products of –

- (a) Tropical Dry Deciduous Forests
(b) Tropical Evergreen Forests
(c) Tropical Thorn Forests
(d) Alpine Forests

U.P.P.C.S. (Mains) 2007

Ans. (a)

See the explanation of the above question.

5. Which one of the following groups of forest is spread over larger area in India?

- (a) Tropical Wet Evergreen
(b) Tropical Dry Deciduous

(c) Tropical Moist Deciduous

(d) Tropical Semi-Evergreen

Uttarakhand P.C.S. (Pre) 2024

Ans. (b)

Tropical dry deciduous forests are found in the largest area in India. They cover 43.85% of the area of all types of forest groups and 8.53% of the country's geographical area. This type of tropical deciduous forest, also known as monsoon forest, is one of the types among tropical humid and dry deciduous forests. Tropical dry deciduous forests are found in those extensive areas of the country where the annual rainfall is approximately 70-100 cm.

6. In Himalayan ranges the types of vegetation changes with altitude due to the following reasons-

1. decrease in temperature
2. changes in rain forms
3. unfertile soil
4. strong winds

Select the correct answer

- (a) 1, 2 and 3 (b) 2, 3 and 4
(c) 1, 3 and 4 (d) 1, 2 and 4
(e) None of these

Chhattisgarh P.C.S. (Pre) 2013

Ans. (a)

If we talk generally, the vegetation decreases with the increasing altitude and the sun facing side of a mountain also has better vegetation growth. Significantly, biodiversity is normally greater in the lower altitudes as compared to the higher altitudes. The major reasons for this change include altitude, drop in temperature, change in rainfall, infertility of soil, lowering of atmospheric pressure and lightening of air. Apart from this, with the increase in altitude, there is a decrease in the land area and changes in climate and latitudinal position also have an effect.

Note - The answer to this question was shown as deleted by the Chhattisgarh Public Service Commission in its revised answer sheet. That is, except for this question, the evaluation was done.

7. Consider the following types of erosion and arrange these in correct sequence in terms of the increasing order of soil loss from the field due to these types of erosion :

- I. Gully erosion II. Splash erosion
III. Rill erosion IV. Sheet erosion

Select the correct answer from the code given below :

Code :

- (a) II, III, IV, I (b) IV, I, III, II
(c) II, IV, III, I (d) III, II, I, IV

U.P. R.O./A.R.O. (Pre) 2023

Ans. (c)

States of soil erosion (increasing order of soil loss)-
Splash erosion, Sheet erosion. Rill erosion - Gully erosion.
Notably, erosion is the process where rocks are broken down by natural forces such as wind or water.

8. Compared to the Eastern Himalayas, the height of the tree-line in the Western Himalayas is:

- (a) More (b) Less
(c) Same (d) Unrelated variable

U.P. P.C.S. (Mains) 2005

Ans. (b)

The tree line is the edge of the habitat at which trees are capable of growing. The value of the natural tree-line in the Eastern Himalayas and Central Himalayas is about 3600-3800m. In the western and northwestern Himalayan region, its value decreases to about 3300-3600m. Therefore, the value of tree-line is less in the Western Himalayas as compared to the Eastern Himalayas.

9. Which is the major resources of the Western Himalayan Region?

- (a) Forest
(b) Metallic Minerals
(c) Non-Metallic Minerals
(d) Atomic Minerals

R.A.S./R.T.S. (Pre) 2010

Ans. (a)

Forests are the main resources of the Western Himalayan region. About 65 percent of the area is covered by forests.

10. Which one of the following tree is abundant in the temperate zone of Western Himalayas?

- (a) Pine (b) Deodar
(c) Silver fir (d) Blue pine

Uttarakhand U.D.A./L.D.A (Pre) 2003

Ans. (b)

The Western Himalayan region extends at an elevation of 1500-2500 meters. Its temperate zone is rich in forests of Deodar, Chir, Pine, other conifers and broad-leaved temperate trees. The trees of Deodar are more abundant in the temperate zone of the Western Himalayas than any other plants.

11. Given below are two statements, in which one is labelled as Assertion (A) and the other as Reason (R) :

Assertion (A) : The northern slopes of the Himalayas have a thicker vegetation cover compared to the southern slopes.

Reason (R) : Southern slopes of Himalayas receive more rainfall than the northern slopes.

Select the correct answer using the code given below :

Code :

- (a) (A) is false but (R) is true.
(b) Both (A) and (R) are true but (R) is not the correct explanation of (A).
(c) Both (A) and (R) are true and (R) is the correct explanation of (A).
(d) (A) is true but (R) is false.

U.P. R.O./A.R.O. (Pre) 2023

Ans. (a)

In the southern slopes of the Himalayas, both rainfall and sunlight are higher compared to the northern slopes. That is why denser vegetation is found on the southern slopes of the Himalayan region compared to the northern slopes.

12. In which one of the following states Cinchona trees does not grow?

- (a) Assam (b) Kerala
(c) Chhattisgarh (d) West Bengal

U.P.P.S.C. (GIC) 2010

Ans. (c)

The Cinchona is an evergreen tropical plant. In India, evergreen forests are found in Tamil Nadu, Karnataka, Kerala, Maharashtra, Assam, Arunachal Pradesh, Meghalaya, Nagaland, Tripura, West Bengal and the Andaman and Nicobar Islands while Chhattisgarh is suitable for tropical dry and monsoon type forests. The Cinchona tree does not grow in Chhattisgarh. It is found in Assam, Kerala and West Bengal.

13. Which one of the following is called 'flame of the forest'?

- (a) Bouhinia variegata
(b) Jacaranda mimosifolia
(c) Butea monosperma
(d) Tectona grandis

R.A.S./R.T.S.(Pre) 2010

Ans. (c)

Butea monosperma is called the "flame of the forest". It is also known as Dhak or Palash. Palash (Butea monosperma) has been declared as the State Flower of Uttar Pradesh.

14. Teak forest is found in which of the following states of India?

- (a) Madhya Pradesh (b) Uttar Pradesh
(c) Jharkhand (d) Karnataka

U.P.P.C.S.(Pre) 2012

Ans. (*)

Among the given options, according to the India State of Forest Report, 2023, teak forest is found in all three states including Madhya Pradesh, Karnataka and Uttar Pradesh. The maximum Teak forest is found in Madhya Pradesh.

15. Which of the following plants have no flowers?

- (a) Jackfruit (b) Fig
(c) Orchid (d) Fern

U.P.P.C.S. (Pre) 1991

Ans. (d)

The fern is a vascular plant that reproduces by spores and has neither seeds nor flowers. These plants are seedless. Thus, option (d) is the correct answer,

16. The alpine vegetation in the western Himalayas is found only up to a height of 3000 metres while in the eastern Himalayas it is found up to a height of 4000 metres. The reason for this variation in the same mountain range is that :

- (a) Eastern Himalayas are higher than the western Himalayas
(b) Eastern Himalayas are nearer to the equator and sea coast than the western Himalayas
(c) Eastern Himalayas get more monsoon rainfall than the western Himalayas
(d) Eastern Himalayan rocks are more fertile than the western Himalayan rocks

I.A.S. (Pre) 1995

Ans. (b)

Since the eastern Himalayas are nearer to the equator and sea coast, the snowline in the eastern Himalayas is at a higher altitude than in the western Himalayas. As a result of this, there is variation in the height of vegetation in the Himalayas.

17. What is the difference between the antelopes Oryx and Chiru?

- (a) Oryx is adapted to live in hot and arid areas, whereas Chiru is adapted to live in steppes and semi-desert areas of cold high mountains
(b) Oryx is poached for its antlers whereas Chiru is poached for its musk

- (c) Oryx exists in western India only whereas Chiru exists in North-East India only
(d) None of the statements (a), (b) and (c) given above is correct

I.A.S. (Pre) 2012

Ans. (a)

Oryx is adapted to live in hot and arid areas like Saudi Arabia, etc. while Chiru is adapted to live in steppes and semi-desert areas of cold high mountains. This is the main difference between the two from the given options.

18. Match the following –

- | | |
|--------------|--------------------------------------|
| (A) Teak | 1. Himalayan Tarai Region |
| (B) Deodar | 2. Central India |
| (C) Sundari | 3. Sunderban |
| (D) Cinchona | 4. High Altitude regions of Himalaya |

Code :

A	B	C	D
(a) 2	4	3	1
(b) 3	2	1	4
(c) 4	1	3	2
(d) 2	3	4	1

U.P.P.C.S. (Pre) 1992

Ans. (a)

Teak: The major expansion of Teak trees is found in India i.e., Madhya Pradesh, Chhattisgarh and Odisha.

Deodar: Cedrus deodara is a species of cedar native to the western Himalayas at an elevation of 1500- 3000 meters.

Sundari : It is the dominant mangrove tree species of the Sundarbans of Bangladesh and India.

Cinchona: The Cinchona is a plant of the Himalayan terai (tarai) region.

19. Sundarbans are which of the following types of forests?

- (a) Tropical dry deciduous forest
(b) Alpine forest
(c) Sub-tropical pine forest
(d) Littoral or Tidal forest

M.P.P.C.S. (Pre) 2022

Ans. (d)

See the explanation of the above question.

20. Which one of the following pairs is correctly matched?

- | | |
|----------------|---------------------|
| (a) Teak | : Jammu and Kashmir |
| (b) Deodar | : Madhya Pradesh |
| (c) Sandalwood | : Kerala |
| (d) Sundari | : West Bengal |

I.A.S. (Pre) 1997

Ans. (d)

See the explanation of the above question.

21. Match List-I with List-II and select the correct answer using the code given below:

List I (Forest)	List II (Region)
A. Tropical moist deciduous	1.Arunachal Pradesh
B. Tropical dry deciduous	2.Sahyadris
C. Alpine	3.Middle Ganga Plain
D. Tropical evergreen	4.Tarai

Code :

- | | A | B | C | D |
|-----|---|---|---|---|
| (a) | 4 | 3 | 1 | 2 |
| (b) | 4 | 2 | 1 | 3 |
| (c) | 1 | 3 | 2 | 4 |
| (d) | 3 | 1 | 4 | 2 |

U.P.P.C.S.(Pre) 2001

Ans. (a)

The description of forest regions found in India is given below:

Tropical moist deciduous forest: These forests are found in areas of moderate rainfall of 100 to 200 cm per annum therefore they are known as monsoon forests. Such areas include a belt running along the eastern slopes of Western Ghats, the northeastern region of the peninsular India and Bhabar and Tarai (Terai) regions.

Tropical dry deciduous forest: The dry deciduous forests are mainly situated in the areas, where annual rainfall ranges from 70- 100 cm. These are spread from central Gangetic plains to southern India.

Alpine Forest: The Alpine forests grow all along the Himalayas at altitudes ranging between 3,000 to 4,900 meters. The expansion of this forest is found in Arunachal Pradesh.

Tropical Evergreen Forests: These are typical rainforests that grow in areas where the annual rainfall exceeds 200 cm. The evergreen forests are found along the Western side of the Western Ghats (The Sahyadri), in the northeast and in the Andaman and Nicobar Islands. Thus, option (a) is the correct answer.

22. Consider the following statements regarding the tropical dry deciduous forest in India:

1. It is found in the regions, where rainfall is between 25 and 75 cm.
2. They occur in an irregular wide strip running from the foot of the Himalayas.

Select the correct answer from the code given below :

Code :

- | | |
|------------|------------|
| (a) 1 only | (b) 2 only |
|------------|------------|

(c) Both 1 and 2

(d) Neither 1 nor 2

Uttarakhand P.C.S. (Pre) 2024

Ans. (d)

Tropical deciduous forests also known as monsoon forests are found in areas where the annual rainfall is between 70 and 200 cm. Based on the availability of water, these forests are divided into tropical humid and dry deciduous forests. Tropical humid deciduous forests are found in areas with 100-200 cm of rainfall. These forests grow in the northeastern states, the foothills of the Himalayas the eastern slopes of the Western Ghats and Odisha.

Tropical dry deciduous forests are found in areas with 70-100 cm of annual rainfall. These forests merge into humid deciduous forests in humid areas and into thorny forests in dry areas. They are found in the more rainfall-prone region of the peninsula. Thus, the statements 1 and 2 given in the context of tropical dry deciduous forest are both incorrect. The commission has considered option (b) as correct in its provisional answer key but it is not correct. The correct option is (d).

23. Match List-I with List-II and select the correct answer by using the codes given below:

List-I	List-II
A. Tropical forest	1. Sunderbans
B. Conifer forest	2. Himachal Pradesh
C. Mangroves	3. Rajasthan
D. Deciduous forest	4. Silent Valley

Code :

- | | A | B | C | D |
|-----|---|---|---|---|
| (a) | 1 | 2 | 4 | 3 |
| (b) | 2 | 1 | 4 | 3 |
| (c) | 1 | 4 | 2 | 3 |
| (d) | 4 | 2 | 1 | 3 |

U.P.P.C.S. (Pre) 2017

Ans. (d)

The correct match of List-I and List-II is as follows :

List - I	List-II
Tropical forest	- Silent Valley
Conifer forest	- Himachal Pradesh
Mangroves	- Sunderbans
Deciduous forest	- Rajasthan

24. Open stunted forests with bushes and small trees having long roots and sharp thorns repeated spines are commonly found in:

- | | |
|-------------------------------|------------------------------|
| (a) Eastern Odisha | (b) North-Eastern Tamil Nadu |
| (c) Siwaliks and Terai region | (d) Western Andhra Pradesh |

I.A.S. (Pre) 2002

Ans. (d)

The above mentioned features are related to tropical thorn forests. These forests are found in areas of low rainfall (less than 70 cm), low humidity (less than 50 percent) and high temperature (25°-30°C). Western Andhra Pradesh is a region of such forests. These forests are found in the north-western parts of the country, including Rajasthan, South Western Punjab, Western Haryana, Kutch and neighboring parts of Saurashtra. Thus, option (d) is the correct answer.

25. Which among the following trees is found at the highest elevation from the sea level ?

- (a) Deodar (b) Pine
(c) Sal (d) Teak

Uttarakhand P.C.S. (Pre) 2002

Ans. (a)

Deodar trees are found at the highest elevation of 1500-3000 meters above sea level in western Himalayas. Details of other plants are as follows:

Chir: It is native to the Himalayas and found at an elevation of between 900 to 1000 m.

Sal: It is mainly found in sub-Himalayan regions, eastern region of Central India and Tamil Nadu.

Teak: These trees are mainly found in Madhya Pradesh, Chhattisgarh, Odisha, Karnataka, Tamil Nadu, Gujarat & Maharashtra.

26. Forest areas have been labeled as 1, 2, 3 and 4 in the rough map:



Among these, those which were threatened in 1997 by a severe epidemic include –

- (a) Teak forests of 3 and 4
(b) Oak forests of 1 and Sal forests of 2
(c) Sal forests of 3
(d) Sandalwood forests of 4

I.A.S. (Pre) 1998

Ans. (c)

The place marked '3' in the map is the Sal forests which were threatened in 1997 by a serious epidemic. These were infected by a tree pest that weakened the trees, resulting in forest loss.

27. Which of the following state's forests are classified as "Subtropical" forests?

- (a) Kerala (b) Madhya Pradesh
(c) Tamil Nadu (d) Karnataka

R.A.S./R.T.S. (Pre) 2003

Ans. (b)

Tropical dry deciduous forest or Sub tropical forests expansion is found in most parts of Madhya Pradesh.

28. Which one of the following is not essentially a species of the Himalayan vegetation?

- (a) Juniper (b) Mahogany
(c) Silver fir (d) Spruce

I.A.S. (Pre) 2008

Ans. (b)

Originally, Mahogany trees are found in tropical regions of North & South America. This tree is abundant in the West Indies, Central America & South America. The rest of the trees are essentially species of Himalayan vegetation.

29. An example of a multipurpose tree used in social forestry is –

- (a) Sheesham (b) Dhak
(c) Eucalyptus (d) Khejri

U.P.P.C.S. (Mains) 2002

Ans. (d)

An example of a multipurpose tree used in social forestry is Khejri. This tree is known as 'king of the desert'. The wood of Khejri is a good source fuel and provides excellent charcoal plus firewood, fodder, green manure and goat-proof thorny fences. Khejri is frost-resistant, drought-resistance and withstands wild temperature extremes. It requires minimal rainfall. That's why Khejri is the preferred plant species for social forestry programs.

30. With reference to 'Victoria Falls in Africa', which of the following statement/s is/are correct?

1. It is located on the border of Zambia and Mozambique.
2. It is situated on Zambezi River.

Select the correct answer using the code given below:
Code "

- (a) Only 1 (b) Neither 1 nor 2
(c) Only 2 (d) Both 1 and 2

U.P. R.O./A.R.O. (Pre) 2023

Ans. (c)

The Victoria Falls, situated on the Zambezi River, lies on the borders of Zambia and Zimbabwe in the African continent. It is also known as 'The Smoke that Thunders'. It was included in the World Heritage List in the year of 1989.

31. Lisa is obtained from the tree of :

- (a) Pine (b) Devdar
(c) Burans (d) Sheesham

Uttarakhand P.C.S. (Mains) 2002

Ans. (a)

Lisa is a significant forest product obtained from pine trees of Uttarakhand. It is also tapped commercially for resin. On distillation, the resin yields an essential oil, commonly known as turpentine oil and non-volatile resin which are used in a wide variety of industries such as paper, soap, cosmetics, paint, varnish, rubber and polish industries. Lisa is the prime resource for self-employment and a base of the rural economy of Uttarakhand.

32. Match List-I (Mangrove) with List-II (State) and select the correct answer from the codes given below the lists:

- | List-I (Mangroves) | List-II (State) |
|--------------------|-------------------|
| A. Achra Ratnagiri | 1. Karnataka |
| B. Coondapur | 2. Kerala |
| C. Pichavaram | 3. Andhra Pradesh |
| D. Vembanad | 4. Maharashtra |
| | 5. Tamil Nadu |

Code :

- | | A | B | C | D |
|-----|---|---|---|---|
| (a) | 2 | 1 | 5 | 4 |
| (b) | 4 | 5 | 3 | 2 |
| (c) | 2 | 5 | 3 | 4 |
| (d) | 4 | 1 | 5 | 2 |

I.A.S. (Pre) 2002

Ans. (d)

The description of mangrove vegetation regions found in India is given below:

Mangrove vegetation areas of Maharashtra: Achra Ratnagiri, Raigad, Devgarh - Vijaydurg, Kundalika Revdanda, Mumbara-Diva, Srivardhan, Vaitarna, Malvan etc.

Mangrove vegetation areas of Karnataka: Coondapur, Honnavar, Karwar, etc.

Mangrove vegetation areas of Kerala : Vembanad, Kannur.

Mangrove vegetation areas of Tamil Nadu: Pichavaram, Muthupet and Ramnad, Pulicat, Kajuveli etc.

Mangrove vegetation areas of West Bengal: Sundarbans

Mangrove vegetation areas of Odisha: Bhitarkanika, Devi, Dhamra, Chilika etc. Thus, option (d) is correct.

33. Which of the following is not correctly matched?

- (a) Sundarbans - West Bengal
(b) Bhitarkanika - Odisha

- (c) Pichavaram - Tamil Nadu
(d) Vembanad - Karnataka

Uttarakhand P.C.S. U.D.A. (Pre) 2003

U.P. Lower Sub. (Pre) 2002

Ans. (d)

See the explanation of the above question.

34. Tropical rainforest in India is found in :

- (a) Assam and Nagaland
(b) Odisha and Chhattisgarh
(c) Uttar Pradesh and Maharashtra
(d) Assam and Kerala

Jharkhand P.C.S. (Mains) 2016

Ans. (d)

The tropical rainforests are found in Kerala and Assam. These forests are confined to the high rainfall areas of the Western Ghats, Lakshadweep, Andaman and Nicobar Islands, upper reaches of Assam and the coast of Tamil Nadu. Significantly they are at their best areas having more than 200cm of rainfall with a short dry season.

35. In India, the State with the largest area under dense deciduous forest cover is :

- (a) Odisha (b) Maharashtra
(c) Madhya Pradesh (d) Chhattisgarh

65th B.P.S.C. (Pre) 2019

Ans. (c)

Madhya Pradesh has the largest area under dense deciduous forest cover.

36. Among the following districts, which one has the larger area under dense deciduous forest cover?

- (a) Paschim Champaran (b) Gaya
(c) Kaimur (d) Nawada
(e) None of the above/More than one of the above.

65th B.P.S.C. (Pre) 2019

Ans. (a)

In the question period and at present, West Champaran district is surrounded by the most dense deciduous forest area. According to the India State of Forest Report, 2023 West Champaran has a total of 831.03 sq. km. of dense forests, while Kaimur has 526 sq. km.

37. 'Khair (Catechu) is a tree of which type of forest?

- (a) Sub- Alpine Forests (b) Cool - Temperate Forests
(c) Sub-Tropical Forests (d) Temperate Forests

Uttarakhand P.C.S. (Pre) 2021

Ans. (c)