

G V V Sharma*

Abstract—This book provides a collection of the international maths olympiad problems in geometry.

1. Construct a right triangle with given hypotenuse c such that the median drawn to the hypotenuse is the geometric mean of the two legs of the triangle.
2. An arbitrary point M is selected in the interior of the segment AB . The squares $AMCD$ and $MBEF$ are constructed on the same side of AB , with the segments AM and MB as their respective bases. The circles circumscribed about these squares, with centers P and Q , intersect at M and also at another point N . Let N_0 denote the point of intersection of the straight lines AF and BC .
 - a) Prove that the points N and N_0 coincide.
 - b) Prove that the straight lines MN pass through a fixed point S independent of the choice of M .
 - c) Find the locus of the midpoints of the segments PQ as M varies between A and B .
3. Two planes, P and Q , intersect along the line p . The point A is given in the plane P , and the point C in the plane Q ; neither of these points lies on the straight line p . Construct an isosceles trapezoid $ABCD$ (with AB parallel to CD) in which a circle can be inscribed, and with vertices B and D lying in the planes P and Q respectively.
4. Consider triangle $P_1P_2P_3$ and a point P within the triangle. Lines P_1P , P_2P , P_3P intersect the opposite sides in points Q_1 , Q_2 , Q_3 respectively. Prove that, of the numbers

$$\frac{P_1P}{PQ_1}, \frac{P_2P}{PQ_2}, \frac{P_3P}{PQ_3}$$

at least one is ≤ 2 and at least one is ≥ 2 .

5. Construct triangle ABC if $AC = b$, $AB = c$ and $\angle AMB = \omega$, where M is the midpoint of segment BC and $\omega < 90^\circ$. Prove that a solution exists if and only if

$$b \tan \frac{\omega}{2} \leq c < b.$$

In what case does the equality hold?

6. Consider a plane ε and three non-collinear points A, B, C on the same side of ε ; suppose the plane determined by these three points is not parallel to ε . In plane ε take three arbitrary points A_0, B_0, C_0 . Let L, M, N be the midpoints of segments AA_0, BB_0, CC_0 ; let G be the centroid of triangle LMN . (We will not consider positions of the points A', B', C' such that the points L, M, N do not form a triangle.) What is the locus of point G as A', B', C' range independently over the plane ε ?
7. Point A and segment BC are given. Determine the locus of points in space which are vertices of right angles with one side passing through A , and the other side intersecting the segment BC .
8. Prove that $\cos \frac{\pi}{7} - \cos \frac{2\pi}{7} + \cos \frac{3\pi}{7} = \frac{1}{2}$
9. Given a triangle ABC , let I be the center of its inscribed circle. The internal bisectors of the angles A, B, C meet the opposite sides in A', B', C' respectively. Prove that $\frac{1}{4} < \frac{AI \cdot BI \cdot CI}{AA' \cdot BB' \cdot CC'} \leq \frac{8}{27}$.
10. Let ABC be a triangle and P an interior point of ABC . Show that at least one of the angles $\angle PAB, \angle PBC, \angle PCA$ is less than or equal to 30° .
11. Consider nine points in space, no four of

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

which are coplanar. Each pair of points is joined by an edge (that is, a line segment) and each edge is either colored blue or red or left uncolored. Find the smallest value of n such that whenever exactly n edges are colored, the set of colored edges necessarily contains a triangle all of whose edges have the same color.

12. In the plane let C be a circle, L a line tangent to the circle C , and M a point on L . Find the locus of all points P with the following property: there exists two points Q, R on L such that M is the midpoint of QR and C is the inscribed circle of triangle PQR .
13. Let D be a point inside acute triangle ABC such that $\angle ADB = \angle ACB + \frac{\pi}{2}$ and $AC \cdot BD = AD \cdot BC$.
 - a) Calculate the ratio $(AB \cdot CD)/(AC \cdot BD)$.
 - b) Prove that the tangents at C to the circumcircles of $\triangle ACD$ and $\triangle BCD$ are perpendicular.
14. ABC is an isosceles triangle with $AB = AC$. Suppose that
 - a) M is the midpoint of BC and O is the point on the line AM such that OB is perpendicular to AB ;
 - b) Q is an arbitrary point on the segment BC different from B and C ;
 - c) E lies on the line AB and F lies on the line AC such that E, Q, F are distinct and collinear.
15. Let A, B, C, D be four distinct points on a line, in that order. The circles with diameters AC and BD intersect at X and Y . The line XY meets BC at Z . Let P be a point on the line XY other than Z . The line CP intersects the circle with diameter AC at C and M , and the line BP intersects the circle with diameter BD at B and N . Prove that the lines AM, DN, XY are concurrent.
16. Let $ABCDEF$ be a convex hexagon with $AB = BC = CD$ and $DE = EF = FA$, such that $\angle BCD = \angle EFA = \frac{\pi}{3}$. Suppose G and H are points in the interior of the hexagon such that $\angle AGB = \angle DHE = \frac{2\pi}{3}$. Prove that $AG + GB + GH + DH + HE \geq CF$.
17. We are given a positive integer r and a rectangular board $ABCD$ with dimensions $|AB| = 20$, $|BC| = 12$. The rectangle is divided into a

grid of 20×12 unit squares. The following moves are permitted on the board: one can move from one square to another only if the distance between the centers of the two squares is \sqrt{r} . The task is to find a sequence of moves leading from the square with A as a vertex to the square with B as a vertex.

- a) Show that the task cannot be done if r is divisible by 2 or 3.
 - b) Prove that the task is possible when $r = 73$.
 - c) Can the task be done when $r = 97$?
18. Let P be a point inside triangle ABC such that
- $$\angle APB - \angle ACB = \angle APC - \angle ABC.$$

Let D, E be the incenters of triangles APB, APC , respectively. Show that AP, BD, CE meet at a point.

19. Let $ABCDEF$ be a convex hexagon such that AB is parallel to DE , BC is parallel to EF , and CD is parallel to FA . Let R_A, R_C, R_E denote the circumradii of triangles FAB, BCD, DEF , respectively, and let P denote the perimeter of the hexagon. Prove that

$$R_A + R_C + R_E \geq \frac{P}{2}.$$

20. In the plane the points with integer coordinates are the vertices of unit squares. The squares are colored alternately black and white (as on a chessboard). For any pair of positive integers m and n , consider a right-angled triangle whose vertices have integer coordinates and whose legs, of lengths m and n , lie along edges of the squares.

Let S_1 be the total area of the black part of the triangle and S_2 be the total area of the white part. Let

$$f(m, n) = |S_1 - S_2|.$$

- a) Calculate $f(m, n)$ for all positive integers m and n which are either both even or both odd.
 - b) Prove that $f(m, n) \leq \frac{1}{2} \max m, n$ for all m and n .
 - c) Show that there is no constant C such that $f(m, n) < C$ for all m and n .
21. The angle at A is the smallest angle of triangle ABC . The points B and C divide the

circumcircle of the triangle into two arcs. Let U be an interior point of the arc between B and C which does not contain A . The perpendicular bisectors of AB and AC meet the line AU at V and W , respectively. The lines BV and CW meet at T . Show that

$$AU = TB + TC.$$

22. In the convex quadrilateral $ABCD$, the diagonals AC and BD are perpendicular and the opposite sides AB and DC are not parallel. Suppose that the point P , where the perpendicular bisectors of AB and DC meet, is inside $ABCD$. Prove that $ABCD$ is a cyclic quadrilateral if and only if the triangles ABP and CDP have equal areas.
23. Let I be the incenter of triangle ABC . Let the incircle of ABC touch the sides BC , CA , and AB at K , L , and M , respectively. The line through B parallel to MK meets the lines LM and LK at R and S , respectively. Prove that angle RIS is acute.
24. Two circles G_1 and G_2 are contained inside the circle G , and are tangent to G at the distinct points M and N , respectively. G_1 passes through the center of G_2 . The line passing through the two points of intersection of G_1 and G_2 meets G at A and B . The lines MA and MB meet G_1 at C and D , respectively.
25. AB is tangent to the circles $CAMN$ and $NMBD$. M lies between C and D on the line CD , and CD is parallel to AB . The chords NA and CM meet at P ; the chords NB and MD meet at Q . The rays CA and DB meet at E . Prove that $PE = QE$.
26. $A_1A_2A_3$ is an acute-angled triangle. The foot of the altitude from A_i is K_i and the incircle touches the side opposite A_i at L_i . The line K_1K_2 is reflected in the line L_1L_2 . Similarly, the line K_2K_3 is reflected in L_2L_3 and K_3K_1 is reflected in L_3L_1 . Show that the three new lines form a triangle with vertices on the incircle.
27. Let ABC be an acute-angled triangle with circumcentre O . Let P on BC be the foot of the altitude from A . Suppose that $\angle BCA \geq \angle ABC + 30^\circ$. Prove that $\angle CAB + \angle COP < 90^\circ$.
28. In a triangle ABC , let AP bisect $\angle BAC$, with P on BC , and let BQ bisect $\angle ABC$, with Q on CA . It is known that $\angle BAC = 60^\circ$ and that $AB + BP = AQ + QB$. What are the possible angles of triangle ABC ?
29. BC is a diameter of a circle center O . A is any point on the circle with $\angle AOC > 60^\circ$. EF is the chord which is the perpendicular bisector of AO . D is the midpoint of the minor arc AB . The line through O parallel to AD meets AC at J . Show that J is the incenter of triangle CEF .
30. $n > 2$ circles of radius 1 are drawn in the plane so that no line meets more than two of the circles. Their centers are O_1, O_2, \dots, O_n . Show that $\sum_{i < j} \frac{1}{O_i O_j} \leq (n-1) \frac{\pi}{4}$.
31. A convex hexagon has the property that for any pair of opposite sides the distance between their midpoints is $\sqrt{3}/2$ times the sum of their lengths. Show that all the hexagon's angles are equal.
32. $ABCD$ is cyclic. The feet of the perpendicular from D to the lines AB , BC , CA are P , Q , R respectively. Show that the angle bisectors of ABC and CDA meet on the line AC if $RP = RQ$.
33. Let ABC be an acute-angled triangle with $AB \neq AC$. The circle with diameter BC intersects the sides AB and AC at M and N respectively. Denote by O the midpoint of the side BC . The bisectors of the angles $\angle BAC$ and $\angle MON$ intersect at R . Prove that the circumcircles of the triangles BMR and CNR have a common point lying on the side BC .
34. In a convex quadrilateral $ABCD$ the diagonal BD does not bisect the angles ABC and CDA . The point P lies inside $ABCD$ and satisfies $\angle PBC = \angle DBA$ and $\angle PDC = \angle BDA$.

Prove that $ABCD$ is a cyclic quadrilateral if and only if $AP = CP$.