

# Geometry through Linear Algebra

G V V Sharma\*




Fig. 1.1




Fig. 1.2

## CONTENTS

1	The Straight Line	1
2	Orthogonality	2
3	Medians of a triangle	3
4	Circle	4

**Abstract**—This textbook introduces linear algebra by exploring Euclidean geometry.

### 1 THE STRAIGHT LINE

1.1 The points  $\mathbf{O} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}$  are as shown in Fig. 1.1. Find the equation of  $OA$ .

**Solution:** Let  $\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$  be any point on  $OA$ . Then, using similar triangles,

$$\frac{x_2}{x_1} = \frac{a_2}{a_1} = m \quad (1.1)$$

$$\Rightarrow x_2 = mx_1 \quad (1.2)$$

where  $m$  is known as the slope of the line. Thus, the equation of the line is

$$\mathbf{x} = \begin{pmatrix} x_1 \\ mx_1 \end{pmatrix} = x_1 \begin{pmatrix} 1 \\ m \end{pmatrix} \quad (1.3)$$

In general, the above equation is written as

$$\mathbf{x} = \begin{pmatrix} x_1 \\ mx_1 \end{pmatrix} = \lambda \begin{pmatrix} 1 \\ m \end{pmatrix} \quad (1.4)$$

1.2 Find the equation of  $AB$  in Fig. 1.2

**Solution:** From Fig. 1.2,

$$\frac{x_2 - a_2}{x_1 - a_1} = \frac{b_2 - a_2}{b_1 - a_1} = m \quad (1.5)$$

$$\Rightarrow x_2 = mx_1 + a_2 - ma_1 \quad (1.6)$$

From (1.6),

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ mx_1 + a_2 - ma_1 \end{pmatrix} \quad (1.7)$$

$$= \mathbf{A} + (x_1 - a_1) \begin{pmatrix} 1 \\ m \end{pmatrix} \quad (1.8)$$

$$= \mathbf{A} + \lambda \begin{pmatrix} 1 \\ m \end{pmatrix} \quad (1.9)$$

1.3 Find the length of  $\mathbf{A}$  in Fig. 1.1

**Solution:** Using Baudhayana's theorem, the

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.




Fig. 1.4

length of the vector  $\mathbf{A}$  is defined as

$$\|\mathbf{A}\| = OA = \sqrt{a_1^2 + a_2^2} = \sqrt{\mathbf{A}^T \mathbf{A}}. \quad (1.10)$$

Also, from (1.4),

$$\|\mathbf{A}\| = \lambda \sqrt{1 + m^2} \quad (1.11)$$

Note that  $\lambda$  is the variable that determines the length of  $\mathbf{A}$ , since  $m$  is constant for all points on the line.

1.4 Find  $\mathbf{A} - \mathbf{B}$ .

**Solution:** See Fig. 1.4. From (1.9), for some  $\lambda$ ,

$$\mathbf{B} = \mathbf{A} + \lambda \begin{pmatrix} 1 \\ m \end{pmatrix} \quad (1.12)$$

$$\Rightarrow \mathbf{A} - \mathbf{B} = -\lambda \begin{pmatrix} 1 \\ m \end{pmatrix}, \quad (1.13)$$

$\mathbf{A} - \mathbf{B}$  is marked in Fig. 1.4.

1.5 Show that  $AB = \|\mathbf{A} - \mathbf{B}\|$

## 2 ORTHOGONALITY

2.1 See Fig. 2.1. In  $\triangle ABC$ ,  $AB \perp BC$ . Show that

$$(\mathbf{A} - \mathbf{B})^T (\mathbf{B} - \mathbf{C}) = 0 \quad (2.1)$$

**Solution:** Using Baudhayana's theorem,

$$\begin{aligned} \|\mathbf{A} - \mathbf{B}\|^2 + \|\mathbf{B} - \mathbf{C}\|^2 &= \|\mathbf{C} - \mathbf{A}\|^2 \quad (2.2) \\ \Rightarrow (\mathbf{A} - \mathbf{B})^T (\mathbf{A} - \mathbf{B}) + (\mathbf{B} - \mathbf{C})^T (\mathbf{B} - \mathbf{C}) \\ &= (\mathbf{C} - \mathbf{A})^T (\mathbf{C} - \mathbf{A}) \\ \Rightarrow 2\mathbf{A}^T \mathbf{B} - 2\mathbf{B}^T \mathbf{B} + 2\mathbf{B}^T \mathbf{C} - 2\mathbf{A}^T \mathbf{C} &= 0 \quad (2.3) \end{aligned}$$

which can be simplified to obtain (2.1).

2.2 Let  $\mathbf{x}$  be any point on  $AB$  in Fig. 2.1. Show that

$$(\mathbf{x} - \mathbf{A})^T (\mathbf{B} - \mathbf{C}) = 0 \quad (2.4)$$




Fig. 2.1

2.3 If  $\mathbf{x}, \mathbf{y}$  are any two points on  $AB$ , show that

$$(\mathbf{x} - \mathbf{y})^T (\mathbf{B} - \mathbf{C}) = 0 \quad (2.5)$$

2.4 In Fig. 2.4,  $BE \perp AC, CF \perp AB$ . Show that  $AD \perp BC$ .




Fig. 2.4

**Solution:** Let  $\mathbf{x}$  be the intersection of  $BE$  and  $CF$ . Then, using (2.5),

$$\begin{aligned} (\mathbf{x} - \mathbf{B})^T (\mathbf{A} - \mathbf{C}) &= 0 \quad (2.6) \\ (\mathbf{x} - \mathbf{C})^T (\mathbf{A} - \mathbf{B}) &= 0 \end{aligned}$$

$$\Rightarrow \mathbf{x}^T (\mathbf{A} - \mathbf{C}) - \mathbf{B}^T (\mathbf{A} - \mathbf{C}) = 0 \quad (2.7)$$

$$\text{and } \mathbf{x}^T (\mathbf{A} - \mathbf{B}) - \mathbf{C}^T (\mathbf{A} - \mathbf{B}) = 0 \quad (2.8)$$

Subtracting (2.8) from (2.7),

$$\mathbf{x}^T (\mathbf{B} - \mathbf{C}) + \mathbf{A}^T (\mathbf{C} - \mathbf{B}) = 0 \quad (2.9)$$

$$\Rightarrow (\mathbf{x}^T - \mathbf{A}^T) (\mathbf{B} - \mathbf{C}) = 0 \quad (2.10)$$

$$\Rightarrow (\mathbf{x} - \mathbf{A})^T (\mathbf{B} - \mathbf{C}) = 0 \quad (2.11)$$

which completes the proof.

### 3 MEDIANS OF A TRIANGLE

3.1 In Fig. 3.1,

$$\frac{AB}{BC} = \frac{\|\mathbf{A} - \mathbf{B}\|}{\|\mathbf{B} - \mathbf{C}\|} = k. \quad (3.1)$$

Show that

$$\frac{\mathbf{A} + k\mathbf{C}}{k + 1} = \mathbf{B}. \quad (3.2)$$

**Solution:** From (1.9),




Fig. 3.1

$$\mathbf{B} = \mathbf{A} + \lambda_1 \begin{pmatrix} 1 \\ m \end{pmatrix}, \quad (3.3)$$

$$\mathbf{B} = \mathbf{C} - \lambda_2 \begin{pmatrix} 1 \\ m \end{pmatrix}.$$

$$\Rightarrow \frac{\|\mathbf{A} - \mathbf{B}\|}{\|\mathbf{B} - \mathbf{C}\|} = \frac{\lambda_1}{\lambda_2} = k \quad (3.4)$$

$$\text{and } \frac{\mathbf{B} - \mathbf{A}}{\lambda_1} = \frac{\mathbf{C} - \mathbf{B}}{\lambda_2} = \begin{pmatrix} 1 \\ m \end{pmatrix}, \quad (3.5)$$

from (3.1). Using (3.4) and (3.4),

$$\mathbf{A} - \mathbf{B} = k(\mathbf{B} - \mathbf{C}) \quad (3.6)$$

resulting in (3.2).

3.2 If  $\mathbf{A}$  and  $\mathbf{B}$  are linearly independent,

$$k_1\mathbf{A} + k_2\mathbf{B} = 0 \Rightarrow k_1 = k_2 = 0 \quad (3.7)$$

3.3  $BE$  and  $CF$  are medians of  $\triangle ABC$  intersecting at  $O$  as shown in Fig. 3.3. Show that

$$\frac{CO}{OF} = \frac{BO}{OE} = 2 \quad (3.8)$$

**Solution:** Let

$$\frac{CO}{OF} = k_1 \quad (3.9)$$

$$\frac{BO}{OE} = k_2 \quad (3.10)$$

Using (3.2),

$$\mathbf{E} = \frac{\mathbf{A} + \mathbf{C}}{2} \quad (3.11)$$

$$\mathbf{F} = \frac{\mathbf{A} + \mathbf{B}}{2} \quad (3.12)$$

and

$$\mathbf{O} = \frac{k_1\mathbf{F} + \mathbf{C}}{k_1 + 1} = \frac{k_1\frac{\mathbf{A} + \mathbf{B}}{2} + \mathbf{C}}{k_1 + 1} \quad (3.13)$$

$$\mathbf{O} = \frac{k_2\mathbf{E} + \mathbf{B}}{k_2 + 1} = \frac{k_2\frac{\mathbf{A} + \mathbf{C}}{2} + \mathbf{B}}{k_2 + 1} \quad (3.14)$$

From (3.13) and (3.14),

$$\frac{k_1\frac{\mathbf{A} + \mathbf{B}}{2} + \mathbf{C}}{k_1 + 1} = \frac{k_2\frac{\mathbf{A} + \mathbf{C}}{2} + \mathbf{B}}{k_2 + 1} \quad (3.15)$$

$$\begin{aligned} \Rightarrow & \left[ \frac{k_1(k_2 + 1)}{2} - \frac{k_2(k_1 + 1)}{2} \right] \mathbf{A} \\ & + \left[ \frac{k_1(k_2 + 1)}{2} - (k_1 + 1) \right] \mathbf{B} \\ & + \left[ (k_2 + 1) - \frac{k_2(k_1 + 1)}{2} \right] \mathbf{C} = 0 \end{aligned} \quad (3.16)$$

resulting in  $k_1 = k_2$ ,

$$k_1^2 - k_1 - 2 = 0 \Rightarrow k_1 = k_2 = 2, \quad (3.17)$$

provided  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are linearly independent. Thus, substituting  $k_1 = 2$  in (3.14),

$$\mathbf{O} = \frac{\mathbf{A} + \mathbf{B} + \mathbf{C}}{3} \quad (3.18)$$

If  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are linearly dependent,

$$\mathbf{A} = \alpha\mathbf{B} + \beta\mathbf{C} \quad (3.19)$$




Fig. 3.3

Note that  $\mathbf{B}, \mathbf{C}$  are linearly independent. Substituting (3.19) in (3.16),

$$\begin{aligned} & \left[ \frac{k_1(k_2+1)}{2} - \frac{k_2(k_1+1)}{2} \right] [\alpha\mathbf{B} + \beta\mathbf{C}] \\ & + \left[ \frac{k_1(k_2+1)}{2} - (k_1+1) \right] \mathbf{B} \\ & + \left[ (k_2+1) - \frac{k_2(k_1+1)}{2} \right] \mathbf{C} = 0 \quad (3.20) \end{aligned}$$

$$\begin{aligned} \implies & (k_1 - k_2)\alpha + k_1k_2 - k_1 - 2 = 0 \\ & (k_1 - k_2)\beta - k_1k_2 + k_2 + 2 = 0 \quad (3.21) \end{aligned}$$

$$\implies (k_1 - k_2)(\alpha + \beta - 1) = 0 \quad (3.22)$$

If  $\alpha + \beta = 1$ ,  $\mathbf{A}, \mathbf{B}, \mathbf{C}$  are collinear according to (3.2) resulting in a contradiction. Hence,  $k_1 = k_2$ , which, upon substitution in (3.21), yields

$$k_1^2 - k_1 - 2 = 0 \implies k_1 = 2. \quad (3.23)$$

#### 4 CIRLE

4.1 A circle with centre at origin  $\mathbf{O}$  and radius  $r$  has the equation

$$\mathbf{x}^T \mathbf{x} = r^2 \quad (4.1)$$

Let  $\mathbf{A}$  and  $\mathbf{B}$  be two points on the circle. Show that

$$(\mathbf{A} - \mathbf{B})^T (\mathbf{A} + \mathbf{B}) = 0 \quad (4.2)$$

4.2 Let  $\mathbf{D}$  be the mid point of  $AB$ . Show that  $OD \perp AB$ .

4.3 Show that the direction vector of the normal to the tangent at  $\mathbf{P}$  is  $\mathbf{P}$ .

4.4 Let  $\mathbf{A}$  be any point on the tangent. Show that  $OP \perp AP$ .

4.5 Show that the angle in a semi-circle is a right angle.