

# The Straight Line

G V V Sharma\*

**Abstract**—Solved problems from JEE mains papers related to 2D lines in coordinate geometry are available in this document. These problems are solved using linear algebra/matrix analysis.

- 1 A straight line through the origin **O** meets the lines

$$(4 \ 3)\mathbf{x} = 10 \quad (1.1)$$

$$(8 \ 6)\mathbf{x} + 5 = 0 \quad (1.2)$$

at **A** and **B** respectively. Find the ratio in which **O** divides **AB**.

**Solution:** Let

$$\mathbf{n} = (4 \ 3) \quad (1.3)$$

Then (1.1) can be expressed as

$$\mathbf{n}^T \mathbf{x} = 10 \quad (1.4)$$

$$2\mathbf{n}^T \mathbf{x} = -5 \quad (1.5)$$

and since **A**, **B** satisfy (1.4) respectively,

$$\mathbf{n}^T \mathbf{A} = 10 \quad (1.6)$$

$$2\mathbf{n}^T \mathbf{B} = -5 \quad (1.7)$$

Let **O** divide the segment **AB** in the ratio  $k : 1$ . Then

$$\mathbf{O} = \frac{k\mathbf{B} + \mathbf{A}}{k + 1} \quad (1.8)$$

$$\therefore \mathbf{O} = \mathbf{0}, \quad (1.9)$$

$$\mathbf{A} = -k\mathbf{B} \quad (1.10)$$

Substituting in (1.6), and simplifying,

$$\mathbf{n}^T \mathbf{B} = \frac{10}{-k} \quad (1.11)$$

$$\mathbf{n}^T \mathbf{B} = \frac{-5}{2} \quad (1.12)$$

resulting in

$$\frac{10}{-k} = \frac{-5}{2} \implies k = 4 \quad (1.13)$$

- 2 The point

$$\mathbf{P} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (2.1)$$

is translated parallel to the line

$$L : (1 \ -1)\mathbf{x} = 4 \quad (2.2)$$

by  $2\sqrt{3}$  units. If the new point **Q** lies in the third quadrant, then find the equation of the line passing through **Q** and perpendicular to **L**.

**Solution:** From (2.2), the direction vector of **L** is

$$\mathbf{m} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (2.3)$$

Thus,

$$\mathbf{Q} = \mathbf{P} + \lambda \mathbf{m} \quad (2.4)$$

However,

$$PQ = 2\sqrt{3} \quad (2.5)$$

$$\implies \|\mathbf{P} - \mathbf{Q}\| = |\lambda| \|\mathbf{m}\| = 2\sqrt{3} \quad (2.6)$$

$$\implies \lambda = \pm \frac{2\sqrt{3}}{\|\mathbf{m}\|} = \pm \sqrt{6} \quad (2.7)$$

$$\therefore \|\mathbf{m}\| = \sqrt{\mathbf{m}^T \mathbf{m}} = \sqrt{2} \quad (2.8)$$

from (2.3). Since **Q** lies in the third quadrant,

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

from (2.4) and (2.7),

$$\mathbf{Q} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} - \sqrt{6} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 - \sqrt{6} \\ 1 - \sqrt{6} \end{pmatrix} \quad (2.9)$$

The equation of the desired line is then obtained as

$$\mathbf{m}^T (\mathbf{x} - \mathbf{Q}) = 0 \quad (2.10)$$

$$\begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = 3 - \sqrt{6} \quad (2.11)$$

- 3 A variable line drawn through the intersection of the lines

$$\begin{pmatrix} 4 & 3 \end{pmatrix} \mathbf{x} = 12 \quad (3.1)$$

$$\begin{pmatrix} 3 & 4 \end{pmatrix} \mathbf{x} = 12 \quad (3.2)$$

meets the coordinate axes at  $\mathbf{A}$  and  $\mathbf{B}$ , then find the locus of the midpoint of  $AB$ .

**Solution:** The intersection of the lines in (3.1) is obtained through the matrix equation

$$\begin{pmatrix} 4 & 3 \\ 3 & 4 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 12 \\ 12 \end{pmatrix} \quad (3.3)$$

by forming the augmented matrix and row reduction as

$$\begin{pmatrix} 4 & 3 & 12 \\ 3 & 4 & 12 \end{pmatrix} \leftrightarrow \begin{pmatrix} 4 & 3 & 12 \\ 0 & 7 & 12 \end{pmatrix} \leftrightarrow \begin{pmatrix} 28 & 0 & 48 \\ 0 & 7 & 12 \end{pmatrix} \\ \leftrightarrow \begin{pmatrix} 7 & 0 & 12 \\ 0 & 7 & 12 \end{pmatrix} \quad (3.4)$$

resulting in

$$\mathbf{C} = \frac{1}{7} \begin{pmatrix} 12 \\ 12 \end{pmatrix} \quad (3.5)$$

Let the  $\mathbf{R}$  be the mid point of  $AB$ . Then,

$$\mathbf{A} = 2 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R} \quad (3.6)$$

$$\mathbf{B} = 2 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{R} \quad (3.7)$$

Let the equation of  $AB$  be

$$\mathbf{n}^T (\mathbf{x} - \mathbf{C}) = 0 \quad (3.8)$$

Since  $\mathbf{R}$  lies on  $AB$ ,

$$\mathbf{n}^T (\mathbf{R} - \mathbf{C}) = 0 \quad (3.9)$$

Also,

$$\mathbf{n}^T (\mathbf{A} - \mathbf{B}) = 0 \quad (3.10)$$

Substituting from (3.6) in (3.10),

$$\mathbf{n}^T \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \mathbf{R} = 0 \quad (3.11)$$

From (3.9) and (3.11),

$$(\mathbf{R} - \mathbf{C}) = k \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \mathbf{R} \quad (3.12)$$

for some constant  $k$ . Multiplying both sides of (3.12) by

$$\mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (3.13)$$

$$\mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (\mathbf{R} - \mathbf{C}) = k \mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \mathbf{R} \\ = k \mathbf{R}^T \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \mathbf{R} = 0 \quad (3.14)$$

$$\therefore \mathbf{R}^T \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \mathbf{R} = 0 \quad (3.15)$$

which can be easily verified for any  $\mathbf{R}$ . from (3.14),

$$\mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (\mathbf{R} - \mathbf{C}) = 0 \\ \Rightarrow \mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R} - \mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{C} = 0 \\ \Rightarrow \mathbf{R}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R} - \mathbf{C}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R} = 0 \quad (3.16)$$

which is the desired locus.

- 4 Two sides of a rhombus are along the lines

$$AB : \begin{pmatrix} 1 & -1 \end{pmatrix} \mathbf{x} + 1 = 0 \quad (4.1)$$

$$AD : \begin{pmatrix} 7 & -1 \end{pmatrix} \mathbf{x} - 5 = 0. \quad (4.2)$$

If its diagonals intersect at

$$\mathbf{P} = \begin{pmatrix} -1 \\ -2 \end{pmatrix}, \quad (4.3)$$

find its vertices.

**Solution:** From (4.1) and (4.2),

$$\begin{pmatrix} 1 & -1 \\ 7 & -1 \end{pmatrix} \mathbf{A} = \begin{pmatrix} -1 \\ 5 \end{pmatrix} \quad (4.4)$$

By row reducing the augmented matrix

$$\begin{pmatrix} 1 & -1 & -1 \\ 7 & -1 & 5 \end{pmatrix} \leftrightarrow \begin{pmatrix} 1 & -1 & -1 \\ 0 & 6 & 12 \end{pmatrix} \leftrightarrow \begin{pmatrix} 1 & -1 & -1 \\ 0 & 1 & 2 \end{pmatrix} \leftrightarrow \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 2 \end{pmatrix} \Rightarrow \mathbf{A} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad (4.5)$$

From (4.3) and (4.5)

$$AP = \|\mathbf{A} - \mathbf{P}\| = 2\sqrt{5} = d(\text{say}) \quad (4.6)$$

The direction vector of  $AP$  is

$$\mathbf{m} = \mathbf{A} - \mathbf{P} = 2 \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (4.7)$$

$$\Rightarrow \|\mathbf{m}\| = 2\sqrt{5} \quad (4.8)$$

Since the direction of  $AP$  is the same as  $AC$ ,

$$\begin{aligned} \mathbf{C} &= \mathbf{P} - d \frac{\mathbf{m}}{\|\mathbf{m}\|} \\ &= -\begin{pmatrix} 1 \\ 2 \end{pmatrix} - 2 \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ -6 \end{pmatrix} \end{aligned} \quad (4.9)$$

Let  $\mathbf{n} \perp \mathbf{m}$ . Then

$$\mathbf{n} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} \quad (4.10)$$

and

$$\begin{aligned} \mathbf{B}, \mathbf{D} &= \mathbf{P} \pm d \frac{\mathbf{n}}{\|\mathbf{n}\|} \\ &= -\begin{pmatrix} 1 \\ 2 \end{pmatrix} \pm 2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} = \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 4 \end{pmatrix} \end{aligned} \quad (4.11)$$

- 5 Let  $k$  be an integer such that the triangle with vertices

$$\begin{pmatrix} k \\ -3k \end{pmatrix}, \begin{pmatrix} 5 \\ k \end{pmatrix}, \begin{pmatrix} -k \\ 2 \end{pmatrix} \quad (5.1)$$

has area 28. Find the orthocentre of this triangle.

- 6 If an equilateral triangle, having centroid at the origin, has a side along the line

$$(1 \ 1)\mathbf{x} = 2, \quad (6.1)$$

then find the area of this triangle.

- 7 Find the locus of the point of intersection of the straight lines

$$(t \ -2)\mathbf{x} - 3t = 0 \quad (7.1)$$

$$(1 \ -2t)\mathbf{x} + 3 = 0 \quad (7.2)$$

- 8 A square, of each side 2, lies above the  $x$ -axis

and has one vertex at the origin. If one of the sides passing through the origin makes an angle  $30^\circ$  with the positive direction of the  $x$ -axis, then find the sum of the  $x$ -coordinates of the vertices of the square.

- 9 Find the locus of the point of intersection of the lines

$$(\sqrt{2} \ -1)\mathbf{x} + 4\sqrt{2}k = 0 \quad (9.1)$$

$$(\sqrt{2}k \ k)\mathbf{x} - 4\sqrt{2} = 0 \quad (9.2)$$

- 10 The sides of a rhombus  $ABCD$  are parallel to the lines

$$(1 \ -1)\mathbf{x} + 2 = 0 \quad (10.1)$$

$$(7 \ -1)\mathbf{x} + 3 = 0. \quad (10.2)$$

If the diagonals of the rhombus intersect at

$$\mathbf{P} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (10.3)$$

and the vertex  $\mathbf{A}$  (different) from the origin is on the  $y$ -axis, then find the ordinate of  $A$ .