

# CONTENTS

<b>1</b>	<b>The Right Angled Triangle</b>	<b>1</b>
1.1	Sum of Angles . . . . .	1
1.2	Budhayana Theorem . . . . .	2
<b>2</b>	<b>Medians of a Triangle</b>	<b>2</b>
2.1	Area of a Triangle . . . . .	2
2.2	Median . . . . .	3
2.3	Similar Triangles . . . . .	5
<b>3</b>	<b>Angle and Perpendicular Bisectors</b>	<b>7</b>
3.1	Angle Bisectors . . . . .	7
3.2	Congruent Triangles . . . . .	7
3.3	Perpendicular Bisectors . . . . .	7
3.4	Perpendiculars from Vertex to Opposite Side . . . . .	8
<b>4</b>	<b>Circle</b>	<b>8</b>
4.1	Chord of a Circle . . . . .	8
4.2	Chords of a circle . . . . .	9
<b>5</b>	<b>Area of a Circle</b>	<b>11</b>
5.1	The Regular Polygon . . . . .	11

**Abstract**—The focus of this text is on investigating the properties of triangles and circles through trigonometry. Several proofs in geometry require a lot of constructions, which may not be easy for all the students in a class. This book provides an alternative approach to geometry which may be useful to those having difficulties with traditional methods. In the process, almost all the basic concepts of trigonometry are covered. Instead of teaching geometry and trigonometry as separate subjects in high school, this textbook shows how to develop a holistic approach for teaching math.

## 1 THE RIGHT ANGLED TRIANGLE

**Definition 1.1.** A right angled triangle looks like Fig. 1.0. with angles  $\angle A, \angle B$  and  $\angle C$  and sides  $a, b$  and  $c$ . The unique feature of this triangle is  $\angle C$  which is defined to be  $90^\circ$ .

**Definition 1.2.** For simplicity, let the greek letter  $\theta = \angle B$ . We have the following definitions.

$$\begin{aligned} \sin \theta &= \frac{a}{c} & \cos \theta &= \frac{b}{c} \\ \tan \theta &= \frac{a}{b} & \cot \theta &= \frac{b}{a} \\ \csc \theta &= \frac{c}{a} & \sec \theta &= \frac{c}{b} \end{aligned} \quad (1.1)$$




Fig. 1.0: Right Angled Triangle




Fig. 1.0: Sum of angles of a triangle

### 1.1 Sum of Angles

**Definition 1.3.** In Fig. 1.0, the sum of all the angles on the top or bottom side of the straight line  $XY$  is  $180^\circ$ .

**Definition 1.4.** In Fig. 1.0, the straight line making an angle of  $90^\circ$  to the side  $AC$  is said to be parallel to the side  $BC$ . Note there is an angle at  $A$  that is equal to  $\theta$ . This is one property of parallel lines. Thus,  $\angle YAZ = 90^\circ$ .

**Problem 1.1.** Show that  $\angle VAZ = 90^\circ - \theta$

**Proof.** Considering the line  $XAZ$ ,

$$\theta + 90^\circ + \angle VAZ = 180^\circ \quad (1.2)$$

$$\Rightarrow \angle VAZ = 90^\circ - \theta \quad (1.3)$$

**Problem 1.2.** Show that  $\angle BAC = 90^\circ - \theta$ .

*Proof.* Consider the line  $VAB$  and use the approach in the previous problem. Note that this implies that  $\angle VAZ = \angle BAC$ . Such angles are known as vertically opposite angles.

**Problem 1.3.** Sum of the angles of a triangle is equal to  $180^\circ$

## 1.2 Budhayana Theorem




Fig. 1.3: Budhayana Theorem

**Problem 1.4.** Using Fig. 1.0, show that

$$\cos \theta = \sin(90^\circ - \theta) \quad (1.4)$$

*Proof.* From Problem 1.2 and (1.1)

$$\cos(90^\circ - \theta) = \frac{b}{c} = \sin \theta \quad (1.5)$$

**Problem 1.5.** Using Fig. 1.3, show that

$$c = a \cos \theta + b \sin \theta \quad (1.6)$$

*Proof.* We observe that

$$BD = a \cos \theta \quad (1.7)$$

$$AD = b \cos(90 - \theta) = b \sin \theta \quad (\text{From } (1.2)) \quad (1.8)$$

Thus,

$$BD + AD = c = a \cos \theta + b \sin \theta \quad (1.9)$$

**Problem 1.6.** From (1.6), show that

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (1.10)$$

*Proof.* Dividing both sides of (1.6) by  $c$ ,

$$1 = \frac{a}{c} \cos \theta + \frac{b}{c} \sin \theta \quad (1.11)$$

$$\Rightarrow \sin^2 \theta + \cos^2 \theta = 1 \quad (\text{from } (1.1)) \quad (1.12)$$

**Problem 1.7.** Using (1.6), show that

$$c^2 = a^2 + b^2 \quad (1.13)$$

(1.13) is known as the Budhayana theorem. It is also known as the Pythagoras theorem.

*Proof.* From (1.6),

$$c = a \frac{a}{c} + b \frac{b}{c} \quad (\text{from } (1.1)) \quad (1.14)$$

$$\Rightarrow c^2 = a^2 + b^2 \quad (1.15)$$

## 2 MEDIANS OF A TRIANGLE

### 2.1 Area of a Triangle

**Definition 2.1.** The area of the rectangle ACBD shown in Fig. 2.0 is defined as  $ab$ . Note that all the angles in the rectangles are  $90^\circ$




Fig. 2.0: Area of a Right Triangle

**Definition 2.2.** The area of the two triangles constituting the rectangle is the same.

**Definition 2.3.** The area of the rectangle is the sum of the areas of the two triangles inside.

**Problem 2.1.** Show that the area of  $\triangle ABC$  is  $\frac{ab}{2}$




Fig. 2.1: Area of a Triangle

*Proof.* From (2.3),

$$ar(ABCD) = ar(ACB) + ar(ADB) \quad (2.1)$$

Also from (2.2),

$$ar(ACB) = ar(ADB) \quad (2.2)$$

From (2.1) and (2.2),

$$2ar(ACB) = ar(ABCD) = ab \text{ (from (2.0))} \quad (2.3)$$

$$\Rightarrow ar(ACB) = \frac{ab}{2} \quad (2.4)$$

**Problem 2.2.** Show that the area of  $\triangle ABC$  in Fig. 2.1 is  $\frac{1}{2}ah$ .

*Proof.* In Fig. 2.1,

$$ar(\triangle ADC) = \frac{1}{2}hy \quad (2.5)$$

$$ar(\triangle ADB) = \frac{1}{2}hx \quad (2.6)$$

Thus,

$$ar(\triangle ABC) = ar(\triangle ADC) + ar(\triangle ADB) \quad (2.7)$$

$$= \frac{1}{2}hy + \frac{1}{2}hx = \frac{1}{2}h(x+y) \quad (2.8)$$

$$= \frac{1}{2}ah \quad (2.9)$$

**Problem 2.3.** Show that the area of  $\triangle ABC$  in Fig. 2.1 is  $\frac{1}{2}ab \sin C$ .

*Proof.* We have

$$ar(\triangle ABC) = \frac{1}{2}ah = \frac{1}{2}ab \sin C \quad (\because h = b \sin C). \quad (2.10)$$

**Problem 2.4.** Show that

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad (2.11)$$

*Proof.* Fig. 2.1 can be suitably modified to obtain

$$ar(\triangle ABC) = \frac{1}{2}ab \sin C = \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B \quad (2.12)$$

Dividing the above by  $abc$ , we obtain

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad (2.13)$$

This is known as the sine formula.

## 2.2 Median

**Definition 2.4.** The line  $AD$  in Fig. 2.4 that divides the side  $a$  in two equal halves is known as the median.




Fig. 2.4: Median of a Triangle

**Problem 2.5.** Show that the median  $AD$  in Fig. 2.4 divides  $\triangle ABC$  into triangles  $ADB$  and  $ADC$  that have equal area.

*Proof.* We have

$$ar(\triangle ADB) = \frac{1}{2} \frac{a}{2} c \sin B = \frac{1}{4} ac \sin B \quad (2.14)$$

$$ar(\triangle ADC) = \frac{1}{2} \frac{a}{2} b \sin C = \frac{1}{4} ab \sin C \quad (2.15)$$

Using the sine formula,  $b \sin C = c \sin B$ ,

$$ar(\triangle ADB) = ar(\triangle ADC) \quad (2.16)$$

**Problem 2.6.**  $BE$  and  $CF$  are the medians in Fig. 2.6. Show that

$$ar(\triangle BFC) = ar(\triangle BEC) \quad (2.17)$$

*Proof.* Since  $BE$  and  $CF$  are the medians,

$$ar(\Delta BFC) = \frac{1}{2}ar(\Delta ABC) \quad (2.18)$$

$$ar(\Delta BEC) = \frac{1}{2}ar(\Delta ABC) \quad (2.19)$$

From the above, we infer that

$$ar(\Delta BFC) = ar(\Delta BEC) \quad (2.20)$$




Fig. 2.6:  $O$  is the Intersection of Two Medians

**Problem 2.7.** The medians  $BE$  and  $CF$  in Fig. 2.6 meet at point  $O$ . Show that

$$\frac{OB}{OE} = \frac{OC}{OF} \quad (2.21)$$

*Proof.* From Problem 2.6,

$$ar(\Delta BFC) = ar(\Delta BEC) \quad (2.22)$$

$$\begin{aligned} \Rightarrow ar(\Delta BOF) + ar(\Delta BOC) \\ = ar(\Delta BOC) + ar(\Delta COE) \end{aligned} \quad (2.23)$$

resulting in

$$ar(\Delta BOF) = ar(\Delta COE) \quad (2.24)$$

Using the sine formula for area of a triangle, the above equation can be expressed as

$$\frac{1}{2}OB \cdot OF \sin \theta = \frac{1}{2}OC \cdot OE \sin \theta \quad (2.25)$$

$$\Rightarrow \frac{OB}{OE} = \frac{OC}{OF} \quad (2.26)$$

**Definition 2.5.** We know that the median of a triangle divides it into two triangles with equal area.

Using this result along with the sine formula for the area of a triangle in Fig. 2.7,

$$\frac{1}{2}AD \sin \theta = \frac{1}{2}AD \sin(180^\circ - \theta) \quad (2.27)$$

$$\Rightarrow \sin \theta = \sin(180^\circ - \theta). \quad (2.28)$$

Note that our geometric definition of  $\sin \theta$  holds only for  $\theta < 90^\circ$ . (2.28) allows us to extend this definition for  $\angle ADC > 90^\circ$ .




Fig. 2.7:  $\sin \theta = \sin(180^\circ - \theta)$

*Proof.* Using the sine formula and (2.8), areas of some triangles in Fig. 2.9 are listed in the following table

Triangle	Area
OFE	$\frac{1}{2}pq \sin \theta$
BOF	$\frac{k}{2}pq \sin \theta$
COE	$\frac{k}{2}pq \sin \theta$
BOC	$\frac{k^2}{2}pq \sin \theta$
BOC	$\frac{1}{2}ar(\Delta ABC)$

where we have used the fact that

$$\sin \angle BOC = \sin(180^\circ - \theta) = \sin \theta \quad (2.33)$$

Since  $BE$  is the median

$$\begin{aligned} ar(\Delta BEC) &= \frac{1}{2}ar(\Delta ABC) \\ &= ar(\Delta BOC) + ar(\Delta COE) \\ ar(\Delta BEA) &= \frac{1}{2}ar(\Delta ABC) \\ &= ar(\Delta AFE) + ar(\Delta BOF) + ar(\Delta FOE) \end{aligned} \quad (2.34)$$

**Problem 2.8.** In Fig. 2.9, show that

$$ar(\Delta AFE) = \frac{1}{4} ar(\Delta ABC) \quad (2.29)$$

*Proof.* We have

$$ar(\Delta AFE) = \frac{1}{2} \frac{b}{2} \frac{c}{2} \sin A = \frac{1}{4} \cdot \frac{1}{2} bc \sin A \quad (2.30)$$

$$= \frac{1}{4} ar(\Delta ABC) \quad (2.31)$$

**Problem 2.9.** Using Fig. 2.9, show that

$$\frac{OB}{OE} = \frac{OC}{OF} = 2 \quad (2.32)$$




Fig. 2.9:  $O$  divides medians in the ratio 2 : 1

Substituting the respective values from the above table,

$$\begin{aligned} \frac{1}{2} ar(\Delta ABC) &= \frac{k^2}{2} pq \sin \theta + \frac{k}{2} pq \sin \theta \\ \frac{1}{2} ar(\Delta ABC) &= \frac{k}{2} pq \sin \theta + \frac{1}{2} pq \sin \theta + \frac{1}{4} ar(\Delta ABC) \end{aligned} \quad (2.35)$$

Simplifying the above,

$$k(k+1) = 2(k+1) \quad (2.36)$$

Since  $k \neq -1$ ,  $k = 2$  and the proof is complete.

### 2.3 Similar Triangles

**Problem 2.10.** In Fig. 2.10, show that

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad (2.37)$$




Fig. 2.10: The cosine formula

*Proof.* From the figure, the first of the following equations

$$a = b \cos C + c \cos B \quad (2.38)$$

$$b = c \cos A + a \cos C \quad (2.39)$$

$$c = b \cos A + a \cos B \quad (2.40)$$

is obvious and the other two can be similarly obtained. The above equations can be expressed in matrix form as

$$\begin{pmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{pmatrix} \begin{pmatrix} \cos A \\ \cos B \\ \cos C \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \quad (2.41)$$

Using the properties of determinants,

$$\cos A = \frac{\begin{vmatrix} a & c & b \\ b & 0 & a \\ c & a & 0 \end{vmatrix}}{\begin{vmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{vmatrix}} = \frac{ab^2 + ac^2 - a^3}{abc + abc} = \frac{b^2 + c^2 - a^2}{2abc} \quad (2.42)$$

**Problem 2.11.** In Fig. 2.11, show that  $EF = \frac{a}{2}$ .

*Proof.* Using the cosine formula for  $\Delta AEF$ ,

$$EF^2 = \left(\frac{b}{2}\right)^2 + \left(\frac{c}{2}\right)^2 - 2\left(\frac{b}{2}\right)\left(\frac{c}{2}\right)\cos A \quad (2.43)$$

$$= \frac{b^2 + c^2 - 2bc \cos A}{4} \quad (2.44)$$

$$= \frac{a^2}{4} \quad (2.45)$$

$$\Rightarrow EF = \frac{a}{2} \quad (2.46)$$




Fig. 2.11: Similar Triangles

**Definition 2.6.** The ratio of sides of triangles  $AEF$  and  $ABC$  is the same. Such triangles are known as similar triangles.

**Problem 2.12.** Show that similar triangles have the same angles.

*Proof.* Use cosine formula and the proof is trivial.

**Problem 2.13.** Show that in Fig. 2.11,  $EF \parallel BC$ .

*Proof.* Since  $\triangle AEF \sim \triangle ABC$ ,  $\angle AEF = \angle ACB$ . Hence the line  $EF \parallel BC$ .

**Problem 2.14.** In Fig. 2.14, the line  $AO$  cuts  $EF$  at  $G$  and is extended to meet the side  $BC$  at  $D$ . Show that

$$\frac{OA}{OD} = 2. \quad (2.47)$$




Fig. 2.14: Similar Triangles and Median

*Proof.* Since  $EF \parallel BC$ ,

$$\triangle AGE \sim \triangle ADC \Rightarrow AG = GD = \frac{AD}{2} \quad (2.48)$$

Also,

$$\triangle OGE \sim \triangle ODB \Rightarrow OD = 2r \quad (2.49)$$

Thus, we have the following relations

$$GD = OG + OD = r + 2r = 3r \quad (2.50)$$

$$AG = GD = 3r \quad (2.51)$$

$$OA = OG + AG = r + 3r = 4r \quad (2.52)$$

Hence

$$\frac{OA}{OD} = \frac{4r}{2r} = 2 \quad (2.53)$$

**Problem 2.15.** In Fig. 2.15,  $BE$  is a median of  $\triangle ABC$  and  $\frac{OA}{OD} = 2$ . Show that  $AD$  is also a median.




Fig. 2.15: Medians meet at a point

*Proof.* Using the cosine formula for  $\triangle ODE$  and  $\triangle OAB$ ,

$$DE^2 = p^2 + t^2 - 2pq \cos \theta \quad (2.54)$$

$$c^2 = (2)p^2 + (2t)^2 - 2(2p)(2q) \cos \theta \quad (2.55)$$

$$\Rightarrow DE = \frac{c}{2}. \quad (2.56)$$

Now using the cosine formula in  $\triangle CDE$  and  $\triangle ABC$ ,

$$\left(\frac{c}{2}\right)^2 = \left(\frac{b}{2}\right)^2 + CD^2 - 2CD\left(\frac{b}{2}\right) \cos C \quad (2.57)$$

$$c^2 = b^2 + a^2 - 2ab \cos C, \quad (2.58)$$

which can be simplified to obtain

$$a^2 - 2ab \cos C = (2CD)^2 - 2(2CD)b \cos C \quad (2.59)$$

$$\begin{aligned} &\Rightarrow (2CD)^2 - 2(2CD)b \cos C + (b \cos C)^2 \\ &= a^2 - 2ab \cos C + (b \cos C)^2 \end{aligned} \quad (2.60)$$

$$\Rightarrow (2CD - b \cos C)^2 = (a - b \cos C)^2 \quad (2.61)$$

$$\Rightarrow 2CD - b \cos C = a - 2 \cos C \quad (2.62)$$

$$\Rightarrow CD = \frac{a}{2} \quad (2.63)$$

Thus,  $AD$  is a median of  $\triangle ABC$ .

**Conclusion:** The medians of a triangle meet at a point.

### 3 ANGLE AND PERPENDICULAR BISECTORS

#### 3.1 Angle Bisectors




Fig. 3.0: Angle bisectors meet at a point

**Definition 3.1.** In Fig. 3.0,  $OB$  divides the  $\angle B$  into half, i.e.

$$\angle OBC = \angle OBA \quad (3.1)$$

$OB$  is known as an angle bisector.

$OB$  and  $OC$  are angle bisectors of angles  $B$  and  $C$ .  $OA$  is joined and  $OD, OF$  and  $OE$  are perpendiculars to sides  $a, b$  and  $c$ .

**Problem 3.1.** Show that  $OD = OE = OF$ .

*Proof.* In  $\triangle ODC$  and  $OEC$ ,

$$OD = OC \sin \frac{C}{2} \quad (3.2)$$

$$OE = OC \sin \frac{C}{2} \quad (3.3)$$

$$\Rightarrow OD = OE. \quad (3.4)$$

Similarly,

$$OD = OF. \quad (3.5)$$

**Problem 3.2.** Show that  $OA$  is the angle bisector of  $\angle A$

*Proof.* In  $\triangle OFA$  and  $OEA$ ,

$$OF = OE \quad (3.6)$$

$$\Rightarrow OA \sin \angle OAF = OA \sin \angle OAE \quad (3.7)$$

$$\Rightarrow \sin \angle OAF = \sin \angle OAE \quad (3.8)$$

$$\Rightarrow \angle OAF = \angle OAE \quad (3.9)$$

which proves that  $OA$  bisects  $\angle A$ . **Conclusion:** The angle bisectors of a triangle meet at a point.

#### 3.2 Congruent Triangles

**Problem 3.3.** Show that in  $\triangle ODC$  and  $OEC$ , corresponding sides and angles are equal.

**Definition 3.2.** Note that  $\triangle ODC$  and  $OEC$  are known as congruent triangles. To show that two triangles are congruent, it is sufficient to show that some angles and sides are equal.

**Problem 3.4.** SSS: Show that if the corresponding sides of three triangles are equal, the triangles are congruent.

**Problem 3.5.** ASA: Show that if two angles and any one side are equal in corresponding triangles, the triangles are congruent.

**Problem 3.6.** SAS: Show that if two sides and the angle between them are equal in corresponding triangles, the triangles are congruent.

**Problem 3.7.** RHS: For two right angled triangles, if the hypotenuse and one of the sides are equal, show that the triangles are congruent.

#### 3.3 Perpendicular Bisectors

**Definition 3.3.** In Fig. 3.8,  $OD \perp BC$  and  $BD = DC$ .  $OD$  is defined as the perpendicular bisector of  $BC$ .

**Problem 3.8.** In Fig. 3.8, show that  $OA = OB = OC$ .

*Proof.* In  $\triangle ODB$  and  $ODC$ , using Budhayana's theorem,

$$\begin{aligned} OB^2 &= OD^2 + BD^2 \\ OC^2 &= OD^2 + DC^2 \end{aligned} \quad (3.10)$$

Since  $BD = DC = \frac{a}{2}$ ,  $OB = OC$ . Similarly, it can be shown that  $OA = OC$ . Thus,  $OA = OB = OC$ .

**Definition 3.4.** In  $\triangle AOB$ ,  $OA = OB$ . Such a triangle is known as an isosceles triangle.




Fig. 3.8: Perpendicular bisectors meet at a point

**Problem 3.9.** Show that  $AF = BF$ .

*Proof.* Trivial using Budhayana's theorem. This shows that  $OF$  is a perpendicular bisector of  $AB$ .  
*Conclusion:* The perpendicular bisectors of a triangle meet at a point.

### 3.4 Perpendiculars from Vertex to Opposite Side

In Fig. 3.10,  $AD \perp BC$  and  $BE \perp AC$ .  $CF$  passes through  $O$  and meets  $AB$  at  $F$ .

**Problem 3.10.** Show that

$$OE = c \cos A \cot C \quad (3.11)$$

*Proof.* In  $\Delta$  s  $AEB$  and  $AEO$ ,

$$AE = c \cos A \quad (3.12)$$

$$OE = AE \tan(90^\circ - C) (\because ADC \text{ is right angled}) \quad (3.13)$$

$$= AE \cot C \quad (3.14)$$

From both the above, we get the desired result.




Fig. 3.10: Perpendiculars from vertex to opposite side meet at a point

**Problem 3.11.** Show that  $\alpha = A$ .

*Proof.* In  $\Delta OEC$ ,

$$CE = a \cos C (\because BEC \text{ is right angled}) \quad (3.15)$$

Hence,

$$\begin{aligned} \tan \alpha &= \frac{CE}{OE} \\ &= \frac{a \cos C}{c \cos A \cot C} \\ &= \frac{a \cos C \sin C}{c \cos A \cos C} \\ &= \frac{a \sin C}{c \cos A} \\ &= \frac{c \sin A}{c \cos A} \left( \because \frac{a}{\sin A} = \frac{c}{\sin C} \right) \\ &= \tan A \end{aligned} \quad (3.16)$$

$$\Rightarrow \alpha = A$$

**Problem 3.12.** Show that  $CF \perp AB$

*Proof.* Consider triangle OFB and the result of the previous problem.  $\because$  the sum of the angles of a triangle is  $180^\circ$ ,  $\angle CFB = 90^\circ$ . *Conclusion:* The perpendiculars from the vertex of a triangle to the opposite side meet at a point.

### 4 CIRCLE

#### 4.1 Chord of a Circle

**Definition 4.1.** Fig. 4.0 represents a circle. The points in the circle are at a distance  $r$  from the centre  $O$ .  $r$  is known as the radius.







Fig. 4.0: Circle Definitions

Fig. 4.1: Angle subtended by chord  $AB$  at the centre  $O$  is twice the angle subtended at  $P$ .

#### 4.2 Chords of a circle

**Definition 4.2.** In Fig. 4.0,  $A$  and  $B$  are points on the circle. The line  $AB$  is known as a chord of the circle.

**Problem 4.1.** In Fig. 4.1 Show that  $\angle OAB = 2\angle APB$ .

*Proof.* In Fig. 4.1, the triangles  $OPA$  and  $OPB$  are isosceles. Hence,

$$\angle OPB = \angle OBP = \theta_1 \quad (4.1)$$

$$\angle OPA = \angle OAP = \theta_2 \quad (4.2)$$

Also,  $\alpha$  and  $\beta$  are exterior angles corresponding to the triangle  $OPB$  and  $OPA$  respectively. Hence

$$\alpha = 2\theta_1 \quad (4.3)$$

$$\beta = 2\theta_2 \quad (4.4)$$

Thus,

$$\angle AOB = \alpha + \beta \quad (4.5)$$

$$= \theta_1 + \theta_2 \quad (4.6)$$

$$= \angle APB \quad (4.7)$$

**Definition 4.3.** The diameter of a circle is the chord that divides the circle into two equal parts. In Fig. 4.2,  $AB$  is the diameter and passes through the centre  $O$

**Problem 4.2.** In Fig. 4.2, show that  $\angle APB = 90^\circ$ .





Fig. 4.2: Diameter of a circle.

**Problem 4.3.** In Fig. 4.3, show that

$$\begin{aligned} \angle ABD &= \angle ACD \\ \angle CAB &= \angle CDB \end{aligned} \quad (4.8)$$

Fig. 4.3:  $PA.PB = PC.PD$ 

*Proof.* Use Problem 4.1.

**Problem 4.4.** In Fig. 4.3, show that the triangles  $PAB$  and  $PBD$  are similar

*Proof.* Trivial using previous problem

**Problem 4.5.** In Fig. 4.3, show that

$$PA.PB = PC.PD \quad (4.9)$$

*Proof.* Since triangles  $PAC$  and  $PBD$  are similar,

$$\frac{PA}{PD} = \frac{PC}{PB} \quad (4.10)$$

$$\Rightarrow PA.PB = PC.PD \quad (4.11)$$

**Definition 4.4.** The line  $PX$  in Fig. 4.6 touches the circle at exactly one point  $P$ . It is known as the tangent to the circle.

**Problem 4.6.**  $OP$  is the perpendicular to the line  $PX$  as shown in the Fig. 4.6. Show that  $OP$  is the shortest distance between the point  $O$  and the line  $PX$ .

*Proof.* Let  $P_1$  be a point on the line  $PX$ . Then  $OPP_1$  is a right angled triangle. Using Budhayana's theorem,

$$\begin{aligned} OP_1^2 &= OP^2 + PP_1^2 \\ \Rightarrow OP_1 &> OP \end{aligned} \quad (4.12)$$





Fig. 4.6: Tangent to a Circle.

Fig. 4.6: Shortest distance from  $O$  to line  $PX$ 

Thus,  $OP$  is the shortest distance between  $O$  and line  $PX$ .

**Problem 4.7.** Show that  $\angle OPX = 90^\circ$

*Proof.* In Fig. 4.6, we can see that  $OP$  is the radius of the circle and the length of all line segments from  $O$  to the line  $PX > r$ . Using the result of the previous problem, it is obvious that  $OP \perp PX$ .

**Problem 4.8.** In Fig. 4.8 show that

$$\angle PCA = \angle PBC \quad (4.13)$$


$O$  is the centre of the circle and  $PC$  is the tangent.

*Proof.* For convenience, greek letters are used for representing certain angles. Since  $\triangle OAC$  is isosceles,

$$2\alpha + 2(\beta - \phi) = 180^\circ \quad (4.14)$$

$$\Rightarrow \alpha + (\beta - \phi) = 90^\circ \quad (4.15)$$

$$\Rightarrow \alpha + \beta = 90^\circ + \phi \quad (4.16)$$

Fig. 4.8:  $PA.PB = PC^2$ .

Since  $\theta$  is an exterior angle for the  $\triangle ABC$ ,

$$\theta = \alpha + \beta \quad (4.17)$$

From both the above equations

$$\theta = 90^\circ + \phi \quad (4.18)$$

Since PC is the tangent,

$$\angle PCB = 90^\circ + \phi = \theta \quad (4.19)$$

Considering the sum of angles in  $\triangle PAC$   $\triangle PBC$ ,

$$\angle P + \theta + \angle PCA = 180^\circ \quad (4.20)$$

$$\angle P + \theta + \alpha = 180^\circ \quad (4.21)$$

Hence,

$$\angle PCA = \alpha \quad (4.22)$$

**Problem 4.9.** In Fig. 4.8, show that the triangles PAC and PBC are similar.

*Proof.* From the previous problem, it is obvious that corresponding angles of both triangles are equal. Hence they are similar.

**Problem 4.10.** Show that  $PA.PB = PC^2$


*Proof.* Since  $\triangle PAC \sim \triangle PBC$ , their sides are in the same ratio. Hence,

$$\frac{PA}{PC} = \frac{PC}{PB} \quad (4.23)$$

$$\Rightarrow PA.PB = PC^2 \quad (4.24)$$

**Problem 4.11.** In Fig. 4.11, show that

$$PA.PB = PC.PD \quad (4.25)$$

Fig. 4.11:  $PA.PB = PC^2$ .

*Proof.* Draw a tangent and use the previous problem.

## 5 AREA OF A CIRCLE

### 5.1 The Regular Polygon

**Definition 5.1.** In Fig. 5.0, 6 congruent triangles are arranged in a circular fashion. Such a figure is known as a regular hexagon. In general,  $n$  number of triangles can be arranged to form a regular polygon.




Fig. 5.0: Polygon Definition

**Definition 5.2.** The angle formed by each of the congruent triangles at the centre of a regular polygon of  $n$  sides is  $\frac{360^\circ}{n}$ .

**Problem 5.1.** Show that the area of a regular polygon is given by

$$\frac{n}{2} r^2 \sin \frac{360^\circ}{n} \quad (5.1)$$




Fig. 5.1: Polygon Area

*Proof.* The triangle that forms the polygon of  $n$  sides is given in Fig. 5.1. Thus,

$$\begin{aligned} \text{ar}(\text{polygon}) &= n \text{ar}(\triangle ABC) \\ &= \frac{n}{2} r^2 \sin \frac{360^\circ}{n} \end{aligned} \quad (5.2)$$

**Problem 5.2.** Using Fig. 5.2, show that

$$\frac{n}{2} r^2 \sin \frac{360^\circ}{n} < \text{area of circle} < n r^2 \tan \frac{180^\circ}{n} \quad (5.3)$$

The portion of the circle visible in Fig. 5.2 is defined to be a sector of the circle.

*Proof.* Note that the circle is squeezed between the inner and outer regular polygons. As we can see from Fig. 5.2, the area of the circle should be in between the areas of the inner and outer polygons. Since

$$\text{ar}(\triangle OAB) = \frac{1}{2} r^2 \sin \frac{360^\circ}{n} \quad (5.4)$$

$$\text{ar}(\triangle OPQ) = 2 \times \frac{1}{2} \times r \tan \frac{360/n}{2} \times r \quad (5.5)$$

$$= r^2 \tan \frac{180^\circ}{n}, \quad (5.6)$$

we obtain (5.3).




Fig. 5.2: Circle Area in between Area of Two Polygons

**Problem 5.3.** Using Fig. 5.3, show that

$$\sin \theta_1 = \sin (\theta_1 + \theta_2) \cos \theta_2 - \cos (\theta_1 + \theta_2) \sin \theta_2 \quad (5.7)$$




Fig. 5.3:  $\sin 2\theta = 2 \sin \theta \cos \theta$

*Proof.* The following equations can be obtained from the figure using the formula for the area of a triangle

$$ar(\Delta ABC) = \frac{1}{2}ac \sin(\theta_1 + \theta_2) \quad (5.8)$$

$$= ar(\Delta BDC) + ar(\Delta ADB) \quad (5.9)$$

$$= \frac{1}{2}cl \sin \theta_1 + \frac{1}{2}al \sin \theta_2 \quad (5.10)$$

$$= \frac{1}{2}ac \sin \theta_1 \sec \theta_2 + \frac{1}{2}a^2 \tan \theta_2 \quad (5.11)$$

( $\because l = a \sec \theta_2$ ). From the above,

$$\Rightarrow \sin(\theta_1 + \theta_2) = \sin \theta_1 \sec \theta_2 + \frac{a}{c} \tan \theta_2 \quad (5.12)$$

$$\Rightarrow \sin(\theta_1 + \theta_2) = \sin \theta_1 \sec \theta_2 + \cos(\theta_1 + \theta_2) \tan \theta_2 \quad (5.13)$$

Multiplying both sides by  $\cos \theta_2$ ,

$$\Rightarrow \sin(\theta_1 + \theta_2) \cos \theta_2 = \sin \theta_1 + \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.14)$$

resulting in

$$\Rightarrow \sin \theta_1 = \sin(\theta_1 + \theta_2) \cos \theta_2 - \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.15)$$

**Problem 5.4.** Prove the following identities

1)

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta. \quad (5.16)$$

2)

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta. \quad (5.17)$$

*Proof.* In (5.7), let

$$\begin{aligned} \theta_1 + \theta_2 &= \alpha \\ \theta_2 &= \beta \end{aligned} \quad (5.18)$$

This gives (5.16). In (5.16), replace  $\alpha$  by  $90^\circ - \alpha$ . This results in

$$\begin{aligned} \sin(90^\circ - \alpha - \beta) \\ &= \sin(90^\circ - \alpha) \cos \beta - \cos(90^\circ - \alpha) \sin \beta \\ \Rightarrow \cos(\alpha + \beta) &= \cos \alpha \cos \beta - \sin \alpha \sin \beta \end{aligned} \quad (5.19)$$

**Problem 5.5.** Using (5.7) and (5.17), show that

$$\sin(\theta_1 + \theta_2) = \sin \theta_1 \cos \theta_2 + \cos \theta_1 \sin \theta_2 \quad (5.20)$$

$$\cos(\theta_1 - \theta_2) = \cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2 \quad (5.21)$$

*Proof.* From (5.7),

$$\sin(\theta_1 + \theta_2) \cos \theta_2 = \sin \theta_1 + \cos(\theta_1 + \theta_2) \sin \theta_2 \quad (5.22)$$

Using (5.17) in the above,

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \sin \theta_1 + (\cos \theta_1 \cos \theta_2 \\ &\quad - \sin \theta_1 \sin \theta_2) \sin \theta_2 \end{aligned} \quad (5.23)$$

which can be expressed as

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \sin \theta_1 + \cos \theta_1 \cos \theta_2 \sin \theta_2 \\ &\quad - \sin \theta_1 \sin^2 \theta_2 \end{aligned} \quad (5.24)$$

Since

$$\sin^2 \theta_2 = 1 - \cos^2 \theta_2, \quad (5.25)$$

we obtain

$$\begin{aligned} \sin(\theta_1 + \theta_2) \cos \theta_2 &= \cos \theta_1 \cos \theta_2 \sin \theta_2 \\ &\quad + \sin \theta_1 \cos^2 \theta_2 \end{aligned} \quad (5.26)$$

resulting in

$$\sin(\theta_1 + \theta_2) = \cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2 \quad (5.27)$$

after factoring out  $\cos \theta_2$ . Using a similar approach, (5.21) can also be proved.

**Problem 5.6.** Show that

$$\sin 2\theta = 2 \sin \theta \cos \theta \quad (5.28)$$

**Problem 5.7.** Show that

$$\cos^2 \frac{180^\circ}{n} < \frac{\text{area of circle}}{nr^2 \tan \frac{180^\circ}{n}} < 1 \quad (5.29)$$

*Proof.* From (5.3) and (5.28),

$$\begin{aligned} \frac{n}{2} r^2 \sin \frac{360^\circ}{n} &< \text{area of circle} \\ &< nr^2 \tan \frac{180^\circ}{n} \\ \Rightarrow nr^2 \sin \frac{180^\circ}{n} \cos \frac{180^\circ}{n} &< \text{area of circle} \\ &< nr^2 \tan \frac{180^\circ}{n} \end{aligned} \quad (5.30)$$

**Problem 5.8.** Show that if

$$\theta_1 < \theta_2, \sin \theta_1 < \sin \theta_2. \quad (5.31)$$

*Proof.* Trivial using the definition and choosing angles  $\theta_1$  and  $\theta_2$  appropriately in a right angled

triangle.

**Problem 5.9.** Show that if

$$\theta_1 < \theta_2, \cos \theta_1 > \cos \theta_2. \quad (5.32)$$

**Problem 5.10.** Show that

$$\sin 0^\circ = 0 \quad (5.33)$$

*Proof.* Follows from (5.32).

**Problem 5.11.** Show that

$$\cos 0^\circ = 1 \quad (5.34)$$

**Problem 5.12.** Show that for large values of  $n$

$$\cos^2 \frac{180^\circ}{n} = 1 \quad (5.35)$$

*Proof.* Follows from previous problem.

**Definition 5.3.** The previous result can be expressed as

$$\lim_{n \rightarrow \infty} \cos^2 \frac{180^\circ}{n} = 1 \quad (5.36)$$

**Problem 5.13.** Show that

$$\text{area of circle} = r^2 \lim_{n \rightarrow \infty} n \tan \frac{180^\circ}{n} \quad (5.37)$$

**Definition 5.4.**

$$\pi = \lim_{n \rightarrow \infty} n \tan \frac{180^\circ}{n} \quad (5.38)$$

Thus, the area of a circle is  $\pi r^2$ .

**Definition 5.5.** The radian is a unit of angle defined by

$$1 \text{ radian} = \frac{360^\circ}{2\pi} \quad (5.39)$$

**Problem 5.14.** Show that the circumference of a circle is  $2\pi r$ .

**Problem 5.15.** Show that the area of a sector with angle  $\theta$  in radians is  $\frac{1}{2}r^2\theta$ .

**Problem 5.16.** Show that

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \quad (5.40)$$