specific mutations and copy number changes, as well as the expression of all genes critically involved in homologous recombination to help us understand how best to treat those patients that have failed PARP inhibition. A better understanding of PARP inhibitor resistance will allow researchers and clinicians to exploit therapeutic liabilities engendered by these adaptive responses and develop rational combination strategies that specifically target or reverse these compensatory signaling pathways.

Acknowledgements

None.

Financial support and sponsorship

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- ■■ of outstanding interest
- Gonzalez-Martin A, Pothuri B, Vergote I, et al., PRIMA/ENGOT-OV26/GOG-3012 Investigators. Niraparib in patients with newly diagnosed advanced ovarian cancer. N Engl J Med 2019; doi: 10.1056/NEJMoa1910962. [Epub

ahead of print]
The PRIMA trial provides data for niraparib use for maintenance therapy in newly diagnosed ovarian cancer, irrespective of BRCA status.

- 2. Coleman RL, Fleming GF, Brady MF, et al. Veliparib with first-line chemother-
- apy and as maintenance therapy in ovarian cancer. N Engl J Med 2019; doi: 10.1056/NEJMoa1909707. [Epub ahead of print]

The VELIA trial provides data for combining veliparib with chemotherapy and then treating with veliparib maintenance monotherapy in newly diagnosed ovarian cancer.

- Ray-Coquard I, Pautier P, Pignata S, et al. Phase III PAOLA-1/ENGOT-0v25 trial: Olaparib plus bevacizumab (bev) as maintenance therapy in patients (pts) with newly diagnosed, advanced ovarian cancer (oc) treated with
- platinum-based chemotherapy (pch) plus bev. ESMO 2019; abstract. LBA2. The PAOLA-1 trial provides data for combining olaparib with bevacizumab for maintenance therapy following upfront treatment with carboplatin, paclitaxel and bevacizumab in newly diagnosed ovarian cancer.
- Sakai W, Swisher EM, Karlan BY, et al. Secondary mutations as a mechanism of cisplatin resistance in BRCA2-mutated cancers. Nature 2008; 451:1116–1120.
- Edwards SL, Brough R, Lord CJ, et al. Resistance to therapy caused by intragenic deletion in BRCA2. Nature 2008; 451:1111-1115.
- Konstantinopoulos PA, Ceccaldi R, Shapiro GI, et al. Homologous recombination deficiency: exploiting the fundamental vulnerability of ovarian cancer. Cancer Discov 2015; 5:1137-1154.
- Norquist B, Wurz KA, Pennil CC, et al. Secondary somatic mutations restoring BRCA1/2 predict chemotherapy resistance in hereditary ovarian carcinomas. J Clin Oncol 2011: 29:3008–3015.
- Patch AM, Christie EL, Etemadmoghadam D, et al. Whole-genome characterization of chemoresistant ovarian cancer. Nature 2015; 521:489–494.
- Barber LJ, Sandhu S, Chen L, et al. Secondary mutations in BRCA2 associated with clinical resistance to a PARP inhibitor. J Pathol 2013; 229:422-429.
- Goodall J, Mateo J, Yuan W, et al. Circulating cell-free DNA to guide prostate cancer treatment with PARP inhibition. Cancer Discov 2017; 7:1006-1017.
- Kondrashova O, Nguyen M, Shield-Artin K, et al., AOCS Study Group. Secondary somatic mutations restoring RAD51C and RAD51D associated with acquired resistance to the PARP inhibitor rucaparib in high-grade ovarian carcinoma. Cancer Discov 2017; 7:984–998.
- Quigley D, Alumkal JJ, Wyatt AW, et al. Analysis of circulating cell-free DNA identifies multiclonal heterogeneity of BRCA2 reversion mutations associated with resistance to PARP inhibitors. Cancer Discov 2017; 7:999–1005.

- Christie EL, Fereday S, Doig K, et al. Reversion of BRCA1/2 germline mutations detected in circulating tumor DNA from patients with high-grade serous ovarian cancer. J Clin Oncol 2017; 35:1274-1280.
- Weigelt B, Comino-Méndez I, de Bruijn I, et al. Diverse BRCA1 and BRCA2 reversion mutations in circulating cell-free DNA of therapy-resistant breast or ovarian cancer. Clin Cancer Res 2017; 23:6708–6720.
- Lin KK, Harrell MI, Oza AM, et al. BRCA reversion mutations in circulating tumor
 DNA predict primary and acquired resistance to the PARP inhibitor Rucaparib in high-grade ovarian carcinoma. Cancer Discov 2019; 9:210-219.

This study provides evidence for detecting BRCA reversion mutations using cfDNA in patients with somatic BRCA mutations.

- Williams RS, Glover JN. Structural consequences of a cancer-causing BRCA1-BRCT missense mutation. J Biol Chem 2003; 278:2630–2635.
- Williams RS, Chasman DI, Hau DD, et al. Detection of protein folding defects caused by BRCA1-BRCT truncation and missense mutations. J Biol Chem 2003; 278:53007 – 53016.
- Lee MS, Green R, Marsillac SM, et al. Comprehensive analysis of missense variations in the BRCT domain of BRCA1 by structural and functional assays. Cancer Res 2010; 70:4880–4890.
- Sy SM, Huen MS, Chen J. PALB2 is an integral component of the BRCA complex required for homologous recombination repair. Proc Natl Acad Sci USA 2009; 106:7155-7160.
- Scully R, Chen J, Plug A. Association of BRCA1 with Rad51 in mitotic and meiotic cells. Cell 1997; 88:265–275.
- Johnson N, Johnson SF, Yao W, et al. Stabilization of mutant BRCA1 protein confers PARP inhibitor and platinum resistance. Proc Natl Acad Sci U S A 2013; 110:17041–17046.
- Drost R, Dhillon KK, van der Gulden H, et al. BRCA1185delAG tumors may acquire therapy resistance through expression of RING-less BRCA1. J Clin Invest 2016; 126:2903–2918.
- 23. Wang Y, Bernhardy AJ, Cruz C, et al. The BRCA1-∆11q alternative splice isoform bypasses germline mutations and promotes therapeutic resistance to PARP inhibition and cisplatin. Cancer Res 2016; 76:2778-2790.
- Cao L, Xu X, Bunting SF, et al. A selective requirement for 53BP1 in the biological response to genomic instability induced by Brca1 deficiency. Mol Cell 2009: 35:534–541.
- 25. Escribano-Diaz C, Orthwein A, Fradet-Turcotte A, et al. A cell cycle-dependent regulatory circuit composed of 53BP1-RIF1 and BRCA1-CtlP controls DNA repair pathway choice. Mol Cell 2013; 49:872-883.
- Ghezraoui H, Oliveira C, Becker JR, et al. 53BP1 cooperation with the REV7-shieldin complex underpins DNA structure-specific NHEJ. Nature 2018; 560:122–127.
- 27. Xu G, Chapman JR, Brandsma I, et al. REV7 counteracts DNA double-strand break resection and affects PARP inhibition. Nature 2017; 521:541-544.
- Zimmermann M, Lottersberger F, Buonomo SB, et al. 53BP1 regulates DSB repair using Rif1 to control 5' end resection. Science 2013; 339:700-704.
- Chapman JR, Barral P, Vannier JB, et al. RIF1 is essential for 53BP1dependent nonhomologous end joining and suppression of DNA doublestrand break resection. Mol Cell 2013; 49:858–871.
- **30.** Feng L, Fong KW, Wang J, *et al.* RIF1 counteracts BRCA1-mediated end resection during DNA repair. J Biol Chem 2013; 288:11135–11143.
- Noordermeer SM, Adam S, Setiaputra D, et al. The Shieldin complex mediates 53BP1-dependent DNA repair. Nature 2018; 560:117–121.
- Dev H, Chiang TW, Lescale C, et al. Shieldin complex promotes DNA endjoining and counters homologous recombination in BRCA1-null cells. Nat Cell Biol 2018: 20:954–965.
- Findlay S, Heath J, Luo VM, et al. SHLD2/FAM35A co-operates with REV7 to coordinate DNA double-strand break repair pathway choice. EMBO J 2018; 37:; pii: e100158.
- Gupta R, Somyajit K, Narita T, et al. DNA repair network analysis reveals Shieldin as a key regulator of NHEJ and PARP inhibitor sensitivity. Cell 2018; 173:972–988.
- Ray Chaudhuri A, Callen E, Ding X, et al. Replication fork stability confers chemoresistance in BRCA-deficient cells. Nature 2016; 535:382-387.
- Lemacon D, Jackson J, Quinet A, et al. MRE11 and EXO1 nucleases degrade reversed forks and elicit MUS81-dependent fork rescue in BRCA2-deficient cells. Nat Commun 2017; 8:860.
- Lai X, Broderick R, Bergoglio V, et al. MUS81 nuclease activity is essential for replication stress tolerance and chromosome segregation in BRCA2-deficient cells. Nat Commun 2017; 8:15983.
- Rondinelli B, Gogola E, Yücel H, et al. EZH2 promotes degradation of stalled replication forks by recruiting MUS81 through histone H3 trimethylation. Nat Cell Biol 2017; 19:1371–1378.
- Taglialatela A, Alvarez S, Leuzzi G, et al. Restoration of replication fork stability in BRCA1- and BRCA2-deficient cells by inactivation of SNF2-family fork remodelers. Mol Cell 2017; 68:414.e8-430.e8.
- Pettitt SJ, Krastev DB, Brandsma I, et al. Genome-wide and high-density CRISPR-Cas9 screens identify point mutations in PARP1 causing PARP inhibitor resistance. Nat Commun 2018; 9:1849.
- Gogola E, Duarte AA, de Ruiter JR, et al. Selective loss of PARG restores PARylation and counteracts PARP inhibitor-mediated synthetic lethality. Cancer Cell 2018; 33:1078–1093.
- Rottenberg S, Nygren AO, Pajic M, et al. Selective induction of chemotherapy resistance of mammary tumors in a conditional mouse model for hereditary breast cancer. Proc Natl Acad Sci USA 2007; 104:12117-12122.

- **43.** Vaidyanathan A, Sawers L, Gannon AL, *et al.* ABCB1 (MDR1) induction defines a common resistance mechanism in paclitaxel- and olaparib-resistant ovarian cancer cells. Br J Cancer 2017; 115:431–441.
- Christie EL, Pattnaik S, Beach J, et al. Multiple ABCB1 transcriptional fusions in drug resistant high-grade serous ovarian and breast cancer. Nat Commun 2019; 10:1295.
- Ordonez LD, Hay T, McEwen R, et al. Rapid activation of epithelial-mesenchymal transition drives PARP inhibitor resistance in Brca2-mutant mammary tumours. Oncotarget 2019; 10:2586–2606.
- 46. Lheureux S, Bruce JP, Burnier JV, et al. Somatic BRCA1/2 recovery as a resistance mechanism after exceptional response to poly (ADP-ribose) polymerase inhibition. J Clin Oncol 2017; 35:1240-1249.
- 47. Park PH, Yamamoto TM, Li H, et al. Amplification of the mutation-carrying BRCA2 allele promotes RAD51 loading and PARP inhibitor resistance in the absence of reversion mutations. Mol Cancer Ther 2019; pii: molcanher. 0256.2019. [Epub ahead of print]
- **48.** Fleury H, Malaquin N, Tu V, et al. Exploiting interconnected synthetic lethal interactions between PARP inhibition and cancer cell reversible senescence. Nat Commun 2019; 10:2556.
- Alotaibi M, Sharma K, Saleh T, et al. Radiosensitization by PARP inhibition in DNA repair proficient and deficient tumor cells: proliferative recovery in senescent cells. Radiat Res 2016; 185:229–245.
- 50. Noordermeer SM, van Attikum H. PARP inhibitor resistance: a tug-of-war inBRCA-mutated cells. Trends Cell Biol 2019; 29:820-834.

Excellent comprehensive review on PARP inhibitor resistance mechanisms.