

MATB42 AMACSS Review Seminar 2023

Alex Teeter

University of Toronto Scarborough


June 1st 2022

Table of Contents

Important Definitions/Operators

Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.



Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

C - set

path: fn
 $c([a, b]) = C$

Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

The image of c under $[a, b]$ is called the curve parametrized by c and is typically denoted C

Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

The image of c under $[a, b]$ is called the curve parametrized by c and is typically denoted C

It is often required that c be one to one and C^1 (meaning all its components are differentiable.) In that case, the curve parametrized by c is called a simple curve.

$$c'(t) = \left(c_1'(t), \dots, c_n'(t) \right)$$


Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

The image of c under $[a, b]$ is called the curve parametrized by c and is typically denoted C



It is often required that c be one to one and C^1 (meaning all its components are differentiable.) In that case, the curve parametrized by c is called a simple curve.

We call $c(a)$ and $c(b)$ the endpoints of the curve. If C is simple and $c(a) = c(b)$, then we call C a simple closed curve.

Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

The image of c under $[a, b]$ is called the curve parametrized by c and is typically denoted C

It is often required that c be one to one and C^1 (meaning all its components are differentiable.) In that case, the curve parametrized by c is called a simple curve.

We call $c(a)$ and $c(b)$ the endpoints of the curve. If C is simple and $c(a) = c(b)$, then we call C a simple closed curve.

Parametrizing Curves and Surfaces

One of the most important things to remember how to do is to be able to parametrize various common curves and surfaces so you may integrate over them.

Parametrizing a Curve

A parametrization of a curve is a map $c : [a, b] \rightarrow \mathbb{R}^n$ (this is commonly called a path).

The image of c under $[a, b]$ is called the curve parametrized by c and is typically denoted C

It is often required that c be one to one and C^1 (meaning all its components are differentiable.) In that case, the curve parametrized by c is called a simple curve.

We call $c(a)$ and $c(b)$ the endpoints of the curve. If C is simple and $c(a) = c(b)$, then we call C a simple closed curve.

Parametrizing Curves and Surfaces

Derivative of a path

Derivative of a path with n components, $c(t) = (c_1(t), \dots, c_n(t))$ is given by

Parametrizing Curves and Surfaces

Derivative of a path

Derivative of a path with n components, $c(t) = (c_1(t), \dots, c_n(t))$ is given by

$$c'(t) = (c'_1(t), \dots, c'_n(t))$$

This is also referred to as the "velocity vector" or the vector tangent to $c(t)$ at time t .

Parametrizing Curves and Surfaces

Derivative of a path

Derivative of a path with n components, $c(t) = (c_1(t), \dots, c_n(t))$ is given by

$$c'(t) = (c'_1(t), \dots, c'_n(t))$$

This is also referred to as the "velocity vector" or the vector tangent to $c(t)$ at time t .

We refer to $\|c'(t)\|$ as the "speed" of the path at time t . It is the length of the velocity vector.

Parametrizing Curves and Surfaces

Derivative of a path

Derivative of a path with n components, $c(t) = (c_1(t), \dots, c_n(t))$ is given by

$$c'(t) = (c'_1(t), \dots, c'_n(t))$$

This is also referred to as the "velocity vector" or the vector tangent to $c(t)$ at time t .

We refer to $\|c'(t)\|$ as the "speed" of the path at time t . It is the length of the velocity vector.

The tangent line to a path at point t_0 is given by

$$\underline{c(t_0) + (t - t_0)c'(t_0)}$$

(Unless $c'(t_0) = 0$, in which case there is no tangent line)



Parametrizing Curves and Surfaces

Derivative of a path

Derivative of a path with n components, $c(t) = (c_1(t), \dots, c_n(t))$ is given by

$$c'(t) = (c'_1(t), \dots, c'_n(t))$$

This is also referred to as the "velocity vector" or the vector tangent to $c(t)$ at time t .

We refer to $\|c'(t)\|$ as the "speed" of the path at time t . It is the length of the velocity vector.

The tangent line to a path at point t_0 is given by

$$c(t_0) + (t - t_0)c'(t_0)$$

(Unless $c'(t_0) = 0$, in which case there is no tangent line)

Parametrizing Curves and Surfaces

We call a path c C^1 if all its component derivatives exist and are differentiable. C^n and C^∞ are what you would expect.

Parametrizing Curves and Surfaces

We call a path c C^1 if all its component derivatives exist and are differentiable. C^n and C^∞ are what you would expect.

Roughly speaking, a path is piecewise C^n if its component paths all satisfy the above condition



Parametrizing Curves and Surfaces

We call a path c C^1 if all its component derivatives exist and are differentiable. C^n and C^∞ are what you would expect.

Roughly speaking, a path is piecewise C^n if its component paths all satisfy the above condition

Parametrizing Curves and Surfaces

Important Examples:

Parametrizing Curves and Surfaces

Important Examples:

The circle of radius R in the plane can be parametrized by $c : [0, 2\pi] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (R\cos(t), R\sin(t))$$



Parametrizing Curves and Surfaces

Important Examples:

The circle of radius R in the plane can be parametrized by $c : [0, 2\pi] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (R\cos(t), R\sin(t))$$

The graph of a function $f : [a, b] \rightarrow \mathbb{R}$ in the plane can be parametrized by $c : [a, b] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (t, f(t))$$



Parametrizing Curves and Surfaces

Important Examples:

The circle of radius R in the plane can be parametrized by $c : [0, 2\pi] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (R\cos(t), R\sin(t))$$

The graph of a function $f : [a, b] \rightarrow \mathbb{R}$ in the plane can be parametrized by $c : [a, b] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (t, f(t))$$

The graph of a line from point (a, b) to (c, d) in the plane can be parametrized by $c : [0, 1] \rightarrow \mathbb{R}^2$ given by

Parametrizing Curves and Surfaces

Important Examples:

The circle of radius R in the plane can be parametrized by $c : [0, 2\pi] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (R\cos(t), R\sin(t))$$

The graph of a function $f : [a, b] \rightarrow \mathbb{R}$ in the plane can be parametrized by $c : [a, b] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (t, f(t))$$

The graph of a line from point (a, b) to (c, d) in the plane can be parametrized by $c : [0, 1] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (a + t(c - a), b + t(d - b))$$

Parametrizing Curves and Surfaces

Important Examples:

The circle of radius R in the plane can be parametrized by $c : [0, 2\pi] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (R\cos(t), R\sin(t))$$

The graph of a function $f : [a, b] \rightarrow \mathbb{R}$ in the plane can be parametrized by $c : [a, b] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (t, f(t))$$


The graph of a line from point (a, b) to (c, d) in the plane can be parametrized by $c : [0, 1] \rightarrow \mathbb{R}^2$ given by

$$c(t) = (a + t(c - a), b + t(d - b))$$

Parametrizing Curves and Surfaces

Sample Question

Parametrize the square with corners at $(0,0)$, $(1,0)$, $(1,1)$, $(0,1)$ with the counterclockwise orientation.



Parametrizing Curves and Surfaces

Sample Question

Parametrize the square with corners at $(0, 0)$, $(1, 0)$, $(1, 1)$, $(0, 1)$ with the counterclockwise orientation.

The square is made up of 4 lines, so this hints at the fact that we will need a piecewise parametrization.

Parametrizing Curves and Surfaces

Sample Question

Parametrize the square with corners at $(0,0)$, $(1,0)$, $(1,1)$, $(0,1)$ with the counterclockwise orientation.

The square is made up of 4 lines, so this hints at the fact that we will need a piecewise parametrization.

The first is a line from $(0,0)$ to $(1,0)$. This can be given easily by $(t, 0), t \in [0, 1]$.

Parametrizing Curves and Surfaces

$$c: [0, 4] \rightarrow \mathbb{R}^2$$

Sample Question

Parametrize the square with corners at $(0, 0)$, $(1, 0)$, $(1, 1)$, $(0, 1)$ with the counterclockwise orientation.

The square is made up of 4 lines, so this hints at the fact that we will need a piecewise parametrization.

The first is a line from $(0, 0)$ to $(1, 0)$. This can be given easily by $(t, 0)$, $t \in [0, 1]$.

Next comes the line from $(1, 0)$ to $(1, 1)$. We have to be more careful here. We want to carry on from where we left off from last parametrization, so we will need to adjust the line equation.



Parametrizing Curves and Surfaces

Sample Question

Parametrize the square with corners at $(0, 0)$, $(1, 0)$, $(1, 1)$, $(0, 1)$ with the counterclockwise orientation.

The square is made up of 4 lines, so this hints at the fact that we will need a piecewise parametrization.

The first is a line from $(0, 0)$ to $(1, 0)$. This can be given easily by $(t, 0)$, $t \in [0, 1]$.

Next comes the line from $(1, 0)$ to $(1, 1)$. We have to be more careful here. We want to carry on from where we left off from last parametrization, so we will need to adjust the line equation.

It can be given by $(1, t - 1)$, $t \in [1, 2]$. (Notice that it would be just $(1, t)$ if we were parametrizing using $[0, 1]$).

Parametrizing Curves and Surfaces

Sample Question

Parametrize the square with corners at $(0, 0)$, $(1, 0)$, $(1, 1)$, $(0, 1)$ with the counterclockwise orientation.

The square is made up of 4 lines, so this hints at the fact that we will need a piecewise parametrization.


The first is a line from $(0, 0)$ to $(1, 0)$. This can be given easily by $(t, 0), t \in [0, 1]$.

Next comes the line from $(1, 0)$ to $(1, 1)$. We have to be more careful here. We want to carry on from where we left off from last parametrization, so we will need to adjust the line equation.

It can be given by $(1, t - 1), t \in [1, 2]$. (Notice that it would be just $(1, t)$ if we were parametrizing using $[0, 1]$).

Parametrizing Curves and Surfaces


We then have the line from $(1, 1)$ to $(0, 1)$, which, again, by adjusting the domain and param is given by $(3 - t, 1), t \in [2, 3]$.



Parametrizing Curves and Surfaces

We then have the line from $(1, 1)$ to $(0, 1)$, which, again, by adjusting the domain and param is given by $(3 - t, 1), t \in [2, 3]$.

By the same idea, the line from $(0, 1)$ to $(0, 0)$ is given by $(0, 4 - t), t \in [3, 4]$.



Parametrizing Curves and Surfaces

We then have the line from $(1, 1)$ to $(0, 1)$, which, again, by adjusting the domain and param is given by $(3 - t, 1), t \in [2, 3]$.


By the same idea, the line from $(0, 1)$ to $(0, 0)$ is given by $(0, 4 - t), t \in [3, 4]$.

We can wrap this all together into the piecewise parametrization

$$c : \underline{[0, 4]} \rightarrow \mathbb{R}^2$$

Given by

$$c(t) = \begin{cases} (t, 0) & \text{if } t \in [0, 1] \\ (1, t - 1) & \text{if } t \in [1, 2] \\ (3 - t, 1) & \text{if } t \in [2, 3] \\ (0, 4 - t) & \text{if } t \in [3, 4] \end{cases}$$



Parametrizing Curves and Surfaces

We also have the analogous 2-dimensional version of parametrizing a curve, which is the parametrization of a surface:



Parametrizing Curves and Surfaces

We also have the analogous 2-dimensional version of parametrizing a curve, which is the parametrization of a surface:

Parametrization of a Surface

A parametrization of a Surface is a function $\Phi : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$. The "Surface" of this parametrization is its image.

$$\Phi(D) = S$$

$$\Phi(u, v) = (x(u, v), y(u, v), z(u, v))$$

$$[a, b] \times [c, d]$$

$$\frac{\partial}{\partial u} \quad \frac{\partial}{\partial v}$$

Analogously, we call Φ C^n if all its component functions are of class C^n wrt u and v .

Parametrizing Curves and Surfaces

T_u and T_v are the component vectors with the partials of Φ in terms of u and v respectively, i.e



Parametrizing Curves and Surfaces

T_u and T_v are the component vectors with the partials of Φ in terms of u and v respectively, i.e

$$\Phi(u_0, v_0) = (x, y, z)$$

$$T_u = \left(\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \right)$$

$$T_v = \left(\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \right)$$

The normal vector to the Surface at point $\Phi(u_0, v_0)$ is given by $n = T_u \times T_v$ at point (u_0, v_0) . Its tangent plane at point (x_0, y_0, z_0) , if $\Phi(u_0, v_0) = (x_0, y_0, z_0)$, is given by

$$n \cdot (x - x_0, y - y_0, z - z_0) = 0$$

Parametrizing Curves and Surfaces

Common Surface Parametrizations:

Parametrizing Curves and Surfaces

Common Surface Parametrizations:

The parametrization of a sphere of radius R centered at the origin can be parametrized by $\Phi : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by

$$\Phi(u, v) = (R \sin(u) \cos(v), R \sin(u) \sin(v), R \cos(u))$$



Parametrizing Curves and Surfaces

Common Surface Parametrizations:

The parametrization of a sphere of radius R centered at the origin can be parametrized by $\Phi : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by

$$\Phi(u, v) = (R\sin(u)\cos(v), R\sin(u)\sin(v), R\cos(u))$$

The parametrization of the torus (donut surface) with hole radius R and tube radius r can be parametrized by $\Phi : [0, 2\pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by




Parametrizing Curves and Surfaces

Common Surface Parametrizations:

The parametrization of a sphere of radius R centered at the origin can be parametrized by $\Phi : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by

$$\Phi(u, v) = (R\sin(u)\cos(v), R\sin(u)\sin(v), R\cos(u))$$

The parametrization of the torus (donut surface) with hole radius R and tube radius r can be parametrized by $\Phi : [0, 2\pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by


$$\Phi(u, v) = ((R + r\cos(u))\cos(v), (R + r\cos(u))\sin(v), r\sin(u))$$

Parametrizing Curves and Surfaces

Common Surface Parametrizations:


The parametrization of a sphere of radius R centered at the origin can be parametrized by $\Phi : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by

$$\Phi(u, v) = (R \sin(u) \cos(v), R \sin(u) \sin(v), R \cos(u))$$

The parametrization of the torus (donut surface) with hole radius R and tube radius r can be parametrized by $\Phi : [0, 2\pi] \times [0, 2\pi] \rightarrow \mathbb{R}^3$ given by

$$\Phi(u, v) = ((R + r \cos(u)) \cos(v), (R + r \cos(u)) \sin(v), r \sin(u))$$

The parametrization of the graph of a scalar function f on \mathbb{R}^2 can be parametrized by $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $\Phi(u, v) = (u, v, f(u, v))$.



Parametrizing Curves and Surfaces

Sample Question: Find the Tangent Plane of a Torus

Given a Torus, where $R = 2$ and $r = 1$, find its tangent plane at point $\Phi(0, 0)$

Parametrizing Curves and Surfaces

Sample Question: Find the Tangent Plane of a Torus

Given a Torus, where $R = 2$ and $r = 1$, find its tangent plane at point $\Phi(0, 0)$

First, we already know that this torus is parametrized by

$$\Phi(u, v) = ((R + r\cos(u))\cos(v), (R + r\cos(u))\sin(v), r\sin(u))$$

Parametrizing Curves and Surfaces

Sample Question: Find the Tangent Plane of a Torus

Given a Torus, where $R = 2$ and $r = 1$, find its tangent plane at point $\Phi(0, 0)$

First, we already know that this torus is parametrized by

$$\Phi(u, v) = ((R + r\cos(u))\cos(v), (R + r\cos(u))\sin(v), r\sin(u))$$

$$= (\overset{R}{\cancel{2}} + \overset{r}{\cancel{1}}\cos(u))\cos(v), (\overset{R}{\cancel{2}} + \overset{r}{\cancel{1}}\cos(u))\sin(v), \overset{r}{\cancel{1}}\sin(u))$$

Parametrizing Curves and Surfaces

Sample Question: Find the Tangent Plane of a Torus

Given a Torus, where $R = 2$ and $r = 1$, find its tangent plane at point $\Phi(0, 0)$

First, we already know that this torus is parametrized by

$$\Phi(u, v) = ((R + r\cos(u))\cos(v), (R + r\cos(u))\sin(v), r\sin(u))$$

$$= ((2 + \cos(u))\cos(v), (2 + \cos(u))\sin(v), \sin(u))$$

Now, we compute T_u and T_v

Parametrizing Curves and Surfaces

Sample Question: Find the Tangent Plane of a Torus

Given a Torus, where $R = 2$ and $r = 1$, find its tangent plane at point $\Phi(0, 0)$

First, we already know that this torus is parametrized by

$$\Phi(u, v) = ((R + r\cos(u))\cos(v), (R + r\cos(u))\sin(v), r\sin(u))$$

$$= ((2 + \cos(u))\cos(v), (2 + \cos(u))\sin(v), \sin(u))$$

Now, we compute T_u and T_v

$$T_u = \left(\frac{\partial((2 + \cos(u))\cos(v))}{\partial u}, \frac{\partial((2 + \cos(u))\sin(v))}{\partial u}, \frac{\partial \sin(u)}{\partial u} \right) = (-\sin(u)\cos(v), -\sin(u)\sin(v), \cos(u))$$

Parametrizing Curves and Surfaces

$$\mathbf{T}_v = (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

Parametrizing Curves and Surfaces

$$= (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

And

Parametrizing Curves and Surfaces

$$= (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

And

$$T_v = \left(\frac{\partial(2 + \cos(u))\cos(v)}{\partial v}, \frac{\partial(2 + \cos(u))\sin(v)}{\partial v}, \frac{\partial\sin(u)}{\partial v} \right) = 0$$

Handwritten notes above the equation:
-2sin(v) - cos(v)sin(u) (under the first term)
2cos(v) + cos(v)cos(u) (under the second term)

Parametrizing Curves and Surfaces

$$= (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

And

$$T_v = \left(\frac{\partial(2 + \cos(u))\cos(v)}{\partial v}, \frac{\partial(2 + \cos(u))\sin(v)}{\partial v}, \frac{\partial\sin(u)}{\partial v} \right)$$

$$\overline{T}_v = (-(2 + \cos(u))\sin(v), (2 + \cos(u))\cos(v), 0)$$

Parametrizing Curves and Surfaces

$$= (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

And

$$T_v = \left(\frac{\partial(2 + \cos(u))\cos(v)}{\partial v}, \frac{\partial(2 + \cos(u))\sin(v)}{\partial v}, \frac{\partial\sin(u)}{\partial v} \right)$$

$$= (-(2 + \cos(u))\sin(v), (2 + \cos(u))\cos(v), 0)$$

Now we compute the normal vector $T_u \times T_v$ at $(0, 0)$.

Parametrizing Curves and Surfaces

$$T_u = (-\cos(v)\sin(u), -\sin(v)\sin(u), \cos(u))$$

And

$$T_v = \left(\frac{\partial(2 + \cos(u))\cos(v)}{\partial v}, \frac{\partial(2 + \cos(u))\sin(v)}{\partial v}, \frac{\partial\sin(u)}{\partial v} \right)$$

$$T_v = (-(2 + \cos(u))\sin(v), (2 + \cos(u))\cos(v), 0)$$

Now we compute the normal vector $T_u \times T_v$ at $(0, 0)$.

Parametrizing Curves and Surfaces

Note that at $(0, 0)$, $T_u = (0, 2, 0)$ and $T_v = (1, 0, 0)$, so

$$n = T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 2 & 0 \\ 1 & 0 & 0 \end{vmatrix}$$

Parametrizing Curves and Surfaces

Note that at $(0,0)$, $T_u = (0, 2, 0)$ and $T_v = (1, 0, 0)$, so

$$\begin{aligned} n = T_u \times T_v &= \begin{vmatrix} i & j & k \\ 0 & 2 & 0 \\ 1 & 0 & 0 \end{vmatrix} \\ &= (0, 0, -2) \end{aligned}$$

$$a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$$

" "

$$(a, b, c)$$

Parametrizing Curves and Surfaces

Note that at $(0, 0)$, $T_u = (0, 2, 0)$ and $T_v = (1, 0, 0)$, so

$$n = T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 2 & 0 \\ 1 & 0 & 0 \end{vmatrix}$$

$$\mathbf{n} = (0, 0, -2)$$

Note that $\Phi(0, 0) = \underline{(2, 0, 1)}$, so the equation of the tangent plane is given by

$$n \cdot (x - 2, y - 0, z - 1) = 0(x - 2) + 0(y - 0) - \underline{2(z - 1)} = -2z + 2$$

at 0, i.e. $-2z + 2 = 0$ or $z = -1$. Thus the equation of the tangent plane at $\Phi(0, 0)$ is $z = -1$.

Parametrizing Curves and Surfaces

Note that at $(0, 0)$, $T_u = (0, 2, 0)$ and $T_v = (1, 0, 0)$, so

$$\begin{aligned} n = T_u \times T_v &= \begin{vmatrix} i & j & k \\ 0 & 2 & 0 \\ 1 & 0 & 0 \end{vmatrix} \\ &= (0, 0, -2) \end{aligned}$$

Note that $\Phi(0, 0) = (2, 0, 1)$, so the equation of the tangent plane is given by


$$n \cdot (x - 2, y - 0, z - 1) = 0(x - 2) + 0(y - z) - 2(z - 1) = -2z + 2$$

at 0, i.e. $-2z + 2 = 0$ or $z = -1$. Thus the equation of the tangent plane at $\Phi(0, 0)$ is $z = -1$.

Vector Fields

Vector Field

A Vector Field is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. This is typically visualized as having the vector assigned to the point \mathbb{R}^n pointing out from that point (though length is scaled back in order to avoid cluttering up the plot)



Vector Fields

$$F : (F_1(x_1, \dots, x_n), \dots, F_n(x_1, \dots, x_n))$$

Vector Field

A Vector Field is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. This is typically visualized as having the vector assigned to the point \mathbb{R}^n pointing out from that point (though length is scaled back in order to avoid cluttering up the plot)

Vector Field Operators

$$\operatorname{div}(F) := \nabla \cdot F = \frac{\partial F_1}{\partial x_1} + \dots + \frac{\partial F_n}{\partial x_n}$$

Vector Fields

$$\nabla = \left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n} \right)$$

Vector Field

A Vector Field is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. This is typically visualized as having the vector assigned to the point \mathbb{R}^n pointing out from that point (though length is scaled back in order to avoid cluttering up the plot)

Vector Field Operators

$$\operatorname{div}(F) := \nabla \cdot F = \frac{\partial F_1}{\partial x_1} + \dots + \frac{\partial F_n}{\partial x_n}$$

$$\operatorname{curl}(F) := \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x_1} & \frac{\partial}{\partial x_2} & \frac{\partial}{\partial x_3} \\ F_1 & F_2 & F_3 \end{vmatrix}$$

Vector Fields

Vector Field

A Vector Field is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. This is typically visualized as having the vector assigned to the point \mathbb{R}^n pointing out from that point (though length is scaled back in order to avoid cluttering up the plot)

Vector Field Operators

$$\operatorname{div}(F) := \nabla \cdot F = \frac{\partial F_1}{\partial x_1} + \cdots + \frac{\partial F_n}{\partial x_n}$$

$$\operatorname{curl}(F) := \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x_1} & \frac{\partial}{\partial x_2} & \frac{\partial}{\partial x_3} \\ F_1 & F_2 & F_3 \end{vmatrix}$$

$$\operatorname{div}(\nabla f)$$

We can also operate on regular scalar functions with the laplacian: $\nabla^2 f := \operatorname{div}(\nabla f)$

Vector Fields

Vector Field

A Vector Field is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. This is typically visualized as having the vector assigned to the point \mathbb{R}^n pointing out from that point (though length is scaled back in order to avoid cluttering up the plot)

Vector Field Operators

$$\operatorname{div}(F) := \nabla \cdot F = \frac{\partial F_1}{\partial x_1} + \cdots + \frac{\partial F_n}{\partial x_n}$$

$$\operatorname{curl}(F) := \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x_1} & \frac{\partial}{\partial x_2} & \frac{\partial}{\partial x_3} \\ F_1 & F_2 & F_3 \end{vmatrix}$$

We can also operate on regular scalar functions with the laplacian: $\nabla^2 f := \operatorname{div}(\nabla f)$

Vector Fields

Important Theorems

For any C^2 function f , $\text{curl}(\nabla f) = 0$ (Gradient has 0 curl for sufficiently nice functions)

Vector Fields

Important Theorems

For any C^2 function f , $\text{curl}(\nabla f) = 0$ (Gradient has 0 curl for sufficiently nice functions)

For any C^2 vector field F , $\text{div}(\text{curl}(F)) = 0$ (the divergence of the curl of any sufficiently nice vector field is 0)

Vector Fields

Important Theorems

For any C^2 function f , $\text{curl}(\nabla f) = 0$ (Gradient has 0 curl for sufficiently nice functions)

For any C^2 vector field F , $\text{div}(\text{curl}(F)) = 0$ (the divergence of the curl of any sufficiently nice vector field is 0)

We will discuss another related theorem soon.

Integration


The Path Integral

The path integral or the integral over the path c on a scalar function f in \mathbb{R}^3 , for $c : [a, b] \rightarrow \mathbb{R}^3$, is given by

Integration

The Path Integral


The path integral or the integral over the path c on a scalar function f in \mathbb{R}^3 , for $c : [a, b] \rightarrow \mathbb{R}^3$, is given by


$$\int_c f ds = \int_a^b f(x(t), y(t), z(t)) \|c'(t)\| dt$$

Integration

The Path Integral

The path integral or the integral over the path c on a scalar function f in \mathbb{R}^3 , for $c : [a, b] \rightarrow \mathbb{R}^3$, is given by

$$\int_c f ds = \int_a^b \underbrace{f(x(t), y(t), z(t))}_{\text{handwritten circle}} \|c'(t)\| dt$$




When $f(x, y, z) = 1$, you get the arclength of the curve parametrized by the path c

Integration

The Path Integral

The path integral or the integral over the path c on a scalar function f in \mathbb{R}^3 , for $c : [a, b] \rightarrow \mathbb{R}^3$, is given by

$$\int_c f ds = \int_a^b f(x(t), y(t), z(t)) \|c'(t)\| dt$$

When $f(x, y, z) = 1$, you get the arclength of the curve parametrized by the path c
If c is a plane curve and f is a scalar function on the plane, then

$$\int_c f(x, y) ds = \int_a^b f(x(t), y(t)) \|c'(t)\| dt$$

Integration

The Path Integral

The path integral or the integral over the path c on a scalar function f in \mathbb{R}^3 , for $c : [a, b] \rightarrow \mathbb{R}^3$, is given by

$$\int_c f ds = \int_a^b f(x(t), y(t), z(t)) \|c'(t)\| dt$$

When $f(x, y, z) = 1$, you get the arclength of the curve parametrized by the path c
If c is a plane curve and f is a scalar function on the plane, then

$$\int_c f(x, y) ds = \int_a^b f(x(t), y(t)) \|c'(t)\| dt$$

Integration

Sample Question

Evaluate the line ~~line~~ ^{path} integral

Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$


Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$

where C is parametrized by $c : [0, \pi] \rightarrow \mathbb{R}^3$, $c(t) = (\cos(t), \sin(t), t)$.



Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$

where C is parametrized by $c : [0, \pi] \rightarrow \mathbb{R}^3$, $c(t) = (\cos(t), \sin(t), t)$.

Well,

Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$

where C is parametrized by $c : [0, \pi] \rightarrow \mathbb{R}^3$, $c(t) = (\cos(t), \sin(t), t)$.

Well,

$$\int_C (xy + z^3) ds = \int_0^\pi (\cos(t)\sin(t) + t^3) \sqrt{\sin^2(t) + \cos^2(t) + 1} dt$$

plug $c(t)$ in

$$\begin{aligned} & -\sin(t) \cos(t) \\ & \frac{d}{dt} \cos(t) = -\sin(t) \\ & \frac{d}{dt} \sin(t) = \cos(t) \\ & \|c'(t)\| \end{aligned}$$

Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$

where C is parametrized by $c : [0, \pi] \rightarrow \mathbb{R}^3$, $c(t) = (\cos(t), \sin(t), t)$.

Well,

$$\begin{aligned} \int_C (xy + z^3) ds &= \int_0^\pi (\cos(t)\sin(t) + t^3) \sqrt{\sin^2(t) + \cos^2(t) + 1} dt \\ &= \int_0^\pi \sqrt{2}\cos(t)\sin(t) + \sqrt{2}t^3 dt \end{aligned}$$

Integration

Sample Question

Evaluate the line integral

$$\int_C (xy + z^3) ds$$

where C is parametrized by $c : [0, \pi] \rightarrow \mathbb{R}^3$, $c(t) = (\cos(t), \sin(t), t)$.

Well,

$$\begin{aligned} \int_C (xy + z^3) ds &= \int_0^\pi (\cos(t)\sin(t) + t^3) \sqrt{\sin^2(t) + \cos^2(t) + 1} dt \\ &= \int_0^\pi \sqrt{2}\cos(t)\sin(t) + \sqrt{2}t^3 dt \end{aligned}$$

Integration

$$\begin{aligned} & \frac{-\cos^2(t) - \sqrt{2}}{2} \\ &= \int_0^\pi \sqrt{2} \cos(t) \sin(t) dt + \int_0^\pi \sqrt{2} t^3 dt \quad \frac{\sqrt{2} + 4}{4} \end{aligned}$$

Integration

$$= \int_0^{\pi} \sqrt{2} \cos(t) \sin(t) dt + \int_0^{\pi} \sqrt{2} t^3 dt$$

$$= \frac{\sqrt{2}}{2} (-\cos^2(\pi) + \cos^2(0)) + \frac{\sqrt{2}}{4} (\pi^4 - 0)$$

Handwritten blue annotations:
- A large blue line is drawn through the entire second equation.
- A blue checkmark is written below the first term of the second equation.
- A blue dot is placed above the plus sign between the two terms.
- A blue circle is drawn around the second term of the second equation.

Integration

$$= \int_0^{\pi} \sqrt{2} \cos(t) \sin(t) dt + \int_0^{\pi} \sqrt{2} t^3 dt$$

$$= \frac{\sqrt{2}}{2} (-\cos^2(\pi) + \cos^2(0)) + \frac{\sqrt{2}}{4} (\pi^4 - 0)$$

$$= \frac{\sqrt{2} \pi^4}{4}$$

Integration

$$\begin{aligned} &= \int_0^{\pi} \sqrt{2} \cos(t) \sin(t) dt + \int_0^{\pi} \sqrt{2} t^3 dt \\ &= \frac{\sqrt{2}}{2} (-\cos^2(\pi) + \cos^2(0)) + \frac{\sqrt{2}}{4} (\pi^4 - 0) \\ &= \frac{\sqrt{2} \pi^4}{4} \end{aligned}$$

Thus

$$\int_C (xy + z^3) ds = \frac{\sqrt{2} \pi^4}{4}$$

Integration

Integral of a Scalar Function over a Surface

If $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, and S is a surface parametrized by Φ , then the scalar integral over S is:


Integration

Integral of a Scalar Function over a Surface

If $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, and S is a surface parametrized by Φ , then the scalar integral over S is:

$$\iint_S f dS = \iint_D f(\Phi(u, v)) \|T_u \times T_v\| du dv$$

Integration



Integral of a Scalar Function over a Surface

If $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, and S is a surface parametrized by Φ , then the scalar integral over S is:

$$\iint_S f dS = \iint_D f(\Phi(u, v)) \|T_u \times T_v\| \underline{du dv}$$

Analogously, if $f = 1$, then the integral gives the surface area of S



Integration

$$F(x, y, z) = y$$


Sample Question

Evaluate the scalar surface integral

$$\int \int (\sqrt{3} \cos(u))^2 + (\sqrt{3} \sin(u))^2 = 3 \int \int y dS$$

where S is the portion of the cylinder $x^2 + y^2 = 3$ that lies between $z = 0$ and $z = 6$

First, we can parametrize this surface through $\Phi(u, v) = (\sqrt{3} \cos(u), \sqrt{3} \sin(u), v)$, where $0 \leq v \leq 6$ and $0 \leq u \leq 2\pi$.



Integration

Sample Question

Evaluate the scalar surface integral

$$\iint y dS$$

where S is the portion of the cylinder $x^2 + y^2 = 3$ that lies between $z = 0$ and $z = 6$

First, we can parametrize this surface through $\Phi(u, v) = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), v)$, where $0 \leq v \leq 6$ and $0 \leq u \leq 2\pi$.

First, we will compute T_u and T_v .

Integration

Sample Question

Evaluate the scalar surface integral

$$\iint y dS$$

where S is the portion of the cylinder $x^2 + y^2 = 3$ that lies between $z = 0$ and $z = 6$

First, we can parametrize this surface through $\Phi(u, v) = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), v)$, where $0 \leq v \leq 6$ and $0 \leq u \leq 2\pi$.

First, we will compute T_u and T_v .

$$T_u = \left(\frac{\partial \sqrt{3}\cos(u)}{\partial u}, \frac{\partial \sqrt{3}\sin(u)}{\partial u}, \frac{\partial v}{\partial u} \right) = (-\sqrt{3}\sin(u), \sqrt{3}\cos(u), 0)$$

Handwritten notes in blue:
Below the first two terms of the vector: $\sqrt{3}\sin(u)$ and $\sqrt{3}\cos(u)$
Above the third term: $\partial v / \partial u$

Integration

Sample Question

Evaluate the scalar surface integral

$$\iint y dS$$

where S is the portion of the cylinder $x^2 + y^2 = 3$ that lies between $z = 0$ and $z = 6$

First, we can parametrize this surface through $\Phi(u, v) = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), v)$, where $0 \leq v \leq 6$ and $0 \leq u \leq 2\pi$.

First, we will compute T_u and T_v .

$$T_u = \left(\frac{\partial \sqrt{3}\cos(u)}{\partial u}, \frac{\partial \sqrt{3}\sin(u)}{\partial u}, \frac{\partial v}{\partial u} \right) = (-\sqrt{3}\sin(u), \sqrt{3}\cos(u), 0)$$

$$T_v = \left(\frac{\partial \sqrt{3}\cos(u)}{\partial v}, \frac{\partial \sqrt{3}\sin(u)}{\partial v}, \frac{\partial v}{\partial v} \right) = (0, 0, 1)$$

Integration

Sample Question

Evaluate the scalar surface integral

$$\iint y dS$$

where S is the portion of the cylinder $x^2 + y^2 = 3$ that lies between $z = 0$ and $z = 6$

First, we can parametrize this surface through $\Phi(u, v) = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), v)$, where $0 \leq v \leq 6$ and $0 \leq u \leq 2\pi$.

First, we will compute T_u and T_v .

$$T_u = \left(\frac{\partial \sqrt{3}\cos(u)}{\partial u}, \frac{\partial \sqrt{3}\sin(u)}{\partial u}, \frac{\partial v}{\partial u} \right) = (-\sqrt{3}\sin(u), \sqrt{3}\cos(u), 0)$$

$$T_v = \left(\frac{\partial \sqrt{3}\cos(u)}{\partial v}, \frac{\partial \sqrt{3}\sin(u)}{\partial v}, \frac{\partial v}{\partial v} \right) = (0, 0, 1)$$

Integration

Now, we compute $T_u \times T_v$.

Integration

Now, we compute $T_u \times T_v$.

$$\mathbf{r}_1 \mathbf{i} - \mathbf{r}_2 \mathbf{j} + \mathbf{r}_3 \mathbf{k} \\ = (F_1, -F_2, F_3)$$

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), 0)$$

Warning!

$$T_u \times T_v \neq T_v \times T_u$$

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (\sqrt{3}\cos(u), \sqrt{3}\sin(u), 0)$$

Lastly, we compute $\|T_u \times T_v\|$.

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (-\sqrt{3}\cos(u), -\sqrt{3}\sin(u), 0)$$

Lastly, we compute $\|T_u \times T_v\|$.

$$\|T_u \times T_v\| = \sqrt{3\cos^2(u) + 3\sin^2(u)} = \sqrt{3}$$

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = \underline{(-\sqrt{3}\cos(u), -\sqrt{3}\sin(u), 0)}$$

Lastly, we compute $\|T_u \times T_v\|$.

$$\|(a, b, c)\| = \sqrt{a^2 + b^2 + c^2}$$

$$\underline{\|T_u \times T_v\|} = \sqrt{\underline{3\cos^2(u) + 3\sin^2(u)}} = \sqrt{3}$$

Pythagorean
identity

Thus,

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (-\sqrt{3}\cos(u), -\sqrt{3}\sin(u), 0)$$

Lastly, we compute $\|T_u \times T_v\|$.

$$\|T_u \times T_v\| = \sqrt{3\cos^2(u) + 3\sin^2(u)} = \sqrt{3}$$

Thus,

$$\iint_S y dS = \iint_D \sqrt{3}\sin(u) \|T_u \times T_v\| dA$$

Handwritten notes: $\times (\phi(u,v))$ and $\approx \sqrt{3}$ are written in blue ink above the integral. The terms $\sqrt{3}\sin(u)$ and $\|T_u \times T_v\|$ in the integrand are circled in blue.

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (-\sqrt{3}\cos(u), -\sqrt{3}\sin(u), 0)$$

Lastly, we compute $\|T_u \times T_v\|$.

$$\|T_u \times T_v\| = \sqrt{3\cos^2(u) + 3\sin^2(u)} = \sqrt{3}$$

Thus,

$$\begin{aligned} \iint_S y dS &= \iint_D \sqrt{3}\sin(u) \|T_u \times T_v\| dA \\ &= \iint_D \sqrt{3}\sin(u) \|T_u \times T_v\| dA \end{aligned}$$

Integration

Now, we compute $T_u \times T_v$.

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 0 & 0 & 1 \\ -\sqrt{3}\sin(u) & \sqrt{3}\cos(u) & 0 \end{vmatrix} = (-\sqrt{3}\cos(u), -\sqrt{3}\sin(u), 0)$$

Lastly, we compute $\|T_u \times T_v\|$.

$$\|T_u \times T_v\| = \sqrt{3\cos^2(u) + 3\sin^2(u)} = \sqrt{3}$$

Thus,

$$\begin{aligned} \iint_S y dS &= \iint_D \sqrt{3}\sin(u) \|T_u \times T_v\| dA \\ &= \iint_D \sqrt{3}\sin(u) \|T_u \times T_v\| dA \end{aligned}$$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$3 \sin(u)$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$$= 3 \int_0^{2\pi} \int_0^6 \sin(u) dv du$$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$$= 3 \int_0^{2\pi} \int_0^6 \sin(u) dv du$$

$$= 18 \int_0^{2\pi} \sin(u) du$$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$$= 3 \int_0^{2\pi} \int_0^6 \sin(u) dv du$$

$$= 18 \int_0^{2\pi} \sin(u) du$$

$$= (-18 \cos(u)) \Big|_0^{2\pi}$$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$$= 3 \int_0^{2\pi} \int_0^6 \sin(u) dv du$$

$$= 18 \int_0^{2\pi} \sin(u) du$$

$$= (-18 \cos(u)) \Big|_0^{2\pi}$$

$$= 0$$

Integration

$$= \int_0^{2\pi} \int_0^6 \sqrt{3} \sin(u) \sqrt{3} dv du$$

$$= 3 \int_0^{2\pi} \int_0^6 \sin(u) dv du$$

$$= 18 \int_0^{2\pi} \sin(u) du$$

$$= (-18 \cos(u)) \Big|_0^{2\pi}$$

$$= 0$$

Integration

Line Integral (The Important One)

If F is a vector field that is continuous on the path $c : [a, b] \rightarrow \mathbb{R}^3$, then the line integral of F along c is given by


$$\int_c F \cdot ds = \int_a^b F(c(t)) \cdot c'(t) dt$$

Integration

Line Integral (The Important One)

If F is a vector field that is continuous on the path $c : [a, b] \rightarrow \mathbb{R}^3$, then the line integral of F along c is given by

$$\int_c F \cdot ds = \int_a^b F(c(t)) \cdot c'(t) dt$$



Theorem of Line Integrals on Gradient Vector Fields

Suppose that $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is C^1 and that $c : [a, b] \rightarrow \mathbb{R}^3$ is a piecewise C^1 path. Then

$$\int_c \nabla f \cdot ds = f(c(b)) - f(c(a))$$

A handwritten blue integral formula: $\int_c \nabla f \cdot ds$. The ∇ is written with a small superscript r (representing \mathbb{R}^3), and the ds has a small superscript r (representing \mathbb{R}^3).

Integration

Line Integral (The Important One)

If F is a vector field that is continuous on the path $c : [a, b] \rightarrow \mathbb{R}^3$, then the line integral of F along c is given by

$$\int_c F \cdot ds = \int_a^b F(c(t)) \cdot c'(t) dt$$

Theorem of Line Integrals on Gradient Vector Fields

Suppose that $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is C^1 and that $c : [a, b] \rightarrow \mathbb{R}^3$ is a piecewise C^1 path. Then

$$\int_c \nabla f \cdot ds = f(c(b)) - f(c(a))$$

Integration .



Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then

Integration

Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then



$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

$= \text{curl}(F)$

Integration

Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then

$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

We can also express this in the language of the vector field operators, in which case it is

Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then

$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

We can also express this in the language of the vector field operators, in which case it is

$$\int_C F \cdot ds = \iint_D \operatorname{curl}(F) \cdot (0, 0, 1) dA$$

Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then

$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

We can also express this in the language of the vector field operators, in which case it is

$$\int_C F \cdot ds = \iint_D \operatorname{curl}(F) \cdot (0, 0, 1) dA$$

Green's Theorem (Important!)

If D is a region that can be decomposed into simple regions, and its boundary is C , and $F = (P(x, y), Q(x, y))$ is a C^1 vector field, then

$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

We can also express this in the language of the vector field operators, in which case it is

$$\int_C F \cdot ds = \iint_D \operatorname{curl}(F) \cdot (0, 0, 1) dA$$

Integration


Sample Question

Evaluate

$$\int_C F \cdot ds$$

where C is the boundary of the triangle with vertices $(-2, -1)$, $(1, 2)$, $(2, -1)$ and $F(x, y) = (2xy + \sin(x^2), 3x^2 + 3^y)$.

The parametrization of C seems too difficult. Bring in Green's Theorem.



Integration

Sample Question

Evaluate

$$\int_C F \cdot ds$$

where C is the boundary of the triangle with vertices $(-2, -1)$, $(1, 2)$, $(2, -1)$ and $F(x, y) = (2xy + \sin(x^2), 3x^2 + 3^y)$.

The parametrization of C seems too difficult. Bring in Green's Theorem.

First we need to compute $\frac{\partial Q}{\partial x}$ and $\frac{\partial P}{\partial y}$

Integration

Sample Question

Evaluate

$$\int_C F \cdot ds$$

where C is the boundary of the triangle with vertices $(-2, -1)$, $(1, 2)$, $(2, -1)$ and $F(x, y) = (2xy + \sin(x^2), 3x^2 + 3y)$.

The parametrization of C seems too difficult. Bring in Green's Theorem.

First we need to compute $\frac{\partial Q}{\partial x}$ and $\frac{\partial P}{\partial y}$

6x

$$\frac{\partial Q}{\partial x} = \frac{\partial(3x^2 + \cancel{3y})}{\partial x} = 6x$$

Integration

Sample Question

Evaluate

$$\int_C F \cdot ds$$

where C is the boundary of the triangle with vertices $(-2, -1)$, $(1, 2)$, $(2, -1)$ and $F(x, y) = (2xy + \sin(x^2), 3x^2 + 3^y)$.

The parametrization of C seems too difficult. Bring in Green's Theorem.

First we need to compute $\frac{\partial Q}{\partial x}$ and $\frac{\partial P}{\partial y}$

$$\frac{\partial Q}{\partial x} = \frac{\partial(3x^2 + 3^y)}{\partial x} = 6x$$

Integration

$$\frac{\partial P}{\partial y} = \frac{\partial(2x + \sin(x^2))}{\partial y} = 2x$$

Integration

$$\frac{\partial P}{\partial y} = \frac{\partial(2xy + \sin(x^2))}{\partial y} = 2x$$

Thus, by Green's Theorem, we have

Integration

$$\frac{\partial P}{\partial y} = \frac{\partial(2xy + \sin(x^2))}{\partial y} = 2x$$

Thus, by Green's Theorem, we have

$$\int_C F \cdot ds = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA = \iint_D 4x dA$$



Integration

$$\frac{\partial P}{\partial y} = \frac{\partial(2xy + \sin(x^2))}{\partial y} = 2x$$

Thus, by Green's Theorem, we have

$$\int_C F \cdot ds = \iint_D 6x - 2x dA = \iint_D 4x dA$$

Now all that's left is to compute $\iint_D 4x dA$.

Integration

$$\frac{\partial P}{\partial y} = \frac{\partial(2xy + \sin(x^2))}{\partial y} = 2x$$

Thus, by Green's Theorem, we have

$$\int_C F \cdot ds = \iint_D 6x - 2x dA = \iint_D 4x dA$$

Now all that's left is to compute $\iint_D 4x dA$.

Integration



The x bounds of the triangular region are bounded above by the line segment between $(1, 2)$ and $(2, -1)$, and below by the line segment between $(-2, -1)$ and $(1, 2)$.

Integration

The x bounds of the triangular region are bounded above by the line segment between $(1, 2)$ and $(2, -1)$, and below by the line segment between $(-2, -1)$ and $(1, 2)$.

The slope of the line segment between $(1, 2)$ and $(2, -1)$ is given by

Integration

The x bounds of the triangular region are bounded above by the line segment between $(1, 2)$ and $(2, -1)$, and below by the line segment between $(-2, -1)$ and $(1, 2)$.

The slope of the line segment between $(1, 2)$ and $(2, -1)$ is given by

$$\text{Slope} = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \frac{-1 - 2}{2 - 1} = -3$$

Integration

The x bounds of the triangular region are bounded above by the line segment between $(1, 2)$ and $(2, -1)$, and below by the line segment between $(-2, -1)$ and $(1, 2)$.

The slope of the line segment between $(1, 2)$ and $(2, -1)$ is given by

$$m = \frac{-1 - 2}{2 - 1} = -3$$

This means the line segment is given by

$$y - y_1 = m(x - x_1)$$

$$y - 2 = -3(x - 1)$$

, or $x = \frac{y}{-3} + \frac{5}{3}$

$$\frac{x}{-3} + \frac{5}{3} = x - 1$$

Integration

The x bounds of the triangular region are bounded above by the line segment between $(1, 2)$ and $(2, -1)$, and below by the line segment between $(-2, -1)$ and $(1, 2)$.

The slope of the line segment between $(1, 2)$ and $(2, -1)$ is given by

$$m = \frac{-1 - 2}{2 - 1} = -3$$

This means the line segment is given by

$$y - 2 = -3(x - 1)$$

, or $x = \frac{y}{-3} + \frac{5}{3}$

Integration

The line segment between $(-2, -1)$ and $(1, 2)$ has slope

Integration

The line segment between $(-2, -1)$ and $(1, 2)$ has slope

$$\frac{2 + 1}{1 + 2} = 1$$

Integration

The line segment between $(-2, -1)$ and $(1, 2)$ has slope

$$\frac{2 + 1}{1 + 2} = 1$$

And so it is given by

$$y + 1 = x + 2$$

or $x = y - 1$

Integration

The line segment between $(-2, -1)$ and $(1, 2)$ has slope


$$\frac{2 + 1}{1 + 2} = 1$$

And so it is given by

$$y + 1 = x + 2$$

or $x = y - 1$

Thus the x region is $y - 1 \leq x \leq \frac{-y}{3} + \frac{5}{3}$



Integration

The line segment between $(-2, -1)$ and $(1, 2)$ has slope

$$\frac{2 + 1}{1 + 2} = 1$$

And so it is given by

$$y + 1 = x + 2$$

or $x = y - 1$

Thus the x region is $y - 1 \leq x \leq \frac{-y}{3} + \frac{5}{3}$

The y region is given by $-1 \leq y \leq 2$

We can finally evaluate the integral!

Integration

$$\iint_D 4x dA = \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy$$

Integration

$$\begin{aligned}\iint_D 4x dA &= \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy \\ &= \int_{-1}^2 2\left(\frac{-y}{3} + \frac{5}{3}\right)^2 - 2(y-1)^2 dy \\ &= \frac{x^2}{9} - \frac{10x}{9} + \frac{25}{9} - \frac{(y-1)^3}{3} + \frac{(y-1)^3}{3} \Big|_{-1}^2\end{aligned}$$

Handwritten notes: $2x^2$ (above the first integral), $-x^2 - 2x + 1$ (above the second integral), and a large blue oval around the final expression.

Integration

$$\iint_D 4x dA = \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy$$

$$= \int_{-1}^2 2\left(\frac{-y}{3} + \frac{5}{3}\right)^2 - 2(y-1)^2 dy$$

$$= 2 \int_{-1}^2 \frac{y^2}{9} - \frac{10y}{9} + \frac{25}{9} - y^2 + 2y - 1 dy$$

Integration

$$\iint_D 4x dA = \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy$$

$$= \int_{-1}^2 2\left(\frac{-y}{3} + \frac{5}{3}\right)^2 - 2(y-1)^2 dy$$

$$= 2 \int_{-1}^2 \frac{y^2}{9} - \frac{10y}{9} + \frac{25}{9} - y^2 + 2y - 1 dy$$

$$= 2 \int_{-1}^2 \left(\frac{-8y^2}{9} + \frac{8y}{9} + \frac{16}{9} \right) dy$$

Integration

$$\begin{aligned}\iint_D 4x dA &= \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy \\&= \int_{-1}^2 2\left(\frac{-y}{3} + \frac{5}{3}\right)^2 - 2(y-1)^2 dy \\&= 2 \int_{-1}^2 \frac{y^2}{9} - \frac{10y}{9} + \frac{25}{9} - y^2 + 2y - 1 dy \\&= 2 \int_{-1}^2 \frac{-8y^2}{9} + \frac{8y}{9} + \frac{16}{9} dy \\&= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right)\end{aligned}$$

Integration

$$\begin{aligned}\iint_D 4x dA &= \int_{-1}^2 \int_{y-1}^{\frac{-y}{3} + \frac{5}{3}} 4x dx dy \\&= \int_{-1}^2 2\left(\frac{-y}{3} + \frac{5}{3}\right)^2 - 2(y-1)^2 dy \\&= 2 \int_{-1}^2 \frac{y^2}{9} - \frac{10y}{9} + \frac{25}{9} - y^2 + 2y - 1 dy \\&= 2 \int_{-1}^2 \frac{-8y^2}{9} + \frac{8y}{9} + \frac{16}{9} dy \\&= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right)\end{aligned}$$

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \end{aligned}$$

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-64}{27} - \frac{8}{27} + \frac{16}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \end{aligned}$$

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-64}{27} - \frac{8}{27} + \frac{16}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-72}{27} + \frac{4}{3} + \frac{16}{3}\right) \end{aligned}$$

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-64}{27} - \frac{8}{27} + \frac{16}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-72}{27} + \frac{4}{3} + \frac{16}{3}\right) \\ &= 2\left(\frac{-72}{27} + \frac{20}{3}\right) \end{aligned}$$

.

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-64}{27} - \frac{8}{27} + \frac{16}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-72}{27} + \frac{4}{3} + \frac{16}{3}\right) \\ &= 2\left(\frac{-72}{27} + \frac{20}{3}\right) \end{aligned}$$

Integration

$$\begin{aligned} &= 2\left(\frac{-8}{9} \int_{-1}^2 y^2 dy + \frac{8}{9} \int_{-1}^2 y dy + \frac{16}{9} \int_{-1}^2 1 dy\right) \\ &= 2\left(\frac{-8(2)^3}{27} + \frac{8}{27}(-1)^3 + \frac{4(2)^2}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-64}{27} - \frac{8}{27} + \frac{16}{9} - \frac{4}{9} + \frac{32}{9} + \frac{16}{9}\right) \\ &= 2\left(\frac{-72}{27} + \frac{4}{3} + \frac{16}{3}\right) \\ &= 2\left(\frac{-72}{27} + \frac{20}{3}\right) \end{aligned}$$

Integration


$$= 2\left(\frac{108}{27}\right)$$

Integration

$$= 2\left(\frac{108}{27}\right)$$

$$= 2(4) = 8$$

Integration

$$\begin{aligned} \int F \cdot ds &\rightarrow \int_D \omega(F) \\ &= 2\left(\frac{108}{27}\right) \int_D F \cdot ds \leftarrow \int_D \omega(F) \\ &= 2(4) = 8 \end{aligned}$$


So $\int_C \vec{f} \cdot d\vec{s} = 8$.

Integration

Surface Integral over a Vector Field

Let \underline{F} be a vector field defined over a surface S parametrized by Φ . Then the surface integral over the vector field F is given by


$$\iint_{\Phi} F \cdot dS = \iint_D F \cdot (T_u \times T_v) du dv$$

Where D is the domain of parametrization.

$$n = \frac{T_u \times T_v}{\|T_u \times T_v\|}$$

$-n$

$$n(0,0,1) = (0,0,1)$$



Integration

Surface Integral over a Vector Field

Let F be a vector field defined over a surface S parametrized by Φ . Then the surface integral over the vector field F is given by

$$\iint_{\Phi} F \cdot dS = \iint_D F \cdot (T_u \times T_v) du dv$$

Where D is the domain of parametrization.



Stokes' Theorem

Let S be an oriented surface, given by a one to one parametrization $\Phi : D \subset \mathbb{R}^2 \rightarrow S$, where D is a region for which Green's Theorem applies. Let ∂S denote the oriented boundary of S and let F be a C^1 vector field on S . Then

$$\iint_S (\nabla \times F) \cdot dS = \int_{\partial S} F \cdot ds$$

Handwritten notes: "= curl(F)" above the first integral and "= 0" to the right of the equation.



Integration

Surface Integral over a Vector Field

Let F be a vector field defined over a surface S parametrized by Φ . Then the surface integral over the vector field F is given by

$$\iint_{\Phi} F \cdot dS = \iint_D F \cdot (T_u \times T_v) du dv$$

Where D is the domain of parametrization.

Stokes' Theorem

Let S be an oriented surface, given by a one to one parametrization $\Phi : D \subset \mathbb{R}^2 \rightarrow S$, where D is a region for which Green's Theorem applies. Let ∂S denote the oriented boundary of S and let F be a C^1 vector field on S . Then

$$\iint_S (\nabla \times F) \cdot dS = \int_{\partial S} F \cdot ds$$

Sample Problem

Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$

where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$


Sample Problem

Evaluate

$$\iint_S \overbrace{\text{curl}(F)}^{\nabla \times F} \cdot dS$$

where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$

This looks too complicated to do directly. Let's use Stokes' Theorem.



Sample Problem


Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$

where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$

This looks too complicated to do directly. Let's use Stokes' Theorem.

We need to determine ∂S if we are going to do this. ∂S is precisely the part of our surface where $z = 0$, or i.e



Sample Problem

Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$

where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$

This looks too complicated to do directly. Let's use Stokes' Theorem.

We need to determine ∂S if we are going to do this. ∂S is precisely the part of our surface where $z = 0$, or i.e

$$0 = 9 - x^2 - y^2$$

Sample Problem


Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$


where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$

This looks too complicated to do directly. Let's use Stokes' Theorem.

We need to determine ∂S if we are going to do this. ∂S is precisely the part of our surface where $z = 0$, or i.e


$$0 = 9 - x^2 - y^2$$

$$x^2 + y^2 = 9$$



Sample Problem

Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$


where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so $n(0, 0, 9) = (0, 0, 1)$ and $F = (x^3 - y, x, xz + e^y z)$

This looks too complicated to do directly. Let's use Stokes' Theorem.

We need to determine ∂S if we are going to do this. ∂S is precisely the part of our surface where $z = 0$, or i.e

$$0 = 9 - x^2 - y^2$$

$$x^2 + y^2 = 9$$



This is the circle of radius 3 in the xy plane.

Sample Problem

Evaluate

$$\iint_S \text{curl}(F) \cdot dS$$

$$\phi(u,v) = (u, v, 9-u^2-v^2)$$

where S is the part of the graph $z = 9 - x^2 - y^2$ above the xy plane oriented so

$n(0,0,9) = (0,0,1)$ and $F = (x^3 - y, x, xz + e^y z)$

$$\phi(0,0) = (0,0,9)$$

This looks too complicated to do directly. Let's use Stokes' Theorem.

We need to determine ∂S if we are going to do this. ∂S is precisely the part of our surface where $z = 0$, or i.e

$$\frac{(2(0), 2(0), 1)}{\sqrt{2(0)^2 + 2(0)^2 + 1^2}} = (0,0,1)$$

$$0 = 9 - x^2 - y^2$$

$$x^2 + y^2 = 9$$

$$T_u = (1, 0, -2u)$$

$$T_v = (0, 1, -2v)$$

$$T_u \times T_v = \begin{vmatrix} i & j & k \\ 1 & 0 & -2u \\ 0 & 1 & -2v \end{vmatrix} = 2ui + 2vj + k = (2u, 2v, 1)$$

This is the circle of radius 3 in the xy plane.

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

$$(3\cos(t))^2 + (3\sin(t))^2 = 9$$

Integration

We can parametrize this by $c(t) = (3\cos(t), \overbrace{3\sin(t)}^{-3\sin(t)}, 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (\overbrace{-3\sin(t)}^{-3\sin(t)}, \overbrace{3\cos(t)}^{3\cos(t)}, 0)$, and that

$$F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$$

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (3\sin(t), -3\cos(t), 0)$, and that

$$F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$$

$$\iint_S \operatorname{curl}(F) \cdot dS = \int_{\partial S} F \cdot dS$$

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (3\sin(t), -3\cos(t), 0)$, and that
 $F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$

$$\iint_S \text{curl}(F) \cdot dS = \int_{\partial S} F \cdot dS$$

$$= \int_0^{2\pi} F(c(t)) \cdot c'(t) dt$$

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (-3\sin(t), 3\cos(t), 0)$, and that

$$F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$$

$$\iint_S \text{curl}(F) \cdot dS = \int_{\partial S} F \cdot dS$$

$$= \int_0^{2\pi} F(c(t)) \cdot c'(t) dt$$

$$= \int_0^{2\pi} -81\cos^3(t)\sin(t) + 9\sin^2(t) + 9\cos^2(t) dt$$

$$\begin{bmatrix} -3\sin(t) \\ 3\cos(t) \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 27\cos^3(t) - 3\sin(t) \\ 3\cos(t) \\ 0 \end{bmatrix}$$

$$= -81\sin(t)\cos^3(t) + 9\sin^2(t) + 9\cos^2(t)$$

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (3\sin(t), -3\cos(t), 0)$, and that

$$F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$$

$$\iint_S \text{curl}(F) \cdot dS = \int_{\partial S} F \cdot dS$$

$$= \int_0^{2\pi} F(c(t)) \cdot c'(t) dt$$

$$= \int_0^{2\pi} 81\cos^3(t)\sin(t) - 9\sin^2(t) - 9\cos^2(t) dt$$

Handwritten in blue: $\frac{81\cos^4(t)}{4} = 0$

$$= \int_0^{2\pi} 81\cos^3(t)\sin(t) dt + \int_0^{2\pi} -9(\cos^2(t) + \sin^2(t)) dt$$

Integration

We can parametrize this by $c(t) = (3\cos(t), 3\sin(t), 0)$, $t \in [0, 2\pi]$

Note then that $c'(t) = (3\sin(t), -3\cos(t), 0)$, and that

$$F(c(t)) = (27\cos^3(t) - 3\sin(t), 3\cos(t), 0)$$

$$\iint_S \text{curl}(F) \cdot dS = \int_{\partial S} F \cdot dS$$

$$= \int_0^{2\pi} F(c(t)) \cdot c'(t) dt$$

$$= \int_0^{2\pi} 81\cos^3(t)\sin(t) \cancel{-} 9\sin^2(t) \cancel{+} 9\cos^2(t) dt$$

$$= \int_0^{2\pi} 81\cos^3(t)\sin(t) dt + \int_0^{2\pi} \cancel{-} 9(\cos^2(t) + \sin^2(t)) dt$$

Integration

$$= \left(\frac{-\cos^4(2\pi)81}{4} + \frac{\cos^4(0)81}{4} \right) + \cancel{-}18\pi$$

$$= 0 \cancel{-} 18\pi$$



$$= \cancel{+}18\pi$$

Thus

$$\iint_S \text{curl}(F) \cdot dS = \cancel{-}18\pi$$

Integration


$$\nabla \times \vec{F} = 0$$



Conservative Vector Fields

If F is a C^1 vector field on \mathbb{R}^3 , except at a finite number of points, then F is called "conservative" if one and hence all of the following equivalent conditions hold:

1. For any oriented simple closed curve, $\int_C F \cdot dR = 0$
2. For any 2 oriented simple curves with the same endpoints, $\int_{C_1} F \cdot dR = \int_{C_2} F \cdot dS$
3. F is the gradient of some function f (we call the function f the potential of F)
4. $\nabla \times F = 0$



Integration

Sample Question

Given $F(x, y, z) = (2xy^2 + 5, 2x^2y - 3y^2, 0)$, compute

$$\int_C F \cdot dr$$

where C is the ellipse given by $\frac{(x-4)^2}{5} + \frac{y^2}{4} = 1$

Integration

Sample Question

Given $F(x, y, z) = (2xy^2 + 5, 2x^2y - 3y^2, 0)$, compute

$$\int_C F \cdot dr$$

where C is the ellipse given by $\frac{(x-4)^2}{5} + \frac{y^2}{4} = 1$

Yikes. This looks too nasty to do directly. However, what if the vector field was conservative? Let's check.

Integration

Sample Question

Given $F(x, y, z) = (2xy^2 + 5, 2x^2y - 3y^2, 0)$, compute

$$\int_C F \cdot dr$$

where C is the ellipse given by $\frac{(x-4)^2}{5} + \frac{y^2}{4} = 1$

Yikes. This looks too nasty to do directly. However, what if the vector field was conservative? Let's check.

Let us compute the curl.


Integration

Sample Question

Given $F(x, y, z) = (2xy^2 + 5, 2x^2y - 3y^2, 0)$, compute

$$\int_C F \cdot dr$$

where C is the ellipse given by $\frac{(x-4)^2}{5} + \frac{y^2}{4} = 1$



Yikes. This looks too nasty to do directly. However, what if the vector field was conservative? Let's check.


Let us compute the curl.

$$\text{curl}(F) = \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xy^2 + 5 & 2x^2y - 3y^2 & 0 \end{vmatrix} = (0, 0, 4xy - 4xy) = (0, 0, 0)$$

Integration

So the vector field IS conservative! Thus

Integration



So the vector field IS conservative! Thus

$$\int_C F \cdot dr = 0$$

Integration

So the vector field IS conservative! Thus

$$\int_C F \cdot dr = 0$$

Since an ellipse is a closed simple curve.

Integration

So the vector field IS conservative! Thus

$$\int_C F \cdot dr = 0$$

Since an ellipse is a closed simple curve.

Integration


little change
on inside
= big change
on outside



Gauss'/The Divergence Theorem

Let W be a symmetric elementary region in space, and ∂W be the surface that bounds it. Then, if F is C^1 , then

$$\iiint_W \operatorname{div}(F) dV = \iint_{\partial W} F \cdot dS$$



Integration

Sample Questions

Evaluate

$$\iint_S \vec{F} \cdot d\vec{S}$$

where $\vec{F} = (2xz, 1 - 4xy^2, 2z - z^2)$ where S is the surface of the solid bounded by $z = 6 - 2x^2 - 2y^2$ and the plane $z = 0$



Integration

Sample Questions

Evaluate

$$\iint_S F \cdot dS$$

where $F = (2xz, 1 - 4xy^2, 2z - z^2)$ where S is the surface of the solid bounded by $z = 6 - 2x^2 - 2y^2$ and the plane $z = 0$

Yeesh. Too hard to evaluate directly. Let's use the divergence theorem.

Integration

Sample Questions

Evaluate

$$\iint_S F \cdot dS$$

where $F = (2xz, 1 - 4xy^2, 2z - z^2)$ where S is the surface of the solid bounded by $z = 6 - 2x^2 - 2y^2$ and the plane $z = 0$

Yeesh. Too hard to evaluate directly. Let's use the divergence theorem.

So, we have

$$\underline{\iint_S F \cdot dS} = \underline{\iiint_E \operatorname{div}(F) dV}$$



Integration

Stokes: $\nabla \times F$
Gauss: F

Sample Questions

Evaluate

$$-2(x^2 + y^2) = -2r^2 \quad \iint_S F \cdot dS$$

where $F = (2xz, 1 - 4xy^2, 2z - z^2)$ where S is the surface of the solid bounded by $z = 6 - 2x^2 - 2y^2$ and the plane $z = 0$

$$z = 6 - r^2$$

Yeesh. Too hard to evaluate directly. Let's use the divergence theorem.

So, we have

$$\iint_S F \cdot dS = \iiint_E \operatorname{div}(F) dV$$

Where E is the region S bounds. Let's set it up in cylindrical coordinates.

Integration

Sample Questions

Evaluate

$$\iint_S F \cdot dS$$

where $F = (2xz, 1 - 4xy^2, 2z - z^2)$ where S is the surface of the solid bounded by $z = 6 - 2x^2 - 2y^2$ and the plane $z = 0$

Yeesh. Too hard to evaluate directly. Let's use the divergence theorem.

So, we have

$$\underline{\iint_S F \cdot dS} = \underline{\iiint_E \operatorname{div}(F) dV}$$

Where E is the region S bounds. Let's set it up in cylindrical coordinates.

Integration

$$z = 6 - 2r^2$$

$$0 \leq z \leq 6 - 2r^2$$

$$6 - 2(\sqrt{3})^2 \\ = 6 - 6 = 0$$

Notice that in cylindrical coordinates, the region is $0 \leq r \leq \sqrt{3}$, $0 \leq \theta \leq 2\pi$,
 $0 \leq z \leq 6 - 2r^2$

$$6 - 2x^2 - 2y^2$$

$$= 6 - 2(x^2 + y^2) = \underline{6 - 2r^2}$$

Notice that in cylindrical coordinates, the region is $0 \leq r \leq \sqrt{3}$, $0 \leq \theta \leq 2\pi$,
 $0 \leq z \leq 6 - 2r^2$

Next, we compute the divergence. It is

Integration

Notice that in cylindrical coordinates, the region is $0 \leq r \leq \sqrt{3}$, $0 \leq \theta \leq 2\pi$,
 $0 \leq z \leq 6 - 2r^2$

Next, we compute the divergence. It is

$$\text{div}(F) = \frac{\partial \overset{F_1}{2xz}}{\partial x} + \frac{\partial \overset{F_2}{1-4xy^2}}{\partial y} + \frac{\partial \overset{F_3}{2z-z^2}}{\partial z} = \cancel{2z} - 8xy + 2 - \cancel{2z} = 2 - 8xy$$

Integration

Notice that in cylindrical coordinates, the region is $0 \leq r \leq \sqrt{3}$, $0 \leq \theta \leq 2\pi$,
 $0 \leq z \leq 6 - 2r^2$

Next, we compute the divergence. It is

$$\operatorname{div}(F) = \frac{\partial 2xz}{\partial x} + \frac{\partial 1 - 4xy^2}{\partial y} + \frac{\partial 2z - z^2}{\partial z} = 2z - 8xy + 2 - 2z = 2 - 8xy$$

At last, we can compute the integral, remembering to pick up an r since we converted E to polar coordinates.

Integration

$$\iiint_E \operatorname{div}(F) dV = \iiint_E 2 - 8xy dV$$

Integration

$$\begin{aligned}\iiint_E \operatorname{div}(F) dV &= \iiint_E 2 - 8xy dV \\ &= \int_0^{2\pi} \int_0^{\sqrt{3}} \int_0^{6-2r^2} \underset{\substack{\text{Jacobian} \\ r}}{r} (2 - 8r^2 \cos(\theta) \sin(\theta)) dz dr d\theta\end{aligned}$$

Integration

$$\begin{aligned}\iiint_E \operatorname{div}(F) dV &= \iiint_E 2 - 8xy dV \\&= \int_0^{2\pi} \int_0^{\sqrt{3}} \int_0^{6-2r^2} r(2 - 8r^2 \cos(\theta) \sin(\theta)) dz dr d\theta \\&= \int_0^{2\pi} \int_0^{\sqrt{3}} \int_0^{6-2r^2} 2r - 8r^3 \cos(\theta) \sin(\theta) dz dr d\theta\end{aligned}$$

Integration

$$\iiint_E \operatorname{div}(F) dV = \iiint_E 2 - 8xy dV$$

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \int_0^{6-2r^2} (2 - 8r^2 \cos(\theta) \sin(\theta)) dz dr d\theta$$

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \int_0^{6-2r^2} 2r - 8r^3 \cos(\theta) \sin(\theta) dz dr d\theta$$

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} 2r - 8r^3 \cos(\theta) \sin(\theta) (6 - 2r^2) dr d\theta$$

Integration

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} (12r - 4r^3 - (48r^3 - 16r^5)\cos(\theta)\sin(\theta)) dr d\theta$$

Integration

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \frac{12r}{6r^2} - \frac{4r^3}{r^4} - (48r^3 - \frac{16r^5}{6}) \cos(\theta) \sin(\theta) dr d\theta$$

$$= \int_0^{2\pi} 6(\sqrt{3})^2 - (\sqrt{3})^4 - (12(\sqrt{3})^4 - \frac{8}{3}(\sqrt{3})^6) \cos(\theta) \sin(\theta) d\theta$$

Integration

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} (12r - 4r^3 - (48r^3 - 16r^5)\cos(\theta)\sin(\theta)) dr d\theta$$

$$= \int_0^{2\pi} (6(\sqrt{3})^2 - (\sqrt{3})^4 - (12(\sqrt{3})^4 - \frac{8}{3}(\sqrt{3})^6)\cos(\theta)\sin(\theta)) d\theta$$

$$= \int_0^{2\pi} (9 - 36\cos(\theta)\sin(\theta)) d\theta$$

Integration

$$= \int_0^{2\pi} \int_0^{\sqrt{3}} 12r - 4r^3 - (48r^3 - 16r^5) \cos(\theta) \sin(\theta) dr d\theta$$

$$= \int_0^{2\pi} 6(\sqrt{3})^2 - (\sqrt{3})^4 - (12(\sqrt{3})^4 - \frac{8}{3}(\sqrt{3})^6) \cos(\theta) \sin(\theta) d\theta$$

$$= \int_0^{2\pi} \boxed{9} - 36 \cos(\theta) \sin(\theta) d\theta$$

~~18 cos²(θ)~~

$$= 9(2\pi) + 18 \cancel{\cos^2(4\pi)} - 9 \cancel{(0)} - 18 \cancel{\cos^2(0)} = 18\pi$$

Integration

Thus,

$$\iint_S F \cdot dS = 18\pi$$

Differential Forms

Differential Forms provide a unified language for all the vector calculus concepts you have learnt thus far.

Differential Forms

Differential Forms provide a unified language for all the vector calculus concepts you have learnt thus far.

They are important to know and understand how to manipulate.

Differential Forms

Differential Forms provide a unified language for all the vector calculus concepts you have learnt thus far.

They are important to know and understand how to manipulate.

They come with 2 fundamental operations: \wedge (the wedge product) and d (The exterior derivative)

Differential Forms

Differential Forms provide a unified language for all the vector calculus concepts you have learnt thus far.

They are important to know and understand how to manipulate.

They come with 2 fundamental operations: \wedge (the wedge product) and d (The exterior derivative)

Differential Forms

Scalars: $F: \mathbb{R}^3 \rightarrow \mathbb{R}$

$k=1,2$ 2 forms $dx \wedge dy$ $dx \wedge dz$ $dy \wedge dz$

Space of k forms

The space of k forms on \mathbb{R}^3 is denoted $\wedge^k(\mathbb{R}^3)$. Some important facts:

1. $\wedge^k(\mathbb{R}^3)$ is a vector space with functions as scalars and will consist of all linear combinations of basic k forms. For $\wedge^1(\mathbb{R}^3)$, it's $f_1 dx + f_2 dy + f_3 dz$, for $\wedge^2(\mathbb{R}^3)$, it's $f_1 dx \wedge dy + f_2 dx \wedge dz + f_3 dy \wedge dz$, and for $\wedge^3(\mathbb{R}^3)$, it's $f_1 dx \wedge dy \wedge dz$

Basic 3-form is just $dx \wedge dy \wedge dz$

Differential Forms

Space of K forms

The space of k forms on \mathbb{R}^3 is denoted $\bigwedge^k(\mathbb{R}^3)$. Some important facts:

1. $\bigwedge^k(\mathbb{R}^3)$ is a vector space with functions as scalars and will consist of all linear combinations of basic k forms. For $\bigwedge^1(\mathbb{R}^3)$, it's $f_1dx + f_2dy + f_3dz$, for $\bigwedge^2(\mathbb{R}^3)$, it's $f_1dx \wedge dy + f_2dx \wedge dz + f_3dy \wedge dz$, and for $\bigwedge^3(\mathbb{R}^3)$, it's $f_1dx \wedge dy \wedge dz$
2. As a sanity check, $\bigwedge^k(\mathbb{R}^3)$ should always be $C(3, k)$ dimensional.

Differential Forms

Space of k forms

The space of k forms on \mathbb{R}^3 is denoted $\bigwedge^k(\mathbb{R}^3)$. Some important facts:

1. $\bigwedge^k(\mathbb{R}^3)$ is a vector space with functions as scalars and will consist of all linear combinations of basic k forms. For $\bigwedge^1(\mathbb{R}^3)$, it's $f_1 dx + f_2 dy + f_3 dz$, for $\bigwedge^2(\mathbb{R}^3)$, it's $f_1 dx \wedge dy + f_2 dx \wedge dz + f_3 dy \wedge dz$, and for $\bigwedge^3(\mathbb{R}^3)$, it's $f_1 dx \wedge dy \wedge dz$
2. As a sanity check, $\bigwedge^k(\mathbb{R}^3)$ should always be $C(3, k)$ dimensional.
3. $\bigwedge^0(\mathbb{R}^3)$ is just the space of scalar functions $f : \mathbb{R}^3 \rightarrow \mathbb{R}$.

$$\bigwedge^k(\mathbb{R}^3) \cong \{0\} \text{ if } k > 3$$

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

► Distributivity: $(dx + dy) \wedge dz = dx \wedge dz + dy \wedge dz$

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

- ▶ Distributivity: $(dx + dy) \wedge dz = dx \wedge dy + dy \wedge dz$
- ▶ Associative: $(dx \wedge dy) \wedge dz = dx \wedge (dy \wedge dz) = dx \wedge dy \wedge dz$

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

- ▶ Distributivity: $(dx + dy) \wedge dz = dx \wedge dy + dy \wedge dz$
- ▶ Associative: $(dx \wedge dy) \wedge dz = dx \wedge (dy \wedge dz) = dx \wedge dy \wedge dz$
- ▶ Anti-Commutative: $dx \wedge dy = (-1)^{kl} dy \wedge dx$

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

- ▶ Distributivity: $(dx + dy) \wedge dz = dx \wedge dz + dy \wedge dz$
- ▶ Associative: $(dx \wedge dy) \wedge dz = dx \wedge (dy \wedge dz) = dx \wedge dy \wedge dz$
- ▶ Anti-Commutative: $dx \wedge dy = (-1)^{kl} dy \wedge dx$
- ▶ Homogeneity wrt functions: $f dx \wedge dy = dx \wedge f dy = f(dx \wedge dy)$

Differential Forms

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

- ▶ Distributivity: $(dx + dy) \wedge dz = dx \wedge dy + dy \wedge dz$
- ▶ Associative: $(dx \wedge dy) \wedge dz = dx \wedge (dy \wedge dz) = dx \wedge dy \wedge dz$
- ▶ Anti-Commutative: $dx \wedge dy = (-1)^{kl} dy \wedge dx$
- ▶ Homogeneity wrt functions: $f dx \wedge dy = dx \wedge f dy = f(dx \wedge dy)$
- ▶ $dx \wedge dx = 0$

F is a 0-form

The Wedge Product

The wedge product takes a k form and an l form and makes a $k + l$ form. It is determined by the following rules:

- ▶ Distributivity: $(dx + dy) \wedge dz = dx \wedge dz + dy \wedge dz$
- ▶ Associative: $(dx \wedge dy) \wedge dz = dx \wedge (dy \wedge dz) = dx \wedge dy \wedge dz$
- ▶ Anti-Commutative: $dx \wedge dy = (-1)^{kl} dy \wedge dx$
- ▶ Homogeneity wrt functions: $f dx \wedge dy = dx \wedge f dy = f(dx \wedge dy)$
- ▶ $dx \wedge dx = 0$
- ▶ $\omega \wedge 0 = 0$ for any k form ω

$\wedge^k(\mathbb{R}^s)$ ← has a 0-form

Differential Forms

The Exterior Derivative

The natural generalization of the derivative to differential forms. Generalizes the curl, grad and div. It is a map $d : \underline{\bigwedge^k(\mathbb{R}^3)} \rightarrow \underline{\bigwedge^{k+1}(\mathbb{R}^3)}$ determined by the following rules:

Differential Forms

The Exterior Derivative

The natural generalization of the derivative to differential forms. Generalizes the curl, grad and div. It is a map $d : \bigwedge^k(\mathbb{R}^3) \rightarrow \bigwedge^{k+1}(\mathbb{R}^3)$ determined by the following rules:

- It is a linear map wrt the vector space structure

$$d(w + \alpha) = dw + d\alpha$$

Differential Forms

The Exterior Derivative

The natural generalization of the derivative to differential forms. Generalizes the curl, grad and div. It is a map $d : \bigwedge^k(\mathbb{R}^3) \rightarrow \bigwedge^{k+1}(\mathbb{R}^3)$ determined by the following rules:

- ▶ It is a linear map wrt the vector space structure
- ▶ For a k form ω and an l form μ , $d(\omega \wedge \mu) = d(\omega) \wedge \mu + (-1)^k \omega \wedge d(\mu)$

Differential Forms

The Exterior Derivative

The natural generalization of the derivative to differential forms. Generalizes the curl, grad and div. It is a map $d : \bigwedge^k(\mathbb{R}^3) \rightarrow \bigwedge^{k+1}(\mathbb{R}^3)$ determined by the following rules:


- ▶ It is a linear map wrt the vector space structure
- ▶ For a k form ω and an l form μ , $d(\omega \wedge \mu) = d(\omega) \wedge \mu + (-1)^k \omega \wedge d(\mu)$
- ▶ $d(d\omega) = 0$ for any k form ω

Differential Forms

The Exterior Derivative

The natural generalization of the derivative to differential forms. Generalizes the curl, grad and div. It is a map $d : \bigwedge^k(\mathbb{R}^3) \rightarrow \bigwedge^{k+1}(\mathbb{R}^3)$ determined by the following rules:

- ▶ It is a linear map wrt the vector space structure
- ▶ For a k form ω and an l form μ , $d(\omega \wedge \mu) = d(\omega) \wedge \mu + (-1)^k \omega \wedge d(\mu)$
- ▶ $d(d\omega) = 0$ for any k form ω
- ▶ For any 0 form f , $d(f) = \frac{\partial f}{\partial x}dx + \frac{\partial f}{\partial y}dy + \frac{\partial f}{\partial z}dz$



Handwritten red arrow pointing from the 0-form rule to a vector of partial derivatives:


$$\left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

Differential Forms

The Generalized Stokes' Theorem

This is the magnum opus of Vector/Manifold Calculus. It generalizes Green's, Gauss' and Stokes' Theorems with the appropriate identifications.

Differential Forms



The Generalized Stokes' Theorem

This is the magnum opus of Vector/Manifold Calculus. It generalizes Green's, Gauss' and Stokes' Theorems with the appropriate identifications.

Let M be an oriented k manifold in \mathbb{R}^n . If ω is a $k - 1$ form, then

Differential Forms

The Generalized Stokes' Theorem

This is the magnum opus of Vector/Manifold Calculus. It generalizes Green's, Gauss' and Stokes' Theorems with the appropriate identifications.

Let M be an oriented k manifold in \mathbb{R}^n . If ω is a $k - 1$ form, then

$$\int_{\partial M} \omega = \int_M d(\omega)$$

Differential Forms

The Generalized Stokes' Theorem

This is the magnum opus of Vector/Manifold Calculus. It generalizes Green's, Gauss' and Stokes' Theorems with the appropriate identifications.

Let M be an oriented k manifold in \mathbb{R}^n . If ω is a $k - 1$ form, then

$$\int_{\partial M} \omega = \int_M d(\omega)$$

Differential Forms

Sample Question

Let $\omega = x^2ydx + xz^2dy + y^3dz$ and $\eta = \cos(y)dx + e^x dy + zdz$. Calculate $\omega \wedge \eta$

Differential Forms

Sample Question


Let $\omega = x^2ydx + xz^2dy + y^3dz$ and $\eta = \cos(y)dx + e^x dy + z dz$. Calculate $\omega \wedge \eta$

$$\omega \wedge \eta = (x^2ydx + xz^2dy + y^3dz) \wedge (\cos(y)dx + e^x dy + z dz)$$

Differential Forms

Sample Question

Let $\omega = x^2ydx + xz^2dy + y^3dz$ and $\eta = \cos(y)dx + e^x dy + zdz$. Calculate $\omega \wedge \eta$

$$\omega \wedge \eta = (x^2ydx + xz^2dy + y^3dz) \wedge (\cos(y)dx + e^x dy + zdz)$$


Distributive
Property of \wedge



$$x^2ydx \wedge (\cos(y)dx + e^x dy + zdz) + xz^2dy \wedge (\cos(y)dx + e^x dy + zdz) + y^3dz \wedge (\cos(y)dx + e^x dy + zdz)$$

Differential Forms

Sample Question

Let $\omega = x^2ydx + xz^2dy + y^3dz$ and $\eta = \cos(y)dx + e^x dy + zdz$. Calculate $\omega \wedge \eta$

$$\omega \wedge \eta = (x^2ydx + xz^2dy + y^3dz) \wedge (\cos(y)dx + e^x dy + zdz)$$

$$x^2ydx \wedge (\cos(y)dx + e^x dy + zdz) + xz^2dy \wedge (\cos(y)dx + e^x dy + zdz) + y^3dz \wedge (\cos(y)dx + e^x dy + zdz)$$

↙ anti-commutativity
OF \wedge

$$-(\cos(y)dx + e^x dy + zdz) \wedge x^2ydx - (\cos(y)dx + e^x dy + zdz) \wedge xz^2dy - (\cos(y)dx + e^x dy + zdz) \wedge y^3dz$$

Differential Forms

Sample Question

Let $\omega = x^2ydx + xz^2dy + y^3dz$ and $\eta = \cos(y)dx + e^x dy + z dz$. Calculate $\omega \wedge \eta$

$$\omega \wedge \eta = (x^2ydx + xz^2dy + y^3dz) \wedge (\cos(y)dx + e^x dy + z dz)$$

$$x^2ydx \wedge (\cos(y)dx + e^x dy + z dz) + xz^2dy \wedge (\cos(y)dx + e^x dy + z dz) + y^3dz \wedge (\cos(y)dx + e^x dy + z dz)$$

$$-(\cos(y)dx + e^x dy + z dz) \wedge x^2ydx - (\cos(y)dx + e^x dy + z dz) \wedge xz^2dy - (\cos(y)dx + e^x dy + z dz) \wedge y^3dz$$

Differential Forms

↓ Distributive Property
+ Homogeneity of $F \subset \Lambda^k$

$$= -e^x x^2 y dy \wedge dx - x^2 y z dz \wedge dx - \cos(y) x z^2 dx \wedge dy - x z^3 dz \wedge dy - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

Differential Forms

$$= -e^x x^2 y dy \wedge dx - x^2 y z dz \wedge dx - \cos(y) x z^2 dx \wedge dy - x z^3 dz \wedge dy - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

anti-symmetry

$$= e^x x^2 y \underline{dx \wedge dy} + x^2 y z dx \wedge dz - \cos(y) x z^2 dx \wedge dy + x z^3 dy \wedge dz - \cos(y) y^3 \underline{dx \wedge dz} - e^x y^3 \underline{dy \wedge dz}$$

Differential Forms

$$= -e^x x^2 y dy \wedge dx - x^2 y z dz \wedge dx - \cos(y) x z^2 dx \wedge dy - x z^3 dz \wedge dy - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

$$= e^x x^2 y dx \wedge dy + x^2 y z dx \wedge dz - \cos(y) x z^2 dx \wedge dy + x z^3 dy \wedge dz - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

✓ because $\wedge^2(\mathbb{R}^3)$
is a vector space

$$(e^x x^2 y - \cos(y) x z^2) \underline{dx} \wedge dy + (x^2 y z - \cos(y) y^3) dx \wedge \underline{dz} + (x z^3 - e^x y^3) dy \wedge dz$$

Differential Forms

$$= -e^x x^2 y dy \wedge dx - x^2 y z dz \wedge dx - \cos(y) x z^2 dx \wedge dy - x z^3 dz \wedge dy - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

$$= e^x x^2 y dx \wedge dy + x^2 y z dx \wedge dz - \cos(y) x z^2 dx \wedge dy + x z^3 dy \wedge dz - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

$$(e^x x^2 y - \cos(y) x z^2) dx \wedge dy + (x^2 y z - \cos(y) y^3) dx \wedge dz + (x z^3 - e^x y^3) dy \wedge dz$$

Thus

$$\omega \wedge \eta = (e^x x^2 y - \cos(y) x z^2) dx \wedge dy + (x^2 y z - \cos(y) y^3) dx \wedge dz + (x z^3 - e^x y^3) dy \wedge dz$$

Differential Forms

$$= -e^x x^2 y dy \wedge dx - x^2 y z dz \wedge dx - \cos(y) x z^2 dx \wedge dy - x z^3 dz \wedge dy - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

$$= e^x x^2 y dx \wedge dy + x^2 y z dx \wedge dz - \cos(y) x z^2 dx \wedge dy + x z^3 dy \wedge dz - \cos(y) y^3 dx \wedge dz - e^x y^3 dy \wedge dz$$

$$(e^x x^2 y - \cos(y) x z^2) dx \wedge dy + (x^2 y z - \cos(y) y^3) dx \wedge dz + (x z^3 - e^x y^3) dy \wedge dz$$

Thus

$$\omega \wedge \eta = (e^x x^2 y - \cos(y) x z^2) dx \wedge dy + (x^2 y z - \cos(y) y^3) dx \wedge dz + (x z^3 - e^x y^3) dy \wedge dz$$

$- dz \wedge dx$

Differential Forms

Sample Question

Let S be the unit sphere with outward pointing normal, and ω be the 2 form
 $(x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx$

Differential Forms

Sample Question

Let S be the unit sphere with outward pointing normal, and ω be the 2 form
 $(x + y + z)dxdy + (3x - 2)dydz + (x - 5y)dzdx$
Evaluate

$$\int_S \omega$$

Yeesh. Too hard to evaluate directly once again. Time to bring out the generalized stokes' Theorem.

Differential Forms



Sample Question

Let S be the unit sphere with outward pointing normal, and ω be the 2 form
 $(x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx$
Evaluate

$$\int_S \omega$$

Yeesh. Too hard to evaluate directly once again. Time to bring out the generalized stokes' Theorem.

Note that the unit sphere is the boundary of the unit ball B , i.e. $\partial B = S^2$

Differential Forms

Sample Question

Let S be the unit sphere with outward pointing normal, and ω be the 2 form
 $(x + y + z)dxdy + (3x - 2)dydz + (x - 5y)dzdx$
Evaluate

$$\int_S \omega$$

Yeesh. Too hard to evaluate directly once again. Time to bring out the generalized stokes' Theorem.

Note that the unit sphere is the boundary of the unit ball B , i.e $\partial B = S^2$

Differential Forms

Next, we need to compute $d\omega$. Note:

$$d\omega = d((x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx)$$

Differential Forms

Next, we need to compute $d\omega$. Note:

$$d\omega = d((x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx)$$

$$= d((x + y + z)dx dy) + d((3x - 2)dy dz) + d((x - 5y)dz dx)$$

Differential Forms

Next, we need to compute $d\omega$. Note:

$$d\omega = d((x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx)$$

$dx \wedge dx = 0$


$$= d((x + y + z)dx dy) + d((3x - 2)dy dz) + d((x - 5y)dz dx)$$

product rule with \wedge

$$= d(x+y+z)dx dy + (x+y+z)d(dx dy) + d(3x-2)dy dz + (3x-2)d(dy dz) + d(x-5y)dz dx + (x-5y)d(dz dx)$$

Differential Forms

Next, we need to compute $d\omega$. Note:

$$d\omega = d((x + y + z)dx dy + (3x - 2)dy dz + (x - 5y)dz dx)$$


$$= d((x + y + z)dx dy) + d((3x - 2)dy dz) + d((x - 5y)dz dx)$$

$$= d(x+y+z)dx dy + (x+y+z)d(dx dy) + d(3x-2)dy dz + (3x-2)d(dy dz) + d(x-5y)dz dx + (x-5y)d(dz dx)$$

$$= dz dx dy + 3 dx dy dz - 5 dy dz dx$$

Differential Forms

$$= \underline{dx dy dz + 3 dx dy dz - 5 dx dy dz}$$

anti-symmetry
of \wedge

Differential Forms

$$= dx dy dz + 3 dx dy dz - 5 dx dy dz$$

$$\underline{\underline{-dx dy dz}}$$

Differential Forms

$$= dx dy dz + 3 dx dy dz - 5 dx dy dz$$

$$- dx dy dz$$

Thus, by the generalized Stokes' Theorem, we have

$$\begin{aligned}\int_S \omega &= \int_{\partial B} \omega \\ &= \int_B d(\omega) \\ &= \iiint_B -1 dx dy dz\end{aligned}$$

$$S = \partial B$$

Differential Forms

This is the negative volume of the unit ball, which is

$$\frac{4\pi r^3}{3}$$

.

Differential Forms

This is the negative volume of the unit ball, which is

$$\frac{4\pi r^3}{3}$$

.
Thus,

$$\int_S \omega = \frac{-4\pi r^3}{3}$$

Differential Forms

This is the negative volume of the unit ball, which is

$$\frac{4\pi \mathbf{1}^3}{3}$$

.
Thus,

$$\int_S \omega = \frac{-4\pi \mathbf{2}^3}{3}$$

Fourier Series

A Fourier approximation is a way of approximating a symmetric function on a certain interval.

Fourier Series

A Fourier approximation is a way of approximating a symmetric function on a certain interval.

Fourier Series

The fourier coefficients of a function defined on $[-\pi, \pi]$ are given by

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(kx) dx$$

$$b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(kx) dx$$

Fourier Series

$$a_0 = \frac{3\pi}{4}$$

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(kx) dx.$$

Sample Question

Calculate the fourier series of the following function:

$$f(x) = \begin{cases} \pi, & \text{if } -\pi \leq x \leq 0 \\ x, & \text{if } 0 \leq x \leq \pi \end{cases}$$

$$\frac{1}{\pi} \left(\int_{-\pi}^0 \pi \cos(kx) dx + \int_0^{\pi} x \cos(kx) dx \right)$$

$$\frac{1}{\pi} \left(\left(\frac{\sin(kx)}{k} \right)_{-\pi}^0 + \int_0^{\pi} \right)$$

Differential Forms

That's all for the review seminar! Any questions?

Differential Forms

That's all for the review seminar! Any questions?