

COMP20003
Algorithms and Data Structures
Recurrences

Nir Lipovetzky
 Department of Computing and
 Information Systems
 University of Melbourne
 Semester 2



Divide and Conquer Algorithms



Mergesort and quicksort are instances of divide-and-conquer algorithms:


- Solve the problem by continually dividing into smaller problems

Other examples?

COMP20003 Algorithms and Data Structures

1-2

Split-solve-join approach:



For problems where the output is a transformation of the input, need to:

- process both sub-problems, and
- join the sub-solutions after processing

COMP20003 Algorithms and Data Structures


4-3

Recurrence for divide and conquer sorting algorithms



One pass through the data reduces problem size by half. Process both halves:

- Operation (process) takes constant time c
- Base case takes time d



COMP20003 Algorithms and Data Structures

1-4

Recurrence for divide and conquer sorting algorithms

One pass through the data reduces problem size by **half**. Process **both halves**

- **Operation** takes constant time **c**
- **Base** case takes time **d**

$$\begin{aligned} T(1) &= d \\ T(n) &= 2T(n/2) + nc \\ &= nc + 2cn/2 + 4cn/4 \dots + n/2 \cdot 2c + nd \\ &= c(n-1)\log n + nd \end{aligned}$$

COMP20003 Algorithms and Data Structures

1-5

Divide and Conquer: Recurrences to Master Theorem

- Most common case:

$$T(n) = 2T(n/2) + n$$

- General case:

$$T(n) = aT(n/b) + f(n)$$

$$f(n) \in \Theta(n^d)$$

- Most common case:

$$T(n) = 2T(n/2) + n$$

$$a=2, b=2, d=1$$

COMP20003 Algorithms and Data Structures

4-6

Master Theorem for Divide and Conquer

- $T(n) = aT(n/b) + f(n)$
 $f(n) \in \Theta(n^d)$
- $T(n)$ closed form varies, depending on whether:
 - $d > \log_b a$ $T(n) \in \Theta(n^d)$
 - $d = \log_b a$ $T(n) \in \Theta(n^d \log n)$
 - $d < \log_b a$ $T(n) \in \Theta(n^{\log_b a})$

COMP20003 Algorithms and Data Structures

4-7

Master Theorem for Divide and Conquer

- $T(n) = aT(n/b) + f(n)$, where
 $a \geq 1, b > 1, n^d$ asymptotically positive
- $T(n)$ closed form varies, depending on whether:
 - $d > \log_b a$ $T(n) \in \Theta(n^d)$
 - $d = \log_b a$ $T(n) \in \Theta(n^d \log n)$
 - $d < \log_b a$ $T(n) \in \Theta(n^{\log_b a})$

COMP20003 Algorithms and Data Structures

4-8

Where do $\Theta()$ solutions to the Master Theorem come from?

$$T(n) = aT(n/b) + f(n), \quad f(n) \in \Theta(n^d)$$

Size of subproblems decreases by b

- So base case reached after $\log_b n$ levels
- Recursion tree $\log_b n$ levels

Branch factor is a

- At k th level, have a^k subproblems

At level k , total work is then

- $a^k * O(n/b^k)^d$
- (#subproblems * cost of solving one)



1-9

Where do $\Theta()$ solutions to the Master Theorem come from?

$$T(n) = aT(n/b) + f(n), \quad f(n) \in \Theta(n^d)$$

• At level k , total work is then

- $a^k * O(n/b^k)^d = O(n^d) * (a/b^d)^k$
 - As k (levels) goes from 0 to $\log_b n$, this is a geometric series, with ratio a/b^d
- $$\sum O(n^d) * (a/b^d)^k$$



COMP20003 Algorithms and Data Structures

1-10

Where do $\Theta()$ solutions to the Master Theorem come from?

$$T(n) = aT(n/b) + f(n), \quad f(n) \in \Theta(n^d)$$

- Geometric series: $O(n^d) * (a/b^d)^k$
 - as k goes from $0 \rightarrow \log_b n$
- Case 1: ratio $a/b^d < 1$
 - $(a/b^d)^k$ gets smaller as k goes from $1 \rightarrow \log n$
 - a/b^d First term is the largest, and is < 1
 - $O(n^d)$




COMP20003 Algorithms and Data Structures

1-11

Example for $a/b^d < 1$

$$T(n) = 2T(n/2) + n^2$$



433-253 Algorithms and Data Structures

4-12

Where do the solutions to the Master Theorem come from?

$$T(n) = aT(n/b) + f(n), \quad f(n) \in \Theta(n^d)$$

- Geometric series: $O(n^d) * (a/b^d)^k$
 - as k goes from $0 \rightarrow \log_b n$
- Case 2: ratio $a/b^d = 1$
 - Series is $O(n^d) + O(n^d) + \dots$
 - For $\log_b n$ levels
 - Sum = $O(n^d \log n)$

COMP20003 Algorithms and Data Structures


1-13

Example for most common case $a/b^d = 1$

$$T(n) = 2T(n/2) + n$$

$$\begin{aligned} T(n) &= 2(2T(n/4) + n/2) + n \\ &= 4T(n/4) + n + n \\ &= 8T(n/8) + n + n + n \end{aligned}$$

....



1 * n
2 * n/2
4 * n/4
....

433-253 Algorithms and Data Structures

4-14

Where do $\Theta()$ solutions to the Master Theorem come from?

$$T(n) = aT(n/b) + f(n), \quad f(n) \in \Theta(n^d)$$


- Geometric series: $O(n^d) * (a/b^d)^k$
 - as k goes from $0 \rightarrow \log_b n$
- Case 3: ratio $a/b^d > 1$
 - $a/b^d > 1 \rightarrow$ series is increasing
 - Sum dominated by last term:
 - $O(n^d)(a/b^d)^{\log(b)n} = n^{\log(b)a}$

COMP20003 Algorithms and Data Structures

1-15

Example for $a/b^d > 1$

$$T(n) = 4T(n/2) + n$$



433-253 Algorithms and Data Structures

4-16



- For more on geometric series, and calculation of closed form, see:
<http://www.youtube.com/watch?v=JJZ-shHiayU>
- 4 minute tutorial from Rose-Hulman Institute of Technology