

Binary Number System

(Continued from previous slide..)

Example

$$\begin{aligned} 10101_2 &= (1 \times 2^4) + (0 \times 2^3) + (1 \times 2^2) + (0 \times 2^1) + (1 \times 2^0) \\ &= 16 + 0 + 4 + 0 + 1 \\ &= 21_{10} \end{aligned}$$

Oct al N um ber Syst em

(Continued from previous slide..)

§ Since there are only 8 digits, 3 bits = 8)
(2³) are sufficient to represent any octal number in binary

Exam ple

$$\begin{aligned} 2057_8 &= (2 \times 8^3) + (0 \times 8^2) + (5 \times 8^1) + (7 \times 8^0) \\ &= 1024 + 0 + 40 + 7 \\ &= 1071_{10} \end{aligned}$$

Hexadecimal Number System

Characteristics

- § A positional number system
- § Has total 16 symbols or digits (0, 1, 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E, F). Hence its base = 16
- § The symbols A, B, C, D, E and F represent the decimal values 10, 11, 12, 13, 14 and 15 respectively
- § The maximum value of a single digit is 15 (one less than the value of the base)

(Continued on next slide)

Hexadecimal Number System

(Continued from previous slide..)

- § Each position of a digit represents a specific power of the base (16)
- § Since there are only 16 digits, 4 bits ($2^4 = 16$) are sufficient to represent any hexadecimal number in binary

Example

$$\begin{aligned} 1AF_{16} &= (1 \times 16^2) + (A \times 16^1) + (F \times 16^0) \\ &= 1 \times 256 + 10 \times 16 + 15 \times 1 \\ &= 256 + 160 + 15 \\ &= 431_{10} \end{aligned}$$

Converting a Number of Another Base to a Decimal Number

Method

- Step 1: Determine the column (positional) value of each digit
- Step 2: Multiply the obtained column values by the digits in the corresponding columns
- Step 3: Calculate the sum of these products

(Continued on next slide)

Converting a Number of Another Base to a Decimal Number

(Continued from previous slide..)

Example

$$4706_8 = ?_{10}$$

$$\begin{aligned} 4706_8 &= 4 \times 8^3 + 7 \times 8^2 + 0 \times 8^1 + 6 \times 8^0 \\ &= 4 \times 512 + 7 \times 64 + 0 + 6 \times 1 \\ &= 2048 + 448 + 0 + 6 \\ &= 2502_{10} \end{aligned}$$

Common values multiplied by the corresponding digits

← Sum of these products

Converting a Decimal Number to a Number of Another Base

(Continued from previous slide..)

Example

$$952_{10} = ?_8$$

Solution:

8	952	Remainder
	<hr/>	
	119	0
	<hr/>	
	14	7
	<hr/>	
	1	6
	<hr/>	
	0	1

$$\text{Hence, } 952_{10} = 1670_8$$

Converting a Number of Some Base to a Number of Another Base

Method

- Step 1: Convert the number to a decimal (base 10)
- Step 2: Convert the decimal number so obtained to the new base number

(Continued on next slide)

Converting a Number of Some Base to a Number of Another Base

(Continued from previous slide..)

Example

$$545_6 = ?_4$$

Solution:

Step 1: Convert from base 6 to base 10

$$\begin{aligned} 545_6 &= 5 \times 6^2 + 4 \times 6^1 + 5 \times 6^0 \\ &= 5 \times 36 + 4 \times 6 + 5 \times 1 \\ &= 180 + 24 + 5 \\ &= 209_{10} \end{aligned}$$

(Continued on next slide)

Converting a Number of Some Base to Another Base

to a Number

(Continued from previous slide..)

Step 2: Convert 209_{10} to base 4

4	209	Remainders
	52	1
	13	0
	3	1
	0	3

Hence, $209_{10} = 3101_4$

So, $545_6 = 209_{10} = 3101_4$

Thus, $545_6 = 3101_4$

Short cut Method for Converting a Binary Number to its Equivalent Octal Number

Method

- Step 1: Divide the digits into groups of three starting from the right
- Step 2: Convert each group of three binary digits to one octal digit using the method of binary to decimal conversion

(Continued on next slide)

Short cut Method for Converting a Binary Number to its Equivalent Octal Number

(Continued from previous slide..)

Example

$$1101010_2 = ?_8$$

Step 1: Divide the binary digits into groups of 3 starting from right

001101010

Step 2: Convert each group into one octal digit

$$001_2 = 0 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = 1$$

$$101_2 = 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = 5$$

$$010_2 = 0 \times 2^2 + 1 \times 2^1 + 0 \times 2^0 = 2$$

$$\text{Hence, } 1101010_2 = 152_8$$

Short cut Method for Converting an Octal Number to its Equivalent Binary Number

Method

Step 1: Convert each octal digit to a 3 digit binary number (the octal digits may be treated as decimal for this conversion)

Step 2: Combine all the binary groups (of 3 number digits each) into a single binary

(Continued on next slide)

Short cut Method for Converting an Octal Number to its Equivalent Binary Number

(Continued from previous slide..)

Example

$$562_8 = ?_2$$

Step 1: Convert each octal digit to 3 binary digits

$$5_8 = 101_2, \quad 6_8 = 110_2, \quad 2_8 = 010_2$$

Step 2: Combine the binary groups

$$562_8 = \begin{array}{ccc} \underline{101} & \underline{110} & \underline{010} \\ 5 & 6 & 2 \end{array}$$

$$\text{Hence, } 562_8 = 101110010_2$$

Short cut Method for Converting a Binary

- Method**
- Step 1: Divide the binary digits into groups of four starting from the right
- Step 2: Combine each group of four binary digits to one hexadecimal digit

(Continued on next slide)

Short cut Method for Converting a Binary Number to its Equivalent Hexadecimal Number

(Continued from previous slide..)

Example

$$111101_2 = ?_{16}$$

Step 1: Divide the binary digits into groups of four starting from the right

0011 1101

Step 2: Convert each group into a hexadecimal digit

$$0011_2 = 0 \times 2^3 + 0 \times 2^2 + 1 \times 2^1 + 1 \times 2^0 = 3_{10} = 3_{16}$$

$$1101_2 = 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = 13_{10} = D_{16}$$

$$\text{Hence, } 111101_2 = 3D_{16}$$

Short cut Method for Converting a Hexadecimal Number to its Equivalent Binary Number

Method

Step 1: Convert the decimal equivalent of each hexadecimal digit to a 4 digit binary number

Step 2: Combine all the resulting binary groups (of 4 digits each) in a single binary number

(Continued on next slide)

Short cut Method for Converting a Hexadecimal Number to its Equivalent Binary Number

(Continued from previous slide..)

Example

$$2AB_{16} = ?_2$$

Step 1: Convert each hexadecimal digit to a 4 digit binary number

$$\begin{aligned} 2_{16} &= 2_{10} = 0010_2 \\ A_{16} &= 10_{10} = 1010_2 \\ B_{16} &= 11_{10} = 1011_2 \end{aligned}$$

Short cut Method for Converting a Hexadecimal Number to its Equivalent Binary Number

(Continued from previous slide..)

Step 2: Combine the binary groups

$$2AB_{16} = \underline{0010} \quad \underline{1010} \quad \underline{1011}$$

2 A B

$$\text{Hence, } 2AB_{16} = 001010101011_2$$

Formation of Fractional Numbers in Binary Number System (Example)

	Binary Point ↓									
Position	4	3	2	1	0	.	-1	-2	-3	-4
Position Value	2^4	2^3	2^2	2^1	2^0		2^{-1}	2^{-2}	2^{-3}	2^{-4}
Quantity Represented	16	8	4	2	1		$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$

(Continued on next slide)

Formation of Fractional Numbers in Binary Number System (Example)

(Continued from previous slide..)

Example

$$\begin{aligned} 110.101_2 &= 1 \times 2^2 + 1 \times 2^1 + 0 \times 2^0 + 1 \times 2^{-1} + 0 \times 2^{-2} + 1 \times 2^{-3} \\ &= 4 + 2 + 0 + 0.5 + 0 + 0.125 \\ &= 6.625_{10} \end{aligned}$$

Formation of Fractional Numbers in Octal Number System (Example)

	Octal Point ↓							
Position	3	2	1	0	.	-1	-2	-3
Position Value	8^3	8^2	8^1	8^0		8^{-1}	8^{-2}	8^{-3}
Quantity Represented	51264		8	1		$\frac{1}{8}$	$\frac{1}{64}$	$\frac{1}{512}$

(Continued on next slide)

Formation of Fractional Numbers in Octal Number System (Example)

(Continued from previous slide..)

Example

$$\begin{aligned} 127.54_8 &= 1 \times 8^2 + 2 \times 8^1 + 7 \times 8^0 + 5 \times 8^{-1} + 4 \times 8^{-2} \\ &= 64 + 16 + 7 + \frac{5}{8} + \frac{4}{64} \\ &= 87 + 0.625 + 0.0625 \\ &= 87.6875_{10} \end{aligned}$$