SILVICULTURE

1.----------the branch of forestry which deals with the establishment, development, care and reproduction of stands of timber
Ans : Silviculture

2 the study of life history and general characteristics of forest trees and crops with particular reference to environmental factors as the basis for the practice of silviculture
Ans : Silvics

3. The natural forest urinfluenced by human activity it may or may not be a climax forest is known as
Ans : Virgin forest

4. To renew a forest crop by natural or artificial means is known as
Ans : Regeneration

5. The renewal of forest crop by self sown seed or by coppice or root suckers is
Ans : Natural regeneration

6. A species which is native to a specified area or region is called
Ans : Indigenous

7. The theory and practice of all the constitutes the creation, conservation and scientific management of forest and utilization of their resources is known as
Ans : Forestry

8. The practice of forestry with the object of obtaining the maximum in volume and quality of products per unit area through the application of the best techniques of silviculture and management is
Ans Intensive forestry

9. The practice of forestry with the object of producing timber and other forest products as a business enterprise is known as
Ans : Commercial forestry

10. The practice of forestry on lands outside the conventional forest area for the benefit of the rural and urban communities is called
Ans : Social forestry

11. The practice of forestry on farmlands is called
Ans : Farm forestry

12. The practice of forestry away from the conventional forest area is
Ans : Extension forestry

13. The plant whose stem is always green and tender and height is usually not more than one meter is
Ans : Herb

14. A shrub is usually not more than meters in height
Ans : 6 meters

15. A large woody perennial plant having a single well defined stem and a more or less definite crown
Ans : Tree

16. The leaves of most conifers are needle shaped and that is why they are called as
Ans Needles

17. The branches originating in clusters from dormant or adventitious buds on the trunk of a tree is
Ans Epicormic branches

18. The condition of the tree with dead upper branches projecting above the green portion of the crown is
Ans : Stagheaded

19. The irregular involutions and swellings on the bole which is just above the basal swell of tree is termed as
Ans : Fluting

20. The roots produced from parts of the plants other than the radicle or its subdivision is called as
Ans : Adventitious roots

21. The special roots produced from its branches and adventitious roots remain suspended in the air till they reached the ground is called as
Ans : Prop roots

22. The adventitious roots which emerge from the butt of a tree above ground level is termed as
Ans : Stilt roots

23. A knee shaped or spike like projection of the roots of swamp tree is known as
Ans : Pneumatophore roots

24, The composite structure or invaded rootlet is called
Ans Mycorrhiza

25. The modified sterns developed from double accessory buds in the axils of cotyledons is termed as
Ans : Lignotubers

26. Lignotubers commonly present in
Ans : Eucalyptus

27. The mutual relationship between microorganisms is termed as
Ans : Symbiosis

28. The science that deals with the time of appearance of characteristics periodic events such as leaf shedding is called
Ans : Phenology

29. The growth in trees is confined only in certain regions is called
Ans : Growing points

30. The width of the growth rings indicates the rate of growth, while the fast growth being indicated by rings wiier than
Ans : 5 mm

31. The diameter at the top of the stem is generally about at the base
Ans : Half the diameter

32. The trees which produce male and female flowers on the same individual is termed as
Ans : Monoecious

33. The trees which produce male and female flowers on different individuals is called as
Ans Dioecious

34. The trees have both unisexual and bisexual flowers is termed as
Ans : Polygamous

35. Pollination of flowers in trees is take place by
Ans : Wind, insects and birds

36. The flowering of one or a few culms in a clump or a few clumps of bamboo in a locality is called as
Ans Sporadic flowering

37. The plants which have naked seeds belong to the subdivision of
Ans : Gymnosperms

38. The plants in which the seed are enclosed in the fruits belong to a sub-division of
Ans : Angiosperms

39. The aggregation of trees occupying a specific area sufficiently uniform in composition is termed as
Ans : Stand

40. The study of plants or animals in relation to their environment is known as
Ans : Ecology

41. The forest which are regenerated from seed are called as
Ans : High forest

42, The forest which are regenerated by some vegetative method are called as
Ans : Coppice forest

43. An uneven aged crop containing theoretically all age classes or gradations is called as
Ans Selection forest

44. A forest composed of almost entirely of one species usually to the extent of not less than 80% is known as
Ans : Pure forest

45. A species of less value than the principal species is termed as
Ans Accessories species

46. Auxiliary species is otherwise called as
Ans : Secondary or subsidiary species

47. A forest which is managed primarily for its produce is called
Ans : Production forest

48. An area constituted under the Indian Forest Act is known as
Ans : Reserved forest
49. An. area which is subject to limited degree of protection under the provisions of chapter IV of the Indian Forest Act is called as
Ans : Protected forest

50. A state forest assigned to a village — community under the proviions of the Indian Forest Act is called as
Ans : Village forest

51. All trees which form the upper most leaf canopy and have their leading shoots free is called as
Ans : Dominant tree

52. The average height of a regular crop is determined by
Ans : Loreys formula

53. A table showing the distribution of stems by diameter classes is called as
Ans : Stand table

54. The uppermost storey of a storyed high forest is
Ans : Overwood

55. The lowest stratum of woody and other vegetation above the ground cover is called as
Ans Under growth

56. All the biotic and abiotic factors of a sited is termed as
Ans : Environment

57. The average weather conditions prevalent in any locality is known as
Ans : Climate

58. The sum of the effective environmental conditions under which an organism lives is known as
Ans : Habitat

59. The maximum energy of solar radiation occurs in the region of
Ans : Green and yellow region

60. Without light, plants become pale yellow and have long thin internodes, a condition known as
Ans Etiolation

61. The estimated light used in photosynthesis is less than
Ans: 2%

62. The relative length of day and night to which the plant is exposed is called as
Ans : Photoperiod

63. The response of the plant to photoperiod is called as
Ans : Photoperiodism

64. Frost occurring on nights with clear sky produced by loss of heat by radiation is called as
Ans : Radiation frost

65. The accumulation to a considerable depth of heavy cold air flowing down into natural depressions from adjoining areas is
Ans : Pool frost

66. A frost produced by cold air brought from elsewhere is called as
Ans : Advective frost

67. A definite relatively localized necrotic lesion primarily of the bark and cambium is known as
Ans : Canker

68. The natural process by which plants become adapted to drought ,cold or heat is known as
Ans : Hardening

69. A month which has less than 50 mm of rainfall is usually referred as
Ans : Dry month

70 is a day on which a minimum of *2.5* mm of rain fall is received
Ans : Rainy day

71. The climate which is modified or adjusted by climatic factors is called
Ans : Bioclimate

72. The direction towards which a slope faces is called as
Ans Aspect

73. The relation of a site to weather conditions especially sun and wind is
Ans Exposure

74. The climate under a plant or other cover is referred as
Ans : Microclimate

75. Ecological influence on characteristics of the soil brought about by its physical and chemical characteristics is
Ans : Edaphic factors

76. A mature soil developed in cool and humid climates by the process of podsolization is called as
Ans : Podsol

77. The relative proportion of the various size groups of individual soil particles is known as
Ans : Soil texture

78. A group of soils having same range in particle size and physical properties based on texture is called as
Ans : Soil class

79. The arrangement of individual soil particles into aggregates of definite size and shape is is known as
Ans : Soil structure

80. The extent to which the gross volume of the soil is unoccupied by solid particles is called as
Ans : Soil porosity

81. A *plant* that can subsist in dry situations is termed as
Ans Xerophyte

82. A plant whose normal habitat is neither very wet nor very dry is called as
Ans : Mesophyte

83. A plant which grows in moist sites is
Ans : Hygrophyte

84. A plant which grows wholly or partially immersed in water is
Ans : Hydrophyte

85, The total capacity of the soil i'or holding cations is referred as
Ans : Cation exchange capacity

86. The ratio of the weight of organic carbon to the weight of total nitrogen in a soil or organic material is called as
Ans : C/N ratio

87, An organism that draws a part or whole of its nourishment from another living organism is

known as -
Ans : Parasites

88. A plant growing on other plant but not nourished by another plant is
Ans : Epiphytes

89. An organism on or within which another organism feeds and develops is termed as_____
Ans : Host

90. The gradual replacement of one community by another in the development of vegetation towards a climax is
Ans : Plant succession

91. The culmination stage in plant succession for a given environment is known as
Ans : Climax

92. Tropical or subtropical grass land containing scattered trees or shrubs is
Ans : Savannah

93. The renewal of a forest crop by natural or artificial means is known as
Ans : Regeneration

94. The renewal of a forest crop by self — sown seed or by coppice or root suckers is known as
Ans : Natural regeneration

95. When regeneration is obtained from seed forms then the crop is called as
Ans : Seedling crop

96. A condition of mature viable seed in which germination is considerably delayed, even though external conditions favour germination is due to
Ans : Seed dormancy

97. The potential capacity of a seed to germinate is called as
Ans : Viability

98. The percentage by number of seeds in a given sample that actually germinate irrespective of time is known as
Ans : Germinative capacity

99. The percentage by number of seeds in a given sample that have germinated upto the time when the rate of germination reaches its peak is known as
Ans : Germinative energy

100. Percentage of the number of the seeds in a sample that develop into seedlings at the end of the growing season is referred as
Ans : Plant percent

101. The shoot portion of seedlings keeps on dying year after year while the root portion remains alive this phenomenon is termed as
Ans : Dying back of seedlings

102. The progressive dying usually backwards from the tip of any portion of the plant is known as
Ans : Die back

103. The period which elapses between the initiation of natural regeneration and the time when it considered safe from adverse influences such as frost, drought or weeds is called as
Ans : Seedling establishment period

104. Seedlings, saplings and poles of species of the over wood that have become established naturally in a forest before regeneration felling's are called as
Ans : Advance growth

105. The coppice shoots arising from the base of the seedlings that have been cut or burnt back is
Ans : Seedling coppice

106. The first stage of regeneration felling under a shelter wood system is
Ans : Seeding felling

107. The removal of the last seed or shelter trees after regeneration has been effected under a shelter wood system is known as
Ans : Final felling

108. A regeneration felling carried out between the seeding felling and the final felling under a

shelter wood system is called as
Ans : Secondary felling

109. The mature crop is removed in a series of operation in which the first stage is seeding felling and the last stage is the final felling is refered as
Ans : Shelterwood system

110. The treatment or handling of slash for reducing hazards from fire, insects or fungi and for providing the seeds with access to thç soil is called as
Ans Slash disposal

111. The time that elapses between successive main fellings on the same area is
Ans : Felling cycle

112. The annual or periodic removal of exploitable trees in an uneven aged forest is known as
Ans : Selection felling

113. A term used to indicate the removal only of certain species of high value is termed as
Ans : Selective felling

114. A survey for the assessment of established and unestablished regeneration generally by sample enumeration is
Ans : Regeneration survey

115. A shoot risng from the root of a woody plant is
Ans : Root sucker

116. Any unrooted portion of a plant used for grafting or budding on a rooted stock is called
Ans : Scion

117. A rooted plant on which a scion is grafted is called
Ans : Stock

11 8. The coppice shoots arising from the base of seedlings that have been Cut cr burnt hcIc ic
Ans : Seedling coppice

119. The coppice arising from the stool or a living stump is called as
Ans : Stool coppice

120. A trees whose stem has been cut off in order to obtain a flush of shoots usually above the height to which the browsing animals can reach is known as
Ans : Pollarding

121. An operation done in a coupe following the main felling is termed as
Ans : Subsidiary felling

122. The removal of less valuable trees in a crop in the interest of better growth of the more valuable individuals is
Ans : Improvement felling

123. A forest crop raised artificially either by sowing or planting is called as
Ans Plantation

124. An area where plants are raised for eventual planting out is known as
Ans : Nursery

125. Restocking of a felled or otherwise cleared woodland by artificial means is
Ans : Reforestation

126. The establishment of a forest by artificial means on an area from which forest vegetation has always or long been absent is
Ans : Afforestation

127. The height increment of 60 cm per annum in the earlier stages of its life and which gives a minimum yield of 10 m3 per hectare per annum in a short rotation of 10 to 15 years.
Ans : Fast growing species

128. The secondary species is mixed with the main species only for a part of the rotation is called as
Ans : Temporary mixture

129. The sowing of seed in drills or single lines is known as
Ans : Line sowing

130. The distance between the plants put out in a plantation or standing in a crop is called as
Ans : Spacing

131. The plantations in which forest crops are raised along with agricultural crops are known as
Ans Taungyas

132. Taungyas is a word
Ans : Burmese

133. Taungyas is otherwise called in tamil
Ans : Kumari

134. In Taungyas each family has hectares of land for cultivation
Ans : 0.8 to 1.6 hectares

135. All cuttings and grafts derived from one tree are collectively called as
Ans : Clone

136. Preparing seedlings in a nursery for planting out by gradually reducing water, shade and shelter resulting in hardening of the plant is called as
Ans : Hardening

137. The killing of the young seedling by certain fungi that cause decay of the stem or roots is called as
Ans Damping off

138. Turning a crop into manure under while green or soon after maturity for the purpose of soil improvement is called as
Ans : Green manuring

139. To mark with stakes the position of plants in a plantation is called as
Ans : Staking out

140. Stump planting is commonly practiced in
Ans : Teak

141. Restocking the blanks in an artificially regenerated area with fresh sowings or plantings is

called as
Ans : Beating up or casuality replacement

142. A crop of trees or shrubs grown to foster the growth of another and more important tree crop in its early stages is known as
Ans Nurse crop

143. A subsidiary crop of low plants introduced in a plantation to afford soil cover between or below the main crop is known as
Ans Cover crop

144. Sowing or planting under an existing stand is
Ans : Under planting

145. A register in which all details of works carried out in a plantation and other points of interest concerning it are recorded from time to time
Ans : Plantation journal

146. Counter trenches are usually made on slopes upto %
Ans 20 %

147. The practice of forestry in all its aspect on farm and village lands generally more or less integrated with other farm operations is called
Ans : Farm forestry

148. An operation carried out for the benefit of a forest crop at any stage of its life between the seedling and the mature stages is called
Ans : Tending operation

149. A tending operation done in the seedling stage in nursery or in a forest crop that involves the removal or cutting back of all weeds is known as
Ans : Weeding

150. A tending operation done in a sapling crop involving the removal or topping of inferior growth including individuals of favoured species is called as
Ans : Cleaning

151. A felling made in an immature stand for the purpose of improving the growth and form of the trees that remain without permanently breaking the canopy is called
Ans : Thinning operation

152. A thinning in which the trees to be cut are selected by some rule of thumb is called
Ans : Mechanical thinning

153. The mechanical thinning is otherwise called as
Ans Stick thinning

154. The mechanical thinning is applied to
Ans : Regular crop

155. The method of thinning in common use that consists in the removal of inferior individuals of a crop is called as
Ans : Ordinary thinning

156. Ordinary thinning is otherwise called as
Ans : German thinning

157. A method in which thinning is primarily directed to the dominant trees in a regular crop is known as
Ans Crown thinning

158. Crown thinning is otherwise called as
Ans : French thinning

159. Free thinning is developed by
Ans : Heck

160. The free thinning is otherwise called as
Ans : Elite thinning

161. The method of thinning which aims at putting as high a proportion as possible of the total potential increment of the area is called as
Ans : Maximum thinning

162. A thinning done in a regular crop in anticipation of suppression is called as
Ans : Advance thinning

163. Advance thinning is developed by
Ans : Craib

164. The stand density index was developed by
Ans : Reineke (1933)

165. The planned interval which elapses between successive thinnings in the same area is called as
Ans : Thinning cycle

166. As an indication in numerical terms the extent to which the crop is thinned is known as
Ans : Thinning intensity

167. The whole set of thinnings carried out in a crop from the earliest stage to maturity is called as
Ans : Thinning regime

168. A method of thinning directed to obtain or maintain selection composition in a crop with all diameter classes adequately represented is known as
Ans : Selection thinning

169. The removal of less valuable trees in a crop in the interest of better growth of the more valuable individuals is called as
Ans : Improvement felling

170. A cutting through bark and outer living layers of wood in a continuous incision all round the bole of a tree is known as
Ans : Girdling

171. The removal of live or dead branches or multiple leaders from standing trees for the improvement of the tree or its timber is known as
Ans : Pruning

172. The natural death and fall of branches of standing trees from such causes as deficiency of light, decay, snow and ice is called as
Ans : Natural pruning

173. The rubbing off the lateral buds to prevent the development of branches as a measure to obtain knot free timber is called as
Ans : Bud pruning

174. CVP index refers to
Ans : Climate, Vegetation and Productivity index

175. A plant which tend to be restricted to calcarious soil is called as
Ans : Calciphile

176. Tropical dry evergreen forests are found in
Ans : Carnatic coast

177. Silvicultural system prevalent during early part of century was
Ans : Clear felling system

178. High forest refers to
Ans : Forests regenerated through seeds

179. Pure forest are those which cover
Ans : 80% of the crop consists of a single species

180. Champion and Seth classified the Indian Forests into major groups
Ans : Sixteen groups

181. Which of the following law is used to estimate the proportionate distribution of trees in selection forests?
Ans : Dc Liocourts law

182. In South India teak was worked under
Ans : Selection system

183. The rotation followed for sandal wood in South India is
Ans : Physical rotation

184. OIeo resins are obtained from
Ans *: Pinus roxburghii*

185. Cutch is obtained from
Ans *: Acacia catechu*

186. Rate of photosynthesis increases with increase in temperature up to
Ans : 30°C

187. Intensity of light on the southern aspects compare to northern aspects will be
Ans : Higher

188. Tetrazolium test for testing viability of seeds is attributable to
Ans : Lakon

189. The exotic tree which has natural hybrids in India is
Ans : Eucalyptus

190. The tree that serves as raw material for the manufacture of pencil is
Ans : Juniperus

191, An example for wood oil is
Ans : *Cedrus deodara*

192. French thinning is otherwise called as
Ans : Crown thinning

193. Maximum thinning was developed by
Ans : Gehrhardt

194. Indian butter tree is
Ans : *Madhuca latifolia*

195. Fruit-tan is obtained from
Ans *Terminalia chebula*

196. Tumbling is the method for
Ans : Extraction of cones

197 Dry strobile is seen in
Ans : Fir

198. 'Chilgoza' is obtained from
Ans *Pinus gerardiana*

199. The leader of the 'Chipko' movement is
Ans : Sunderlal Baguhuna

200. World Forest Day is celebrated on
Ans : 21st March

201. Epicormic branches originate from
Ans : Adventious buds

202. Prop roots is a common phenomenon in
Ans : Ficus

203. Based on the methods of regeneration, forests are classified as forests
Ans : High and coppice

204. Trees which do not form part of the upper most leaf canopy but the leading of shoots,which are not definitely overtopped by the neighbouring trees are called as
Ans : Dominated

205. The mean annual temperature of montane sub-tropical forest ranged between
Ans : 17°C to 24°C

206. A soil formed from parent material *in situ* is called as
Ans : Sedentary soil

207. The term 'forest succession' was first used by
Ans : Thoreau

208. Soil host properties are influenced mainly by parental material is called
Ans : Endodynamorphic

209. An organism on or within which another organism feeds and develops is called as
Ans : Host

210. The C:N ratio considered suitable for measuring nitrate status of the soil is
Ans : 10 : 1

211. Nitrogn fixing non-leguminous tree species is
Ans : Casuarina

212. A forest fire spreading slowly over the ground with low flame is termed as
Ans : Creeping fire

213. The productive capacity of Indian Forest is
Ans : 0.5 m3/ha

214. The climax stage of estuarine succession is
Ans : Evergreen forest

215. If the canopy density is between 0.75 and 1.0, then the forest is referred as
Ans Dense

216. The Latin name for Andaman Padauk tree is
Ans *: Plerocarpus da!bergioides*

217. The Latin name for golden bamboo is
Ans : *Bambusa vulgaris*

218. Nitrogen fixing ability of Casuarina is attributable to
Ans : Frankia

219. Site index is the relation between
Ans : Age and height

220. In north eastern hill region, shifting cultivation is called as
Ans : Jhum

221. The clear felling or clear strip system is suitable for
Ans Light demanders

222. The poly climax theory in plant succession was developed by
Ans : Tansley

223. Fungal roots are otherwise called
Ans : Mycorrhiza

224. The Father of Indian Forestry is
Ans : Dietrich Brandis

225. A method of silvicultural procedure worked out in accordance with accepted sets of silvicultural principles by which crops constituting forests are tended, harvested and replaced by new crops of distinctive forms is called as
Ans : Silvicultural system

226. The silvicultural system in which the regeneration is normally of seedling origin either natural or artificial or combination and where the rotation is generally long is termed as
Ans : High forest system

227. The silvicultural systems in which the crop originates mainly from coppice and where the rotation of the coppice is short is
Ans : Coppice system

228. A change from one silvicultural system to another is necessitated and this is usually referred to as
Ans : Conversion

229. The period during which the change from one silvicultural system to another is effected as
Ans : Conversion period

230 a silvicultural system in which equal or equi — productive areas of mature crop are successively clear felled in one operation
Ans : Clear felling system

231. A vigorous tree usually of bad form occupying more space than its future value warrants and threatening potentially better neighbours is called as
Ans : Wolf tree

232. A plantation map showing areas suitable for different species is
Ans : Suitability map

233. The clear strip system is also called as
Ans : Progressive strip system

234. A subdivision of felling series formed with the object of regulating fellings in some special manner is
Ans : Cutting section

35. The regeneration is obtained under shelter of the over wood such system are called as
Ans : Shelter wood system

236. In shelter wood system over wood in two or more operation these are also known as
Ans : System of successive regeneration fellings

237. The canopy is uniformly opened up over the whole area of a compartment to obtain uniform

regeneration
Ans : Uniform system

238. The uniform system is also called as
Ans : Shelterwood compartment or compartment system

239. A felling made with a view to inviting or assisting regeneration under a shelterwood system is
Ans : Regeneration felling

240. Opening the canopy of a mature stand to provide conditions for securing regeneration from the seed of trees retained for the purpose is called
Ans : Seeding felling

241. The first stage of regeneration felling under a shelterwood system is
Ans : Seeding felling

242. A regeneration felling carried, out between the seeding felling and the final felling is
Ans : Secondary felling

243. The removal of the last seed or shelter trees after regeneration has been effected under a shelter wood system is called
Ans : Final felling

244. The interval between the seeding felling and final felling on a particular area under one of the shelterwood systems is called
Ans : Regeneration interval

245. The period required to regenerate the whole of a periodic block is known as
Ans : Regeneration period

246. A felling made under a high forest system usually towards the end of the rotation with the object of creating conditions favourable to seed production and natural regeneration is called as
Ans : Preparatory fellings

247. A silvicultural system in which regeneration fellings are carried out uniformally all ove the compartment in scattered groups is called as
Ans : Group system

248. The group system is developed by
Ans : Karl gayer

249. The group system also known as
Ans : Gayers or Bavarian femelschlag

250. In group system canopy has a wavy profile usually known as
Ans : Gayers line

251. A silvicultural system in which regeneration felling are done in the form of strips successively from one side of the compartment is called as
Ans Shelterwood strip system

252. In wagners blendersaumschlag system in which regeneration feflings are carried out in narrow strips extending in direction and advancing from
Ans East to west direction and north to south

253, The strip and group system is a modification of the system
Ans : Strip and group system

254. The wedge system is developed by
Ans : Eberhard

255. The regeneration fellings are on the pattern of group system but as the regeneration period is long and the crop produced is uneven aged or irregular is termed as
Ans : Irregular shelterwood system

256. A silvicultural system in which the crop to be regenerated is opened up irregularly and the crop is uneven aged is called as
Ans : Indian irregular shelterwood system

257. The Indian irregular shelterwood system is otherwise called as
Ans : Punjab shelterwood system

258. A silvicultural system in which fellings and regeneration is distributed over the whole area and all age of trees are found mixed together is called as
Ans : Selection system

259. The time that elapses between successive main fellings on the same area is known as
Ans : Felling cycle

260. •The group selection system is suitable for
Ans : Light demanders

261. In two storeyed high forest system a crop of trees in which the canopy can be differentiated into
Ans : Two strata

262. It is not silvicultural system as it neither aims at regenerating the crop nor producing a crop of distinctive characteristics is known as
Ans : Improvement felling

263. The improvement felling is applied to
Ans : Mixed uneven aged forests

264. In some situations improvement felling is combined with selection felling is referred as
Ans : Selection cum improvement felling

265. The high forest is developed from
Ans : Seed

266. The silvicultural system in which the new crop is originates mainly from stool coppice is called as
Ans : Coppice system

267. In coppice system generally the rotation of the coppice is
Ans : Short

268. A silvicultural system based on stool coppice in which the old crop is clear felled without any reservation of trees is called as
Ans : Simple coppice system

269. In coppice system the number of shoots is reduced only to one per stool the special operation known as
Ans : Singling out operation

270. The coppice of two rotation system is modification of
Ans : Simple coppice system

271. The shelterwood coppice system is modification of system
Ans : Simple coppice system

272. A silvicultural system in which fellings are carried out on the principles of selection system but regeneration is obtained by
Ans Coppice

273. The tree stem has been cut off in order to obtain a flush of shoots is called as
Ans : Pollarding

274. Deodar is managed under
Ans : Shelter wood system

275. Scientific name of toon
Ans *: Toona ciliala*

276. *Eucalyptus* is a
Ans : Good coppicer

277. Tendu is a source of
Ans : Beedi Leaf

278. Sandal is commonly affected by
Ans : Spike disease

279. *Terminalias* belong to the family
Ans : Combretaceae

280. Ligno tuber is present in
Ans : Eucalyptus

281. *Prosopis juflflora* is native of
Ans Mexico

282. “Mimosin” is present in
Ans : Leucaena

283. An important Multi-purpose tree in coastal area
Ans : Casuarina

284. Which is the family of Bamboo
Ans : Poaceae

285. Casuarina root nodules are formed in association with
Ans : Frankia

286. *Shorea robusta* is susceptible to
Ans : Drought

287. *Santalum album* is a
Ans : Partial root parasite

288. *Mesuaferrea* belongs to
Ans : Guttiferae

289. Most suitable Silvicultural system of deodar in hilly terrain for control of soil erosion is
Ans Selection system

290. The Silviculture of Indian Trees book was written by
Ans : Banerjee

291. Neem is found to occur wild in India in
Ans : Western ghats

292. The scientific name of nagasampigo is
Ans *Mesuaferrea*

293. Fragrant flowers characterized for
Ans : *Michelia champaca*

294, Very strong durable wood is
Ans : *Mesuaferrea*

295. Commonest companion tree of Sal is
Ans : *Buchanania latifolia*

296. Iron wood of Malabar is
Ans : *Hopea parviflora*

297. Native of *Swietenia mahagoni* is
Ans : West Indies

298. *Dalbergia sissoo* grows most typically in
Ans : Alluvial

299. Forest of *Acacia arabica* are frequently termed as
Ans : Babul meadow

300. Calorific value of wood is highest in
Ans : *Casuarina equisettfolla*

301. The growing points consist of cells which have the capacity to divide and give rise to new cells.
Ans : Meristematic cells

302. Kail is common name of
Ans : *Pinus wallichiana*

303. Example for presence of root nodules in trees
Ans : *C. equistifolia*

304. The meristematic cells found in the apices of shoot and root are called as
Ans : Apical meristems

305. *Vitex negundo* is belongs to family
Ans Verbenaceae

306. *Hopea parviflora* belongs to the family
Ans : Dipterocarpaceae

307. *Scientific* name of yellow bamboo is
Ans *: Bambusa vulgaris*

308. Sal is a
Ans Good coppicer

309. *Dalbergia sissoo* is
Ans : Frost hardy

3 10. *Atteva fabriciella* is a common pest of
Ans *: Ailanthus excelsa*

311. Silk *cotton* tree belongs to the family
Ans : Malvaceae

312 is a good *pollarding* ability tree
Ans *: Hardwickia binata*

313, Santalin is present in
Ans *: Pterocarpus santalinus*

314. Mesquite bean is the common name for
Ans *: Prosopisjuhflora*

315, *Azadirachta indica* belongs to family
Ans *:* Meliaceae

3 16. *Eucalyptus tereticornis* belongs to family of
Ans *:* Myrtaceae

317. Dry teak forest is associated with
Ans *: Terminalia tomentosa*

318. *Gmelina arborea* belongs to family
Ans *:* Verbenaceae

319. *Pinus wallichiana* is a
Ans : 5 needle pine

320. Origin of *Mangfera indica* is
Ans *:* India

321. Indian *Copal* tree is
Ans *: Vateria indica*

322. *Terminalia arfuna* belongs to family of
Ans : Combretaceae

323. Karanj *oil* is extracted from
Ans: *Pongamia pinnata*

324. CVP *index* on forest productivity does not include
Ans : Edaphic factors

325. Presence of chloroxylon in moist deciduous forest is
Ans : Pre climax

326. Total *geographical* area in India is
Ans : 328.7 m.ha.

327. Red sanders occur on soil
Ans : Quartzite

328. *Lignotubers* present in
Ans : Eucalyptus

329. The *direction* towards which a slope faces is called
Ans : Aspect

330. The *climate* of small area is
Ans : Micro climate

331. Scientific name of rose wood is
Ans : *Dalbergia 1atfo!oia*

332. Non-legume which fixes atmospheric nitrogen is
Ans : Frankia

333. Epiphytes is
Ans : *Ficus bengalensis*

334. Sal borer is
Ans *: Haplocerambyx spinicornis*

335. Example of annual flowering *bamboo* is
Ans : *Ochiandra stridula*

336. The time which elapses between formation to final felling is called
Ans : Rotation

337. Stilt roots present in
Ans : Mangroves

338. *Cedrus deodara* is
Ans : Dioecious

339. Teak pollination occurs by
Ans : Insects

340, Afternoon shower is also known as
Ans : instability rain

341. Teak grows on soil
Ans : Lime rich soil

342. The pH of alkali soil is
Ans : > 8.5

343. Plants grows in moist site *is* called
Ans : Hygrophytes

344. Scientific name of mesquite is
Ans : *Prosopisjuliflora*

345. Teak defoliator is
Ans *: Hybleae purea*

346. Tree seed every year is
Ans : Teak and shisham

347. Scientific name of Iron wood is
Ans : *Xylia xylocarpa*

348. Lianas are
Ans : Woody climber

349 is a Evergreen tree
Ans : Neem

350. Regeneration obtained through seeds is called
Ans : High forest

351. Restocking of degraded areas are known as
Ans : Reforestation

352. Sal grows well in
Ans : Quartzite soil

353. The gradual replacement of one plant community by the another is known as
Ans : Succession

354. Configuration of land surface is
Ans : Topography

355. Forest conservation act was implemented in the year
Ans : 1980

356. FSI is located at
Ans : Dehradun

357. Who proposed mono climax theory?
Ans : Clements

358. The up and down movement of wind in valley is known as
Ans : Dadu

359. Indian forest act was implemented in the year
Ans : 1927

360. Who is known as Father of Indian Forestry?
Ans : Sir Deidritch Brandis

361. Who is known as first silviculturist of India?
Ans : Troup (1909)

362. First forest policy of Independent India was introduced in
Ans : 1952

363. Plant community involved in succession is called
Ans : Sere

364. Culmination stage in forest development is known as
Ans : Climax

365. Apical meristerm is responsible for
Ans : Height growth

366. Lateral meristerm is responsible for
Ans Diameter growth

367 is the distance north or south of equator
Ans Latitude

368. Elevation of an object from the known level is called
Ans : Altitude

369 % of light is used for photosynthesis in plants
Ans : Less than 2%

370. Nematophores are common in
Ans Mangrove sp.

371. The first species to grow in a site is called as
Ans Pioneer sp.

372. Who proposed Mosaic theory?
Ans : Aubrevilles

373. To move nursery stock from one part of nursery to another in order to obtain young trees suitable for planting out is called as
Ans : Transplant

374. How many hot spots are there in India?
Ans : 2

375. hA/C 1 denotes forest type
Ans : Shola forest

376. Earth's albedo is
Ans : 42 %

377. Symbiotic bacteria is
Ans Rhizobium

378. Amount of sunlight actually used in photosynthesis is
Ans : 2 per cent

379. The type of frost common in Dehradun valley is
Ans : Pool frost

380. Pneumatophore is special root type present in
Ans : Roots of swamp tree

381. Chilling of air below the freezing point is termed as
Ans : Frost

382. Controlled Burning is a
Ans : Cultural operation

383. Indian biodiversity contribution to world biodiversity is
Ans 7%

384. Stage immediately preceding the climax stage in forest succession is known as
Ans : Pre climax

385. Stage more exacting the climax stage in forest succession is called as
Ans : Post climax

386. The revised classification of forest type was proposed by s
Ans : Champion and Seth

387. How many forest types are there in the world?
Ans : 8

388. How many major forest types are there in India?
Ans : 5

389. Change in colour of leaf to yellow due to inadequate light is known as
Ans : Etiolation

390. The relative productive capacity of the species is known as
Ans Site quality

391. Who coined the word Ecology?
Ans : Hackel

392. First biosphere reserve in India is
Ans : Nilgiri Biosphere reserve

393. Vanamahotsava started in the year
Ans 1952

394. Scientific name of red gum is
Ans *: Eucalyptus tereticornis*

395. First National Park in India is
Ans Hailey/Corbett

396. Scientific name of Spruce is
Ans : *Spicea smithiana*

397, First bird sanctuary in India is
Ans : Vedanthangal

398. The regeneration fellings which are distributed over the whole felling series is called as
Ans : Selection system

399.. Salix in Kashmir valley is worked under
Ans : Pollard system

400, Integration of commodity production with demands on forests to provide other value is called as
Ans : Multiple use management

401, An appropriate silvicultural system should be
Ans : Optimizing the yield, improve the quality and shorten the investment period

402. RIL stands for
Ans : Reduced impact logging

403. Yellow or Golden Bamboo is called as
Ans *: Bamboosa vulgaris*

404. Species which is highly suitable for making cooperage is
Ans : Quercus spp

405. What percentage of regeneration of a sample plot could be taken for a successful regeneration of that area?
Ans : 60 -70%

406. A series of thinning carried out in the life of a teak plantation is called as
Ans : Thinning regime

407. Palmrosa oil is a
Ans : Grass oil

408. Commercial cultivation of *Saussuria lappa* is found in the state of
Ans Himachal Pradesh

409. The national afforestation programme is being implemented through a tier system
Ans : 2

410. *Eucalyptus globules* plantation in the Nilgiri hills was commenced in the year
Ans : 1843

411. Introduction of teak in Srilanka was done by Van Rhede in the year
Ans 1680

412. *Dendrocalamus strictus* is a good associate of
Ans : Teak

413. Institute of Tropical forestry is situated at
Ans : Puerto Rica

414. The beef wood is also known as
Ans : Casuarina

415. Origin of Eucalyptus is
Ans : Australia

416. The family of *Acacia catechu* is
Ans : Leguminaceae

417. Teak has a
Ans : Large inflorescence

418. *Sesbania grandiflora* is a
Ans : Leguminous tree

419. As per Benthem and Hookers' classification, there are non-standard exceptions to the family name ending 'aceae'.
Ans : 8

420. The natural system of classification is based on
Ans : form

421. The flower lacking either stamen or pistil is called as
Ans : Incomplete flower

422. The largest family among the Indian flora is
Ans : Asteracea

423. The origin of teak is
Ans : Burma

424. The arrangement of ovules within the ovary is termed as
Ans : Placentation

425. Ergrostis ergrostis is an example for
Ans : Tautonym

426. A pair of appendages that present at the point of attachment of a leaf to the stem is
Ans : Stipules

427. Thick and fleshy root without any particular shape is
Ans : Tuberous

428. Sucking roots of parasite is known as
Ans : Haustoria

429. The specimen or other element designated by the author or used by him as the nomenclatural type is called
Ans biotype

430, Father of Botany is
Ans : Theophrastus

431. If all the petals are free, the condition is called
Ans : Polypetale

432. The alternative name of Guttiferae is
Ans : Clusiaceae

433. *Acacia nilotica* belongs to the family
Ans Mimoceae

434. Peeling of bark is a characteristic feature of
Ans : Myrtaceae

435. *Santalum album* is found in
Ans : Dry deciduous forests

436. Name the Indian Rubber plant
Ans *: Ficus elastica*

437. Pellucid dots are characteristic of the family
Ans : Myrtaceae

438. The clear felled strips progress successively in one direction from one end of the forest to another is known as
Ans : Clear strip system

439. The length and breadth of standard nursery bed is
Ans : 12 m length and 1.2 m width

440. The cotyledons are pushed upwards by the rapid elongation of the hypocotyls is called type of germination.
Ans : Epigeous

441. The cotyledons are remain in the soil or just on its surface of earth is called type of germination
Ans : Hypogeous

442. The stem is induced to rooting, when it is still attached to the parent plant is called as
Ans : Layering

443. The technique of connecting two pieces of living plant tissue together so that they will unite and form a functional plant is called as
Ans: B

444. The micropropogation is synonymous of
Ans : in-vitro propogation

445, A technique of propagating plants under a controlled environment using aseptic plant cells. tissues in a artificial growing medium is called as
Ans : Micropropagation

446. The selection forest is otherwise called as
Ans All aged forest

447. The trees are felled in small group stand not as scattered single trees of the typical selection system is called as
Ans : Group selection system

448. The improvement fellings is usually applied to forest
Ans : Mixed uneven aged forest

449. The fellings are carried out on the principle of selection system but regeneration is obtained by coppice and the resultant crop is uneven aged is
Ans Coppice selection system

450. The silvicultural system which is adopted for bamboo forest is
Ans : Cuims selection system

451. Sampling intensity in regeneration survey is
Ans : 2-4 %

452, The branch of forestry aims to get maximum income per unit area is called as
Ans : Intensive forestry

453. Felling debris such as branches, leaves and twigs are called as
Ans Slash

454. Improvement felling is one of the opertion
Ans Tending

455. Two types of layering is and
Ans : air layering and soil layering

456. The succession found in extremely dry location is
Ans : Xerarch succession

457. Monkey puzzle tree is
Ans *: Araucaria imbricata*

458. The wavy profile of the canopy obtained in group system is called as
Ans : Gayer's line

459. Floating periodic block is also known as
Ans : Single periodic block

460. The felling area in selection system is
Ans : Coupe

461. The new crop produced under clear felling system is
Ans : Absolutely even aged

462. The wedge shape of strips are produced in system
Ans : Eberhard Wedge system

463. The character of the new crop produced in bamboo forest is
Ans : Uneven aged

464. A felling area, usually one of an annual series is called as
Ans : Coupe

465. The method of raising forest plantation in combination with agricultural crops is called as
Ans : Taungya cultivation

466. Mycorrhizal symbiosis helps plant to provide
Ans : Phosphorus

467. Which is a known as a very good soil for the all type of plants is
Ans Sandy loam

468. *Populous deltoids* is introduced in India. The origin of this species is which country
Ans : United kingdom

469. Mean annual temperature of tropical zone is about
Ans : Over 24°C

470. The crown cover of green leaves and branches formed by the crowns of all trees in a forest is
Ans Canopy

471. The portion of trees in a stand below the overstory is
Ans understory

472. Cheapest method of site preparation is
Ans : Prescribed burning

473 World’s first Teak Forest was planted in 1842 in Kerala at place.
Ans : Nilambur

474 The Native Place of Redwood trees is
Ans : California

475 In aerial seeding seeds are very effective.
Ans : Pellatised seeds

476. Which among these practices is an important tool for regeneration of moist sal

Ans : Weeding

477, Thinning of valuable timber trees in a mixed forest plantation is called as
Ans : Selection thinning

478. What helps in protecting forests through promoting responsible and better management practices is
Ans : Forest certification

479 type of rotation is most favourable for naural reproduction?
Ans : Silvicultural rotation

480 system is called as Andaman’s canopy lifting shelter wood system.
Ans : Indian irregular shelter wood system

481 The system in which regeneration is obtained from seed is system.
Ans : High forest

482 is also known as single periodic block.
Ans : Floating periodic block

483. The National Afforestation and Eco-Development Board (NAEB), set up in Year.
Ans 1992

484 In shelterwood strip system felling is done against the direction of
Ans : Wind

485 is a function of height, diameter and the form of the tree.
Ans : Volume growth

486 *Acacia Lenticularis* is found in the state of
Ans : Sikkim

487 System is a compromise between selection system and shelter wood group system.
Ans : Irregular shelter wood

488. *Celtis australis* is found in type of forest.
Ans : Western Himalaya forests

489. The flowering of trees is affected by factors.
Ans : Internal and external factors

490 Trees have mostly type of flower.
Ans : Bisexual flowers

491 indigenous poplar species.
Ans *: Populus ciliata*

492. *Acacia albida* is tree.
Ans : Multipurpose tree

493. The name Kannimara is given as the word
Ans : Virgin

494. Scientific name of kadamba
Ans *: IVeolamarckia cadamba*

495. Tectona grandis belong to the family of
Ans : Lamiaceae

496. Lagerstroemia microcarpa is an endemic tree of
Ans : The Western Ghats

497. The world’s first Teak Museum is located at place.
Ans : Nilambur in the Malappuram district of Kerala

498. Nilambur is renowned for the oldest teak plantation in the world is planted by

Ans H.V. Conolly

499 Teak is one of the world’s largest living teak tree found in Parambikulam.
Ans : Kannimara

500. Kannimara teak tree has been awarded by the Government of India.
Ans Mahavriksha Puraskar