


Adv. Macro: Heterogenous Agent Models

Jeppe Druedahl & Patrick Moran 2022







• Teachers: Jeppe Druedahl and Patrick Moran

- Teachers: Jeppe Druedahl and Patrick Moran
- Central economic questions:
  - 1. What explains the level and dynamics of heterogeneity/inequality?
  - 2. What role does heterogeneity play for understanding consumption-saving dynamics in partial equilibrium?
  - 3. What role does heterogeneity play for understanding business-cycle fluctuations in general equilibrium?

- Teachers: Jeppe Druedahl and Patrick Moran
- Central economic questions:
  - 1. What explains the level and dynamics of heterogeneity/inequality?
  - 2. What role does heterogeneity play for understanding consumption-saving dynamics in partial equilibrium?
  - 3. What role does heterogeneity play for understanding business-cycle fluctuations in general equilibrium?
- Central technical method: Programming in Python

Prerequisite: Intro. to Programming and Numerical Analysis

**Complicated:** Close to the research frontier

- Teachers: Jeppe Druedahl and Patrick Moran
- Central economic questions:
  - 1. What explains the level and dynamics of heterogeneity/inequality?
  - 2. What role does heterogeneity play for understanding consumption-saving dynamics in partial equilibrium?
  - 3. What role does heterogeneity play for understanding business-cycle fluctuations in general equilibrium?
- Central technical method: Programming in Python
   Prerequisite: Intro. to Programming and Numerical Analysis
   Complicated: Close to the research frontier
- Plan for today:
  - 1. More about the course
  - 2. Dynamic programming theory
  - 3. Dynamic programming practice

### Model components:

- 1. Optimizing individual agents (households + firms)
- 2. Idiosyncratic and aggregate risk
- 3. Information flows (who knows what when  $\Rightarrow$  often everything)
- 4. Market clearing (Walras vs. search-and-match)

### Model components:

- 1. Optimizing individual agents (households + firms)
- 2. Idiosyncratic and aggregate risk
- 3. Information flows (who knows what when  $\Rightarrow$  often everything)
- 4. Market clearing (Walras vs. search-and-match)

#### Insurance:

 $Complete 
ightarrow idiosyncratic risk insured away \sim representative agent Incomplete 
ightarrow$  agents need to self-insure

### Model components:

- 1. Optimizing individual agents (households + firms)
- 2. Idiosyncratic and aggregate risk
- 3. Information flows (who knows what when  $\Rightarrow$  often everything)
- 4. Market clearing (Walras vs. search-and-match)

#### Insurance:

 $Complete 
ightarrow idiosyncratic risk insured away \sim representative agent Incomplete 
ightarrow$  agents need to self-insure

### Heterogeneity:

Ex ante in preferences, abilities etc.

Ex post after realization of idiosyncratic shocks

### Model components:

- 1. Optimizing individual agents (households + firms)
- 2. Idiosyncratic and aggregate risk
- 3. Information flows (who knows what when  $\Rightarrow$  often everything)
- 4. Market clearing (Walras vs. search-and-match)

#### Insurance:

 $Complete 
ightarrow idiosyncratic risk insured away \sim representative agent Incomplete 
ightarrow$  agents need to self-insure

### Heterogeneity:

Ex ante in preferences, abilities etc.

Ex post after realization of idiosyncratic shocks

• HANC: Heterogeneous Agent Neo-Classical model

### Model components:

- 1. Optimizing individual agents (households + firms)
- 2. Idiosyncratic and aggregate risk
- 3. Information flows (who knows what when  $\Rightarrow$  often everything)
- 4. Market clearing (Walras vs. search-and-match)

#### Insurance:

 $Complete 
ightarrow idiosyncratic risk insured away \sim representative agent Incomplete 
ightarrow agents need to self-insure$ 

### Heterogeneity:

Ex ante in preferences, abilities etc.

Ex post after realization of idiosyncratic shocks

- HANC: Heterogeneous Agent Neo-Classical model
- HANK: Heterogeneous Agent New Keynesian model (i.e. include price and wage setting frictions)

- **Lectures:** Thursday 10-13
  - 2 hours of »normal« lecture
  - 1 hour of active problem solving (no exercise classes)

- Lectures: Thursday 10-13
  - 2 hours of »normal« lecture
  - 1 hour of active problem solving (no exercise classes)

#### Content:

- 1. Explanation of computational methods
- 2. Discussion of research papers
- Examples of code for central mechanisms (you should run the notebook codes simultaneously)

- Lectures: Thursday 10-13
  - 2 hours of »normal« lecture
  - 1 hour of active problem solving (no exercise classes)

### Content:

- 1. Explanation of computational methods
- 2. Discussion of research papers
- Examples of code for central mechanisms (you should run the notebook codes simultaneously)

#### Material:

Web: sites.google.com/view/numeconcph-advmacrohet/ Git: github.com/numeconcopenhagen/adv-macro-het

- Lectures: Thursday 10-13
  - 2 hours of »normal« lecture
  - 1 hour of active problem solving (no exercise classes)

### Content:

- 1. Explanation of computational methods
- 2. Discussion of research papers
- Examples of code for central mechanisms (you should run the notebook codes simultaneously)

#### Material:

Web: sites.google.com/view/numeconcph-advmacrohet/ Git: github.com/numeconcopenhagen/adv-macro-het

#### Code:

- 1. We provide code you will build upon
- 2. Based on the GEModelTools package

• Individual assignments (hand-in on Absalon)

- Individual assignments (hand-in on Absalon)
  - Assignment based on a HANC-model)
     Deadline: 14th of October (must be approved before exam)

- Individual assignments (hand-in on Absalon)
  - 1. Assignment based on a HANC-model)
    - Deadline: 14th of October (must be approved before exam)
  - 2. Assignment based on a **HANK** -model
    - Deadline: 25th of November (must be approved before exam)

- Individual assignments (hand-in on Absalon)
  - 1. Assignment based on a **HANC-model**)

<u>Deadline</u>: 14th of October (must be approved before exam)

2. Assignment based on a **HANK** -model

<u>Deadline</u>: 25th of November (*must be approved before exam*)

3. **Free assignment** with essay on a relevant model extension of own choice and a simple implementation

Deadline for proposal: 9th of December

Deadline for peer feedback: 16th of December (exam requirement)

- Individual assignments (hand-in on Absalon)
  - 1. Assignment based on a HANC-model)

<u>Deadline</u>: 14th of October (*must be approved before exam*)

- 2. Assignment based on a HANK -model
  - <u>Deadline</u>: 25th of November (*must be approved before exam*)
- Free assignment with essay on a relevant model extension of own choice and a simple implementation

Deadline for proposal: 9th of December

Deadline for peer feedback: 16th of December (exam requirement)

All feedback can be used to improve assignments before the exam

- Individual assignments (hand-in on Absalon)
  - 1. Assignment based on a HANC-model)

<u>Deadline</u>: 14th of October (*must be approved before exam*)

- 2. Assignment based on a **HANK** -model
  - <u>Deadline</u>: 25th of November (*must be approved before exam*)
- 3. **Free assignment** with essay on a relevant model extension of own choice and a simple implementation

Deadline for proposal: 9th of December

Deadline for peer feedback: 16th of December (exam requirement)

- All feedback can be used to improve assignments before the exam
- Exam:
  - 1. Hand-in 3×assignments
  - 2. 48 hour take-home: Programming of new extension
    - + analysis of model + interpretation of results

### **Python**

- Assumed knowledge: From Introduction to Programming and Numerical Analysis you are assumed to know the basics of
  - 1.1 Python
  - 1.2 JupyterLab
  - 1.3 VSCode
  - 1.4 git
- 2. Updated Python: Install (or re-install) newest Anaconda
- 3. Packages: pip install quantecon, EconModel, consav
- 4. **GEMoodel tools:** 
  - 4.1 Clone the GEModelTools repository
  - 4.2 Locate repository in command prompt
  - 4.3 Run pip install -e .

# Course plan

See CoursePlan.pdf

# Knowledge

- 1. Account for, formulate and interpret precautionary saving models
- 2. Account for stochastic and non-stochastic simulation methods
- Account for, formulate and interpret general equilibrium models with ex ante and ex post heterogeneity, idiosyncratic and aggregate risk, and with and without pricing frictions
- 4. Discuss the difference between the stationary equilibrium, the transition path and the dynamic equilibrium
- Discuss the relationship between various equilibrium concepts and their solution methods
- Identify and account for methods for analyzing the dynamic distributional effects of long-run policy (e.g. taxation and social security) and short-run policy (e.g. monetary and fiscal policy)

### Skills

- 1. Solve precautionary saving problems with dynamic programming and simulate behavior with stochastic and non-stochastic techniques
- 2. Solve general equilibrium models with ex ante and ex post heterogeneity, idiosyncratic and aggregate risk, and with and without pricing frictions (stationary equilibrium, transition path, dynamic equilibrium)
- 3. Analyze dynamics of income and wealth inequality
- 4. Analyze transitional and permanent structural changes (e.g. inequality trends and the long-run decline in the interest rate)
- Analyze the dynamic distributional effects of long-run policy (e.g. taxation and social security) and short-run policy (e.g. monetary and fiscal policy)

### Competencies

- Independently formulate, discuss and assess research on both the causes and effects of heterogeneity and risk for both long-run and short-run outcomes
- 2. Discuss and assess the importance of how heterogeneity and risk is modeled for questions about both long-run and short-run dynamics

**Dynamic Programming** 

■ Budget constraint for  $t \in \{0, 1, ..., T-1\}$ 

$$assets_t = (1 + return rate) \times assets_{t-1} + wage \times productivity_t - consumption_t$$

$$a_t = (1 + r)a_{t-1} + wz_t - c_t$$

■ Budget constraint for  $t \in \{0, 1, ..., T-1\}$ 

$$\mathsf{assets}_t = (1 + \mathsf{return}\,\mathsf{rate}) \times \mathsf{assets}_{t-1} + \mathsf{wage} \times \mathsf{productivity}_t - \mathsf{consumption}_t$$

$$\mathsf{a}_t = (1 + r)\mathsf{a}_{t-1} + \mathsf{wz}_t - \mathsf{c}_t$$

• Borrowing constraint:  $a_t \ge 0$ 

■ Budget constraint for  $t \in \{0, 1, ..., T-1\}$ 

$$\mathsf{assets}_t = (1 + \mathsf{return}\,\mathsf{rate}) \times \mathsf{assets}_{t-1} + \mathsf{wage} \times \mathsf{productivity}_t - \mathsf{consumption}_t \\ a_t = (1 + r)a_{t-1} + wz_t - c_t$$

- Borrowing constraint:  $a_t \ge 0$
- Static problem:
  - 1. **Information:**  $z_t$  is known for all t
  - 2. Target: Discounted utility,  $\sum_{t=0}^{T-1} \beta^t u(c_t)$ ,  $\beta > 0$
  - 3. **Behavior:** Choose  $c_0, c_1, \ldots, c_{T-1}$  simultaneously
  - 4. **Solution:** Sequence of consumption *choices*  $c_0, c_1, \ldots, c_{T-1}$

■ Budget constraint for  $t \in \{0, 1, ..., T-1\}$ 

$$\mathsf{assets}_t = (1 + \mathsf{return}\,\mathsf{rate}) \times \mathsf{assets}_{t-1} + \mathsf{wage} \times \mathsf{productivity}_t - \mathsf{consumption}_t \\ a_t = (1 + r)a_{t-1} + wz_t - c_t$$

- Borrowing constraint:  $a_t \ge 0$
- Static problem:
  - 1. **Information:**  $z_t$  is known for all t
  - 2. Target: Discounted utility,  $\sum_{t=0}^{T-1} \beta^t u(c_t)$ ,  $\beta > 0$
  - 3. **Behavior:** Choose  $c_0, c_1, \ldots, c_{T-1}$  simultaneously
  - 4. **Solution:** Sequence of consumption *choices*  $c_0, c_1, \ldots, c_{T-1}$
- Dynamic programming:
  - 1. **Information:**  $z_t$  is revealed period-by-period
  - 2. Target: Expected discounted utility,  $\sum_{t=0}^{T-1} \beta^t \mathbb{E}_t[u(c_t)], \ \beta > 0$
  - 3. Behavior: Choose  $c_t$  sequentially as information is revealed
  - 4. **Solution:** Sequence of consumption functions,  $c_t^{\star}(z_t, a_{t-1})$

### Static solution: IBC

Substitution implies Intertemporal Budget Constraint (IBC)

$$a_{T-1} = (1+r)a_{T-2} + wz_{T-1} - c_{T-1}$$

$$= (1+r)^2 a_{T-3} + (1+r)wz_{T-2} - (1+r)c_{T-1} + wz_{T-1} - c_{T-1}$$

$$= (1+r)^T a_{-1} + \sum_{t=0}^{T-1} (1+r)^{T-1-t} (wz_t - c_t)$$

### Static solution: IBC

Substitution implies Intertemporal Budget Constraint (IBC)

$$a_{T-1} = (1+r)a_{T-2} + wz_{T-1} - c_{T-1}$$

$$= (1+r)^2 a_{T-3} + (1+r)wz_{T-2} - (1+r)c_{T-1} + wz_{T-1} - c_{T-1}$$

$$= (1+r)^T a_{-1} + \sum_{t=0}^{T-1} (1+r)^{T-1-t} (wz_t - c_t)$$

• Use **terminal condition**  $a_{T-1} = 0$  (equality due utility max.)

$$(1+r)^{-(T-1)}a_{T-1}=0 \Leftrightarrow b_0+h_0-\sum_{t=0}^{T-1}(1+r)^{-t}c_t=0$$

where 
$$b_0 = (1+r)a_{-1}$$
 and  $h_0 \equiv \sum_{t=0}^{T-1} (1+r)^{-t} w z_t$ 

# Static solution: FOC and consumption function

$$\mathcal{L} = \sum_{t=0}^{T-1} \beta^t \frac{c_t^{1-\rho}}{1-\rho} + \lambda \left[ \sum_{t=0}^{T-1} (1+r)^{-t} c_t - b_0 - h_0 \right]$$

First order conditions:

$$\forall t : 0 = \beta^t c_t^{-\rho} - \lambda (1+r)^{-t} \Leftrightarrow c_t^{-\rho} = \beta (1+r) c_{t+1}^{-\rho}$$

# Static solution: FOC and consumption function

$$\mathcal{L} = \sum_{t=0}^{T-1} \beta^t \frac{c_t^{1-\rho}}{1-\rho} + \lambda \left[ \sum_{t=0}^{T-1} (1+r)^{-t} c_t - b_0 - h_0 \right]$$

First order conditions:

$$\forall t : 0 = \beta^t c_t^{-\rho} - \lambda (1+r)^{-t} \Leftrightarrow c_t^{-\rho} = \beta (1+r) c_{t+1}^{-\rho}$$

Insert Euler into IBC to get consumption choice

$$\sum_{t=0}^{T-1} (1+r)^{-t} (\beta(1+r))^{t/\rho} c_0 = b_0 + h_0 \Leftrightarrow$$

$$c_0 = \frac{1 - (\beta(1+r))^{1/\rho}/(1+r)}{1 - ((\beta(1+r))/(1+r))^T} (b_0 + h_0)$$

### Static solution: FOC and consumption function

$$\mathcal{L} = \sum_{t=0}^{T-1} \beta^t \frac{c_t^{1-\rho}}{1-\rho} + \lambda \left[ \sum_{t=0}^{T-1} (1+r)^{-t} c_t - b_0 - h_0 \right]$$

First order conditions:

$$\forall t : 0 = \beta^t c_t^{-\rho} - \lambda (1+r)^{-t} \Leftrightarrow c_t^{-\rho} = \beta (1+r) c_{t+1}^{-\rho}$$

Insert Euler into IBC to get consumption choice

$$\sum_{t=0}^{T-1} (1+r)^{-t} (\beta(1+r))^{t/\rho} c_0 = b_0 + h_0 \Leftrightarrow$$

$$c_0 = \frac{1 - (\beta(1+r))^{1/\rho}/(1+r)}{1 - ((\beta(1+r))/(1+r))^T} (b_0 + h_0)$$

• Question: Is this the solution correct?

• In words: An optimal policy has the property that whatever the initial state and initial decision are, the remaining decisions must constitute an optimal policy with regard to the state resulting from the first decision. (See Bellman, 1957, Chap. III.3.)

- In words: An optimal policy has the property that whatever the initial state and initial decision are, the remaining decisions must constitute an optimal policy with regard to the state resulting from the first decision. (See Bellman, 1957, Chap. III.3.)
- In math:

- In words: An optimal policy has the property that whatever the initial state and initial decision are, the remaining decisions must constitute an optimal policy with regard to the state resulting from the first decision. (See Bellman, 1957, Chap. III.3.)
- In math:
  - 1. Value function,  $v_t$ : Defined recursively from

$$v_t(z_t, a_{t-1}) = \max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

with  $v_T(\bullet) = 0$ .

- In words: An optimal policy has the property that whatever the initial state and initial decision are, the remaining decisions must constitute an optimal policy with regard to the state resulting from the first decision. (See Bellman, 1957, Chap. III.3.)
- In math:
  - 1. Value function,  $v_t$ : Defined recursively from

$$v_t(z_t, a_{t-1}) = \max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

with 
$$v_T(\bullet) = 0$$
.

2. **Policy function,**  $c_t^{\star}$ : Is the same as

$$c_t^*(z_t, a_{t-1}) = \arg\max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

#### Vocabulary

$$v_t(z_t, a_{t-1}) = \max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

- 1. State variables:  $z_t$  and  $a_{t-1}$
- 2. Control variable:  $c_t$
- 3. Continuation value:  $\beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$
- 4. **Parameters:** r, w, and stuff in  $u(\bullet)$

#### Timing of shocks

• Realization of shocks: First in the period before choices are made

#### Timing of shocks

- Realization of shocks: First in the period before choices are made
- Beginning-of-period value function (before realization):

$$\underline{v}_t(z_{t-1}, a_{t-1}) = \mathbb{E}_{t-1}[v_t(z_t, a_{t-1})]$$

#### Timing of shocks

- Realization of shocks: First in the period before choices are made
- Beginning-of-period value function (before realization):

$$\underline{v}_t(z_{t-1}, a_{t-1}) = \mathbb{E}_{t-1}[v_t(z_t, a_{t-1})]$$

End-of-period value function (after realization):

$$v_t(z_t, a_{t-1}) = \max_{c_t} u(c_t) + \beta \underline{v}_{t+1}(z_t, a_{t-1})$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

#### Numerical value function iteration - basics

 Discretization: All state variable belong to discrete monotonically increasing sets 

 = grids,

$$\begin{split} z_t &\in \mathcal{G}_z = \{z^0, z^1, \dots, z^{\#z-1}\} \\ a_t &\in \mathcal{G}_a = \{a^0, a^1, \dots, a^{\#_a-1}\} \end{split}$$

#### Numerical value function iteration - basics

 Discretization: All state variable belong to discrete monotonically increasing sets 

 = grids,

$$z_t \in \mathcal{G}_z = \{z^0, z^1, \dots, z^{\#z-1}\}\$$
  
 $a_t \in \mathcal{G}_a = \{a^0, a^1, \dots, a^{\#a-1}\}\$ 

• Transition probabilities:  $\pi_{i_z-,i_z}=\Pr[z_t=z^{i_z}\,|\,z_t=z^{i_{z-}}]$ 

#### Numerical value function iteration - basics

 Discretization: All state variable belong to discrete monotonically increasing sets 

 = grids,

$$z_t \in \mathcal{G}_z = \{z^0, z^1, \dots, z^{\#z-1}\}\$$
  
 $a_t \in \mathcal{G}_a = \{a^0, a^1, \dots, a^{\#a-1}\}\$ 

- Transition probabilities:  $\pi_{i_z...,i_z} = \Pr[z_t = z^{i_z} \, | \, z_t = z^{i_{z-}}]$
- Linear interpolation (function approximation):
  - 1. Assume  $\underline{v}_{t+1}$  is known on  $\mathcal{G}_z \times \mathcal{G}_a$  (tensor product)
  - 2. Evaluate  $\underline{v}_{t+1}(z^{i_z}, a)$  for arbitrary a by

$$egin{aligned} & \underline{\check{\mathbf{v}}}_{t+1}(\mathbf{z}^{i_{\mathbf{z}}}, \mathbf{a}) = \underline{\mathbf{v}}_{t+1}(\mathbf{z}^{i_{\mathbf{z}}}, \mathbf{a}^{\iota}) + \omega_{i} \frac{\mathbf{a} - \mathbf{a}^{\iota}}{\mathbf{a}^{\iota+1} - \mathbf{a}^{\iota}} \\ & \omega_{i} \equiv \frac{\mathbf{v}_{t+1}(\mathbf{z}^{i_{\mathbf{z}}}, \mathbf{a}^{\iota+1}) - \mathbf{v}_{t+1}(\mathbf{z}^{\iota_{\mathbf{z}}}, \mathbf{a}^{\iota})}{\mathbf{a}^{\iota+1} - \mathbf{a}^{\iota}} \\ & \iota \equiv \operatorname{largest} \ i_{\mathbf{a}} \in \{0, 1, \dots, \#_{\mathbf{a}} - 2\} \ \operatorname{such that} \ \mathbf{a}^{i_{\mathbf{a}}} \leq \mathbf{a} \end{aligned}$$

# **Deriving transition probabilities**

• **Specification:** Assume  $\log z_t$  follows the AR(1) process

$$\log z_{t+1} = \rho_z \log z_t + \psi_{t+1}, \ \psi_{t+1} \sim \mathcal{N}(\mu_{\psi}, \sigma_{\psi})$$

where  $\mu_{\psi}$  is used to ensure  $\mathbb{E}[\mathsf{z}_t] = 1$ 

• Literature: Tauchen (1986), Tauchen and Hussey (1991) and Rouwenhorst (1995) develops method for deriving  $\mathcal{G}_z$  and  $\pi_{i_z,i_z}$  given  $\rho_z$  and  $\sigma_{\psi}$ , but we don't care about the details here

#### Numerical value function iteration - loops

Beginning-of-period value function:

$$\underline{v}_{t}(z^{i_{z-}}, a^{i_{a-}}) = \sum_{i_{z}=0}^{\#_{z}-1} \pi_{i_{z-}, i_{z}} v_{t}(z^{i_{z}}, a^{i_{a-}})$$

#### Numerical value function iteration - loops

Beginning-of-period value function:

$$\underline{v}_t(z^{i_{z-}}, a^{i_{s-}}) = \sum_{i_z=0}^{\#_z-1} \pi_{i_{z-}, i_z} v_t(z^{i_z}, a^{i_{s-}})$$

End-of-period value-of-choice:

$$v_t(z^{i_z}, a^{i_{s-}}|c_t) = u(c_t) + \beta \sum_{i_{z+}=0}^{\#_z-1} \pi_{i_z, i_{z+}} \check{v}_{t+1}(z^{i_{z+1}}, a_t)$$

$$a_t = (1+r)a^{i_{s-}} + wz^{i_z} - c_t$$

#### Numerical value function iteration - loops

Beginning-of-period value function:

$$\underline{v}_t(z^{i_{z-}}, a^{i_{s-}}) = \sum_{i_z=0}^{\#_z-1} \pi_{i_{z-}, i_z} v_t(z^{i_z}, a^{i_{s-}})$$

End-of-period value-of-choice:

$$v_t(z^{i_z}, a^{i_{s-}}|c_t) = u(c_t) + \beta \sum_{i_{z+}=0}^{\#_z-1} \pi_{i_z, i_{z+}} \breve{v}_{t+1}(z^{i_{z+1}}, a_t)$$
 $a_t = (1+r)a^{i_{s-}} + wz^{i_z} - c_t$ 

- Nested loops:
  - 1. **Outer loop:** Backwards in time from t = T 1 (note  $\underline{v}_T$  is known)
  - 2. **Inner loop:** For each grid point in  $\mathcal{G}_z \times \mathcal{G}_a$  find  $c_t^*(z_t, a_{t-1})$  and therefore  $v_t^*(z_t, a_{t-1})$  with a numerical optimizer

#### In practice

- Example-notebook: Introduction.ipynb
  - 1. Introduces EconModel package
  - 2. Show implementation of solution and simulation methods

#### **Numerical Monte Carlo simulation**

• Initial distribution: Draw  $z_{i,-1}$  and  $a_{i,-1}$  for  $i \in \{0,1,\dots,N-1\}$ 

#### **Numerical Monte Carlo simulation**

- Initial distribution: Draw  $z_{i,-1}$  and  $a_{i,-1}$  for  $i \in \{0,1,\ldots,N-1\}$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Draw  $z_{it}$  given transition probabilities
  - 2. Use linear interpolation to evaluate

$$c_{it} = \breve{c}_t^*(z_{it}, a_{it-1})$$
  
 $a_{it} = (1+r)a_{it-1} + wz_{it} - c_{it}$ 

#### **Numerical Monte Carlo simulation**

- Initial distribution: Draw  $z_{i,-1}$  and  $a_{i,-1}$  for  $i \in \{0,1,\ldots,N-1\}$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Draw  $z_{it}$  given transition probabilities
  - 2. Use linear interpolation to evaluate

$$c_{it} = \breve{c}_{t}^{\star}(z_{it}, a_{it-1})$$
  
 $a_{it} = (1+r)a_{it-1} + wz_{it} - c_{it}$ 

- Review:
  - Pro: Simple to implement
  - Con: Computationally costly and introduces randomness

■ Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$ 

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_z=0}^{\#_z-1} \pi_{i_z, i_z} \underline{D}_t(z^{i_z}, a^{i_{a-}})$$

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_{z-}=0}^{\#_z-1} \pi_{i_{z-}, i_z} \underline{D}_t(z^{i_{z-}}, a^{i_{a-}})$$

2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$ 

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv \textit{histogram}$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_z = 0}^{\#_z - 1} \pi_{i_z = ,i_z} \underline{D}_t(z^{i_z}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv \textit{histogram}$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_{z-}=0}^{\#_z-1} \pi_{i_{z-}, i_z} \underline{D}_t(z^{i_{z-}}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do
  - 3.1 Find  $\iota \equiv \text{largest } i_a \in \{0,1,\ldots,\#_a-2\} \text{ such that } a^{i_a} \leq a_t^\star(z^{i_z},a^{i_{a-}})$

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_z = 0}^{\#_z - 1} \pi_{i_z, i_z} \underline{D}_t(z^{i_z}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do
  - 3.1 Find  $\iota \equiv$  largest  $i_a \in \{0,1,\ldots,\#_a-2\}$  such that  $a^{i_a} \leq a_t^\star(z^{i_z},a^{i_{a-}})$
  - 3.2 Calculate  $\omega=rac{a^{\iota+1}-a^*(z^{i_z},a^{\prime a-})}{a^{\iota+1}-a^{\iota}}\in[0,1]$

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_{z-}=0}^{\#_z-1} \pi_{i_{z-}, i_z} \underline{D}_t(z^{i_{z-}}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do
  - 3.1 Find  $\iota \equiv \text{largest } i_a \in \{0,1,\ldots,\#_a-2\}$  such that  $a^{i_a} \leq a_t^\star(z^{i_z},a^{i_a-})$
  - 3.2 Calculate  $\omega=rac{a^{\iota+1}-a^*(z^{i_z},a^{i_{a^-}})}{a^{\iota+1}-a^{\iota}}\in[0,1]$
  - 3.3 Increment  $\underline{m{D}}_{t+1}(z^{i_z},a^\iota)$  with  $\omega m{D}_t(z^{i_z},a^{i_{\vartheta-}})$

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_{z-}=0}^{\#_z-1} \pi_{i_{z-}, i_z} \underline{D}_t(z^{i_{z-}}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do
  - 3.1 Find  $\iota \equiv$  largest  $i_a \in \{0,1,\ldots,\#_a-2\}$  such that  $a^{i_a} \leq a_t^\star(z^{i_z},a^{i_{a-}})$
  - 3.2 Calculate  $\omega=rac{a^{\iota+1}-a^*(z^{i_z},a^{i_{a^-}})}{a^{\iota+1}-a^{\iota}}\in[0,1]$
  - 3.3 Increment  $\underline{\boldsymbol{D}}_{t+1}(\boldsymbol{z}^{i_z}, \boldsymbol{a}^{\iota})$  with  $\omega \boldsymbol{D}_t(\boldsymbol{z}^{i_z}, \boldsymbol{a}^{i_{a-}})$
  - 3.4 Increment  $\underline{\textbf{\textit{D}}}_{t+1}(z^{i_{\it{z}}},a^{\iota+1})$  with  $(1-\omega)\textbf{\textit{D}}_t(z^{i_{\it{z}}},a^{i_{\it{a}-}})$

- Initial distribution: Choose  $\underline{\mathcal{D}}_0(z_{-1}, a_{-1})$ , which is defined on  $\mathcal{G}_z \times \mathcal{G}_a$  and sum to  $1 \equiv histogram$
- **Simulation:** Forwards in time from t = 0 and in each time period
  - 1. Distribute stochastic mass: For each  $i_z$  and  $i_{a-}$  calculate

$$D_t(z^{i_z}, a^{i_{a-}}) = \sum_{i_{z-}=0}^{\#_z-1} \pi_{i_{z-}, i_z} \underline{D}_t(z^{i_{z-}}, a^{i_{a-}})$$

- 2. Initial zero mass: Set  $\underline{D}_{t+1}(z^{i_z}, a^{i_a}) = 0$  for all  $i_{z+}$  and  $i_a$
- 3. Distribute endogenous mass: For each  $i_z$  and  $i_{a-}$  do

3.1 Find 
$$\iota \equiv \text{largest } i_a \in \{0,1,\ldots,\#_a-2\}$$
 such that  $a^{i_a} \leq a_t^{\star}(z^{i_z},a^{i_{a-}})$ 

3.2 Calculate 
$$\omega=rac{a^{\iota+1}-a^*(z^{j_z},a^{j_a-})}{a^{\iota+1}-a^{\iota}}\in[0,1]$$

- 3.3 Increment  $\underline{D}_{t+1}(z^{i_z}, a^{\iota})$  with  $\omega D_t(z^{i_z}, a^{i_{a-}})$
- 3.4 Increment  $\underline{\boldsymbol{D}}_{t+1}(\boldsymbol{z}^{i_z}, \boldsymbol{a}^{\iota+1})$  with  $(1-\omega)\boldsymbol{D}_t(\boldsymbol{z}^{i_z}, \boldsymbol{a}^{i_{\vartheta-}})$
- Review:
  - 1. Pro: Computationally efficient and no randomness
  - 2. Con: Introduces a non-continuous distribution

#### Side-note: Matrix formulation

• The histogram method can be written in **matrix form**:

$$oldsymbol{D}_t = \Pi_z' \underline{oldsymbol{D}}_t \ \underline{oldsymbol{D}}_{t+1} = \Lambda_t oldsymbol{D}_t$$

where

 $\underline{\textbf{\textit{D}}}_{t}$  is vector of length  $\#_{z} \times \#_{a}$ 

 $\boldsymbol{D}_t$  is vector of length  $\#_z \times \#_a$ 

 $\Pi_z'$  is derived from the  $\pi_{i_z,i_z}$ 's

 $\Lambda_t'$  is derived from the  $\iota$ 's and  $\omega$ 's

Note: Example showed in notebook.

#### **Infinite horizon:** $T \to \infty$ ?

$$v_t(z_t, a_{t-1}) = \max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)]$$
  
s.t.  $a_t = (1+r)a_{t-1} + wz_t - c_t \ge 0$ 

- Contraction mapping result: If  $\beta$  is low enough (strong enough impatience) then the value and policy function converge to  $v(z_t, a_{t-1})$  and  $c^*(z_t, a_{t-1})$  for large enough T
- Maximum upper limit for  $\beta$ :  $\frac{1}{1+r}$
- In practice: Solve backwards until value and policy functions does not change anymore (given some tolerance)

# EGM

# **Euler-equation from variation argument**

- Case I: If  $c_t^{-\rho} > \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ : Increase  $c_t$  by  $\Delta > 0$ , and lower  $c_{t+1}$  by (1+r)
  - 1. **Feasible:** Yes, if  $a_t > 0$
  - 2. Utility change:  $\left(c_{t}^{-\rho}\right)+\beta\left(-(1+r)\right)\mathbb{E}_{t}\left[c_{t+1}^{-\rho}\right]>0$

# **Euler-equation from variation argument**

- Case I: If  $c_t^{-\rho} > \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ : Increase  $c_t$  by  $\Delta > 0$ , and lower  $c_{t+1}$  by (1+r)
  - 1. **Feasible:** Yes, if  $a_t > 0$
  - 2. Utility change:  $\left(c_{t}^{-\rho}\right)+\beta\left(-(1+r)\right)\mathbb{E}_{t}\left[c_{t+1}^{-\rho}\right]>0$
- Case II: If  $c_t^{-\rho} < \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ : Lower  $c_t$  by  $\Delta > 0$ , and increase  $c_{t+1}$  by (1+r)
  - 1. Feasible: Yes (always)
  - 2. Utility change:  $\left(-c_t^{-\rho}\right) + \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right] > 0$

# **Euler-equation from variation argument**

- Case I: If  $c_t^{-\rho} > \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ : Increase  $c_t$  by  $\Delta > 0$ , and lower  $c_{t+1}$  by (1+r)
  - 1. **Feasible:** Yes, if  $a_t > 0$
  - 2. Utility change:  $\left(c_{t}^{-\rho}\right)+\beta\left(-(1+r)\right)\mathbb{E}_{t}\left[c_{t+1}^{-\rho}\right]>0$
- Case II: If  $c_t^{-\rho} < \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ : Lower  $c_t$  by  $\Delta > 0$ , and increase  $c_{t+1}$  by (1+r)
  - 1. Feasible: Yes (always)
  - 2. Utility change:  $\left(-c_{t}^{-\rho}\right)+\beta(1+r)\mathbb{E}_{t}\left[c_{t+1}^{-\rho}\right]>0$
- Conclusion: By contradiction
  - 1. Constrained:  $a_t = 0$  and  $c_t^{-\rho} \geq \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$ , or
  - 2. Unconstrained:  $a_t > 0$  and  $c_t^{-\rho} = \beta(1+r)\mathbb{E}_t\left[c_{t+1}^{-\rho}\right]$

# Endogenous grid-point method (EGM)

Alternative to value function iteration:

1. Calculate post-decision marginal value of cash:

$$q(z^{i_z}, a^{i_s}) = \sum_{i_{r+}=0}^{\#_z-1} \pi_{i_z, i_{z+}} c_+ (z^{i_{z+}}, a^{i_s})^{-\sigma}$$

# Endogenous grid-point method (EGM)

Alternative to value function iteration:

1. Calculate post-decision marginal value of cash:

$$q(z^{i_z}, a^{i_s}) = \sum_{i_{r+1}=0}^{\#_z-1} \pi_{i_z, i_{z+}} c_+ (z^{i_{z+}}, a^{i_s})^{-\sigma}$$

2. Invert Euler-equation:

$$c(z^{i_z}, a^{i_a}) = (\beta(1+r)q(z^{i_z}, a^{i_a}))^{-\frac{1}{\sigma}}$$

# Endogenous grid-point method (EGM)

Alternative to value function iteration:

1. Calculate post-decision marginal value of cash:

$$q(z^{i_z}, a^{i_a}) = \sum_{i_{z_+}=0}^{\#_z-1} \pi_{i_z,i_{z_+}} c_+(z^{i_{z_+}}, a^{i_a})^{-\sigma}$$

2. Invert Euler-equation:

$$c(z^{i_z}, a^{i_a}) = (\beta(1+r)q(z^{i_z}, a^{i_a}))^{-\frac{1}{\sigma}}$$

3. Endogenous cash-on-hand:

$$m(z^{i_z}, a^{i_a}) = a^{i_a} + c_+(z^{i_z}, a^{i_a})$$

# Endogenous grid-point method (EGM)

Alternative to value function iteration:

1. Calculate post-decision marginal value of cash:

$$q(z^{i_z}, a^{i_a}) = \sum_{i_{z_+}=0}^{\#_z-1} \pi_{i_z, i_{z_+}} c_+ (z^{i_{z_+}}, a^{i_a})^{-\sigma}$$

2. Invert Euler-equation:

$$c(z^{i_z}, a^{i_a}) = (\beta(1+r)q(z^{i_z}, a^{i_a}))^{-\frac{1}{\sigma}}$$

3. Endogenous cash-on-hand:

$$m(z^{i_z}, a^{i_a}) = a^{i_a} + c_+(z^{i_z}, a^{i_a})$$

4. Consumption function:  $c^*(z^{i_z}, a^{i_{z-}}) = \text{interpolation of function}$  from  $m(z^{i_z}, :)$  to  $c(z^{i_z}, :)$  at  $m = (1 + r)a^{i_{z-}} + wz^{i_z}$ 

# Exercises

## **Exercises: Model extensions**

#### Three exercises for you to do:

- 1. Ensure the stationary distribution is found in the simulation
- 2. Make some borrowing allowed, b
- 3. Introduce transitory shock,  $\xi_t$  (hardest)

#### • Full extended model:

$$\begin{aligned} v_t(z_t, a_{t-1}) &= \max_{c_t} u(c_t) + \beta \mathbb{E}_t[v_{t+1}(z_{t+1}, a_t)] \\ \text{s.t.} \\ a_t &= (1+r)a_{t-1} + wz_t - c_t + \xi_t \\ \log z_{t+1} &= \rho_z \log z_t + \psi_{t+1}, \ \mathbb{E}[z_t] = 1, \mathsf{Var}[\psi_t] = \sigma_\psi^2 \\ \xi_t &\sim \mathcal{N}(0, \sigma_\xi^2) \\ a_t &\geq -b \end{aligned}$$

### Extra: Gauss-Hermite I

• General problem: How can we calculate

$$\mathbb{E}(f(x)) = \int f(x)g(x)dx$$

- $f: \mathbb{R} \to \mathbb{R}$  some function
- g(x) is the probability distribution function (PDF) for x and G(x) is the CDF

### Extra: Gauss-Hermite I

General problem: How can we calculate

$$\mathbb{E}(f(x)) = \int f(x)g(x)dx$$

- $f: \mathbb{R} \to \mathbb{R}$  some function
- g(x) is the probability distribution function (PDF) for x and G(x) is the CDF
- General solution: Turn it into a discrete sum

$$\mathbb{E}(f(x)) \approx \sum_{i=1}^{S} \omega_i f(x_i)$$

### Extra: Gauss-Hermite I

General problem: How can we calculate

$$\mathbb{E}(f(x)) = \int f(x)g(x)dx$$

- $f: \mathbb{R} \to \mathbb{R}$  some function
- g(x) is the probability distribution function (PDF) for x and G(x) is the CDF
- General solution: Turn it into a discrete sum

$$\mathbb{E}(f(x)) \approx \sum_{i=1}^{S} \omega_i f(x_i)$$

• How to choose S and the *nodes*  $(x_i)$  and *weights*  $(\omega_i)$ ? Answer: Guassian quadrature

### Extra: Gauss-Hermite II

• Gauss-Hermite quadrature uses that

$$\int_{-\infty}^{\infty} f(x)e^{-x^2}dx = \sum_{i=1}^{S} \omega_i f(x_i) + \frac{S!\sqrt{\pi}}{s^S(2S)!}f^{(2S)}(\epsilon)$$

for some  $\epsilon$  and where the  $(x_i, \omega_i)$ 's can be easily found

### Extra: Gauss-Hermite II

Gauss-Hermite quadrature uses that

$$\int_{-\infty}^{\infty} f(x)e^{-x^2}dx = \sum_{i=1}^{S} \omega_i f(x_i) + \frac{S!\sqrt{\pi}}{s^5(2S)!}f^{(2S)}(\epsilon)$$

for some  $\epsilon$  and where the  $(x_i, \omega_i)$ 's can be easily found

• Well behaved function: For  $S \to \infty$  we have

$$\int_{-\infty}^{\infty} f(x)e^{-x^2}dx \approx \sum_{i=1}^{S} \omega_i f(x_i)$$

## Extra: Gauss-Hermite II

Gauss-Hermite quadrature uses that

$$\int_{-\infty}^{\infty} f(x)e^{-x^2}dx = \sum_{i=1}^{S} \omega_i f(x_i) + \frac{S!\sqrt{\pi}}{s^S(2S)!}f^{(2S)}(\epsilon)$$

for some  $\epsilon$  and where the  $(x_i, \omega_i)$ 's can be easily found

• Well behaved function: For  $S \to \infty$  we have

$$\int_{-\infty}^{\infty} f(x)e^{-x^2}dx \approx \sum_{i=1}^{S} \omega_i f(x_i)$$

■ Example: Random normal variable:  $Y \sim \mathcal{N}(\mu, \sigma^2)$  so that

$$\mathbb{E}[f(Y)] = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{\infty} f(y)e^{-\frac{(y-\mu)^2}{2\sigma^2}} dy$$
$$\approx \frac{1}{\sqrt{\pi}} \sum_{i=1}^{S} \omega_i f(\sqrt{2}\sigma x_i + \mu)$$

**Summary** 

## Summary and next week

#### Today:

- 1. Introduction to course
- 2. Dynamic programming in theory
- 3. Dynamic programming in practice
- Next week: More on consumption-saving models and precautionary savings in partial equilibrium

#### Homework:

- 1. Work on completing the model extension exercise
- 2. Read: TBA