
SF²Bench: Evaluating Data-Driven Models for Compound Flood Forecasting in South Florida

Xu Zheng¹ Chaohao Lin¹ Sipeng Chen¹ Zhuomin Chen¹ Jimeng Shi¹
Wei Cheng² Jayantha Obeysekera¹ Jason Liu¹ Dongsheng Luo¹

¹Florida International University ²NEC Laboratories America

{xzhen019, clin027, schen131, zchen051, jshi008}@fiu.edu

{weicheng}@nec-labs.com, {jobeysk, liux, dluo}@fiu.edu

Abstract

Forecasting compound floods presents a significant challenge due to the *intricate interplay of meteorological, hydrological, and oceanographic factors*. Analyzing compound floods has become more critical as the global climate increases flood risks. Traditional physics-based methods, such as the Hydrologic Engineering Center’s River Analysis System, are often time-inefficient. Machine learning has recently demonstrated promise in both modeling accuracy and computational efficiency. However, the scarcity of comprehensive datasets currently hinders systematic analysis. Existing water-related datasets are often limited by a sparse network of monitoring stations and incomplete coverage of relevant factors. To address this challenge, we introduce SF²Bench, a comprehensive time series collection on compound floods in South Florida, which integrates four key factors: tide, rainfall, groundwater, and human management activities (gate and pump controlling). This integration allows for a more detailed analysis of the individual contributions of these drivers to compound flooding and informs the development of improved flood forecasting approaches. To comprehensively evaluate the potential of various modeling paradigms, we assess the performance of six categories of methods, encompassing Multilayer Perceptrons, Convolutional Neural Networks, Recurrent Neural Networks, Graph Neural Networks, Transformers, and Large Language Models. We verified the impact of different key features on flood forecasting through experiments. Our analysis examines temporal and spatial aspects, providing insights into the influence of historical data and spatial dependencies. The varying performance across these approaches underscores the diverse capabilities of each in capturing complex temporal and spatial dependencies inherent in compound floods. By making the code and data publicly available¹, we aim to foster collaboration between the machine learning and environmental science communities, driving advancements in real-world flood forecasting solutions.

1 Introduction

Floods are among the most common and hazardous natural events, causing environmental damage [80], catastrophic loss of life [29], and property damage [8]. Compared with single-driver flood events, such as fluvial floods and pluvial floods [20], compound floods, occurring when two or more distinct flood drivers coincide in space or time [60], pose greater challenges for prediction and prevention, making it an important research topic in environmental science. Recent research indicates a rise in both the frequency and scale of compound floods due to global climate change [65, 71, 24]. Therefore, understanding the underlying causes of compound floods is both critical and urgent.

¹<https://github.com/AslanDing/SFBench>

Accurate and explainable compound flood models can support decision-making in water management, thereby minimizing damage to human life and infrastructure.

Classical physics-based methods predict the water stage by solving complex partial differential equations (PDE) [52, 81], such as the Hydrologic Engineering Center’s River Analysis System (HEC-RAS) [9]. Despite their accuracy and explainability, the extensive data requirements of physics-based methods, including high-resolution terrain data, reservoir characteristics, canal networks, and river geometries [59, 83], limit their widespread applications. The rapid development of machine learning (ML) has led to the application of data-centric methods, which utilize deep learning (DL) models for flood prediction and prevention. Researchers employ Convolutional Neural Networks (CNNs) [36], Long Short-Term Memory networks (LSTMs) [25], Graph Neural Networks (GNNs) [31], and Transformers [64] to uncover the underlying principles of compound flood. However, existing methods [2, 57, 46, 63, 62, 43] largely focus on temporal causality, often underestimating the complex interplay of factors. Most prominently, compound floods have garnered increasing attention due to their capacity to analyze multiple influencing factors [6, 65, 76, 50, 33]. Nevertheless, existing datasets [30, 57, 63] often contain limited factors, hindering a systematic analysis. For example, LamaH-CE [34] provides hydrological and topological data for the Danube River basin but lacks other important factors, such as rainfall.

Developing a new dataset and benchmark for systematic analysis of compound floods presents several challenges, including the diversity of relevant factors like meteorological drivers and tides, the need for long-term data, and the spatial distribution of data collection. To mitigate these challenges, previous studies have often focused on local regions with limited factors, such as Haikou City in [76]. However, the limited spatial scope of such datasets restricts their representativeness and hinders generalization to other regions.

In this paper, we introduce SF²Bench, a comprehensive time series dataset for compound floods in South Florida. South Florida’s intricate waterway system, encompassing rivers, canals, reservoirs, and extensive coastlines, presents substantial challenges for both the prediction and attribution analysis of compound floods. Moreover, its unique confluence of low-lying topography, converging flood drivers such as hurricanes and sea-level rise, urban development, porous geology, and frequent compound flood renders it an unparalleled area for studying the dynamics of compound flooding. According to [26], the key factors for compound flooding in the South Florida region include sea level, rainfall, river discharge, groundwater table, storm surge, and waves. Moreover, human management activities, such as water flow control, represent another crucial factor for comprehensive flood analysis. To provide a representative and comprehensive dataset, we consider the multiple key factors: water level, sea level, groundwater level, rainfall, and human management activities on hydraulic structures (gates and pumps), as shown in Figure 1. We compiled time series data from 2,452 monitoring stations across counties, spanning from 1985 to 2024. To the best of our knowledge, SF²Bench represents the first comprehensive dataset for compound flood analysis incorporating such a range of driving factors.

To validate data-centric AI methods for compound flood forecasting, we benchmark a wide range of forecasting methods using SF²Bench, including Multilayer Perceptrons (MLPs), Recurrent Neural Networks(RNNs), CNNs, GNNs, Transformers, and Large Language Models(LLM)-based approaches. Our observations indicate that MLPs and Transformers exhibit advantages in terms of MAE and MSE metrics, while MLPs and GNNs demonstrate better performance in extreme flood events. The benchmark results highlight the varying degrees of effectiveness of each method in capturing the complex temporal and spatial dependencies inherent in compound flooding. Furthermore, we conduct experiments to demonstrate the individual and combined effectiveness of the different factors included in SF²Bench across various model architectures. Finally, we discuss potential strategies for improving flood forecasting performance by leveraging both spatial and temporal information.




Figure 1: The schematic diagram of compound flood. The key factors include rainfall, sea, groundwater, and human control. The figure is assisted by OpenAI SORA.

2 Related Work

2.1 Flood Dataset

Monitoring floods presents a significant challenge due to their unpredictable nature and potentially devastating consequences. Existing flood datasets can be broadly categorized into satellite image datasets [56, 17, 47, 53, 77, 7] and time series monitoring datasets [76, 30, 2, 57, 34, 18, 12]. Satellite image datasets utilize remote sensing to capture surface water extent [4]. While effective in delineating flood-affected areas, this type of dataset [77, 7] often lacks crucial temporal dynamics and information on the underlying hydrological and meteorological factors that drive flood formation, limiting its utility for in-depth modeling. Time series monitoring datasets, on the other hand, typically utilize fixed monitoring stations to record hydrological-related data such as soil moisture, water level, and temperature. A prominent example within this category is the CAMELS-x family of datasets [1, 3, 16, 12, 18]. For instance, CAMELS-BR [12] encompasses data from 3,679 gauges across Brazil. LamaH-CE [34] provides daily and hourly time series data from 882 gauges, including runoff, meteorological variables, and catchment attributes. In [33], LamaH-CE is used as a benchmark for flood forecasting, primarily focusing on temporal and spatial aspects. However, these datasets primarily focus on general hydrological modeling. Analyzing compound floods, as highlighted in [26], necessitates detailed data on rainfall, water levels, and groundwater, which are often limited in existing time series datasets.

2.2 Machine Learning for Forecasting

The task of forecasting time series data presents inherent complexities and high dimensionality. Recent advancements in time series forecasting have been significantly propelled by a data-centric approach [85], underscoring the critical role of extensive, high-quality data in training robust models. Deep learning methodologies, with their powerful representation learning capabilities, have shown considerable promise in this domain. Based on their architectural designs, deep learning methods applied to time series forecasting can be categorized as follows: **MLP-based models** [14, 84, 69, 38, 37] leverage the capabilities of multilayer perceptrons for analyzing temporal sequences. **RNN-based methods** [35, 58, 67, 54, 27] are widely adopted in time series forecasting due to their inherent ability to model temporal dependencies within sequential data. **CNN-based methods** [68, 15, 45, 72, 66] employ convolutional operations to extract hierarchical features from time series data, enabling effective learning of underlying patterns and trends. **GNN-based methods** [74, 75, 11, 40, 79, 10] utilize graph structures to model intricate relationships between different time series variables, enhancing forecasting accuracy. **Transformer-based methods** [70, 41, 48, 88, 39, 73, 86] have demonstrated remarkable performance in capturing long-range dependencies and complex temporal dynamics within time series data. **LLM-based methods** [28, 51, 89, 21] explore the application of prompting and reprogramming techniques to align time series data with text embeddings for forecasting tasks.

3 The SF²Bench Dataset

3.1 Overview

SF²Bench comprises a time series data collection from 2,452 monitoring stations across a 67,349 km² area in South Florida, sourced from the South Florida Water Management District (SFWMD)². The dataset spans the period from 1985 to 2024 and is divided into 8 temporal splits. This dataset incorporates key factors that play critical roles in compound floods [26], including water level³, rainfall, groundwater level, and human control data for pumps and gates. Notably, sea level data is inherently included within the water level at certain monitoring stations due to their direct connection to the sea. It is specifically collected for benchmarking data-driven forecasting approaches in the context of compound flood analysis. In Table 1, we provide a comparison with other datasets. Compared to CAMELS-x [1, 3, 16, 12, 18], SF²Bench focuses on a region particularly susceptible to flooding, making it more relevant for compound flood analysis. In comparison to Bangladesh-

²<https://www.sfwmd.gov/>

³Water stage and water level are used interchangeably.

Table 1: Comparison with other datasets for flood forecasting. N/A indicates Not Available. * denotes datasets sharing the cell of Other Attributions. Climatic indices refer to statistical data about climate, such as daily rainfall, potential evapotranspiration, and temperature time series. Land cover attributes describe the physical material on the Earth’s surface (e.g., the ratio of woodland). Soil attributes are characteristics of the soil (e.g., porosity and soil depth). Geological attributes refer to features of the Earth’s surface (e.g., geologic class and subsurface porosity). Anthropogenic influences encompass activities such as water abstraction and discharges. Other Catchment attributes include location, area, topographic data, and so forth.

Dataset	Time Span	Interval	Type	Gauges	Area(km ²)	Public	Other Attributions
DarlingFlood [2]	1900-2018	Daily	Flow	12	3.5×10^4	No	Rainfall
SekongFlood [2]	1981-2013	Daily	Flow	8	2.8×10^4	No	Rainfall
BangladeshFlood [57]	1979-2013	Daily	Stage	24	1.5×10^5	No	N/A
Qi River [61]	1979-2020	Hour	Flow	7	7.1×10^3	No	Rainfall
Tunxi basins [61]	1981-2007	Hour	Flow	12	N/A	No	Rainfall
CAMELS* [1]	1989-2009	Daily	Flow	671	1.0×10^4	Yes	Climatic Indices
CAMELS-CL* [3]	1913-2018	Daily	Flow	516	N/A	Yes	Land Cover Attributes
CAMELS-GB* [16]	1970-2015	Daily	Flow	671	2.1×10^5	Yes	Soil Attributes
CAMELS-BR* [12]	1925-2024	Daily	Flow	4,025	N/A	Yes	Geological Attributes
CAMELS-AUS* [18]	1951-2014	Daily	Flow	107	6.9×10^5	Yes	Anthropogenic Influences
LamaH-CE* [34]	1951-2014	Daily & Hour	Flow	859	1.7×10^5	Yes	Other Catchment Attributes
SF ² Bench	1985-2024	Hour	Stage	2,452	6.7×10^4	Yes	Rainfall, Groundwater, Human Control

Flood [57], SF²Bench offers a more comprehensive set of driving factors relevant to compound flooding, considering both temporal and spatial dimensions.

3.2 Preprocessing

Data Collection. The data for SF²Bench was collected from DBHYDRO ⁴, an environmental database maintained by the SFWMD that stores a wide range of hydrologic, meteorologic, hydrogeologic, and water quality data. We initially collected data from 3731 monitoring stations and subsequently screened them based on data availability and relevance to flood analysis, resulting in a final set of 2,452 valid stations. This included 993 water stage monitoring stations, 349 rainwater monitoring stations, 582 groundwater level monitoring stations, 99 pump stations, and 429 gates. The different types of data and their physical meanings are summarized in Table 2. To capture fine-grained temporal dynamics, all raw data is collected at their native ‘breakpoint’ frequency, resulting in a high (up to second-level) temporal resolution for each recorded value. However, this breakpoint frequency collection method results in inconsistent data across stations, with some recording data at much higher frequencies than others. In addition, some monitoring stations have missing data for certain periods, while others provide data only for a limited duration, such as one year.

Table 2: The summary of the data type in SF²Bench.

Type	Stations	Unit	Description
Water	993	Feet	Water Stage
Groundwater	582	Feet	Stage of Groundwater
Rainfall	349	Inches	Rainfall
Pump	99	RPM	Rotational Speed
Gate	429	Feet	Opening Level

Data Processing. To provide an AI-ready dataset, we introduce data processing to unify the format. As previously mentioned, the data collection ranges vary across different monitoring stations, making it challenging to standardize all data to a uniform length. To balance the number of time series and temporal length, we divided the data into eight splits. Each time-series data is sampled with an hourly interval for all splits. During this processing, for each hourly interval, we first compute the mean of the available data points within that interval. Subsequently, we address missing values using interpolation methods. According to the characteristics of the data, we have two interpolation methods: linear interpolation and zero data filling. For water stage and groundwater level, we regard them as continuous variables and apply linear interpolation to fill missing values. For the other

⁴<https://apps.sfwmd.gov/dbhydroInsights/%23/homepage>




Figure 2: (a) Monitor stations location distribution. (b) Flood observation location distribution. (c) Temporal patterns of key features over a year. In (a) and (b), we highlight three interest parts: Orlando, Fort Myers, and Miami. In (c), the x-axis is the number of days in one year.

variables, namely rainfall and control data (pumps and gates), we treat them as discrete events and fill missing values with zero. The detailed information for each split is presented in Appendix Table 7.

3.3 Qualitative Analysis

To provide an intuitive insight of SF²Bench, we provide visualization results from spatial and temporal aspects. More detailed information are provided in the Appendix A.

Spatial Distribution. Figure 2a illustrates the spatial distribution of all monitoring stations, which are primarily located around the intricate river system of South Florida, radiating outwards from Lake Okeechobee. The geographically staggered distribution of hydrological, groundwater, and rainwater monitoring stations enables more effective spatial analysis. For example, the water level at a specific location is likely correlated with rainfall in the surrounding area and the local groundwater level (representing the soil's water storage capacity). We also highlight the observed flood locations from 2020 to 2023 and in 2008 in Figure 2b provided by SFWMD [49], which are predominantly concentrated in urban areas.

Temporal Pattern Visualization. Figure 2c illustrates the average annual temporal patterns across all monitoring stations, using data from split S7 as a representative example. From this pattern, we observe a strong correlation between groundwater level and water level data. The water level generally rises from approximately the 150th to the 300th day of the year, followed by a gradual decrease. According to the National Weather Service (NWS)⁵, Florida's climate is characterized by distinct dry and rainy seasons, with the latter typically spanning from May to October. This aligns well with the observed data patterns in our dataset. Taking rainfall as a reference, we note that increases in water level tend to correspond with rainfall events, such as the rainfall peak after the 300th day and the subsequent rise in water level. In addition, human control activities on hydraulic structures (e.g., pumps and gates) appear to influence water level changes in response to rainfall. We can infer that human intervention has played a role in mitigating potential flooding.

4 Forecasting Benchmarks

4.1 Problem Definition

Given water stage time series data $\mathbf{X} \in \mathbb{R}^{N \times L}$ from N water monitoring stations, the forecasting task is to predict the water stage values for the next T time steps, denoted as $\mathbf{Y} \in \mathbb{R}^{N \times T}$, using a fixed look-back window of length L . We also have access to additional time series information,

⁵<https://www.weather.gov/tbw/TBWTimeClimoQuickReference>

including groundwater levels $\mathbf{X}_w \in \mathbb{R}^{N_w \times L}$ (from N_w stations), rainfall $\mathbf{X}_r \in \mathbb{R}^{N_r \times L}$ (from N_r stations), pump control data $\mathbf{X}_p \in \mathbb{R}^{N_p \times L}$ (from N_p stations), gate control data $\mathbf{X}_g \in \mathbb{R}^{N_g \times L}$ (from N_g stations), and location information for the monitoring stations. For the primary benchmark experiments aimed at fair comparison across different models, we mainly consider the water stage data as the supervised data. However, we acknowledge that incorporating the additional information (groundwater levels, rainfall, pump and gate control data, and location) has the potential to further enhance forecasting performance.

4.2 Metric

We follow standard time series forecasting practices by using the Mean Absolute Error (MAE) and Mean Squared Error (MSE) as our primary evaluation metrics. To better assess the performance of our models in real-world applications, particularly for extreme flood events, we also employ the Symmetric Extremal Dependence Index (SEDI) [22, 78], as suggested by [23]. By selecting quantile thresholds (e.g., the 95th and 5th percentiles of the observed values), SEDI classifies each time stamp as belonging to either a normal or an extreme case and then calculates the hit rate of this classification. A higher SEDI value indicates better performance in predicting extreme events. The formulation of SEDI is as follows:

$$\text{SEDI}(p) = \frac{|\hat{\mathbf{Y}} < V_{1-\frac{p}{2}} \& \mathbf{Y} < V_{1-\frac{p}{2}}| + |\hat{\mathbf{Y}} > V_{\frac{p}{2}} \& \mathbf{Y} > V_{\frac{p}{2}}|}{|\mathbf{Y} < V_{1-\frac{p}{2}}| + |\mathbf{Y} > V_{\frac{p}{2}}|}, \quad (1)$$

where $|\cdot|$ means the number of the true values satisfying the condition, $\hat{\mathbf{Y}}$ is the forecasting results, p is the quantile of the threshold, and $V_{1-\frac{p}{2}}, V_{\frac{p}{2}}$ are the lower and upper threshold of top and worst $\frac{p}{2}$ percent, respectively.

4.3 Methods

We benchmark six categories of time series forecasting architectures: MLP, CNN, RNN, GNN, Transformer, and LLM. We select two advanced methods as representative examples for each of these categories. Additionally, for the first four classical architectures, we also implement a basic foundational architecture. The specific advanced methods we evaluate are: MLP: NLinear [84], TSMixer [14], CNN: ModernTCN [45], TimesNet [72], RNN: DeepAR [58], DilatedRNN [13], GNN: FourierGNN [79], StemGNN [11], Transformer: PatchTST [48], iTransformer [41], LLM: GPT4TS [89], AutoTimes [42]. We follow the source code of NeuralForecast⁶ for the implementation of these approaches. The summary of these methods can be found in Appendix B.2.

4.4 Experiment Setup

Due to the memory and training time limitations associated with GPUs, applying some methods, particularly LLM-based approaches [51, 28, 89, 42], to the entire dataset is challenging. To facilitate a fair comparison, we conduct our experiments using two setups: evaluation on three specific areas of interest and evaluation on the entire dataset. The three areas were selected based on the flood observation data presented in Figure 2. Figure 2b visualizes these flood locations alongside the selected areas. We report the performance of all evaluated methods within these areas of interest. For the entire dataset, we only report the results of the basic foundational techniques. For each data split, the last year’s data is used for testing, the data of the second-to-last year is used for validation, and the remaining preceding data is used for training. In the benchmark, we maintain a consistent lookback window of two days, and we evaluate prediction windows of one, three, five, and seven days. The detailed experimental setup, including software and hardware platforms, is provided in the Appendix B.

Table 3: Average results of basic methods on the whole dataset. The best and second results are shown in bold font and underlined, respectively.

Metric	MLP	LSTM	TCN	GCN
\downarrow MAE	0.2328	0.3302	0.3491	<u>0.3053</u>
\downarrow MSE	0.3902	2.2772	<u>0.6807</u>	1.0673
\uparrow SEDI(10%)	0.6853	0.5751	0.5167	<u>0.6137</u>
\uparrow SEDI(5%)	0.5970	0.4516	0.3904	<u>0.5213</u>
\uparrow SEDI(1%)	0.4107	0.1828	0.1842	<u>0.3134</u>

⁶<https://github.com/Nixtla/neuralforecast>

Table 4: Benchmark of average MAE&MSE results on three interest areas across 8 splits. The best and second results are shown in bold font and underlined, respectively.

Metric	T	MLP			CNN			Transformer	
		MLP	TSMixer	NLinear	TCN	ModernTCN	TimesNet	iTransformer	PatchTST
\downarrow MAE	1D	0.0788	0.0928	0.0817	0.1792	0.0798	0.0983	<u>0.0756</u>	0.0741
	3D	0.1351	0.1442	0.1373	0.2281	0.1441	0.1528	0.1314	0.1316
	5D	0.1764	0.1816	0.1769	0.2697	0.1891	0.1927	<u>0.1722</u>	0.1719
	7D	0.2063	0.2126	0.2082	0.3088	0.2248	0.2267	0.2041	0.2044
	Avg.	0.1492	0.1578	0.1510	0.2465	0.1594	0.1676	<u>0.1458</u>	0.1455
\downarrow MSE	1D	<u>0.0521</u>	0.0648	0.0556	0.3722	0.0681	0.0704	0.0253	0.0531
	3D	0.1029	0.1132	<u>0.1105</u>	0.4647	0.2443	0.1358	0.1112	0.1132
	5D	0.1432	0.1566	0.1547	0.4173	0.2808	0.1829	0.1583	0.1566
	7D	0.1707	0.1903	0.1841	0.5386	0.2723	0.2241	0.1911	0.1903
	Avg.	0.1172	0.1283	<u>0.1262</u>	0.4482	0.2164	0.1533	0.1284	0.1283
Metric		RNN			GNN			LLM	
		LSTM	DeepAR	DilatedRNN	GCN	FourierGNN	StemGNN	GPT4TS	AutoTimes
\downarrow MAE	1D	0.1182	0.1178	0.0919	0.1696	0.0921	0.1332	0.1256	0.0846
	3D	0.1821	0.1837	0.1573	0.2006	0.1503	0.2181	0.1521	0.1362
	5D	0.2232	0.2247	0.2022	0.2504	0.1930	0.3153	0.1911	0.1752
	7D	0.2576	0.2596	0.2374	0.2799	0.2280	0.3570	0.2247	0.2062
	Avg.	0.1953	0.1964	0.1722	0.2251	0.1658	0.2559	0.1734	0.1505
\downarrow MSE	1D	0.1339	0.1230	0.1033	7.2299	0.0768	0.1632	0.0966	0.0584
	3D	0.1985	0.1954	0.1672	1.5645	0.1416	0.2543	0.1410	0.1125
	5D	0.2348	0.2389	0.2341	2.1341	0.2071	0.4189	0.1847	<u>0.1522</u>
	7D	0.2873	0.2691	0.2591	1.0374	0.2125	0.4716	0.2245	<u>0.1823</u>
	Avg.	0.2136	0.2066	0.1909	2.9915	0.1595	0.3270	0.1617	0.1263

Table 5: Benchmark of average SEDI results on three interest areas across 8 splits. The best and second results are shown in bold font and underlined, respectively.

Metric		MLP			CNN			Transformer	
		MLP	TSMixer	NLinear	TCN	ModernTCN	TimesNet	iTransformer	PatchTST
\uparrow SEDI(10%)	0.6897	0.6144	0.6278	0.5311	0.6067	0.5829	0.6286	0.6296	
	0.5834	0.4942	0.5111	0.3706	0.4846	0.4589	0.5079	0.5086	
	0.3666	0.2480	0.2767	0.1387	0.2512	0.2222	0.2690	0.2685	
Metric		RNN			GNN			LLM	
		LSTM	DeepAR	DilatedRNN	GCN	FourierGNN	StemGNN	GPT4TS	AutoTimes
\uparrow SEDI(10%)	0.6097	0.5989	0.6164	0.6179	<u>0.6623</u>	0.5507	0.5581	0.6239	
	0.4702	0.4643	0.5014	0.5138	<u>0.5511</u>	0.4270	0.4640	0.5015	
	0.1680	0.1478	0.2499	0.2828	<u>0.3217</u>	0.1690	0.2268	0.2568	

4.5 Results & Observations

Overall Performance. Table 4 presents the average MSE and MAE results across the three interest areas across eight data splits. The performance-leading methods include PatchTST, iTransformer, MLP, NLinear, and AutoTimes. PatchTST and iTransformer demonstrate the best performance according to the MAE results. However, MLP, NLinear, and AutoTimes exhibit the best performance in terms of the MSE metric. A smaller MAE and a larger MSE suggest that a method accurately predicts the majority of data points, but occasionally produces abnormal prediction values. Conversely, a larger MAE and a smaller MSE indicate a relatively stable method that might not be highly accurate for most points. Table 5 demonstrates the performance of the models on extreme cases. Surprisingly, FourierGNN achieves the second-best performance in SEDI, despite not showing superiority in MSE and MAE. The detailed results for each split are provided in the Appendix C. Furthermore, Table 3 presents the average results of the basic methods on the entire dataset. The detailed results are available in the Appendix C.1. Among the four basic methods, the performance of MLP remains

Table 6: Input factors ablation study on S6. All, G, R, and C represent all factors, groundwater, rainfall, and human control(pump and gate). The best and second results are shown in bold font and underlined, respectively.

Metric	method	w/ All	w/o G	w/o R	w/o C	w/o GR	w/o RC	w/o WC	w/o WRC
\downarrow MAE	iTransformer	0.1406	0.1407	0.1406	<u>0.1405</u>	0.1411	0.1410	0.1402	0.1411
	PatchTST	0.1376	0.1391	0.1378	<u>0.1377</u>	0.1398	0.1396	0.1385	0.1421
	TSMixer	0.1596	<u>0.1419</u>	0.2523	0.2111	0.1418	0.2807	0.1423	0.1422
	NLinear	0.1546	0.1577	0.1483	0.1540	0.1485	<u>0.1450</u>	0.1579	0.1435
	TimesNet	0.1642	0.1599	0.1662	0.1650	0.1592	0.1656	0.1578	<u>0.1580</u>
\downarrow MSE	iTransformer	0.0953	0.0960	0.0949	0.0954	0.0957	0.0956	0.0949	0.0962
	PatchTST	<u>0.0917</u>	0.0932	0.0927	0.0916	0.0950	0.0938	0.0923	0.0971
	TSMixer	0.1080	0.0946	0.4061	0.2698	0.0946	0.6697	<u>0.0943</u>	0.0941
	NLinear	0.0992	0.1011	0.0966	0.0984	0.0973	<u>0.0954</u>	0.1006	0.0947
	TimesNet	0.1188	0.1154	0.1226	0.1202	0.1145	0.1237	0.1111	<u>0.1123</u>

the best, followed by GCN and TCN, which is consistent with the MSE and MAE results in Table 4. Overall, our observations indicate that for this task, MLP, transformer-based models (PatchTST and iTransformer), and LLM-based models (AutoTimes and GPT4TS) performed outstandingly in terms of MAE and MSE, while the GNN-based model (FourierGNN) showed astonishing performance on the SEDI metric. In this context, the prediction results for extreme cases are particularly important as they reflect the predictive ability for flood occurrences. Notably, we did not observe strong correlations between the MAE/MSE results and the SEDI results, underscoring the necessity of reporting multiple evaluation metrics. Additionally, we found no clear relationship between model performance and the number of trainable parameters, as evidenced by NLinear(9K) achieving comparable results to AutoTimes(4M) despite a significant difference in trainable parameter count.

Ablation of Factors. To verify and demonstrate the influence of different input factors, we conduct an ablation study using five methods. These methods contains two settings: channel-independent (including NLinear and PatchTST) and channel-dependent (including iTransformer, TimesNet, and TSMixer). For the channel-independent methods, we used all available factors as input. In contrast, for the channel-dependent methods, we primarily used the water stage data as the supervised target, while other factors were considered as potential additional inputs. As shown in Table 6, NLinear and TSMixer achieve comparable results when the input is limited to only the water stage data, suggesting that the inclusion of other factors does not significantly improve their performance. For iTransformer, PatchTST, and TimesNet, we observe performance improvements when additional information is provided, highlighting the potential benefit of incorporating multi-faceted data for this forecasting task.

Moreover, for iTransformer and PatchTST, we find excluding groundwater information (denoted as “w/o G” and “w/o GR”) results in larger MSE and MAE errors compared to settings where other factors are excluded (e.g., “w/o R” for without rainfall, “w/o C” for without control data, “w/o RC”, and “w/o WC”). Moreover, for iTransformer and TimesNet, providing only groundwater information as the additional input leads to the best performance among the ablation settings, suggesting that the groundwater is a particularly informative factor for these models. We also observe similar results on the SEDI(10%) metric, provided in Appendix C.2.




Figure 3: The different interest areas with a radius for ablation study on spatial information.

Impact of Temporal and Spatial Information. To provide insight into the impact of temporal input length and spatial information, we conduct ablation studies respectively. We first select an interest area as the anchor area, where the water stages are regarded as the forecasting target. As shown in Figure 3, R is the radius of the anchor area. Then, we incorporate information from surrounding stations by incrementally increasing the radius of the interest area. In these experiments, we consider radius scale factors of 1, 1.2, 1.4, 1.6, and 1.8. The MAE, MSE, and SEDI results, presented in Figures 4, show that iTransformer, PatchTST, and TSMixer experience a performance improvement as the input




Figure 4: Study on spatial information impact.




Figure 5: Study on temporal information impact.

area expands. This indicates the effectiveness of incorporating additional spatial information for the forecasting task. We provide the detailed results in Appendix C.2.

Furthermore, maintaining the anchor area as the forecasting target, we evaluate the impact of temporal input length by considering a range of durations: 6 hours, 12 hours, and 1 to 6 days. As shown in Figure 5, we observe that PatchTST, TSMixer, and iTransformer generally show improved performance with increasing input length. However, for iTransformer, performance begins to decrease beyond an input length of 1 day. A potential reason for this phenomenon is that longer input sequences require a larger amount of training data to effectively learn the underlying patterns. The detailed results are available in Appendix C.2.

Comparing these two strategies, we find that both can enhance task performance. Increasing spatial input generally leads to a relatively stable, albeit limited, improvement. In contrast, extending the temporal input length can yield more substantial gains, particularly for models like TSMixer, where a significant reduction in MSE is observed.

5 Conclusion

In this paper, we introduce SF²Bench, a new dataset collected for comprehensive compound flood analysis, aiming to foster collaboration between the machine learning and environmental science communities. SF²Bench comprehensively covers the majority of South Florida and integrates five key factors: water level, sea level, groundwater table, rainfall, and human control activities. We evaluate six types of time series forecasting approaches on this dataset and observe that different architectures exhibit distinct advantages. Furthermore, our ablation studies on input factors reveal that groundwater level is a particularly effective predictor compared to other information sources. Additionally, we conduct experiments to explore the effectiveness of increasing spatial and temporal information, and the results demonstrate that both strategies improve forecasting performance for this task.

Limitations. While SF²Bench provides five key factors, it currently lacks explicit topological linkage information between monitoring stations due to the intricate nature of South Florida’s water system. Although we provide some flood observation data, the locations of these observations may not directly correspond to our monitoring stations. Therefore, this information is provided in a separate file rather than being integrated into the time series data.

References

- [1] Nans Addor, Andrew J Newman, Naoki Mizukami, and Martyn P Clark. The camels data set: catchment attributes and meteorology for large-sample studies. *Hydrology and Earth System Sciences*, 21(10):5293–5313, 2017.
- [2] Kasuni E. Adikari, Sangam Shrestha, Dhanika T. Ratnayake, Aakanchya Budhathoki, S. Mo-hanasundaram, and Matthew N. Dailey. Evaluation of artificial intelligence models for flood and drought forecasting in arid and tropical regions. *Environmental Modelling & Software*, 144:105136, 2021.
- [3] Camila Alvarez-Garreton, Pablo A Mendoza, Juan Pablo Boisier, Nans Addor, Mauricio Galleguillos, Mauricio Zambrano-Bigiarini, Antonio Lara, Cristóbal Puelma, Gonzalo Cortes, Rene Garreaud, et al. The camels-cl dataset: catchment attributes and meteorology for large sample studies–chile dataset. *Hydrology and Earth System Sciences*, 22(11):5817–5846, 2018.
- [4] Donato Amitrano, Gerardo Di Martino, Alessio Di Simone, and Pasquale Imperatore. Flood detection with sar: A review of techniques and datasets. *Remote Sensing*, 16(4), 2024.
- [5] Shaojie Bai, J Zico Kolter, and Vladlen Koltun. An empirical evaluation of generic convolutional and recurrent networks for sequence modeling. *arXiv preprint arXiv:1803.01271*, 2018.
- [6] Emanuele Bevacqua, Douglas Maraun, Michalis Ioannis Voudoukas, Evangelos Voukouvalas, Mathieu Vrac, Lorenzo Mentaschi, and Martin Widmann. Higher probability of compound flooding from precipitation and storm surge in europe under anthropogenic climate change. *Science advances*, 5(9):eaaw5531, 2019.
- [7] Derrick Bonafilia, Beth Tellman, Tyler Anderson, and Erica Issenberg. Sen1floods11: A georeferenced dataset to train and test deep learning flood algorithms for sentinel-1. In *Proceedings of the IEEE/CVF conference on computer vision and pattern recognition workshops*, pages 210–211, 2020.
- [8] Samuel D Brody, Sammy Zahran, Praveen Maghelal, Himanshu Grover, and Wesley E Highfield. The rising costs of floods: Examining the impact of planning and development decisions on property damage in florida. *Journal of the American Planning Association*, 73(3):330–345, 2007.
- [9] Gary W Brunner. Hec-ras river analysis system. hydraulic reference manual. version 1.0. 1997.
- [10] Wanlin Cai, Yuxuan Liang, Xianggen Liu, Jianshuai Feng, and Yuankai Wu. Msgnet: Learning multi-scale inter-series correlations for multivariate time series forecasting. *arXiv preprint arXiv:2401.00423*, 2023.
- [11] Defu Cao, Yujing Wang, Juanyong Duan, Ce Zhang, Xia Zhu, Conguri Huang, Yunhai Tong, Bixiong Xu, Jing Bai, Jie Tong, and Qi Zhang. Spectral temporal graph neural network for multivariate time-series forecasting. In *Proceedings of the 34th International Conference on Neural Information Processing Systems*, NIPS ’20, Red Hook, NY, USA, 2020. Curran Associates Inc.
- [12] Vinícius BP Chagas, Pedro LB Chaffe, Nans Addor, Fernando M Fan, Ayan S Fleischmann, Rodrigo CD Paiva, and Vinícius A Siqueira. Camels-br: hydrometeorological time series and landscape attributes for 897 catchments in brazil. *Earth System Science Data*, 12(3):2075–2096, 2020.
- [13] Shiyu Chang, Yang Zhang, Wei Han, Mo Yu, Xiaoxiao Guo, Wei Tan, Xiaodong Cui, Michael Witbrock, Mark A Hasegawa-Johnson, and Thomas S Huang. Dilated recurrent neural networks. *Advances in neural information processing systems*, 30, 2017.
- [14] Si-An Chen, Chun-Liang Li, Sercan O Arik, Nathanael Christian Yoder, and Tomas Pfister. TSMixer: An all-MLP architecture for time series forecast-ing. *Transactions on Machine Learning Research*, 2023.

- [15] Mingyue Cheng, Jiqian Yang, Tingyue Pan, Qi Liu, and Zhi Li. Convtimenet: A deep hierarchical fully convolutional model for multivariate time series analysis. *arXiv preprint arXiv:2403.01493*, 2024.
- [16] Gemma Coxon, Nans Addor, John P Bloomfield, Jim Freer, Matt Fry, Jamie Hannaford, Nicholas JK Howden, Rosanna Lane, Melinda Lewis, Emma L Robinson, et al. Camels-gb: hydrometeorological time series and landscape attributes for 671 catchments in great britain. *Earth System Science Data*, 12(4):2459–2483, 2020.
- [17] F. Dottori, L. Alferi, A. Bianchi, J. Skoien, and P. Salamon. A new dataset of river flood hazard maps for europe and the mediterranean basin. *Earth System Science Data*, 14(4):1549–1569, 2022.
- [18] K. J. A. Fowler, S. C. Acharya, N. Addor, C. Chou, and M. C. Peel. Camels-aus: hydrometeorological time series and landscape attributes for 222 catchments in australia. *Earth System Science Data*, 13(8):3847–3867, 2021.
- [19] Jean-Yves Franceschi, Aymeric Dieuleveut, and Martin Jaggi. Unsupervised scalable representation learning for multivariate time series. *Advances in neural information processing systems*, 32, 2019.
- [20] Joshua Green, Ivan D Haigh, Niall Quinn, Jeff Neal, Thomas Wahl, Melissa Wood, Dirk Eilander, Marleen de Ruiter, Philip Ward, and Paula Camus. A comprehensive review of coastal compound flooding literature. *arXiv preprint arXiv:2404.01321*, 2024.
- [21] Nate Gruver, Marc Finzi, Shikai Qiu, and Andrew Gordon Wilson. Large language models are zero-shot time series forecasters. In *Proceedings of the 37th International Conference on Neural Information Processing Systems, NIPS ’23*, Red Hook, NY, USA, 2023. Curran Associates Inc.
- [22] Tao Han, Zhenghao Chen, Song Guo, Wanghan Xu, and Lei Bai. Cra5: Extreme compression of era5 for portable global climate and weather research via an efficient variational transformer. *arXiv preprint arXiv:2405.03376*, 2024.
- [23] Tao Han, Song Guo, Zhenghao Chen, Wanghan Xu, and Lei Bai. How far are today’s time-series models from real-world weather forecasting applications? *arXiv preprint arXiv:2406.14399*, 2024.
- [24] Yukiko Hirabayashi, Roobavannan Mahendran, Sujan Koirala, Lisako Konoshima, Dai Yamazaki, Satoshi Watanabe, Hyungjun Kim, and Shinjiro Kanae. Global flood risk under climate change. *Nature climate change*, 3(9):816–821, 2013.
- [25] Sepp Hochreiter and Jürgen Schmidhuber. Long short-term memory. *Neural computation*, 9(8):1735–1780, 1997.
- [26] R. Jane, L. Cadavid, J. Obeysekera, and T. Wahl. Multivariate statistical modelling of the drivers of compound flood events in south florida. *Natural Hazards and Earth System Sciences*, 20(10):2681–2699, 2020.
- [27] Sheo Yon Jhin, Seojin Kim, and Noseong Park. Addressing prediction delays in time series forecasting: A continuous gru approach with derivative regularization. In *Proceedings of the 30th ACM SIGKDD Conference on Knowledge Discovery and Data Mining*, pages 1234–1245, 2024.
- [28] Ming Jin, Shiyu Wang, Lintao Ma, Zhixuan Chu, James Y. Zhang, Xiaoming Shi, Pin-Yu Chen, Yuxuan Liang, Yuan-Fang Li, Shirui Pan, and Qingsong Wen. Time-LLM: Time series forecasting by reprogramming large language models. In *The Twelfth International Conference on Learning Representations*, 2024.
- [29] Sebastiaan N Jonkman and Johannes K Vrijling. Loss of life due to floods. *Journal of flood risk management*, 1(1):43–56, 2008.
- [30] Syed Kabir, Sandhya Patidar, Xilin Xia, Qiuahua Liang, Jeffrey Neal, and Gareth Pender. A deep convolutional neural network model for rapid prediction of fluvial flood inundation. *Journal of Hydrology*, 590:125481, 2020.

- [31] Thomas N Kipf and Max Welling. Semi-supervised classification with graph convolutional networks. *arXiv preprint arXiv:1609.02907*, 2016.
- [32] Thomas N. Kipf and Max Welling. Semi-supervised classification with graph convolutional networks. In *International Conference on Learning Representations*, 2017.
- [33] Nikolas Kirschstein and Yixuan Sun. The merit of river network topology for neural flood forecasting. In *International Conference on Machine Learning*, pages 24713–24725. PMLR, 2024.
- [34] Christoph Klingler, Karsten Schulz, and Mathew Herrnegger. Lamahl large-sample data for hydrology and environmental sciences for central europe. *Earth System Science Data Discussions*, 2021:1–46, 2021.
- [35] Guokun Lai, Wei-Cheng Chang, Yiming Yang, and Hanxiao Liu. Modeling long-and short-term temporal patterns with deep neural networks. In *The 41st international ACM SIGIR conference on research & development in information retrieval*, pages 95–104, 2018.
- [36] Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, 1998.
- [37] Shengsheng Lin, Weiwei Lin, Xinyi Hu, Wentai Wu, Ruichao Mo, and Haocheng Zhong. Cyclenet: Enhancing time series forecasting through modeling periodic patterns. In A. Globerson, L. Mackey, D. Belgrave, A. Fan, U. Paquet, J. Tomczak, and C. Zhang, editors, *Advances in Neural Information Processing Systems*, volume 37, pages 106315–106345. Curran Associates, Inc., 2024.
- [38] Shengsheng Lin, Weiwei Lin, Wentai Wu, Haojun Chen, and Junjie Yang. SparseTSF: Modeling long-term time series forecasting with *1k* parameters. In Ruslan Salakhutdinov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett, and Felix Berkenkamp, editors, *Proceedings of the 41st International Conference on Machine Learning*, volume 235 of *Proceedings of Machine Learning Research*, pages 30211–30226. PMLR, 21–27 Jul 2024.
- [39] Shizhan Liu, Hang Yu, Cong Liao, Jianguo Li, Weiyao Lin, Alex X. Liu, and Schahram Dustdar. Pyraformer: Low-complexity pyramidal attention for long-range time series modeling and forecasting. In *International Conference on Learning Representations*, 2022.
- [40] Yijing Liu, Qinxian Liu, Jian-Wei Zhang, Haozhe Feng, Zhongwei Wang, Zihan Zhou, and Wei Chen. Multivariate time-series forecasting with temporal polynomial graph neural networks. In S. Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, *Advances in Neural Information Processing Systems*, volume 35, pages 19414–19426. Curran Associates, Inc., 2022.
- [41] Yong Liu, Tengge Hu, Haoran Zhang, Haixu Wu, Shiyu Wang, Lintao Ma, and Mingsheng Long. itransformer: Inverted transformers are effective for time series forecasting. In *The Twelfth International Conference on Learning Representations*, 2024.
- [42] Yong Liu, Guo Qin, Xiangdong Huang, Jianmin Wang, and Mingsheng Long. Autotimes: Autoregressive time series forecasters via large language models. In *The Thirty-eighth Annual Conference on Neural Information Processing Systems*, 2024.
- [43] Zichuan Liu, Tianchun Wang, Jimeng Shi, Xu Zheng, Zhuomin Chen, Lei Song, Wenqian Dong, Jayantha Obeysekera, Farhad Shirani, and Dongsheng Luo. Timex++: Learning time-series explanations with information bottleneck. In *International Conference on Machine Learning*, pages 32062–32082. PMLR, 2024.
- [44] Ilya Loshchilov and Frank Hutter. Decoupled weight decay regularization. In *International Conference on Learning Representations*, 2019.
- [45] Donghao Luo and Xue Wang. Moderntcn: A modern pure convolution structure for general time series analysis. In *The twelfth international conference on learning representations*, pages 1–43, 2024.

- [46] Scott Miau and Wei-Hsi Hung. River flooding forecasting and anomaly detection based on deep learning. *Ieee Access*, 8:198384–198402, 2020.
- [47] Fabio Montello, Edoardo Arnaudo, and Claudio Rossi. Mmflood: A multimodal dataset for flood delineation from satellite imagery. *IEEE Access*, 10:96774–96787, 2022.
- [48] Yuqi Nie, Nam H Nguyen, Phanwadee Sinthong, and Jayant Kalagnanam. A time series is worth 64 words: Long-term forecasting with transformers. In *The Eleventh International Conference on Learning Representations*, 2023.
- [49] Office of Resilience, South Florida Water Management District. Resiliency and flood protection, 2025.
- [50] Agnieszka I Olbert, Sogol Moradian, Stephen Nash, Joanne Comer, Bartosz Kazmierczak, Roger A Falconer, and Michael Hartnett. Combined statistical and hydrodynamic modelling of compound flooding in coastal areas-methodology and application. *Journal of Hydrology*, 620:129383, 2023.
- [51] Zijie Pan, Yushan Jiang, Sahil Garg, Anderson Schneider, Yuriy Nevmyvaka, and Dongjin Song. ²ip-llm: Semantic space informed prompt learning with llm for time series forecasting. In *Forty-first International Conference on Machine Learning*, 2024.
- [52] Claudio Paniconi and Mario Putti. Physically based modeling in catchment hydrology at 50: Survey and outlook. *Water Resources Research*, 51(9):7090–7129, 2015.
- [53] Katerina Papagiannaki, Olga Petrucci, Michalis Diakakis, Vassiliki Kotroni, Luigi Aceto, Cinzia Bianchi, Rudolf Brázdil, Miquel Grimalt Gelabert, Moshe Inbar, Abdullah Kahraman, et al. Developing a large-scale dataset of flood fatalities for territories in the euro-mediterranean region, ffem-db. *Scientific data*, 9(1):166, 2022.
- [54] Yao Qin, Dongjin Song, Haifeng Chen, Wei Cheng, Guofei Jiang, and Garrison W Cottrell. A dual-stage attention-based recurrent neural network for time series prediction. In *Proceedings of the Twenty-Sixth International Joint Conference on Artificial Intelligence*, pages 2627–2633. International Joint Conferences on Artificial Intelligence Organization, 2017.
- [55] Alec Radford, Jeffrey Wu, Rewon Child, David Luan, Dario Amodei, Ilya Sutskever, et al. Language models are unsupervised multitask learners. *OpenAI blog*, 1(8):9, 2019.
- [56] Maryam Rahnemoonfar, Tashnim Chowdhury, Argho Sarkar, Debyrat Varshney, Masoud Yari, and Robin Roberson Murphy. Floodnet: A high resolution aerial imagery dataset for post flood scene understanding. *IEEE Access*, 9:89644–89654, 2021.
- [57] Jannatul Ferdous Ruma, Mohammed Sarfaraz Gani Adnan, Ashraf Dewan, and Rashedur M. Rahman. Particle swarm optimization based lstm networks for water level forecasting: A case study on bangladesh river network. *Results in Engineering*, 17:100951, 2023.
- [58] David Salinas, Valentin Flunkert, Jan Gasthaus, and Tim Januschowski. Deepar: Probabilistic forecasting with autoregressive recurrent networks. *International journal of forecasting*, 36(3):1181–1191, 2020.
- [59] Christopher C Sampson, Andrew M Smith, Paul D Bates, Jeffrey C Neal, Lorenzo Alfieri, and Jim E Freer. A high-resolution global flood hazard model. *Water resources research*, 51(9):7358–7381, 2015.
- [60] Antonia Sebastian. Chapter 7 - compound flooding. In Samuel Brody, Yoonjeong Lee, and Baukje Bee Kothuis, editors, *Coastal Flood Risk Reduction*, pages 77–88. Elsevier, 2022.
- [61] Pingping Shao, Jun Feng, Jiamin Lu, Pengcheng Zhang, and Chenxin Zou. Data-driven and knowledge-guided denoising diffusion model for flood forecasting. *Expert Systems with Applications*, 244:122908, 2024.
- [62] Jimeng Shi, Vitalii Stebliankin, Zhaonan Wang, Shaowen Wang, and Giri Narasimhan. Graph transformer network for flood forecasting with heterogeneous covariates. *arXiv preprint arXiv:2310.07631*, 2023.

- [63] Jimeng Shi, Zeda Yin, Arturo Leon, Jayantha Obeysekera, and Giri Narasimhan. Fidlar: Forecast-informed deep learning architecture for flood mitigation. *arXiv preprint arXiv:2402.13371*, 2024.
- [64] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.
- [65] Thomas Wahl, Shaleen Jain, Jens Bender, Steven D Meyers, and Mark E Luther. Increasing risk of compound flooding from storm surge and rainfall for major us cities. *Nature Climate Change*, 5(12):1093–1097, 2015.
- [66] Huiqiang Wang, Jian Peng, Feihu Huang, Jince Wang, Junhui Chen, and Yifei Xiao. Micn: Multi-scale local and global context modeling for long-term series forecasting. In *The eleventh international conference on learning representations*, 2023.
- [67] Jingyuan Wang, Ze Wang, Jianfeng Li, and Junjie Wu. Multilevel wavelet decomposition network for interpretable time series analysis. In *Proceedings of the 24th ACM SIGKDD international conference on knowledge discovery & data mining*, pages 2437–2446, 2018.
- [68] Shiyu Wang, Jiawei Li, Xiaoming Shi, Zhou Ye, Baichuan Mo, Wenze Lin, Shengtong Ju, Zhixuan Chu, and Ming Jin. Timemixer++: A general time series pattern machine for universal predictive analysis. *arXiv preprint arXiv:2410.16032*, 2024.
- [69] Shiyu Wang, Haixu Wu, Xiaoming Shi, Tengge Hu, Huakun Luo, Lintao Ma, James Y. Zhang, and JUN ZHOU. Timemixer: Decomposable multiscale mixing for time series forecasting. In *The Twelfth International Conference on Learning Representations*, 2024.
- [70] Yuxuan Wang, Haixu Wu, Jiaxiang Dong, Guo Qin, Haoran Zhang, Yong Liu, Yunzhong Qiu, Jianmin Wang, and Mingsheng Long. Timexer: Empowering transformers for time series forecasting with exogenous variables. In *The Thirty-eighth Annual Conference on Neural Information Processing Systems*, 2024.
- [71] Oliver EJ Wing, William Lehman, Paul D Bates, Christopher C Sampson, Niall Quinn, Andrew M Smith, Jeffrey C Neal, Jeremy R Porter, and Carolyn Kousky. Inequitable patterns of us flood risk in the anthropocene. *Nature Climate Change*, 12(2):156–162, 2022.
- [72] Haixu Wu, Tengge Hu, Yong Liu, Hang Zhou, Jianmin Wang, and Mingsheng Long. Timesnet: Temporal 2d-variation modeling for general time series analysis. In *The Eleventh International Conference on Learning Representations*, 2023.
- [73] Haixu Wu, Jiehui Xu, Jianmin Wang, and Mingsheng Long. Autoformer: decomposition transformers with auto-correlation for long-term series forecasting. In *Proceedings of the 35th International Conference on Neural Information Processing Systems*, pages 22419–22430, 2021.
- [74] Zonghan Wu, Shirui Pan, Guodong Long, Jing Jiang, Xiaojun Chang, and Chengqi Zhang. Connecting the dots: Multivariate time series forecasting with graph neural networks. In *Proceedings of the 26th ACM SIGKDD international conference on knowledge discovery & data mining*, pages 753–763, 2020.
- [75] Zonghan Wu, Shirui Pan, Guodong Long, Jing Jiang, and Chengqi Zhang. Graph wavenet for deep spatial-temporal graph modeling, 2019.
- [76] Kui Xu, Yunchao Zhuang, Lingling Bin, Chenyue Wang, and Fuchang Tian. Impact assessment of climate change on compound flooding in a coastal city. *Journal of Hydrology*, 617:129166, 2023.
- [77] Qingsong Xu, Yilei Shi, Jie Zhao, and Xiao Xiang Zhu. Floodcastbench: A large-scale dataset and foundation models for flood modeling and forecasting. *Scientific Data*, 12(1):431, 2025.
- [78] Wanghan Xu, Kang Chen, Tao Han, Hao Chen, Wanli Ouyang, and Lei Bai. Extremecast: Boosting extreme value prediction for global weather forecast. *arXiv preprint arXiv:2402.01295*, 2024.

- [79] Kun Yi, Qi Zhang, Wei Fan, Hui He, Liang Hu, Pengyang Wang, Ning An, Longbing Cao, and Zhendong Niu. Fouriergnn: rethinking multivariate time series forecasting from a pure graph perspective. In *Proceedings of the 37th International Conference on Neural Information Processing Systems*, NIPS ’23, Red Hook, NY, USA, 2023. Curran Associates Inc.
- [80] Jie Yin, Yao Gao, Ruishan Chen, Dapeng Yu, Robert Wilby, Nigel Wright, Yong Ge, Jeremy Bricker, Huili Gong, and Mingfu Guan. Flash floods: why are more of them devastating the world’s driest regions? *Nature*, 615(7951):212–215, 2023.
- [81] Zeda Yin, Linglong Bian, Beichao Hu, Jimeng Shi, and Arturo S Leon. Physic-informed neural network approach coupled with boundary conditions for solving 1d steady shallow water equations for riverine system. In *World Environmental and Water Resources Congress 2023*, pages 280–288, 2023.
- [82] Zhihan Yue, Yujing Wang, Juanyong Duan, Tianmeng Yang, Congrui Huang, Yunhai Tong, and Bixiong Xu. Ts2vec: Towards universal representation of time series. In *Proceedings of the AAAI conference on artificial intelligence*, volume 36, pages 8980–8987, 2022.
- [83] Shuaihong Zang, Zhijia Li, Ke Zhang, Cheng Yao, Zhiyu Liu, Jingfeng Wang, Yingchun Huang, and Sheng Wang. Improving the flood prediction capability of the xin’ anjiang model by formulating a new physics-based routing framework and a key routing parameter estimation method. *Journal of Hydrology*, 603:126867, 2021.
- [84] Ailing Zeng, Muxi Chen, Lei Zhang, and Qiang Xu. Are transformers effective for time series forecasting? In *Proceedings of the AAAI conference on artificial intelligence*, volume 37, pages 11121–11128, 2023.
- [85] Daochen Zha, Zaid Pervaiz Bhat, Kwei-Herng Lai, Fan Yang, Zhimeng Jiang, Shaochen Zhong, and Xia Hu. Data-centric artificial intelligence: A survey. *ACM Computing Surveys*, 57(5):1–42, 2025.
- [86] Haoyi Zhou, Shanghang Zhang, Jieqi Peng, Shuai Zhang, Jianxin Li, Hui Xiong, and Wancai Zhang. Informer: Beyond efficient transformer for long sequence time-series forecasting. In *The Thirty-Fifth AAAI Conference on Artificial Intelligence, AAAI 2021, Virtual Conference*, volume 35, pages 11106–11115. AAAI Press, 2021.
- [87] Haoyi Zhou, Shanghang Zhang, Jieqi Peng, Shuai Zhang, Jianxin Li, Hui Xiong, and Wancai Zhang. Informer: Beyond efficient transformer for long sequence time-series forecasting. In *Proceedings of the AAAI conference on artificial intelligence*, volume 35, pages 11106–11115, 2021.
- [88] Tian Zhou, Ziqing Ma, Qingsong Wen, Xue Wang, Liang Sun, and Rong Jin. Fedformer: Frequency enhanced decomposed transformer for long-term series forecasting. In *International conference on machine learning*, pages 27268–27286. PMLR, 2022.
- [89] Tian Zhou, Peisong Niu, Xue Wang, Liang Sun, and Rong Jin. One fits all: power general time series analysis by pretrained lm. In *Proceedings of the 37th International Conference on Neural Information Processing Systems*, NIPS ’23, Red Hook, NY, USA, 2023. Curran Associates Inc.

Contents

1	Introduction	1
2	Related Work	3
2.1	Flood Dataset	3
2.2	Machine Learning for Forecasting	3
3	The SF²Bench Dataset	3
3.1	Overview	3
3.2	Preprocessing	4
3.3	Qualitative Analysis	5
4	Forecasting Benchmarks	5
4.1	Problem Definition	5
4.2	Metric	6
4.3	Methods	6
4.4	Experiment Setup	6
4.5	Results & Observations	7
5	Conclusion	9
A	Detailed information of SF²Bench	17
B	Experimental Details	18
B.1	Data Preprocessing	18
B.2	Methods	18
B.3	Platform	19
C	Detailed Results	20
C.1	Detailed Benchmark Results	20
C.2	Detailed Other Results	20

A Detailed information of SF²Bench

We provide the detailed information about the number of monitor stations in each split in Table 7. In each split, the number of water monitor stations is more than other kinds of features, which means the water level feature is the main information and others are additional parts.

Table 7: The summary of the all splits in SF²Bench

Splits	Time Span	Interval	Water	Groundwater	Rainfall	Pump	Gate
S0	1985-1990	1 Hour	159	40	143	17	82
S1	1990-1995	1 Hour	227	36	139	18	104
S2	1995-2000	1 Hour	332	44	170	26	94
S3	2000-2005	1 Hour	402	178	227	31	107
S4	2005-2010	1 Hour	518	296	254	48	172
S5	2010-2015	1 Hour	585	333	216	65	256
S6	2015-2020	1 Hour	670	317	186	85	300
S7	2020-2024	1 Hour	716	352	194	89	329

We also provide the geographical distribution information of monitor stations in different splits in Figure 6. As time goes by, the number of monitoring stations gradually increases, and in terms of spatial distribution, the locations of monitoring stations remain consistent.




Figure 6: The location distribution in each split.

In addition, we visualize the temporal pattern of features over a year in each split. As shown in Figure 7, 8, 9, and 10, the temporal patterns in 8 splits are similar and consistent. From the view of the climate, the dry and rainy seasons are highly consistent but the intensity of rainfall in different splits is different. These patterns that are highly aligned with the actual situation demonstrate the quality of the dataset.

B Experimental Details

B.1 Data Preprocessing

Normalization. To make the data have a zero mean and unit variance, we follow [19, 87, 82] using z-score to normalize the time series data. For the time series, whose variance is less than 1E-4, we regard its variance as one to avoid the variable overflow(inf or NaN). For forecasting tasks, all the report metrics are based on the normalized data.

Timestamp Features. Because of some methods, such as NLinear, we do not consider extracting the timestamp feature as part of the input. For those methods that are capable of handling the timestamp feature, we ignore this part. In our code, we also provide timestamp features extracted by following [87, 82] for further works.

B.2 Methods

MLP. We implement a classical three-layer MLP with ReLU as the activation function. The input layer dimension is determined by the input sequence length, and the output layer dimension corresponds to the forecast horizon. The hidden layer dimension is 64. For training, the learning rate is 1×10^{-4} , weight decay is 1×10^{-6} , and the batch size is 64. The model is trained for 15 epochs.

NLinear [84]. This is a simple linear model that treats each time series independently, modeling future values using a linear transformation of the most recent input values. Implementation is based on the NeuralForecast library⁷. The learning rate is 1×10^{-4} , weight decay is 1×10^{-6} , batch size is 64, and training is performed for 50 epochs.

TSMixer [14]. Inspired by MLP-Mixer models from vision tasks, TSMixer is a neural network architecture for time series forecasting. It alternately applies MLPs along the time and feature axes, learning dependencies across both dimensions without requiring attention mechanisms or complex sequence modeling. Implementation follows the NeuralForecast default settings. The architecture includes 2 mixing layers, and the second feed-forward layer has 64 units. The learning rate is 1×10^{-4} , batch size is 32, and the model is trained for 10 epochs.

TCN [5]. It incorporates causal convolutions, ensuring predictions depend only on current and past inputs, thus preserving temporal order. They also utilize dilated convolutions to efficiently capture long-range dependencies by expanding the receptive field without significantly increasing layers. Our implementation follows the official code⁸ using the popular channel-wise setting. It employs a three-layer backbone with a kernel size of 3 and a fixed dilation of 1. The learning rate is 1×10^{-3} , weight decay is 1×10^{-7} , batch size is 256, and training is performed for 50 epochs.

ModernTCN [45]. ModernTCN introduces a streamlined, fully convolutional architecture that aims to simplify design while enhancing performance. It incorporates components like depth-wise separable convolutions and Gated Linear Units (GLUs) to efficiently capture local and long-range temporal dependencies. Implementation follows the long-term forecasting settings from the source code⁹, using the Weather dataset hyperparameters as defaults. The learning rate is 1×10^{-4} , batch size is 256, and the model is trained for 100 epochs.

TimesNet [72]. TimesNet models temporal variations in a two-dimensional space by reshaping time series data into a pseudo-image format and applying 2D convolutional techniques. This enables it to capture both short-term dynamics and long-term dependencies. Implementation uses the Time-Series-Library¹⁰, with default hyperparameters from the long-term forecasting setting for the Weather dataset. The learning rate is 1×10^{-4} , batch size is 32, and training is performed for 10 epochs.

LSTM [25]. Our Long Short-Term Memory (LSTM) implementation is a two-layer model with a

⁷<https://github.com/Nixtla/neuralforecast>

⁸<https://github.com/locuslab/TCN/tree/master>

⁹<https://github.com/luodhhh/ModernTCN>

¹⁰<https://github.com/thuml/Time-Series-Library>

hidden dimension of 32. The learning rate is 1×10^{-3} , weight decay is 1×10^{-6} , batch size is 64, and the model is trained for 50 epochs.

DeepAR [58]. DeepAR is a global model trained on multiple related time series, which aids generalization, especially for series with limited history. It employs an RNN architecture to predict future values by modeling the conditional distribution of the next value given past observations. Implementation is based on the NeuralForecast library. The learning rate is 1×10^{-3} , batch size is 64, and training is performed for 20 epochs.

DilatedRNN [13]. Dilated RNNs exponentially expand their receptive field by stacking layers with different dilation factors, allowing efficient capture of short- and long-range patterns without a drastic increase in parameters. This makes them suitable for forecasting tasks with wide-ranging temporal dependencies. Implementation is based on the NeuralForecast library. The learning rate is 1×10^{-3} , batch size is 64, and training is performed for 40 epochs.

GCN [32]. The GCN architecture consists of two layers with a hidden dimension of 32. The graph topology is derived from location information using Delaunay triangulation¹¹. The learning rate is 1×10^{-4} , weight decay is 1×10^{-5} , batch size is 32, and the model is trained for 50 epochs.

FourierGNN [79]. FourierGNN leverages graph neural networks and Fourier transforms to capture temporal and inter-variable dependencies. Time series variables are treated as graph nodes, with edges representing their relationships. Fourier transforms project data into the frequency domain to model periodic and long-range dependencies. Implementation follows the source code¹². The learning rate is 1×10^{-5} , batch size is 32, and training is performed for 100 epochs.

StemGNN [11]. StemGNN is designed to capture both temporal (via temporal convolutions) and spatial (via spectral graph convolutions) dependencies in time-series data, learning smooth representations over the graph structure and dynamic patterns. Implementation follows the source code¹³. The learning rate is 1×10^{-4} , batch size is 32, and the model is trained for 50 epochs.

iTransformer [41]. The iTransformer uses an encoder-decoder structure where the encoder processes the sequence in reverse order. This allows the decoder to predict future values based on this processed representation, enhancing focus on relevant temporal sequences while mitigating the computational cost of traditional transformers. Implementation is based on the NeuralForecast library. The learning rate is 1×10^{-4} , batch size is 32, and training is performed for 10 epochs.

PatchTST [48]. PatchTST is a Transformer-based architecture employing patching and channel independence. Time series are divided into patches, which are transformed into tokens and processed by a transformer model to capture local and global dependencies via self-attention. This is particularly useful for long-term forecasting. Implementation is based on the NeuralForecast library. The learning rate is 1×10^{-4} , batch size is 128, and training is performed for 100 epochs.

GPT4TS [89]. GPT4TS treats time series as a language, leveraging pretrained language models (LLMs) to learn temporal patterns. Time series data is tokenized for LLM processing, enabling zero-shot or few-shot generalization. Implementation follows the source code¹⁴, using long-term forecasting settings for the Weather dataset as default hyperparameters. The default language model is GPT-2 [55]. The learning rate is 1×10^{-4} , batch size is 64, and training is performed for 10 epochs.

AutoTimes [42]. AutoTimes projects time series segments into the embedding space of language tokens, leveraging the autoregressive capabilities of LLMs for forecasting. By training the model to predict subsequent time series segments given preceding ones, AutoTimes generates multi-step forecasts. Implementation follows the source code¹⁵, with GPT-2 [55] as the language model. The learning rate is 5×10^{-4} , batch size is 64, and training is performed for 10 epochs.

For all benchmark experiments, AdamW [44] is used as the optimizer, and the loss function is Mean Squared Error (MSE).

B.3 Platform

All experiments are conducted on two Linux machines, one with 8 NVIDIA A100 GPUs, each with 40GB of memory, and another with 4 RTX 4090 GPUs. We used Python 3.12.9 and Pytorch 2.6.0 to construct our project.

¹¹<https://docs.scipy.org/doc/scipy/reference/generated/scipy.spatial.Delaunay.html>

¹²<https://github.com/aikunyi/FourierGNN>

¹³<https://github.com/microsoft/StemGNN>

¹⁴<https://github.com/DAMO-DI-ML/NeurIPS2023-One-Fits-All>

¹⁵<https://github.com/thuml/AutoTimes/tree/main>

C Detailed Results

C.1 Detailed Benchmark Results

In Section 4.5, we report the average of our benchmark results. In this section, we report the detailed results, including MSE, MAE, and three SEDI values on different prediction lengths, on three interest areas(Orlando, Miami, and Fort Myers) from Table 8 to 37. The best average results are presented in bold font, while the second-best are underlined. Furthermore, from Table 38 to 42, we report the detailed results of basic methods on the whole dataset. To provide qualitative analysis, from Figure 11 to 16, we demonstrate the case visualization of each method on the S7 split in the Orlando area.

C.2 Detailed Other Results

To provide more information about the ablation study, we report the detailed results of the factor ablation study in Table 43 to 45. We report the average results on three interest areas on the S6 split. For the spatial and temporal information ablation study, the results are available from Table 46 to 49.




Figure 7: The temporal pattern of features of S0 and S1.




Figure 8: The temporal pattern of features of S2 and S3.




Figure 9: The temporal pattern of features of S4 and S5.




Figure 10: The temporal pattern of features of S6 and S7.




Figure 11: Cases visualization of MLP-based architecture methods on S7(Orlando area).




Figure 12: Cases visualization of CNN-based architecture methods on S7(Orlando area). Modern. means ModernTCN.




Figure 13: Cases visualization of Transformer-based architecture methods on S7(Orlando area). iTrans. means iTransformer.




Figure 14: Cases visualization of RNN-based architecture methods on S7(Orlando area). Dilated. means DilatedRNN.




Figure 15: Cases visualization of GNN-based architecture methods on S7(Orlando area). Fourier. means FourierGNN.




Figure 16: Cases visualization of LLM-based architecture methods on S7(Orlando area). Dilated. means DilatedRNN.

Table 8: Benchmark(MAE) on part 1(Orlando area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.0915	0.1229	0.0679	0.0810	0.0716	0.0691	0.0598	0.0846	0.0810
	3D	0.1424	0.1928	0.1167	0.1437	0.1174	0.1154	0.1086	0.1314	0.1335
	5D	0.1746	0.2472	0.1607	0.1844	0.1503	0.1529	0.1418	0.1718	0.1730
	7D	0.1919	0.2899	0.1848	0.2244	0.1750	0.1793	0.1753	0.1970	0.2022
	Avg.	0.1501	<u>0.2132</u>	0.1325	0.1584	<u>0.1286</u>	0.1292	0.1214	0.1462	0.1474
MLP	1D	0.1046	0.1355	0.0812	0.0846	0.0767	0.0778	0.1184	0.1015	0.0975
	3D	0.1494	0.2140	0.1252	0.1378	0.1185	0.1219	0.1500	0.1458	0.1453
	5D	0.1769	0.2673	0.1595	0.1783	0.1496	0.1547	0.1751	0.1798	0.1802
	7D	0.1938	0.3093	0.1892	0.2120	0.1765	0.1819	0.2098	0.2096	0.2103
	Avg.	0.1562	0.2315	0.1388	0.1532	0.1303	0.1341	0.1633	0.1592	0.1583
NLinear	1D	0.0980	0.1279	0.0777	0.0820	0.0771	0.0741	0.0640	0.0929	0.0867
	3D	0.1475	0.2109	0.1234	0.1390	0.1207	0.1211	0.1079	0.1415	0.1390
	5D	0.1835	0.2663	0.1584	0.1814	0.1527	0.1541	0.1422	0.1763	0.1769
	7D	0.1964	0.3085	0.1891	0.2161	0.1803	0.1814	0.1711	0.2065	0.2062
	Avg.	0.1563	0.2284	0.1371	0.1546	0.1327	0.1327	0.1213	0.1543	0.1522
TCN	1D	0.1887	0.1417	0.2145	0.3830	0.1322	0.0889	0.1164	0.1913	0.1821
	3D	0.1588	0.2302	0.1857	0.4058	0.1968	0.1359	0.1471	0.2127	0.2091
	5D	0.2110	0.3028	0.2139	0.4483	0.2128	0.1928	0.2024	0.2579	0.2552
	7D	0.2135	0.3379	0.2605	0.5715	0.2686	0.2052	0.2102	0.2695	0.2921
	Avg.	0.1930	0.2532	0.2186	0.4522	0.2026	0.1557	0.1690	0.2329	0.2346
CNN	1D	0.0893	0.1238	0.0721	0.0760	0.0704	0.0661	0.0588	0.0802	0.0796
	3D	0.1486	0.2113	0.1366	0.1450	0.1212	0.1162	0.1073	0.1324	0.1398
	5D	0.1886	0.2685	0.1798	0.1887	0.1632	0.1537	0.1426	0.1697	0.1818
	7D	0.2140	0.3228	0.2240	0.2310	0.1936	0.1833	0.1728	0.2002	0.2177
	Avg.	0.1601	0.2316	0.1531	0.1602	0.1371	<u>0.1298</u>	0.1204	<u>0.1456</u>	0.1547
TimesNet	1D	0.1068	0.1436	0.0805	0.0909	0.0919	0.0925	0.0787	0.1051	0.0987
	3D	0.1558	0.2190	0.1252	0.1515	0.1382	0.1402	0.1254	0.1582	0.1517
	5D	0.1813	0.2745	0.1637	0.1933	0.1735	0.1766	0.1578	0.1958	0.1896
	7D	0.2034	0.3222	0.2025	0.2309	0.2102	0.2146	0.1886	0.2263	0.2248
	Avg.	0.1618	0.2398	0.1430	0.1666	0.1534	0.1560	0.1376	0.1714	0.1662
Transformer	1D	0.0865	0.1227	0.0645	0.0730	0.0710	0.0700	0.0629	0.0889	0.0799
	3D	0.1395	0.1963	0.1103	0.1313	0.1193	0.1195	0.1058	0.1376	0.1325
	5D	0.1717	0.2546	0.1471	0.1723	0.1534	0.1528	0.1401	0.1775	0.1712
	7D	0.1889	0.2962	0.1801	0.2068	0.1833	0.1830	0.1708	0.2048	0.2017
	Avg.	<u>0.1466</u>	0.2175	<u>0.1255</u>	0.1459	0.1317	0.1313	<u>0.1199</u>	0.1522	<u>0.1463</u>
PatchTST	1D	0.0843	0.1148	0.0638	0.0678	0.0711	0.0693	0.0617	0.0858	0.0773
	3D	0.1365	0.1922	0.1113	0.1269	0.1149	0.1199	0.1046	0.1362	0.1303
	5D	0.1677	0.2451	0.1471	0.1728	0.1489	0.1569	0.1409	0.1718	0.1689
	7D	0.1866	0.2927	0.1787	0.2054	0.1786	0.1859	0.1679	0.2019	0.1997
	Avg.	0.1438	0.2112	0.1252	0.1432	0.1284	0.1330	0.1188	0.1489	0.1441

Table 9: Benchmark(MAE) on part 1(Orlando area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.1275	0.1790	0.0935	0.1714	0.1044	0.1032	0.0865	0.1211	0.1233
	3D	0.1652	0.2751	0.1431	0.2458	0.1518	0.1528	0.1481	0.1647	0.1808
	5D	0.1972	0.3297	0.1801	0.2977	0.1871	0.1891	0.1784	0.1985	0.2197
	7D	0.2139	0.3517	0.2277	0.3351	0.2198	0.2143	0.1978	0.2266	0.2484
	Avg.	0.1759	0.2839	0.1611	0.2625	0.1658	0.1648	0.1527	0.1777	0.1931
RNN	1D	0.1209	0.1855	0.0946	0.1726	0.1024	0.0971	0.0881	0.1171	0.1223
	3D	0.1667	0.2677	0.1488	0.2465	0.1534	0.1620	0.1421	0.1736	0.1826
	5D	0.1942	0.3110	0.1907	0.3237	0.1889	0.1828	0.1770	0.2045	0.2216
	7D	0.2146	0.3522	0.2210	0.3619	0.2122	0.2197	0.2123	0.2317	0.2532
	Avg.	0.1741	0.2791	0.1638	0.2762	0.1642	0.1654	0.1548	0.1817	0.1949
DilatedRNN	1D	0.0939	0.1317	0.0700	0.1552	0.0836	0.0748	0.0672	0.1006	0.0971
	3D	0.1499	0.2318	0.1249	0.2031	0.1339	0.1318	0.1224	0.1457	0.1554
	5D	0.1794	0.2793	0.1627	0.2897	0.1678	0.1635	0.1520	0.1809	0.1969
	7D	0.2110	0.3329	0.1959	0.3061	0.1990	0.1945	0.1931	0.1478	0.2225
	Avg.	0.1586	0.2439	0.1384	0.2385	0.1461	0.1411	0.1337	0.1437	0.1680
GCN	1D	0.1003	0.2139	0.1034	0.3710	0.1030	0.1005	0.0775	0.0962	0.1457
	3D	0.1543	0.3137	0.1411	0.3384	0.1524	0.1517	0.1208	0.1392	0.1889
	5D	0.1862	0.3741	0.1665	0.4935	0.1893	0.1818	0.1610	0.1732	0.2407
	7D	0.2034	0.4156	0.1912	0.5967	0.2153	0.2073	0.1927	0.2024	0.2781
	Avg.	0.1610	0.3293	0.1505	0.4499	0.1650	0.1603	0.1380	0.1528	0.2134
GNN	1D	0.0985	0.1324	0.0796	0.1337	0.0863	0.0929	0.0672	0.0927	0.0979
	3D	0.1445	0.2034	0.1178	0.2733	0.1295	0.1246	0.1125	0.1392	0.1556
	5D	0.1746	0.2644	0.1524	0.3672	0.1712	0.1640	0.1513	0.1719	0.2021
	7D	0.1918	0.3028	0.1936	0.4107	0.1992	0.2019	0.1871	0.2068	0.2367
	Avg.	0.1523	0.2257	0.1358	0.2962	0.1466	0.1459	0.1295	0.1527	0.1731
StemGNN	1D	0.1270	0.1782	0.0786	0.6896	0.1005	0.0833	0.0741	0.0969	0.1785
	3D	0.1717	0.3602	0.1536	0.6742	0.1778	0.1720	0.1498	0.1715	0.2539
	5D	0.2081	0.4895	0.2141	1.4591	0.2144	0.2729	0.2258	0.2412	0.4156
	7D	0.2477	0.5122	0.2577	1.4988	0.2552	0.3307	0.2486	0.2419	0.4491
	Avg.	0.1886	0.3850	0.1760	1.0804	0.1870	0.2147	0.1746	0.1879	0.3243
GPT4TS	1D	0.1050	0.1372	0.0833	0.0934	0.0905	0.0898	0.0778	0.1106	0.0984
	3D	0.1530	0.2169	0.1221	0.1502	0.1335	0.1380	0.1222	0.1547	0.1488
	5D	0.1817	0.2650	0.1565	0.1928	0.1666	0.1723	0.1559	0.1948	0.1857
	7D	0.1967	0.3105	0.1930	0.2294	0.1969	0.2054	0.1962	0.2224	0.2188
	Avg.	0.1591	0.2324	0.1387	0.1664	0.1469	0.1514	0.1380	0.1706	0.1629
LLM	1D	0.0967	0.1369	0.0747	0.0770	0.0768	0.0784	0.0660	0.0963	0.0878
	3D	0.1453	0.2057	0.1196	0.1296	0.1164	0.1203	0.1099	0.1409	0.1360
	5D	0.1733	0.2664	0.1499	0.1751	0.1498	0.1534	0.1411	0.1739	0.1729
	7D	0.1902	0.3029	0.1814	0.2113	0.1780	0.1823	0.1691	0.2032	0.2023
	Avg.	0.1514	0.2280	0.1314	0.1482	0.1302	0.1336	0.1215	0.1536	0.1497

Table 10: Benchmark(MSE) on part 1(Orlando area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.1472	0.1254	0.0376	0.0610	0.0379	0.0275	0.0248	0.0523	0.0642
	3D	0.2661	0.2014	0.0621	0.1232	0.0681	0.0536	0.0465	0.0913	0.1140
	5D	0.3569	0.2609	0.0875	0.1714	0.0999	0.0772	0.0670	0.1226	0.1554
	7D	0.2963	0.3124	0.1097	0.2215	0.1199	0.0975	0.0871	0.1498	0.1743
	Avg.	0.2666	0.2250	0.0742	0.1443	0.0814	0.0640	0.0564	0.1040	0.1270
MLP	1D	0.1649	0.1283	0.0423	0.0696	0.0380	0.0312	0.0661	0.0633	0.0755
	3D	0.2915	0.2214	0.0679	0.1255	0.0694	0.0575	0.0862	0.1000	0.1274
	5D	0.1769	0.2903	0.0936	0.1673	0.0981	0.0812	0.1010	0.1325	0.1626
	7D	0.2817	0.3469	0.1190	0.2048	0.1248	0.1031	0.1384	0.1644	0.1854
	Avg.	0.2688	0.2467	0.0807	0.1418	<u>0.0826</u>	0.0682	0.0979	0.1150	0.1377
NLinear	1D	0.1468	0.1294	0.0449	0.0678	0.0405	0.0308	0.0291	0.0580	0.0684
	3D	0.2830	0.2223	0.0699	0.1257	0.0720	0.0575	0.0510	0.0974	0.1223
	5D	0.4103	0.2915	0.0950	0.1689	0.1013	0.0812	0.0722	0.1300	0.1688
	7D	0.3001	0.3481	0.1204	0.2080	0.1291	0.1029	0.0924	0.1617	0.1828
	Avg.	0.2851	0.2478	0.0825	<u>0.1426</u>	0.0857	<u>0.0681</u>	0.0612	0.1118	0.1356
TCN	1D	0.3257	0.1416	0.1462	0.8044	0.1152	0.0344	0.0567	0.1260	0.2188
	3D	0.2475	0.2295	0.1342	0.9118	0.1694	0.0597	0.0731	0.1423	0.2459
	5D	0.2906	0.3102	0.1367	0.6579	0.1763	0.1011	0.1119	0.1884	0.2466
	7D	0.3160	0.3587	0.1805	1.3613	0.2556	0.1105	0.1087	0.2119	0.3629
	Avg.	0.2950	0.2600	0.1494	0.9338	0.1791	0.0764	0.0876	0.1671	0.2686
CNN	1D	0.1391	0.1432	0.0407	0.0778	0.0392	0.0279	0.0255	0.0520	0.0682
	3D	0.2746	0.2966	0.1185	0.1814	0.0799	0.0575	0.0505	0.0921	0.1439
	5D	0.3575	0.3305	0.1278	0.2059	0.1303	0.0850	0.0723	0.1257	0.1794
	7D	0.3067	0.4557	0.1821	0.2844	0.1657	0.1096	0.0950	0.1579	0.2196
	Avg.	0.2695	0.3065	0.1173	0.1874	0.1038	0.0700	0.0608	0.1069	0.1528
TimesNet	1D	0.1770	0.1473	0.0429	0.0742	0.0546	0.0449	0.0394	0.0761	0.0821
	3D	0.3474	0.2545	0.0758	0.1626	0.1049	0.0829	0.0737	0.1365	0.1548
	5D	0.3436	0.3207	0.1069	0.2093	0.1475	0.1193	0.0939	0.1907	0.1915
	7D	0.3053	0.4134	0.1426	0.2522	0.2020	0.1648	0.1283	0.2267	0.2294
	Avg.	0.2933	0.2840	0.0921	0.1746	0.1273	0.1030	0.0838	0.1575	0.1644
Transformer	1D	0.1277	0.1337	0.0357	0.0674	0.0382	0.0314	0.0295	0.0619	0.0657
	3D	0.2760	0.2269	0.0688	0.1343	0.0755	0.0627	0.0529	0.1057	0.1254
	5D	0.3581	0.3195	0.1004	0.1846	0.1081	0.0855	0.0769	0.1483	0.1727
	7D	0.2999	0.3668	0.1288	0.2222	0.1384	0.1129	0.1003	0.1810	0.1938
	Avg.	0.2654	0.2617	0.0834	0.1521	0.0901	0.0731	0.0649	0.1242	0.1394
PatchTST	1D	0.1315	0.1368	0.0370	0.0608	0.0385	0.0325	0.0321	0.0593	0.0661
	3D	0.3008	0.2298	0.0845	0.1276	0.0713	0.0645	0.0535	0.1041	0.1295
	5D	0.3504	0.2951	0.1222	0.1736	0.1042	0.0946	0.0777	0.1368	0.1693
	7D	0.3145	0.3605	0.1365	0.2170	0.1412	0.1198	0.0968	0.1708	0.1946
	Avg.	0.2743	0.2555	0.0951	0.1448	0.0888	0.0778	0.0650	0.1177	0.1399

Table 11: Benchmark(MSE) on part 1(Orlando area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.2408	0.2451	0.0505	0.2788	0.0655	0.0496	0.0435	0.0846	0.1323
	3D	0.2765	0.3219	0.0784	0.3653	0.1097	0.0768	0.0837	0.1162	0.1786
	5D	0.3053	0.3853	0.1085	0.4750	0.1569	0.1011	0.0953	0.1525	0.2225
	7D	0.3058	0.4057	0.1427	0.5133	0.1855	0.1226	0.1103	0.1818	0.2459
	Avg.	0.2821	0.3395	0.0950	0.4081	0.1294	0.0875	0.0832	0.1338	0.1948
RNN	1D	0.2406	0.2367	0.0480	0.2190	0.0615	0.0435	0.0386	0.0782	0.1208
	3D	0.2720	0.3088	0.0793	0.4084	0.1117	0.0783	0.0652	0.1212	0.1806
	5D	0.3033	0.3441	0.1072	0.5899	0.1537	0.1003	0.0883	0.1564	0.2304
	7D	0.3046	0.4056	0.1356	0.5250	0.1639	0.1221	0.1070	0.1800	0.2430
	Avg.	0.2801	0.3238	0.0925	0.4356	0.1227	0.0860	0.0748	0.1340	0.1937
DilatedRNN	1D	0.1818	0.1370	0.0372	0.3297	0.0496	0.0356	0.0304	0.0860	0.1109
	3D	0.2517	0.2674	0.0689	0.3124	0.0887	0.0645	0.0722	0.1150	0.1551
	5D	0.2822	0.3308	0.1004	0.6438	0.1338	0.0915	0.0916	0.1423	0.2270
	7D	0.3125	0.3989	0.1242	0.5068	0.1702	0.1159	0.1257	0.0714	0.2282
	Avg.	0.2571	0.2835	0.0827	0.4482	0.1106	0.0769	0.0800	0.1037	0.1803
GCN	1D	0.1465	0.5157	0.0443	1.1053	0.0519	0.0353	0.0275	0.0518	0.2473
	3D	0.2526	1.6031	0.0656	0.7137	0.0944	0.0654	0.0494	0.0891	0.3667
	5D	0.3055	1.9044	0.0867	1.1121	0.1266	0.0895	0.0730	0.1184	0.4770
	7D	0.2687	2.2856	0.1071	1.4113	0.1651	0.1087	0.0928	0.1471	0.5733
	Avg.	0.2433	1.5772	<u>0.0759</u>	1.0856	0.1095	0.0747	0.0607	0.1016	0.4161
GNN	1D	0.1619	0.1324	0.0412	0.0901	0.0442	0.0326	0.0275	0.0575	0.0734
	3D	0.2714	0.2058	0.0649	0.2944	0.0746	0.0560	0.0489	0.0941	0.1388
	5D	0.3066	0.2721	0.0891	0.4409	0.1106	0.0806	0.0705	0.1238	0.1868
	7D	0.2766	0.3139	0.1154	0.4880	0.1381	0.1065	0.0919	0.1528	0.2104
	Avg.	0.2541	0.2310	0.0777	0.3283	0.0919	0.0689	0.0597	0.1071	0.1523
StemGNN	1D	0.2290	0.1703	0.0403	1.1688	0.0652	0.0410	0.0325	0.0723	0.2274
	3D	0.2703	0.4534	0.0903	1.2569	0.1323	0.0896	0.0743	0.1263	0.3117
	5D	0.3017	0.7711	0.1305	3.3143	0.1663	0.1663	0.1417	0.2080	0.6500
	7D	0.3225	0.7901	0.1768	3.4285	0.1979	0.2323	0.1489	0.1864	0.6854
	Avg.	0.2809	0.5462	0.1095	2.2921	0.1404	0.1323	0.0994	0.1483	0.4686
GPT4TS	1D	0.1984	0.1399	0.0456	0.0882	0.0542	0.0431	0.0387	0.0935	0.0877
	3D	0.3883	0.2385	0.0811	0.1648	0.0986	0.0788	0.0690	0.1343	0.1567
	5D	0.3668	0.3102	0.0997	0.2037	0.1304	0.1091	0.0870	0.1876	0.1868
	7D	0.2928	0.3700	0.1378	0.2503	0.1622	0.1494	0.1319	0.2233	0.2147
	Avg.	0.3116	0.2646	0.0911	0.1768	0.1114	0.0951	0.0816	0.1597	0.1615
LLM	1D	0.1531	0.1507	0.0401	0.0671	0.0390	0.0310	0.0287	0.0623	0.0715
	3D	0.3072	0.2333	0.0683	0.1278	0.0686	0.0572	0.0508	0.0978	0.1264
	5D	0.3527	0.2975	0.0916	0.1700	0.0992	0.0811	0.0715	0.1319	0.1619
	7D	0.2913	0.3428	0.1181	0.2061	0.1265	0.1031	0.0916	0.1612	0.1801
	Avg.	0.2761	0.2561	0.0795	0.1427	0.0833	<u>0.0681</u>	<u>0.0606</u>	0.1133	0.1350

Table 12: Benchmark(SEDI(10%)) on part 1(Orlando area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.8143	0.9334	0.9011	0.8780	0.6133	0.7659	0.9351	0.7772	0.8273
	3D	0.7599	0.8673	0.8592	0.9286	0.5201	0.7173	0.9040	0.7097	0.7833
	5D	0.6435	0.8095	0.8142	0.8702	0.4517	0.6956	0.8571	0.6573	0.7249
	7D	0.6022	0.8056	0.7759	0.8158	0.4198	0.6622	0.8516	0.5969	0.6912
	Avg.	0.7050	0.8540	0.8376	0.8731	0.5012	0.7103	0.8870	0.6853	0.7567
MLP	1D	0.7483	0.8690	0.8917	0.9141	0.5971	0.7394	0.7901	0.7191	0.7836
	3D	0.5937	0.7493	0.8103	0.8420	0.4798	0.6766	0.7626	0.6334	0.6935
	5D	0.5167	0.6755	0.7538	0.7783	0.4139	0.6214	0.7289	0.5685	0.6321
	7D	0.4706	0.6241	0.7042	0.7236	0.3734	0.5854	0.6798	0.5107	0.5840
	Avg.	0.5823	0.7295	0.7900	0.8145	0.4660	0.6557	0.7403	0.6079	0.6733
NLinear	1D	0.7729	0.8803	0.8951	0.9171	0.5984	0.7844	0.9259	0.7302	0.8130
	3D	0.6007	0.7581	0.8166	0.8390	0.4784	0.6961	0.8550	0.6410	0.7106
	5D	0.5198	0.6821	0.7576	0.7721	0.4129	0.6535	0.7956	0.5750	0.6461
	7D	0.4731	0.6290	0.7063	0.7159	0.3701	0.6155	0.7512	0.5173	0.5973
	Avg.	0.5916	0.7374	0.7939	0.8110	0.4649	0.6873	0.8319	0.6159	0.6918
TCN	1D	0.5634	0.8939	0.7243	0.7307	0.4343	0.7156	0.8132	0.5039	0.6724
	3D	0.5462	0.8010	0.6694	0.7131	0.3092	0.7072	0.7778	0.3687	0.6116
	5D	0.4606	0.7638	0.6291	0.6572	0.2804	0.6123	0.7136	0.3351	0.5565
	7D	0.4598	0.7014	0.6092	0.5674	0.2201	0.5762	0.5692	0.3620	0.5082
	Avg.	0.5075	0.7900	0.6580	0.6671	0.3110	0.6528	0.7184	0.3924	0.5872
CNN	1D	0.7912	0.8737	0.8857	0.8915	0.6023	0.7692	0.9240	0.7395	0.8096
	3D	0.6031	0.7637	0.7893	0.8041	0.4970	0.6746	0.8469	0.6370	0.7020
	5D	0.4960	0.6839	0.7201	0.7403	0.4277	0.6142	0.7898	0.5668	0.6298
	7D	0.4357	0.6233	0.6667	0.6818	0.3869	0.5726	0.7499	0.5076	0.5781
	Avg.	0.5815	0.7362	0.7654	0.7794	0.4785	0.6576	0.8277	0.6127	0.6799
TimesNet	1D	0.7224	0.8584	0.8701	0.8789	0.5727	0.7000	0.9025	0.7044	0.7762
	3D	0.5820	0.7493	0.7919	0.7959	0.4368	0.6345	0.7953	0.5956	0.6727
	5D	0.4866	0.6771	0.7336	0.7370	0.3854	0.5657	0.7453	0.5191	0.6062
	7D	0.4083	0.6298	0.6863	0.6808	0.3325	0.5181	0.7137	0.4662	0.5545
	Avg.	0.5498	0.7286	0.7705	0.7732	0.4318	0.6045	0.7892	0.5713	0.6524
Transformer	1D	0.8049	0.8881	0.9039	0.9057	0.5997	0.7663	0.9234	0.7186	0.8138
	3D	0.6433	0.7820	0.8294	0.8350	0.4807	0.6836	0.8500	0.6270	0.7164
	5D	0.5540	0.7202	0.7733	0.7744	0.4121	0.6302	0.7956	0.5481	0.6510
	7D	0.5043	0.6631	0.7177	0.7288	0.3558	0.5843	0.7486	0.4997	0.6003
	Avg.	0.6266	0.7634	0.8061	0.8110	0.4621	0.6661	0.8294	0.5983	0.6954
PatchTST	1D	0.8082	0.8872	0.8947	0.9082	0.5937	0.7683	0.9141	0.7252	0.8125
	3D	0.6607	0.7851	0.8203	0.8352	0.4994	0.6789	0.8537	0.6293	0.7203
	5D	0.5672	0.7188	0.7626	0.7659	0.4175	0.6272	0.7914	0.5625	0.6516
	7D	0.5081	0.6574	0.7167	0.7186	0.3708	0.5887	0.7506	0.5049	0.6020
	Avg.	0.6361	0.7621	0.7986	0.8070	0.4704	0.6658	0.8274	0.6055	0.6966

Table 13: Benchmark(SEDI(10%)) on part 1(Orlando area) stations(RNN, GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.7709	0.8270	0.8825	0.8781	0.5134	0.7182	0.9039	0.7302	0.7780
	3D	0.6027	0.8070	0.8162	0.8833	0.4448	0.5963	0.8482	0.6473	0.7057
	5D	0.5247	0.7396	0.7646	0.7848	0.3667	0.6529	0.7852	0.5842	0.6503
	7D	0.4409	0.6678	0.7459	0.7427	0.3836	0.6063	0.7591	0.4874	0.6042
	Avg.	0.5848	0.7604	0.8023	0.8222	0.4271	0.6434	0.8241	0.6123	0.6846
RNN	1D	0.7411	0.8286	0.9006	0.9469	0.5732	0.7033	0.9113	0.7681	0.7966
	3D	0.6168	0.7816	0.7859	0.7710	0.4175	0.6046	0.8704	0.6535	0.6877
	5D	0.5086	0.7014	0.7872	0.8217	0.3716	0.6180	0.7106	0.6096	0.6411
	7D	0.4686	0.7151	0.7627	0.8162	0.3346	0.6131	0.6993	0.5065	0.6145
	Avg.	0.5838	0.7567	0.8091	0.8390	0.4242	0.6348	0.7979	0.6344	0.6850
DilatedRNN	1D	0.8033	0.9139	0.8972	0.9067	0.6086	0.7706	0.9528	0.7437	0.8246
	3D	0.6053	0.8119	0.7907	0.8376	0.4752	0.5884	0.8004	0.6515	0.6951
	5D	0.5254	0.7325	0.7443	0.8017	0.4259	0.6454	0.8208	0.6270	0.6654
	7D	0.4664	0.7255	0.7337	0.7261	0.4031	0.6275	0.7647	0.5970	0.6305
	Avg.	0.6001	0.7960	0.7915	0.8180	0.4782	0.6580	0.8347	0.6548	0.7039
GCN	1D	0.8018	0.7777	0.8496	0.7072	0.5680	0.7274	0.9103	0.7172	0.7574
	3D	0.6240	0.7415	0.8169	0.6969	0.5115	0.6977	0.8412	0.6788	0.7011
	5D	0.5324	0.7147	0.7394	0.6223	0.4217	0.6488	0.8378	0.6075	0.6406
	7D	0.4949	0.6795	0.6887	0.5567	0.3824	0.6114	0.7941	0.5809	0.5986
	Avg.	0.6133	0.7283	0.7737	0.6458	0.4709	0.6713	0.8458	0.6461	0.6744
GNN	1D	0.8505	0.8952	0.9035	0.8991	0.5899	0.7981	0.9211	0.7834	0.8301
	3D	0.6929	0.8059	0.8177	0.7987	0.5029	0.6747	0.8521	0.6853	0.7288
	5D	0.6012	0.7985	0.7753	0.7296	0.4659	0.6222	0.8349	0.6308	0.6823
	7D	0.5372	0.7731	0.7619	0.6927	0.4257	0.6830	0.8133	0.5499	0.6546
	Avg.	0.6705	0.8182	0.8146	0.7800	0.4961	0.6945	0.8554	0.6624	0.7240
StemGNN	1D	0.6812	0.8950	0.8559	0.4137	0.5486	0.7748	0.9207	0.7578	0.7310
	3D	0.5934	0.7234	0.8229	0.3851	0.4289	0.6430	0.8470	0.6376	0.6352
	5D	0.5151	0.5492	0.6765	0.0829	0.3765	0.4472	0.7608	0.5663	0.4968
	7D	0.4576	0.5168	0.6141	0.0549	0.3011	0.4146	0.8487	0.5028	0.4638
	Avg.	0.5618	0.6711	0.7423	0.2341	0.4138	0.5699	0.8443	0.6161	0.5817
GPT4TS	1D	0.7588	0.8697	0.8790	0.8738	0.5594	0.7173	0.9066	0.6863	0.7814
	3D	0.5816	0.7515	0.8012	0.7983	0.4484	0.6322	0.8277	0.5967	0.6797
	5D	0.4917	0.6840	0.7480	0.7384	0.3970	0.5830	0.7591	0.5241	0.6157
	7D	0.4388	0.6309	0.6941	0.6878	0.3582	0.5430	0.7099	0.4672	0.5662
	Avg.	0.5677	0.7340	0.7805	0.7746	0.4408	0.6189	0.8008	0.5686	0.6607
AutoTimes	1D	0.7957	0.8638	0.8896	0.9167	0.5985	0.7343	0.9218	0.7225	0.8054
	3D	0.6299	0.7622	0.8190	0.8520	0.4892	0.6712	0.8516	0.6389	0.7142
	5D	0.5392	0.6780	0.7622	0.7817	0.4135	0.6243	0.7982	0.5804	0.6472
	7D	0.4895	0.6363	0.7204	0.7309	0.3710	0.5879	0.7553	0.5180	0.6012
	Avg.	0.6136	0.7351	0.7978	0.8203	0.4680	0.6544	0.8317	0.6149	0.6920

Table 14: Benchmark(SEDI(5%)) on part 1(Orlando area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.5128	0.9250	0.7735	0.8437	0.4047	0.7105	0.8731	0.6329	0.7095
	3D	0.5032	0.8575	0.7227	0.9190	0.3281	0.6590	0.8035	0.5460	0.6674
	5D	0.4580	0.7990	0.6804	0.8465	0.2779	0.6314	0.7573	0.5308	0.6227
	7D	0.4174	0.8105	0.6355	0.7808	0.2684	0.5818	0.7227	0.4304	0.5809
	Avg.	0.4729	0.8480	0.7030	0.8475	<u>0.3198</u>	<u>0.6457</u>	0.7892	<u>0.5350</u>	0.6451
MLP	1D	0.5204	0.8556	0.7619	0.8916	0.3911	0.7094	0.7193	0.6167	0.6833
	3D	0.3386	0.7182	0.6688	0.7902	0.3201	0.6698	0.6612	0.4533	0.5775
	5D	0.2734	0.6303	0.6004	0.7015	0.2664	0.5025	0.6127	0.3537	0.4926
	7D	0.2257	0.5757	0.5444	0.6277	0.2313	0.4536	0.5548	0.2847	0.4372
	Avg.	0.3395	0.6950	0.6439	0.7528	0.3022	0.5838	0.6370	0.4271	0.5477
NLinear	1D	0.5802	0.8689	0.7743	0.9014	0.4122	0.7091	0.8529	0.6285	0.7159
	3D	0.3752	0.7312	0.6721	0.7904	0.3451	0.6185	0.7482	0.4648	0.5932
	5D	0.3456	0.6392	0.6011	0.6967	0.2872	0.5471	0.6727	0.3603	0.5187
	7D	0.2322	0.5818	0.5408	0.6184	0.2505	0.4979	0.6121	0.2912	0.4531
	Avg.	0.3833	0.7053	0.6471	0.7517	0.3238	0.5932	0.7215	0.4362	0.5702
TCN	1D	0.3567	0.8742	0.2636	0.6239	0.2001	0.6360	0.4088	0.4724	0.4795
	3D	0.4252	0.7796	0.3238	0.6102	0.1573	0.4839	0.4081	0.3764	0.4456
	5D	0.2808	0.7319	0.2196	0.5901	0.1484	0.3628	0.4501	0.3326	0.3895
	7D	0.3380	0.6245	0.2753	0.4466	0.1223	0.3825	0.2879	0.2875	0.3456
	Avg.	0.3502	0.7526	0.2706	0.5677	0.1570	0.4663	0.3887	0.3672	0.4150
CNN	1D	0.5274	0.8480	0.7496	0.8700	0.3762	0.7081	0.8590	0.6339	0.6965
	3D	0.3525	0.7260	0.6128	0.7416	0.3183	0.5902	0.7544	0.4769	0.5716
	5D	0.2805	0.6333	0.5298	0.6535	0.2708	0.5110	0.6824	0.3742	0.4919
	7D	0.2334	0.5539	0.4603	0.5819	0.2352	0.4597	0.6208	0.3082	0.4317
	Avg.	0.3484	0.6903	0.5881	0.7118	0.3001	0.5673	0.7292	0.4483	0.5479
TimesNet	1D	0.4528	0.8341	0.7355	0.8507	0.3561	0.5952	0.8295	0.5725	0.6533
	3D	0.3185	0.7133	0.6279	0.7271	0.2782	0.5261	0.7186	0.4477	0.5447
	5D	0.2639	0.6133	0.5616	0.6561	0.2305	0.4305	0.6461	0.3400	0.4678
	7D	0.2184	0.5629	0.4993	0.5816	0.1851	0.3748	0.5961	0.2652	0.4104
	Avg.	0.3134	0.6809	0.6061	0.7039	0.2625	0.4816	0.6976	0.4064	0.5190
Transformer	1D	0.5441	0.8814	0.7818	0.8940	0.3864	0.7065	0.8510	0.5962	0.7052
	3D	0.3974	0.7637	0.6710	0.7924	0.3104	0.5943	0.7502	0.4656	0.5931
	5D	0.3080	0.6876	0.6000	0.7083	0.2453	0.5257	0.6786	0.3554	0.5136
	7D	0.2514	0.6234	0.5337	0.6420	0.2047	0.4701	0.6208	0.2896	0.4545
	Avg.	0.3752	0.7390	0.6466	0.7592	0.2867	0.5742	0.7252	0.4267	0.5666
PatchTST	1D	0.5491	0.8715	0.7655	0.8953	0.3878	0.7004	0.8402	0.6004	0.7013
	3D	0.3800	0.7584	0.6663	0.7948	0.3210	0.5907	0.7519	0.4659	0.5911
	5D	0.3052	0.6788	0.5937	0.6959	0.2631	0.5219	0.6719	0.3734	0.5130
	7D	0.2502	0.6018	0.5339	0.6317	0.2196	0.4734	0.6156	0.3054	0.4539
	Avg.	0.3711	0.7276	0.6399	0.7544	0.2979	0.5716	0.7199	0.4363	0.5648

Table 15: Benchmark(SEDI(5%)) on part 1(Orlando area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.4017	0.8107	0.7419	0.8659	0.3198	0.6701	0.8120	0.5639	0.6483
	3D	0.3042	0.7513	0.6543	0.8644	0.2781	0.5208	0.6626	0.3957	0.5539
	5D	0.2563	0.7351	0.5817	0.7538	0.2252	0.4925	0.6287	0.3346	0.5010
	7D	0.2557	0.6068	0.5389	0.7051	0.2231	0.4831	0.5848	0.2670	0.4581
	Avg.	0.3045	0.7260	0.6292	0.7973	0.2615	0.5416	0.6720	0.3903	0.5403
RNN	1D	0.5253	0.8201	0.7607	0.9462	0.3488	0.6711	0.8081	0.5904	0.6838
	3D	0.3304	0.7414	0.6407	0.7018	0.2825	0.5496	0.6188	0.4553	0.5401
	5D	0.2763	0.6725	0.5825	0.8152	0.2105	0.5123	0.5172	0.3650	0.4939
	7D	0.2556	0.6896	0.5741	0.8037	0.2025	0.4243	0.4710	0.3234	0.4680
	Avg.	0.3469	0.7309	0.6395	0.8167	0.2611	0.5393	0.6038	0.4335	0.5465
DilatedRNN	1D	0.5340	0.8917	0.7632	0.8782	0.3685	0.7112	0.8792	0.6419	0.7085
	3D	0.3663	0.8079	0.6105	0.7865	0.2975	0.5017	0.6871	0.4803	0.5672
	5D	0.3059	0.6907	0.5751	0.7343	0.2560	0.5750	0.6966	0.4549	0.5361
	7D	0.3052	0.6703	0.5421	0.6376	0.2480	0.5455	0.6476	0.4271	0.5029
	Avg.	0.3779	0.7652	0.6227	0.7591	0.2925	0.5833	0.7276	0.5010	0.5787
GCN	1D	0.5137	0.7241	0.7046	0.6410	0.3511	0.7351	0.8413	0.5751	0.6357
	3D	0.3818	0.6910	0.6424	0.6378	0.3101	0.7135	0.7400	0.4640	0.5726
	5D	0.3292	0.6557	0.5711	0.5477	0.2499	0.6674	0.7444	0.4156	0.5226
	7D	0.3021	0.6169	0.5142	0.4740	0.2346	0.5327	0.6955	0.4517	0.4777
	Avg.	0.3817	0.6719	0.6081	0.5751	0.2864	0.6622	0.7553	0.4766	0.5522
GNN	1D	0.5676	0.8926	0.7800	0.8434	0.3851	0.7068	0.8439	0.6727	0.7115
	3D	0.4629	0.7868	0.6564	0.7028	0.3275	0.5714	0.7651	0.5707	0.6054
	5D	0.3950	0.7889	0.6028	0.6333	0.2748	0.5038	0.7352	0.5055	0.5549
	7D	0.3873	0.7530	0.6115	0.5815	0.2572	0.5880	0.7235	0.4659	0.5460
	Avg.	0.4532	0.8053	0.6626	0.6903	0.3112	0.5925	0.7669	0.5537	0.6045
StemGNN	1D	0.3886	0.8590	0.7176	0.2562	0.3499	0.7175	0.8579	0.6320	0.5973
	3D	0.3247	0.6710	0.6642	0.1607	0.2578	0.5487	0.7490	0.5132	0.4862
	5D	0.2763	0.4268	0.4857	0.0234	0.2251	0.3075	0.6028	0.3641	0.3390
	7D	0.2189	0.3938	0.3786	0.0456	0.1851	0.2623	0.7196	0.3385	0.3178
	Avg.	0.3021	0.5877	0.5615	0.1215	0.2545	0.4590	0.7324	0.4619	0.4351
GPT4TS	1D	0.4904	0.8517	0.7383	0.8528	0.3567	0.6286	0.8333	0.5737	0.6657
	3D	0.3418	0.7131	0.6424	0.7412	0.2897	0.5298	0.7352	0.4407	0.5542
	5D	0.2811	0.6474	0.5588	0.6647	0.2451	0.4542	0.6619	0.3371	0.4813
	7D	0.2508	0.5786	0.4827	0.6002	0.2173	0.4100	0.5996	0.2547	0.4243
	Avg.	0.3410	0.6977	0.6055	0.7147	0.2772	0.5056	0.7075	0.4016	0.5314
LLM	1D	0.5235	0.8448	0.7499	0.9034	0.3786	0.6736	0.8310	0.5961	0.6876
	3D	0.3733	0.7387	0.6568	0.8177	0.3245	0.5828	0.7471	0.4660	0.5884
	5D	0.3023	0.6336	0.5923	0.7174	0.2638	0.5145	0.6738	0.3625	0.5075
	7D	0.2343	0.5856	0.5382	0.6401	0.2301	0.4620	0.6092	0.3000	0.4499
	Avg.	0.3584	0.7007	0.6343	0.7696	0.2993	0.5582	0.7153	0.4311	0.5584

Table 16: Benchmark(SEDI(1%)) on part 1(Orlando area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.2398	0.8330	0.3903	0.7439	0.2199	0.2365	0.4548	0.1809	0.4124
	3D	0.1994	0.7910	0.2967	0.7730	0.2051	0.1198	0.3723	0.1528	0.3638
	5D	0.1508	0.7647	0.2900	0.7580	0.1927	0.1346	0.3785	0.1380	0.3509
	7D	0.1877	0.7879	0.2579	0.6627	0.1935	0.1097	0.3831	0.1107	0.3367
	Avg.	0.1944	0.7942	0.3087	0.7344	0.2028	0.1502	0.3972	0.1456	0.3660
MLP	1D	0.2154	0.7569	0.3670	0.7701	0.2092	0.1228	0.2195	0.1553	0.3520
	3D	0.1462	0.5898	0.2661	0.5455	0.1401	0.0936	0.1884	0.0939	0.2580
	5D	0.1104	0.4818	0.1939	0.4161	0.1069	0.0697	0.1647	0.0636	0.2009
	7D	0.1078	0.3947	0.1464	0.3318	0.0871	0.0548	0.1533	0.0456	0.1652
	Avg.	0.1450	0.5558	0.2433	0.5159	0.1358	0.0852	0.1815	0.0896	0.2440
NLinear	1D	0.2205	0.7772	0.3813	0.7807	0.2302	0.2270	0.3975	0.1560	0.3963
	3D	0.1508	0.6056	0.2785	0.5478	0.1460	0.1937	0.3198	0.0888	0.2914
	5D	0.1156	0.4949	0.2062	0.4135	0.1131	0.1686	0.2773	0.0607	0.2313
	7D	0.1104	0.4048	0.1576	0.3231	0.0919	0.1536	0.2289	0.0453	0.1895
	Avg.	0.1493	0.5706	0.2559	0.5163	0.1453	0.1857	0.3059	0.0877	0.2771
TCN	1D	0.1520	0.7471	0.1639	0.2160	0.0801	0.0385	0.0000	0.0404	0.1797
	3D	0.1481	0.6252	0.1179	0.3218	0.0634	0.0303	0.0232	0.0397	0.1712
	5D	0.1935	0.5904	0.0941	0.2732	0.0516	0.0161	0.0496	0.0063	0.1593
	7D	0.1863	0.3808	0.1512	0.1816	0.0468	0.0121	0.0149	0.0185	0.1240
	Avg.	0.1700	0.5859	0.1318	0.2481	0.0605	0.0243	0.0219	0.0262	0.1586
CNN	1D	0.1937	0.7645	0.3466	0.7558	0.2153	0.2077	0.3743	0.1827	0.3801
	3D	0.1202	0.6078	0.2371	0.4858	0.1331	0.1410	0.2984	0.1105	0.2667
	5D	0.0829	0.5308	0.1742	0.3875	0.1018	0.1315	0.2742	0.0778	0.2201
	7D	0.0804	0.4283	0.1325	0.3070	0.0795	0.0623	0.2074	0.0606	0.1698
	Avg.	0.1193	0.5829	0.2226	0.4840	0.1324	0.1356	0.2886	0.1079	0.2592
TimesNet	1D	0.1451	0.7329	0.3521	0.6753	0.1455	0.1175	0.3185	0.1265	0.3267
	3D	0.0951	0.5803	0.2451	0.4443	0.1134	0.0911	0.2359	0.0766	0.2352
	5D	0.0651	0.4805	0.1894	0.3933	0.0900	0.0482	0.2077	0.0577	0.1915
	7D	0.0372	0.3977	0.1499	0.2984	0.0631	0.0399	0.1878	0.0400	0.1518
	Avg.	0.0856	0.5479	0.2341	0.4528	0.1030	0.0742	0.2375	0.0752	0.2263
Transformer	1D	0.2066	0.7969	0.3789	0.7930	0.2251	0.1355	0.3441	0.1508	0.3789
	3D	0.1275	0.6825	0.2843	0.5750	0.1318	0.0913	0.2617	0.0926	0.2808
	5D	0.0961	0.5919	0.2149	0.4326	0.0930	0.0727	0.2305	0.0576	0.2237
	7D	0.0714	0.4919	0.1664	0.3504	0.0721	0.0542	0.1959	0.0448	0.1809
	Avg.	0.1254	0.6408	0.2611	0.5378	0.1305	0.0884	0.2581	0.0864	0.2661
PatchTST	1D	0.2149	0.7937	0.3601	0.8041	0.2110	0.1291	0.3226	0.1593	0.3744
	3D	0.1327	0.6565	0.2770	0.5851	0.1429	0.0968	0.2652	0.0913	0.2809
	5D	0.0915	0.5578	0.2143	0.4411	0.1047	0.0734	0.2243	0.0673	0.2218
	7D	0.0757	0.4552	0.1661	0.3639	0.0799	0.0570	0.1975	0.0518	0.1809
	Avg.	0.1287	0.6158	0.2544	0.5485	0.1346	0.0891	0.2524	0.0924	0.2645

Table 17: Benchmark(SEDI(1%)) on part 1(Orlando area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.1231	0.5677	0.3286	0.5730	0.1160	0.1235	0.2804	0.1423	0.2818
	3D	0.0722	0.3377	0.2315	0.4083	0.1017	0.0801	0.1284	0.0580	0.1772
	5D	0.0462	0.4222	0.1899	0.1858	0.0782	0.0618	0.2246	0.0612	0.1587
	7D	0.0583	0.2870	0.0658	0.3472	0.0747	0.0526	0.1112	0.0411	0.1297
	Avg.	0.0749	0.4036	0.2040	0.3786	0.0927	0.0795	0.1862	0.0756	0.1869
RNN	1D	0.1138	0.6157	0.3092	0.4319	0.1276	0.1392	0.2927	0.1144	0.2681
	3D	0.0954	0.3621	0.1573	0.2978	0.0974	0.1047	0.1110	0.0291	0.1568
	5D	0.0588	0.3228	0.1519	0.2054	0.0821	0.0691	0.0985	0.0716	0.1325
	7D	0.1016	0.2944	0.0939	0.0666	0.0793	0.0690	0.1249	0.0787	0.1135
	Avg.	0.0924	0.3987	0.1781	0.2504	0.0966	0.0955	0.1568	0.0734	0.1677
DilatedRNN	1D	0.2111	0.7473	0.3720	0.7648	0.1895	0.1384	0.3324	0.2115	0.3709
	3D	0.1417	0.6452	0.2337	0.5917	0.1539	0.0764	0.1583	0.1111	0.2640
	5D	0.1170	0.5410	0.1941	0.4338	0.1278	0.0712	0.2353	0.1309	0.2314
	7D	0.0923	0.4478	0.1716	0.3167	0.1039	0.0766	0.1639	0.0805	0.1817
	Avg.	0.1405	0.5953	0.2428	0.5267	0.1438	0.0906	0.2225	0.1335	0.2620
GCN	1D	0.2025	0.5785	0.3234	0.4610	0.2208	0.1729	0.3612	0.1966	0.3146
	3D	0.1393	0.5720	0.2439	0.4175	0.0475	0.1439	0.3082	0.1379	0.2513
	5D	0.1185	0.5251	0.1685	0.3352	0.0296	0.1351	0.3633	0.1096	0.2231
	7D	0.1255	0.4695	0.1648	0.2308	0.0269	0.1249	0.3558	0.1684	0.2083
	Avg.	0.1465	0.5363	0.2252	0.3611	0.0812	0.1442	0.3471	0.1531	0.2493
GNN	1D	0.2074	0.7982	0.3768	0.7408	0.2155	0.1250	0.3068	0.1698	0.3675
	3D	0.1567	0.7499	0.2746	0.6374	0.1849	0.1035	0.2813	0.1395	0.3160
	5D	0.1326	0.7111	0.2177	0.5194	0.1881	0.1060	0.2724	0.1089	0.2820
	7D	0.1830	0.6629	0.2348	0.4327	0.1766	0.1203	0.3300	0.1289	0.2836
	Avg.	0.1699	0.7305	0.2760	0.5826	0.1913	0.1137	0.2976	0.1368	0.3123
StemGNN	1D	0.1717	0.4968	0.3191	0.0206	0.1620	0.1239	0.2858	0.2005	0.2225
	3D	0.1410	0.3176	0.2135	0.0176	0.1152	0.0530	0.1569	0.1344	0.1437
	5D	0.1395	0.0319	0.1679	0.0013	0.0691	0.0051	0.1753	0.0661	0.0820
	7D	0.1056	0.0337	0.1386	0.0003	0.0784	0.0036	0.2534	0.0487	0.0828
	Avg.	0.1395	0.2200	0.2098	0.0100	0.1062	0.0464	0.2179	0.1124	0.1328
GPT4TS	1D	0.1732	0.7775	0.3626	0.7129	0.1662	0.1221	0.3125	0.1272	0.3443
	3D	0.1047	0.6083	0.2679	0.4717	0.1214	0.0922	0.2501	0.0771	0.2492
	5D	0.0698	0.5106	0.1923	0.3910	0.0886	0.0498	0.2192	0.0558	0.1971
	7D	0.0579	0.4151	0.1366	0.3045	0.0733	0.0356	0.2041	0.0334	0.1576
	Avg.	0.1014	0.5779	0.2399	0.4700	0.1124	0.0749	0.2465	0.0734	0.2370
LLM	1D	0.1985	0.7427	0.3622	0.8053	0.2213	0.1268	0.3335	0.1362	0.3658
	3D	0.1179	0.6341	0.2763	0.5863	0.1497	0.0970	0.2677	0.0801	0.2761
	5D	0.0788	0.5089	0.2057	0.4334	0.1039	0.0723	0.2216	0.0601	0.2106
	7D	0.0714	0.4392	0.1579	0.3310	0.0873	0.0559	0.1895	0.0444	0.1721
	Avg.	0.1167	0.5812	0.2505	0.5390	0.1406	0.0880	0.2531	0.0802	0.2562

Table 18: Benchmark(MAE) on part 2(Miami area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.0739	0.0969	0.1538	0.1234	0.1172	0.1069	0.1168	0.1548	0.1180
	3D	0.1588	0.1621	0.2502	0.2129	0.1952	0.1638	0.1855	0.2459	0.1968
	5D	0.2147	0.2079	0.3080	0.2637	0.2568	0.1984	0.2268	0.3070	0.2479
	7D	0.2650	0.2436	0.3517	0.3046	0.2989	0.2236	0.2595	0.3454	0.2865
	Avg.	0.1781	0.1776	0.2659	0.2262	0.2170	0.1732	0.1971	0.2633	0.2123
MLP	1D	0.0932	0.1147	0.1777	0.1381	0.1322	0.1152	0.1279	0.2124	0.1389
	3D	0.1637	0.1770	0.2667	0.2123	0.2056	0.1680	0.1926	0.2856	0.2089
	5D	0.2179	0.2209	0.3239	0.2646	0.2615	0.2014	0.2349	0.3376	0.2578
	7D	0.2621	0.2558	0.3663	0.3055	0.3081	0.2258	0.2663	0.3779	0.2960
	Avg.	0.1842	0.1921	0.2836	0.2301	0.2269	0.1776	0.2054	0.3034	0.2254
NLinear	1D	0.0857	0.1055	0.1672	0.1274	0.1207	0.1085	0.1191	0.1484	0.1228
	3D	0.1603	0.1709	0.2606	0.2059	0.1988	0.1642	0.1882	0.2414	0.1988
	5D	0.2153	0.2167	0.3190	0.2608	0.2560	0.1984	0.2311	0.3010	0.2498
	7D	0.2603	0.2526	0.3618	0.3022	0.3031	0.2237	0.2635	0.3457	0.2891
	Avg.	0.1804	0.1864	0.2772	0.2241	0.2196	0.1737	0.2005	0.2591	0.2151
TCN	1D	0.1747	0.1498	0.1937	0.1790	0.1882	0.1957	0.1439	0.7995	0.2531
	3D	0.2303	0.1938	0.2971	0.2766	0.2811	0.2347	0.2049	0.9442	0.3328
	5D	0.3770	0.2350	0.3711	0.3214	0.3427	0.2733	0.2633	0.8945	0.3848
	7D	0.3407	0.2960	0.3925	0.3659	0.3887	0.3089	0.2950	1.0090	0.4246
	Avg.	0.2807	0.2186	0.3136	0.2857	0.3002	0.2531	0.2268	0.9118	0.3488
CNN	1D	0.0777	0.1049	0.1764	0.1385	0.1134	0.1034	0.1154	0.1565	0.1233
	3D	0.1611	0.1813	0.2938	0.2362	0.1982	0.1669	0.1906	0.2810	0.2136
	5D	0.2289	0.2449	0.3621	0.3098	0.2576	0.2038	0.2351	0.3383	0.2726
	7D	0.2809	0.2903	0.4201	0.3565	0.3054	0.2314	0.2702	0.3758	0.3163
	Avg.	0.1871	0.2054	0.3131	0.2603	0.2186	0.1764	0.2028	0.2879	0.2315
TimesNet	1D	0.0962	0.1269	0.1984	0.1537	0.1505	0.1309	0.1520	0.1927	0.1502
	3D	0.1684	0.1943	0.2871	0.2323	0.2249	0.1854	0.2122	0.2767	0.2227
	5D	0.2212	0.2409	0.3516	0.2849	0.2861	0.2200	0.2587	0.3347	0.2748
	7D	0.2722	0.2761	0.3937	0.3329	0.3326	0.2409	0.2915	0.3775	0.3147
	Avg.	0.1895	0.2095	0.3077	0.2509	0.2485	0.1943	0.2286	0.2954	0.2406
Transformer	1D	0.0731	0.0925	0.1477	0.1178	0.1129	0.1024	0.1104	0.1409	0.1122
	3D	0.1446	0.1603	0.2490	0.1975	0.1901	0.1593	0.1807	0.2334	0.1894
	5D	0.2057	0.2052	0.3095	0.2518	0.2498	0.1938	0.2248	0.3007	0.2427
	7D	0.2513	0.2460	0.3529	0.2933	0.2968	0.2193	0.2555	0.3480	0.2829
	Avg.	0.1687	0.1760	0.2648	0.2151	0.2124	0.1687	0.1928	0.2558	0.2068
PatchTST	1D	0.0691	0.0926	0.1472	0.1158	0.1114	0.1012	0.1104	0.1394	0.1109
	3D	0.1419	0.1610	0.2449	0.1975	0.1927	0.1590	0.1828	0.2376	0.1897
	5D	0.2002	0.2094	0.3042	0.2524	0.2504	0.1943	0.2251	0.2999	0.2420
	7D	0.2463	0.2461	0.3499	0.2947	0.3005	0.2200	0.2574	0.3444	0.2824
	Avg.	0.1644	0.1773	0.2615	0.2151	0.2138	0.1686	0.1939	0.2553	0.2062

Table 19: Benchmark(MAE) on part 2(Miami area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.1150	0.1373	0.2165	0.1678	0.1691	0.1378	0.1531	0.2537	0.1688
	3D	0.2287	0.2081	0.3074	0.2574	0.2650	0.2002	0.2272	0.3706	0.2581
	5D	0.2646	0.2556	0.3832	0.3241	0.3190	0.2378	0.2694	0.4339	0.3109
	7D	0.3326	0.2989	0.4425	0.3728	0.3855	0.2579	0.2909	0.4743	0.3569
	Avg.	0.2352	0.2250	0.3374	0.2805	0.2847	0.2084	0.2352	0.3831	0.2737
RNN	1D	0.1196	0.1356	0.2067	0.1678	0.1659	0.1410	0.1546	0.2557	0.1684
	3D	0.2086	0.2169	0.3113	0.2615	0.2693	0.2047	0.2301	0.3624	0.2581
	5D	0.2839	0.2586	0.3710	0.3123	0.3166	0.2299	0.2790	0.4224	0.3092
	7D	0.3404	0.2992	0.4478	0.3695	0.3657	0.2598	0.3007	0.4670	0.3562
	Avg.	0.2381	0.2276	0.3342	0.2778	0.2794	0.2088	0.2411	0.3769	0.2730
DilatedRNN	1D	0.0781	0.1134	0.1706	0.1359	0.1309	0.1155	0.1217	0.2111	0.1347
	3D	0.1724	0.1856	0.2843	0.2268	0.2393	0.1769	0.2149	0.3164	0.2271
	5D	0.2463	0.2351	0.3483	0.2967	0.3012	0.2133	0.2548	0.3866	0.2853
	7D	0.3312	0.2732	0.4052	0.3472	0.3469	0.2460	0.2917	0.4338	0.3344
	Avg.	0.2070	0.2018	0.3021	0.2516	0.2546	0.1879	0.2208	0.3370	0.2454
GCN	1D	0.0910	0.1254	0.1880	0.1761	0.1626	0.1374	0.1528	0.9648	0.2498
	3D	0.1718	0.1904	0.2843	0.2485	0.2410	0.1928	0.2173	0.5350	0.2601
	5D	0.2460	0.2340	0.3390	0.3061	0.3085	0.2319	0.2571	0.7279	0.3313
	7D	0.2998	0.2700	0.3911	0.3452	0.3508	0.2609	0.2914	0.6410	0.3563
	Avg.	0.2022	0.2050	0.3006	0.2690	0.2657	0.2057	0.2297	0.7172	0.2994
GNN	1D	0.0825	0.1072	0.1604	0.1365	0.1226	0.1165	0.1234	0.2444	0.1367
	3D	0.1593	0.1722	0.2527	0.2059	0.2024	0.1734	0.1925	0.3399	0.2123
	5D	0.2109	0.2189	0.3114	0.2658	0.2568	0.2123	0.2320	0.4031	0.2639
	7D	0.2896	0.2506	0.3590	0.3121	0.3067	0.2357	0.2624	0.4222	0.3048
	Avg.	0.1856	0.1872	0.2709	0.2301	0.2221	0.1845	0.2026	0.3524	0.2294
StemGNN	1D	0.0859	0.1404	0.1762	0.1382	0.1444	0.1204	0.1252	0.3656	0.1620
	3D	0.2047	0.2609	0.2958	0.2616	0.2763	0.2432	0.2227	0.4821	0.2809
	5D	0.2874	0.3173	0.3743	0.3535	0.4289	0.2839	0.2765	0.5247	0.3558
	7D	0.3603	0.3402	0.4218	0.4172	0.4253	0.3211	0.3283	0.6507	0.4081
	Avg.	0.2346	0.2647	0.3170	0.2926	0.3187	0.2422	0.2382	0.5058	0.3017
GPT4TS	1D	0.0931	0.1264	0.1860	0.1431	0.1392	0.1304	0.1510	0.1974	0.1458
	3D	0.1639	0.1959	0.2833	0.2283	0.2281	0.1895	0.2159	0.2917	0.2246
	5D	0.2256	0.2429	0.3469	0.2834	0.2897	0.2195	0.2535	0.3389	0.2751
	7D	0.2728	0.2842	0.3941	0.3275	0.3342	0.2466	0.2878	0.3960	0.3179
	Avg.	0.1888	0.2124	0.3026	0.2456	0.2478	0.1965	0.2271	0.3060	0.2408
LLM	1D	0.0863	0.1029	0.1665	0.1318	0.1270	0.1140	0.1312	0.1587	0.1273
	3D	0.1530	0.1663	0.2545	0.2069	0.2008	0.1668	0.1926	0.2450	0.1982
	5D	0.2065	0.2109	0.3079	0.2619	0.2556	0.2017	0.2342	0.3065	0.2481
	7D	0.2552	0.2469	0.3515	0.3009	0.3029	0.2267	0.2654	0.3482	0.2872
	Avg.	0.1752	0.1817	0.2701	0.2254	0.2216	0.1773	0.2058	0.2646	0.2152

Table 20: Benchmark(MSE) on part 2(Miami area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.0315	0.0629	0.1405	0.0774	0.0602	0.0525	0.0787	0.1295	0.0792
	3D	0.0808	0.1311	0.2777	0.1506	0.1289	0.0935	0.1471	0.2762	0.1607
	5D	0.1300	0.1819	0.3613	0.2040	0.1912	0.1197	0.1905	0.3950	0.2217
	7D	0.1783	0.2232	0.4256	0.2480	0.2456	0.1399	0.2230	0.4617	0.2682
	Avg.	0.1052	0.1498	0.3012	0.1700	0.1565	0.1014	0.1599	0.3156	0.1824
MLP	1D	0.0397	0.0747	0.1695	0.0877	0.0697	0.0578	0.0868	0.2450	0.1039
	3D	0.0932	0.1486	0.3241	0.1611	0.1419	0.0985	0.1605	0.3833	0.1889
	5D	0.1460	0.2061	0.4276	0.2192	0.2111	0.1263	0.2114	0.4777	0.2532
	7D	0.1968	0.2537	0.5054	0.2679	0.2786	0.1475	0.2509	0.5544	0.3069
	Avg.	0.1189	0.1708	0.3566	0.1840	0.1753	0.1075	0.1774	0.4151	0.2132
NLinear	1D	0.0374	0.0687	0.1605	0.0830	0.0639	0.0554	0.0816	0.1269	0.0847
	3D	0.0924	0.1442	0.3186	0.1581	0.1376	0.0974	0.1568	0.2825	0.1734
	5D	0.1450	0.2026	0.4224	0.2174	0.2065	0.1253	0.2083	0.3881	0.2395
	7D	0.1962	0.2511	0.5021	0.2666	0.2743	0.1471	0.2484	0.4696	0.2944
	Avg.	0.1177	0.1666	0.3509	0.1813	0.1706	0.1063	0.1738	0.3168	0.1980
TCN	1D	0.0976	0.0963	0.1756	0.1090	0.1377	0.1694	0.0984	5.8002	0.8355
	3D	0.1318	0.1480	0.3271	0.2042	0.2212	0.2113	0.1527	7.0577	1.0567
	5D	0.3490	0.2009	0.4521	0.2532	0.3033	0.2986	0.2079	5.1319	0.8996
	7D	0.2479	0.2662	0.4783	0.2929	0.3587	0.2911	0.2391	6.7494	1.1155
	Avg.	0.2066	0.1779	0.3583	0.2148	0.2552	0.2426	0.1745	6.1848	0.9768
CNN	1D	0.0395	0.0838	0.2180	0.1137	0.0630	0.0539	0.0818	0.3165	0.1213
	3D	0.1168	0.1894	0.4307	0.2323	0.1464	0.1036	0.1678	2.9814	0.5460
	5D	0.1869	0.3223	0.5611	0.3355	0.2198	0.1352	0.2221	2.7558	0.5923
	7D	0.2499	0.4475	0.7525	0.3950	0.2877	0.1613	0.2656	1.4768	0.5045
	Avg.	0.1483	0.2608	0.4906	0.2691	0.1792	0.1135	0.1843	1.8826	0.4411
TimesNet	1D	0.0453	0.0899	0.1964	0.1033	0.0884	0.0700	0.1090	0.1729	0.1094
	3D	0.1100	0.1839	0.3598	0.1995	0.1694	0.1154	0.1857	0.3311	0.2069
	5D	0.1666	0.2634	0.4952	0.2586	0.2486	0.1484	0.2477	0.4500	0.2848
	7D	0.2283	0.3207	0.5949	0.3303	0.3296	0.1674	0.2892	0.5382	0.3498
	Avg.	0.1376	0.2145	0.4116	0.2230	0.2090	0.1253	0.2079	0.3731	0.2377
Transformer	1D	0.0337	0.0645	0.1413	0.0798	0.0614	0.0536	0.0766	0.1210	0.0790
	3D	0.0880	0.1424	0.3086	0.1614	0.1361	0.0964	0.1512	0.2755	0.1700
	5D	0.1486	0.2002	0.4253	0.2206	0.2052	0.1266	0.2082	0.3973	0.2415
	7D	0.2032	0.2658	0.4903	0.2697	0.2727	0.1490	0.2408	0.5022	0.2992
	Avg.	0.1184	0.1682	0.3414	0.1828	0.1688	0.1064	0.1692	0.3240	0.1974
PatchTST	1D	0.0326	0.0667	0.1392	0.0789	0.0615	0.0525	0.0781	0.1213	0.0789
	3D	0.0872	0.1459	0.2950	0.1614	0.1389	0.0963	0.1586	0.2825	0.1707
	5D	0.1402	0.2096	0.3952	0.2208	0.2068	0.1249	0.2112	0.3960	0.2381
	7D	0.1899	0.2624	0.4758	0.2686	0.2763	0.1473	0.2494	0.4766	0.2933
	Avg.	0.1125	0.1711	0.3263	0.1824	0.1709	0.1053	0.1743	0.3191	0.1952

Table 21: Benchmark(MSE) on part 2(Miami area) stations(RNN,GNN,LLM)

Method	Metric	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.0575	0.1123	0.2405	0.1158	0.1092	0.0775	0.1224	1.1111	0.2433
	3D	0.1531	0.1869	0.3854	0.2070	0.2091	0.1206	0.1944	1.4368	0.3617
	5D	0.1983	0.2390	0.4905	0.2732	0.2894	0.1479	0.2368	1.3811	0.4070
	7D	0.2734	0.2901	0.5826	0.3426	0.3981	0.1677	0.2615	1.8380	0.5192
	Avg.	0.1706	0.2071	0.4248	0.2346	0.2515	0.1284	0.2038	1.4417	0.3828
RNN	1D	0.0611	0.1108	0.2390	0.1248	0.1152	0.0790	0.1230	0.9006	0.2192
	3D	0.1276	0.1894	0.3864	0.2041	0.2173	0.1226	0.1953	1.3494	0.3490
	5D	0.1973	0.2385	0.4877	0.2623	0.2918	0.1463	0.2344	1.4248	0.4104
	7D	0.2820	0.3010	0.5995	0.3402	0.3564	0.1667	0.2657	1.4515	0.4704
	Avg.	0.1670	0.2099	0.4282	0.2328	0.2452	0.1286	0.2046	1.2816	0.3622
DilatedRNN	1D	0.0366	0.0808	0.1810	0.0931	0.0823	0.0635	0.0944	0.8270	0.1823
	3D	0.1137	0.1710	0.3481	0.1894	0.2205	0.1151	0.1746	1.0864	0.3023
	5D	0.1970	0.2275	0.4660	0.2738	0.3055	0.1469	0.2277	1.3925	0.4046
	7D	0.2899	0.2742	0.5486	0.3313	0.3679	0.1781	0.2746	1.3743	0.4549
	Avg.	0.1593	0.1884	0.3859	0.2219	0.2441	0.1259	0.1928	1.1701	0.3360
GCN	1D	0.0333	0.0964	0.1750	0.1817	0.1209	0.0909	0.0948	170.0996	21.3616
	3D	0.0829	0.1800	0.3166	0.2245	0.1894	0.1255	0.1616	32.4988	4.2224
	5D	0.1407	0.2346	0.4015	0.2996	0.2709	0.1500	0.2053	44.7123	5.8019
	7D	0.2021	0.2599	0.4775	0.3266	0.3373	0.1828	0.2379	17.1457	2.3962
	Avg.	0.1147	0.1927	0.3426	0.2581	0.2296	0.1373	0.1749	66.1141	8.4455
GNN	1D	0.0339	0.0703	0.1442	0.0825	0.0631	0.0563	0.0809	0.6083	0.1425
	3D	0.0851	0.1359	0.2816	0.1522	0.1320	0.0988	0.1493	0.9670	0.2502
	5D	0.1331	0.1885	0.3656	0.2117	0.1916	0.1290	0.1920	1.6289	0.3800
	7D	0.2053	0.2240	0.4330	0.2525	0.2511	0.1483	0.2250	1.0897	0.3536
	Avg.	0.1144	0.1547	0.3061	0.1747	0.1594	0.1081	0.1618	1.0735	0.2816
StemGNN	1D	0.0391	0.1055	0.1724	0.0912	0.0873	0.0665	0.0898	1.2556	0.2384
	3D	0.1283	0.2207	0.3498	0.2164	0.2680	0.1644	0.1796	1.5368	0.3830
	5D	0.2199	0.2945	0.4663	0.3239	0.5381	0.2094	0.2414	1.6949	0.4985
	7D	0.3149	0.3582	0.5362	0.3986	0.4688	0.2426	0.3031	2.0180	0.5801
	Avg.	0.1755	0.2447	0.3812	0.2575	0.3406	0.1707	0.2035	1.6263	0.4250
GPT4TS	1D	0.0473	0.0949	0.1828	0.0977	0.0819	0.0735	0.1124	0.1921	0.1103
	3D	0.1163	0.1995	0.3605	0.2009	0.1762	0.1251	0.1953	0.3751	0.2186
	5D	0.1908	0.2827	0.4853	0.2619	0.2654	0.1534	0.2487	0.4691	0.2947
	7D	0.2394	0.3739	0.5955	0.3207	0.3350	0.1826	0.2946	0.5944	0.3670
	Avg.	0.1484	0.2377	0.4060	0.2203	0.2146	0.1337	0.2127	0.4077	0.2477
LLM	1D	0.0375	0.0718	0.1582	0.0879	0.0687	0.0581	0.0900	0.1373	0.0887
	3D	0.0889	0.1438	0.3015	0.1598	0.1410	0.0984	0.1615	0.2986	0.1742
	5D	0.1406	0.2011	0.3952	0.2175	0.2057	0.1265	0.2115	0.4012	0.2374
	7D	0.1944	0.2496	0.4659	0.2653	0.2754	0.1485	0.2510	0.4759	0.2908
	Avg.	0.1154	0.1666	0.3302	0.1826	0.1727	0.1079	0.1785	0.3283	0.1978

Table 22: Benchmark(SEDI10) on part 2(Miami area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.8741	0.8301	0.8503	0.8179	0.8890	0.5135	0.4864	0.7414	0.7503
	3D	0.8258	0.7763	0.8076	0.7682	0.8627	0.4220	0.4150	0.6270	0.6881
	5D	0.7627	0.7409	0.7604	0.7023	0.8228	0.3802	0.3853	0.6126	0.6459
	7D	0.7108	0.7069	0.7572	0.6765	0.8028	0.3550	0.3867	0.5522	0.6185
	Avg.	0.7933	0.7636	0.7939	0.7412	0.8443	0.4177	0.4183	0.6333	0.6757
MLP	1D	0.8506	0.7672	0.8000	0.7572	0.8709	0.4446	0.4225	0.5891	0.6878
	3D	0.7622	0.6598	0.7016	0.6497	0.7979	0.3370	0.3233	0.4826	0.5893
	5D	0.6958	0.5955	0.6414	0.5820	0.7421	0.2694	0.2707	0.4073	0.5255
	7D	0.6433	0.5499	0.5962	0.5360	0.6957	0.2335	0.2379	0.3550	0.4810
	Avg.	0.7380	0.6431	0.6848	0.6312	0.7767	0.3211	0.3136	0.4585	0.5709
NLinear	1D	0.8486	0.7786	0.8039	0.7660	0.8697	0.4931	0.4452	0.7005	0.7132
	3D	0.7606	0.6705	0.7059	0.6597	0.7969	0.3753	0.3424	0.5532	0.6081
	5D	0.6961	0.6019	0.6452	0.5928	0.7418	0.3265	0.2868	0.4605	0.5439
	7D	0.6445	0.5544	0.6000	0.5465	0.6962	0.2868	0.2522	0.3980	0.4973
	Avg.	0.7374	0.6514	0.6887	0.6412	0.7761	0.3704	0.3316	0.5280	0.5906
TCN	1D	0.7593	0.7900	0.8204	0.6993	0.7827	0.2702	0.4436	0.2651	0.6038
	3D	0.7316	0.6984	0.7447	0.5739	0.7257	0.1869	0.3902	0.1932	0.5306
	5D	0.5618	0.6040	0.6806	0.5003	0.6732	0.1566	0.3651	0.1776	0.4649
	7D	0.6365	0.5666	0.6778	0.4988	0.5719	0.1185	0.2944	0.1443	0.4386
	Avg.	0.6723	0.6647	0.7309	0.5681	0.6884	0.1831	0.3733	0.1951	0.5095
CNN	1D	0.8450	0.7649	0.7755	0.7226	0.8710	0.4506	0.4383	0.6989	0.6959
	3D	0.7464	0.6562	0.6524	0.5939	0.7859	0.3206	0.3270	0.5369	0.5774
	5D	0.6737	0.5673	0.5828	0.5158	0.7293	0.2661	0.2691	0.4358	0.5050
	7D	0.6180	0.5240	0.5309	0.4738	0.6825	0.2327	0.2278	0.3797	0.4587
	Avg.	0.7208	0.6281	0.6354	0.5766	0.7672	0.3175	0.3156	0.5128	0.5592
TimesNet	1D	0.8231	0.7314	0.7478	0.7043	0.8409	0.3910	0.3584	0.6258	0.6528
	3D	0.7335	0.6302	0.6612	0.6004	0.7609	0.2651	0.2711	0.4914	0.5517
	5D	0.6745	0.5599	0.5876	0.5382	0.7072	0.2241	0.2204	0.4053	0.4896
	7D	0.6158	0.5245	0.5430	0.4907	0.6482	0.1888	0.1921	0.3477	0.4438
	Avg.	0.7117	0.6115	0.6349	0.5834	0.7393	0.2672	0.2605	0.4675	0.5345
Transformer	1D	0.8637	0.8038	0.8207	0.7756	0.8762	0.4669	0.4545	0.7247	0.7233
	3D	0.7686	0.7074	0.7133	0.6568	0.8054	0.3441	0.3459	0.5810	0.6153
	5D	0.6971	0.6270	0.6584	0.5889	0.7435	0.2774	0.2809	0.4665	0.5424
	7D	0.6451	0.5751	0.6083	0.5437	0.6964	0.2396	0.2535	0.3979	0.4949
	Avg.	0.7436	0.6783	0.7002	0.6412	0.7803	0.3320	0.3337	<u>0.5425</u>	0.5940
PatchTST	1D	0.8595	0.7994	0.8269	0.7725	0.8746	0.4772	0.4750	0.7220	0.7259
	3D	0.7678	0.6914	0.7225	0.6534	0.7956	0.3561	0.3503	0.5633	0.6125
	5D	0.7022	0.6194	0.6605	0.5880	0.7365	0.3018	0.2951	0.4659	0.5462
	7D	0.6513	0.5667	0.6103	0.5407	0.6906	0.2540	0.2564	0.4046	0.4968
	Avg.	0.7452	0.6692	0.7051	0.6386	0.7743	<u>0.3473</u>	0.3442	0.5389	0.5954

Table 23: Benchmark(SEDI10) on part 2(Miami area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.8360	0.7782	0.8055	0.7590	0.8905	0.4015	0.3896	0.6608	0.6901
	3D	0.7519	0.6710	0.6999	0.6089	0.8233	0.2542	0.2837	0.5149	0.5760
	5D	0.6692	0.6430	0.6611	0.5801	0.7650	0.2298	0.2401	0.3879	0.5220
	7D	0.6402	0.5991	0.6269	0.4682	0.7060	0.1930	0.2364	0.3669	0.4796
	Avg.	0.7243	0.6728	0.6984	0.6041	0.7962	0.2696	0.2874	0.4826	0.5669
RNN	1D	0.8152	0.7886	0.7825	0.7384	0.8582	0.3910	0.4268	0.6741	0.6843
	3D	0.7450	0.6511	0.7208	0.6443	0.8068	0.2397	0.2744	0.5008	0.5729
	5D	0.6943	0.6334	0.6518	0.5570	0.7574	0.2019	0.2405	0.4519	0.5235
	7D	0.6155	0.5923	0.6195	0.5072	0.7319	0.2241	0.2245	0.3572	0.4840
	Avg.	0.7175	0.6664	0.6936	0.6117	0.7886	0.2642	0.2916	0.4960	0.5662
DilatedRNN	1D	0.8692	0.7880	0.8197	0.7892	0.8793	0.4684	0.4362	0.6942	0.7180
	3D	0.7796	0.7056	0.7306	0.6319	0.7575	0.2916	0.3150	0.5490	0.5951
	5D	0.6999	0.6585	0.6645	0.5788	0.7220	0.2488	0.2843	0.4543	0.5389
	7D	0.6469	0.5807	0.5970	0.5429	0.6725	0.2202	0.2308	0.3934	0.4856
	Avg.	0.7489	0.6832	0.7029	0.6357	0.7578	0.3072	0.3166	0.5227	0.5844
GCN	1D	0.8394	0.7752	0.8086	0.7437	0.8631	0.4585	0.4685	0.6737	0.7038
	3D	0.7913	0.7114	0.7760	0.6987	0.8134	0.4180	0.3846	0.5844	0.6472
	5D	0.7726	0.6603	0.7433	0.6416	0.8037	0.3545	0.3369	0.5404	0.6067
	7D	0.7195	0.6370	0.7368	0.6266	0.7624	0.2961	0.3282	0.5247	0.5789
	Avg.	0.7807	0.6960	0.7662	0.6777	0.8107	0.3818	0.3795	0.5808	0.6342
GNN	1D	0.8842	0.8339	0.8446	0.8034	0.8794	0.4924	0.4661	0.5660	0.7213
	3D	0.8071	0.7862	0.7878	0.6988	0.8383	0.3740	0.3840	0.5319	0.6510
	5D	0.7275	0.6700	0.7674	0.6396	0.8200	0.3434	0.3735	0.5033	0.6056
	7D	0.7208	0.6275	0.7505	0.6695	0.7939	0.3157	0.3493	0.4627	0.5863
	Avg.	0.7849	0.7294	0.7876	0.7028	0.8329	0.3814	0.3932	0.5160	0.6410
StemGNN	1D	0.8646	0.7785	0.7954	0.7146	0.8769	0.4439	0.4379	0.4597	0.6714
	3D	0.7744	0.6132	0.7012	0.6006	0.7568	0.2438	0.2951	0.3788	0.5455
	5D	0.6814	0.6045	0.6612	0.5584	0.5665	0.2083	0.2529	0.3115	0.4806
	7D	0.6648	0.5171	0.6013	0.4415	0.5642	0.1738	0.2100	0.2598	0.4291
	Avg.	0.7463	0.6283	0.6898	0.5788	0.6911	0.2674	0.2990	0.3525	0.5316
GPT4TS	1D	0.8216	0.7340	0.7701	0.7331	0.8414	0.3715	0.3565	0.6124	0.6551
	3D	0.7389	0.6277	0.6603	0.6071	0.7518	0.2678	0.2655	0.4756	0.5493
	5D	0.6746	0.5605	0.5939	0.5327	0.6894	0.2279	0.2255	0.3964	0.4876
	7D	0.6139	0.5161	0.5541	0.4812	0.6419	0.1929	0.1982	0.3275	0.4407
	Avg.	0.7123	0.6096	0.6446	0.5885	0.7311	0.2650	0.2614	0.4530	0.5332
AutoTimes	1D	0.8574	0.7975	0.8157	0.7598	0.8726	0.4113	0.4080	0.7040	0.7033
	3D	0.7652	0.6858	0.7175	0.6466	0.8032	0.3067	0.3215	0.5668	0.6017
	5D	0.7002	0.6266	0.6601	0.5837	0.7408	0.2576	0.2676	0.4668	0.5379
	7D	0.6453	0.5683	0.6149	0.5348	0.6967	0.2213	0.2319	0.4040	0.4896
	Avg.	0.7420	0.6695	0.7021	0.6312	0.7783	0.2992	0.3073	0.5354	0.5831

Table 24: Benchmark(SEDI5) on part 2(Miami area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.8203	0.8018	0.8453	0.6927	0.8843	0.3386	0.3853	0.6361	0.6755
	3D	0.7427	0.7459	0.8080	0.6251	0.8496	0.2768	0.3320	0.5376	0.6147
	5D	0.6747	0.7152	0.7456	0.5607	0.8084	0.2658	0.3134	0.5075	0.5739
	7D	0.6180	0.6833	0.7520	0.5447	0.7774	0.2555	0.3284	0.4584	0.5522
	Avg.	0.7139	0.7366	0.7878	0.6058	0.8299	0.2842	0.3398	0.5349	0.6041
MLP TSMixer	1D	0.7829	0.7231	0.7853	0.6187	0.8603	0.2823	0.3224	0.4521	0.6034
	3D	0.6518	0.5942	0.6704	0.5094	0.7743	0.2085	0.2223	0.3548	0.4982
	5D	0.5559	0.5210	0.5950	0.4429	0.7069	0.1762	0.1753	0.2891	0.4328
	7D	0.4809	0.4667	0.5379	0.3967	0.6516	0.1424	0.1484	0.2437	0.3835
	Avg.	0.6179	0.5763	0.6471	0.4919	0.7483	0.2023	0.2171	0.3349	0.4795
NLinear	1D	0.7842	0.7341	0.7857	0.6272	0.8583	0.3155	0.3398	0.5837	0.6286
	3D	0.6549	0.6065	0.6750	0.5297	0.7740	0.2379	0.2452	0.4378	0.5201
	5D	0.5606	0.5291	0.6000	0.4519	0.7081	0.2114	0.1966	0.3530	0.4513
	7D	0.4881	0.4734	0.5439	0.4077	0.6537	0.1884	0.1690	0.2964	0.4026
	Avg.	0.6220	0.5858	0.6512	0.5041	0.7485	0.2383	0.2377	0.4177	0.5007
TCN	1D	0.6732	0.7264	0.8085	0.4636	0.6769	0.1780	0.3211	0.1701	0.5022
	3D	0.6003	0.6072	0.7341	0.3332	0.6076	0.1251	0.2746	0.1106	0.4241
	5D	0.3759	0.4967	0.6449	0.3185	0.5094	0.0967	0.2207	0.1031	0.3457
	7D	0.3713	0.4688	0.6172	0.2808	0.4117	0.0658	0.1851	0.0710	0.3089
	Avg.	0.5052	0.5748	0.7012	0.3490	0.5514	0.1164	0.2504	0.1137	0.3952
CNN ModernTCN	1D	0.7753	0.7152	0.7470	0.5845	0.8521	0.2784	0.3290	0.5653	0.6058
	3D	0.6425	0.5891	0.6141	0.4597	0.7544	0.1988	0.2123	0.4098	0.4851
	5D	0.5456	0.4885	0.5393	0.3821	0.6921	0.1645	0.1691	0.3149	0.4120
	7D	0.4749	0.4345	0.4788	0.3437	0.6387	0.1410	0.1413	0.2653	0.3648
	Avg.	0.6096	0.5568	0.5948	0.4425	0.7343	0.1957	0.2129	0.3888	0.4669
TimesNet	1D	0.7487	0.6768	0.7300	0.5653	0.8128	0.2235	0.2528	0.4910	0.5626
	3D	0.6236	0.5582	0.6225	0.4521	0.7272	0.1473	0.1642	0.3592	0.4568
	5D	0.5369	0.4827	0.5380	0.3935	0.6590	0.1184	0.1226	0.2851	0.3920
	7D	0.4596	0.4353	0.4911	0.3486	0.5869	0.1043	0.0991	0.2359	0.3451
	Avg.	0.5922	0.5383	0.5954	0.4399	0.6965	0.1484	0.1597	0.3428	0.4391
Transformer iTransformer	1D	0.7985	0.7670	0.8047	0.6444	0.8568	0.2787	0.3549	0.6033	0.6385
	3D	0.6767	0.6518	0.6871	0.5192	0.7699	0.1998	0.2785	0.4604	0.5304
	5D	0.5748	0.5559	0.6193	0.4492	0.6981	0.1652	0.1791	0.3529	0.4493
	7D	0.5036	0.4938	0.5666	0.4040	0.6433	0.1402	0.1705	0.2874	0.4012
	Avg.	0.6384	0.6171	0.6694	0.5042	0.7420	0.1960	0.2457	0.4260	0.5049
Transformer PatchTST	1D	0.7948	0.7663	0.8131	0.6364	0.8546	0.2859	0.3648	0.6001	0.6395
	3D	0.6706	0.6408	0.6990	0.5158	0.7641	0.2161	0.2558	0.4454	0.5260
	5D	0.5757	0.5538	0.6241	0.4486	0.6977	0.1833	0.2059	0.3516	0.4551
	7D	0.5016	0.4906	0.5665	0.4029	0.6450	0.1581	0.1751	0.2979	0.4047
	Avg.	0.6357	0.6129	0.6757	0.5009	0.7404	0.2108	0.2504	0.4238	0.5063

Table 25: Benchmark(SEDI5) on part 2(Miami area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.7782	0.7372	0.7975	0.6088	0.8707	0.2226	0.2661	0.5371	0.6023
	3D	0.6193	0.6102	0.6635	0.4559	0.7919	0.1437	0.1772	0.4233	0.4856
	5D	0.5323	0.5690	0.6237	0.3969	0.7194	0.1253	0.1418	0.3040	0.4266
	7D	0.4947	0.4718	0.5666	0.3301	0.6305	0.1009	0.1329	0.2593	0.3734
	Avg.	0.6061	0.5971	0.6628	0.4479	0.7531	0.1481	0.1795	0.3809	0.4720
RNN	1D	0.7506	0.7444	0.7702	0.5797	0.8298	0.2470	0.2639	0.5466	0.5915
	3D	0.6252	0.6021	0.6902	0.4647	0.7750	0.1426	0.1678	0.4273	0.4869
	5D	0.5361	0.5584	0.5951	0.3839	0.7114	0.1120	0.1259	0.3603	0.4229
	7D	0.4436	0.5078	0.5647	0.3391	0.6956	0.0967	0.1104	0.2599	0.3772
	Avg.	0.5889	0.6032	0.6551	0.4419	0.7530	0.1496	0.1670	0.3985	0.4696
DilatedRNN	1D	0.8054	0.7393	0.8027	0.6504	0.8471	0.3023	0.3125	0.5724	0.6290
	3D	0.6871	0.6505	0.7110	0.5126	0.7126	0.1749	0.2102	0.4345	0.5117
	5D	0.5834	0.5993	0.6388	0.4463	0.6757	0.1499	0.1713	0.3450	0.4512
	7D	0.5413	0.4900	0.5612	0.4197	0.6188	0.1372	0.1388	0.2847	0.3990
	Avg.	0.6543	0.6198	0.6784	0.5072	0.7136	0.1911	0.2082	0.4092	0.4977
GCN	1D	0.7502	0.7320	0.8148	0.6321	0.8439	0.3394	0.3757	0.5786	0.6333
	3D	0.6899	0.6528	0.7953	0.5605	0.7889	0.2891	0.3176	0.5118	0.5757
	5D	0.6957	0.5923	0.7658	0.5245	0.7818	0.2388	0.2643	0.4678	0.5414
	7D	0.6440	0.5577	0.7625	0.4949	0.7402	0.2244	0.2587	0.4532	0.5169
	Avg.	0.6949	0.6337	0.7846	0.5530	0.7887	0.2729	0.3041	0.5029	0.5668
GNN	1D	0.8279	0.7998	0.8378	0.6750	0.8669	0.3249	0.3500	0.4899	0.6465
	3D	0.7233	0.7394	0.7814	0.5680	0.8201	0.2450	0.2873	0.4436	0.5760
	5D	0.6198	0.6098	0.7593	0.5051	0.7966	0.2324	0.2902	0.4006	0.5267
	7D	0.6588	0.5544	0.7447	0.5131	0.7637	0.2154	0.2739	0.4022	0.5158
	Avg.	0.7074	0.6758	0.7808	0.5653	0.8118	0.2544	0.3003	0.4340	0.5662
StemGNN	1D	0.8015	0.7321	0.7884	0.5807	0.8594	0.2851	0.3385	0.3684	0.5943
	3D	0.6608	0.5265	0.7032	0.4503	0.7169	0.1409	0.1868	0.2969	0.4603
	5D	0.5775	0.5009	0.6416	0.4189	0.3915	0.1244	0.1569	0.2450	0.3821
	7D	0.4858	0.4275	0.5752	0.2864	0.4690	0.0999	0.1371	0.1949	0.3345
	Avg.	0.6314	0.5468	0.6771	0.4341	0.6092	0.1626	0.2048	0.2763	0.4428
GPT4TS	1D	0.7424	0.6726	0.7407	0.5721	0.8175	0.2057	0.2580	0.4738	0.5604
	3D	0.6382	0.5471	0.6227	0.4534	0.7177	0.1509	0.1622	0.3482	0.4550
	5D	0.5484	0.4662	0.5447	0.3914	0.6360	0.1230	0.1265	0.2813	0.3897
	7D	0.4653	0.4102	0.4972	0.3462	0.5821	0.1040	0.1057	0.2184	0.3411
	Avg.	0.5986	0.5240	0.6013	0.4408	0.6883	0.1459	0.1631	0.3304	0.4366
LLM	1D	0.7923	0.7578	0.8113	0.6238	0.8590	0.2414	0.3024	0.5828	0.6213
	3D	0.6587	0.6310	0.6989	0.5032	0.7737	0.1776	0.2137	0.4392	0.5120
	5D	0.5670	0.5597	0.6312	0.4430	0.7028	0.1544	0.1690	0.3459	0.4466
	7D	0.4857	0.4895	0.5722	0.3936	0.6512	0.1312	0.1424	0.2904	0.3945
	Avg.	0.6259	0.6095	0.6784	0.4909	0.7467	0.1761	0.2069	0.4146	0.4936

Table 26: Benchmark(SEDI1) on part 2(Miami area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.6392	0.7486	0.7968	0.4636	0.7523	0.1510	0.1849	0.4622	0.5248
	3D	0.5460	0.6948	0.7434	0.4294	0.6635	0.1429	0.1752	0.3631	0.4698
	5D	0.5736	0.6829	0.6508	0.3551	0.6734	0.1527	0.1643	0.3667	0.4524
	7D	0.5641	0.6625	0.7139	0.4331	0.6920	0.1401	0.1617	0.3547	0.4653
	Avg.	0.5807	0.6972	0.7263	0.4203	0.6953	0.1467	0.1715	0.3867	0.4781
MLP	1D	0.5899	0.6011	0.6559	0.3407	0.6954	0.1186	0.1495	0.2268	0.4222
	3D	0.4351	0.4299	0.4696	0.2279	0.5486	0.0778	0.0867	0.1369	0.3016
	5D	0.3255	0.3379	0.3531	0.1776	0.4431	0.0599	0.0690	0.1006	0.2333
	7D	0.2522	0.2693	0.2787	0.1494	0.3661	0.0524	0.0697	0.0791	0.1896
	Avg.	0.4007	0.4096	0.4393	0.2239	0.5133	0.0772	0.0937	0.1359	0.2867
NLinear	1D	0.6032	0.6324	0.6721	0.3915	0.7185	0.1412	0.1508	0.3811	0.4613
	3D	0.4433	0.4556	0.4838	0.2745	0.5680	0.1052	0.1023	0.2296	0.3328
	5D	0.3408	0.3553	0.3639	0.2278	0.4626	0.0956	0.0801	0.1734	0.2624
	7D	0.2637	0.2851	0.2905	0.1992	0.3820	0.0851	0.0702	0.1435	0.2149
	Avg.	0.4128	0.4321	0.4526	0.2732	0.5328	0.1068	0.1008	0.2319	0.3179
TCN	1D	0.1324	0.4480	0.7226	0.2066	0.2128	0.0640	0.1211	0.0506	0.2448
	3D	0.0330	0.3732	0.5677	0.1723	0.1441	0.0387	0.1129	0.0265	0.1835
	5D	0.0258	0.2706	0.4580	0.1274	0.1506	0.0312	0.1130	0.0283	0.1506
	7D	0.0571	0.2566	0.4746	0.1193	0.1263	0.0128	0.0770	0.0104	0.1418
	Avg.	0.0621	0.3371	0.5557	0.1564	0.1584	0.0367	0.1060	0.0289	0.1802
CNN	1D	0.6055	0.5939	0.6605	0.3129	0.7173	0.1196	0.1522	0.3788	0.4426
	3D	0.4336	0.4005	0.4627	0.1837	0.5687	0.0721	0.0722	0.2224	0.3020
	5D	0.3390	0.2918	0.3464	0.1267	0.4670	0.0555	0.0590	0.1442	0.2287
	7D	0.2672	0.2162	0.2717	0.1025	0.3878	0.0519	0.0415	0.1163	0.1819
	Avg.	0.4113	0.3756	0.4353	0.1814	0.5352	0.0748	0.0812	0.2154	0.2888
TimesNet	1D	0.5404	0.5278	0.6337	0.2851	0.6289	0.0906	0.0988	0.2989	0.3880
	3D	0.4211	0.3841	0.4394	0.1624	0.5016	0.0434	0.0522	0.1626	0.2708
	5D	0.3299	0.2943	0.3291	0.1157	0.3647	0.0295	0.0269	0.1104	0.2001
	7D	0.2249	0.2508	0.2616	0.0851	0.3262	0.0234	0.0230	0.0944	0.1612
	Avg.	0.3791	0.3643	0.4160	0.1621	0.4554	0.0467	0.0502	0.1666	0.2550
Transformer	1D	0.6163	0.6752	0.7437	0.3883	0.7228	0.1268	0.1668	0.4121	0.4815
	3D	0.4566	0.5015	0.5470	0.2322	0.5716	0.0745	0.0972	0.2676	0.3435
	5D	0.3503	0.3820	0.4225	0.1779	0.4512	0.0560	0.0565	0.1561	0.2566
	7D	0.2829	0.2964	0.3415	0.1450	0.3636	0.0446	0.0921	0.1163	0.2103
	Avg.	0.4265	0.4638	0.5137	0.2359	0.5273	0.0755	0.1031	0.2380	0.3230
PatchTST	1D	0.6112	0.6690	0.7429	0.3695	0.7238	0.1261	0.1691	0.4179	0.4787
	3D	0.4618	0.4940	0.5613	0.2286	0.5741	0.0822	0.0949	0.2473	0.3430
	5D	0.3572	0.3782	0.4444	0.1734	0.4708	0.0668	0.0695	0.1708	0.2664
	7D	0.2791	0.3030	0.3527	0.1388	0.3865	0.0589	0.0600	0.1358	0.2143
	Avg.	0.4273	0.4610	0.5253	0.2276	0.5388	0.0835	0.0984	0.2429	0.3256

Table 27: Benchmark(SEDI1) on part 2(Miami area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.5173	0.5951	0.6803	0.2523	0.4421	0.0731	0.0896	0.3249	0.3718
	3D	0.3565	0.3467	0.4334	0.1171	0.2033	0.0380	0.0398	0.1530	0.2110
	5D	0.3267	0.2902	0.3336	0.0565	0.2339	0.0143	0.0343	0.1030	0.1740
	7D	0.2477	0.2608	0.4109	0.0553	0.0981	0.0078	0.0051	0.0914	0.1471
	Avg.	0.3621	0.3732	0.4646	0.1203	0.2443	0.0333	0.0422	0.1681	0.2260
RNN	1D	0.4712	0.6332	0.6489	0.2446	0.4433	0.0756	0.1161	0.3283	0.3701
	3D	0.3647	0.3977	0.4097	0.0820	0.1804	0.0310	0.0387	0.1553	0.2074
	5D	0.2339	0.2331	0.2853	0.0355	0.1330	0.0220	0.0048	0.1158	0.1329
	7D	0.1074	0.2155	0.2837	0.0609	0.0813	0.0165	0.0063	0.1074	0.1099
	Avg.	0.2943	0.3699	0.4069	0.1058	0.2095	0.0363	0.0415	0.1767	0.2051
DilatedRNN	1D	0.6372	0.5916	0.7119	0.4065	0.7022	0.1088	0.1333	0.3922	0.4605
	3D	0.5222	0.4714	0.6236	0.2306	0.4730	0.0741	0.0689	0.2576	0.3402
	5D	0.3632	0.4556	0.5025	0.1754	0.4427	0.0560	0.0513	0.1841	0.2789
	7D	0.3681	0.3293	0.4419	0.1630	0.4273	0.0491	0.0411	0.1600	0.2475
	Avg.	0.4727	0.4620	0.5700	0.2439	0.5113	0.0720	0.0737	0.2485	0.3317
GCN	1D	0.5814	0.5680	0.7359	0.3744	0.6712	0.1554	0.1852	0.4307	0.4628
	3D	0.5052	0.5117	0.7286	0.2892	0.5979	0.1691	0.1555	0.3875	0.4181
	5D	0.4462	0.4254	0.6934	0.2584	0.5692	0.1573	0.1310	0.3559	0.3796
	7D	0.4309	0.3783	0.6980	0.2593	0.5114	0.1639	0.1560	0.3369	0.3668
	Avg.	0.4909	0.4709	0.7140	0.2953	0.5874	0.1614	0.1569	0.3777	0.4068
GNN	1D	0.6570	0.6977	0.7677	0.4359	0.7147	0.1492	0.1707	0.3647	0.4947
	3D	0.5526	0.6366	0.6920	0.3083	0.6153	0.1285	0.1482	0.2883	0.4212
	5D	0.4619	0.4865	0.6791	0.2633	0.5934	0.1286	0.1450	0.2778	0.3794
	7D	0.6103	0.4415	0.7151	0.3320	0.6158	0.1261	0.1514	0.2791	0.4089
	Avg.	0.5705	0.5656	0.7135	0.3349	0.6348	0.1331	0.1538	0.3025	0.4261
StemGNN	1D	0.5891	0.5290	0.7186	0.3042	0.6594	0.1308	0.1546	0.2231	0.4136
	3D	0.3107	0.2996	0.6110	0.1650	0.4861	0.0484	0.0712	0.1342	0.2658
	5D	0.3170	0.2849	0.5104	0.1400	0.1212	0.0405	0.0469	0.1600	0.2026
	7D	0.2300	0.2427	0.3921	0.0846	0.1728	0.0315	0.0513	0.0838	0.1611
	Avg.	0.3617	0.3391	0.5580	0.1735	0.3599	0.0628	0.0810	0.1503	0.2608
GPT4TS	1D	0.5577	0.5380	0.6412	0.2952	0.6627	0.0716	0.0904	0.2589	0.3895
	3D	0.4106	0.3726	0.4585	0.1692	0.4833	0.0402	0.0416	0.1690	0.2681
	5D	0.3439	0.2834	0.3346	0.1284	0.3308	0.0317	0.0291	0.1154	0.1997
	7D	0.2543	0.2254	0.2716	0.0894	0.2808	0.0250	0.0233	0.0790	0.1561
	Avg.	0.3917	0.3548	0.4265	0.1706	0.4394	0.0421	0.0461	0.1556	0.2533
LLM	1D	0.5904	0.6595	0.7269	0.3583	0.6964	0.1035	0.1381	0.3728	0.4557
	3D	0.4307	0.4794	0.5365	0.2296	0.5553	0.0609	0.0708	0.2425	0.3257
	5D	0.3386	0.3883	0.4334	0.2038	0.4503	0.0524	0.0534	0.1652	0.2607
	7D	0.2539	0.2968	0.3465	0.1403	0.3743	0.0457	0.0386	0.1488	0.2056
	Avg.	0.4034	0.4560	0.5108	0.2330	0.5191	0.0656	0.0752	0.2323	0.3119

Table 28: Benchmark(MAE) on part 3(Fort Myers area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.	
MLP	1D	0.0251	0.0246	0.0207	0.0463	0.0302	0.0628	0.0505	0.0380	0.0373	
	3D	0.0526	0.0607	0.0523	0.0777	0.0603	0.1228	0.0977	0.0757	0.0750	
	5D	0.0839	0.0940	0.0814	0.1102	0.0958	0.1610	0.1305	0.1105	0.1084	
	7D	0.1005	0.1202	0.1013	0.1327	0.1143	0.1891	0.1558	0.1281	0.1303	
	Avg.	0.0655	0.0749	0.0639	0.0917	0.0752	<u>0.1339</u>	<u>0.1086</u>	0.0881	0.0877	
MLP	TSMixer	1D	0.0247	0.0327	0.0265	0.0477	0.0353	0.0684	0.0548	0.0461	0.0420
	3D	0.0503	0.0683	0.0560	0.0821	0.0665	0.1226	0.0987	0.0811	0.0782	
	5D	0.0734	0.0989	0.0827	0.1093	0.0899	0.1596	0.1307	0.1108	0.1069	
	7D	0.0946	0.1254	0.1081	0.1330	0.1100	0.1873	0.1561	0.1368	0.1314	
	Avg.	0.0608	0.0813	0.0683	0.0930	0.0754	0.1345	0.1101	0.0937	0.0896	
NLinear	1D	0.0184	0.0243	0.0200	0.0427	0.0307	0.0615	0.0480	0.0377	0.0354	
	3D	0.0463	0.0613	0.0516	0.0798	0.0636	0.1196	0.0947	0.0752	0.0740	
	5D	0.0705	0.0931	0.0796	0.1089	0.0880	0.1577	0.1279	0.1055	0.1039	
	7D	0.0933	0.1207	0.1053	0.1334	0.1078	0.1861	0.1545	0.1322	0.1292	
	Avg.	0.0571	0.0749	0.0641	0.0912	0.0725	0.1312	0.1063	0.0877	0.0856	
TCN	1D	0.0577	0.0505	0.0318	0.0970	0.1129	0.0920	0.2015	0.1758	0.1024	
	3D	0.0968	0.1382	0.0689	0.1048	0.1373	0.1937	0.1485	0.2512	0.1424	
	5D	0.1406	0.1002	0.0931	0.1558	0.1816	0.2363	0.1885	0.2557	0.1690	
	7D	0.2256	0.1780	0.1241	0.1633	0.2116	0.2240	0.2846	0.2676	0.2098	
	Avg.	0.1302	0.1167	0.0795	0.1302	0.1608	0.1865	0.2058	0.2376	0.1559	
CNN	1D	0.0221	0.0265	0.0205	0.0443	0.0278	0.0654	0.0491	0.0357	0.0364	
	3D	0.0567	0.0702	0.0510	0.0861	0.0603	0.1332	0.1010	0.0719	0.0788	
	5D	0.0871	0.1048	0.0834	0.1215	0.0883	0.1774	0.1384	0.1015	0.1128	
	7D	0.1202	0.1362	0.1056	0.1499	0.1094	0.2057	0.1686	0.1280	0.1405	
	Avg.	0.0715	0.0845	0.0651	0.1005	0.0714	0.1454	0.1143	0.0843	0.0921	
TimesNet	1D	0.0199	0.0288	0.0242	0.0513	0.0445	0.0809	0.0651	0.0528	0.0459	
	3D	0.0467	0.0651	0.0558	0.0875	0.0724	0.1426	0.1164	0.0866	0.0841	
	5D	0.0703	0.0970	0.0838	0.1148	0.0988	0.1852	0.1484	0.1127	0.1139	
	7D	0.0923	0.1284	0.1121	0.1438	0.1158	0.2145	0.1759	0.1417	0.1406	
	Avg.	0.0573	0.0798	0.0690	0.0993	0.0829	0.1558	0.1264	0.0984	0.0961	
Transformer	iTransformer	1D	0.0190	0.0253	0.0192	0.0383	0.0278	0.0641	0.0482	0.0345	0.0346
	3D	0.0469	0.0617	0.0487	0.0726	0.0576	0.1244	0.0959	0.0706	0.0723	
	5D	0.0709	0.0948	0.0766	0.1023	0.0834	0.1633	0.1331	0.0977	0.1028	
	7D	0.0914	0.1226	0.1041	0.1275	0.1048	0.1889	0.1596	0.1214	0.1275	
	Avg.	0.0571	0.0761	<u>0.0621</u>	0.0852	0.0684	0.1352	0.1092	0.0811	0.0843	
PatchTST	1D	0.0183	0.0241	0.0177	0.0381	0.0288	0.0637	0.0481	0.0349	0.0342	
	3D	0.0464	0.0654	0.0545	0.0765	0.0588	0.1242	0.1007	0.0714	0.0747	
	5D	0.0762	0.0951	0.0761	0.1057	0.0835	0.1651	0.1354	0.1014	0.1048	
	7D	0.0984	0.1266	0.1032	0.1319	0.1053	0.1955	0.1608	0.1257	0.1309	
	Avg.	0.0598	0.0778	0.0629	<u>0.0881</u>	<u>0.0691</u>	0.1371	0.1112	<u>0.0833</u>	0.0862	

Table 29: Benchmark(MAE) on part 3(Fort Myers area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.0486	0.0447	0.0393	0.0640	0.0465	0.1120	0.0839	0.0606	0.0625
	3D	0.0895	0.0878	0.0788	0.1048	0.0828	0.1775	0.1299	0.1091	0.1075
	5D	0.1246	0.1247	0.1167	0.1344	0.1172	0.2057	0.1590	0.1281	0.1388
	7D	0.1621	0.1584	0.1496	0.1725	0.1389	0.2203	0.1823	0.1567	0.1676
	Avg.	0.1062	0.1039	0.0961	0.1189	0.0964	0.1789	0.1388	0.1136	0.1191
RNN	1D	0.0569	0.0461	0.0381	0.0632	0.0486	0.1023	0.0763	0.0694	0.0626
	3D	0.0859	0.0950	0.0851	0.1180	0.0949	0.1640	0.1403	0.0993	0.1103
	5D	0.1086	0.1219	0.1137	0.1424	0.1388	0.2005	0.1742	0.1466	0.1433
	7D	0.1496	0.1542	0.1549	0.1745	0.1467	0.2247	0.2012	0.1485	0.1693
	Avg.	0.1003	0.1043	0.0980	0.1245	0.1073	0.1729	0.1480	0.1159	0.1214
DilatedRNN	1D	0.0394	0.0268	0.0277	0.0441	0.0472	0.0691	0.0596	0.0372	0.0439
	3D	0.0726	0.0714	0.0679	0.0978	0.0648	0.1390	0.1186	0.0837	0.0895
	5D	0.1100	0.1028	0.0887	0.1257	0.0966	0.2000	0.1578	0.1128	0.1243
	7D	0.1312	0.1500	0.1175	0.1499	0.1234	0.2363	0.1870	0.1478	0.1554
	Avg.	0.0883	0.0877	0.0754	0.1043	0.0830	0.1611	0.1308	0.0954	0.1033
GCN	1D	0.0543	0.0596	0.0264	0.3354	0.1422	0.0789	0.0946	0.1151	0.1133
	3D	0.0858	0.0942	0.0598	0.3643	0.1756	0.1443	0.1374	0.1593	0.1526
	5D	0.1006	0.1192	0.0851	0.3859	0.2002	0.1767	0.1713	0.1952	0.1793
	7D	0.1196	0.1408	0.1141	0.4079	0.2197	0.2033	0.2068	0.2315	0.2055
	Avg.	0.0901	0.1034	0.0713	0.3734	0.1845	0.1508	0.1525	0.1753	0.1627
GNN	1D	10.0224	0.0295	0.0262	0.0462	0.0370	0.0732	0.0543	0.0436	0.0415
	3D	0.0548	0.0673	0.0538	0.0829	0.0784	0.1336	0.1087	0.0841	0.0830
	5D	0.0830	0.0978	0.0814	0.1153	0.0967	0.1668	0.1511	0.1109	0.1129
	7D	0.1064	0.1236	0.1114	0.1452	0.1335	0.2015	0.1823	0.1354	0.1424
	Avg.	0.0666	0.0796	0.0682	0.0974	0.0864	0.1438	0.1241	0.0935	0.0949
StemGNN	1D	0.0636	0.0445	0.0335	0.0635	0.0586	0.1001	0.0599	0.0489	0.0591
	3D	0.1114	0.1001	0.0788	0.1273	0.1028	0.1524	0.1421	0.1408	0.1195
	5D	0.1690	0.1688	0.1022	0.1698	0.1352	0.1980	0.2366	0.2161	0.1745
	7D	0.2094	0.2090	0.1719	0.2016	0.1625	0.2325	0.2514	0.2730	0.2139
	Avg.	0.1384	0.1306	0.0966	0.1406	0.1148	0.1707	0.1725	0.1697	0.1417
GPT4TS	1D	0.0209	0.0287	0.0261	0.0482	0.0407	0.0740	0.0596	0.7614	0.1324
	3D	0.0467	0.0638	0.0553	0.0845	0.0730	0.1431	0.1105	0.0869	0.0830
	5D	0.0702	0.0946	0.0832	0.1141	0.0980	0.1823	0.1455	0.1120	0.1125
	7D	0.0909	0.1231	0.1105	0.1392	0.1160	0.2087	0.1715	0.1397	0.1375
	Avg.	0.0572	0.0776	0.0688	0.0965	0.0819	0.1520	0.1218	0.2750	0.1163
AutoTimes	1D	0.0228	0.0282	0.0199	0.0429	0.0321	0.0680	0.0542	0.0407	0.0386
	3D	0.0475	0.0615	0.0486	0.0760	0.0638	0.1240	0.0993	0.0751	0.0745
	5D	0.0692	0.0965	0.0737	0.1075	0.0892	0.1632	0.1315	0.1056	0.1045
	7D	0.0921	0.1218	0.1038	0.1304	0.1083	0.1900	0.1573	0.1285	0.1290
	Avg.	0.0579	0.0770	0.0615	0.0892	0.0734	0.1363	0.1106	0.0875	0.0867

Table 30: Benchmark(MSE) on part 3(Fort Myers area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.0036	0.0046	0.0052	0.0165	0.0137	0.0368	0.0139	0.0095	0.0130
	3D	0.0156	0.0159	0.0199	0.0359	0.0318	0.0888	0.0374	0.0252	0.0338
	5D	0.0295	0.0300	0.0345	0.0538	0.0486	0.1253	0.0552	0.0426	0.0524
	7D	0.0461	0.0442	0.0494	0.0729	0.0635	0.1537	0.0701	0.0562	0.0695
	Avg.	0.0237	0.0237	0.0272	0.0448	0.0394	0.1011	0.0441	0.0334	0.0422
MLP	1D	0.0047	0.0056	0.0070	0.0182	0.0163	0.0420	0.0159	0.0116	0.0152
	3D	0.0176	0.0174	0.0217	0.0381	0.0353	0.0983	0.0403	0.0280	0.0371
	5D	0.0348	0.0315	0.0372	0.0561	0.0522	0.1410	0.0605	0.0452	0.0573
	7D	0.0548	0.0465	0.0531	0.0740	0.0679	0.1735	0.0778	0.0624	0.0763
	Avg.	0.0280	0.0253	0.0297	0.0466	0.0429	0.1137	0.0486	0.0368	0.0465
NLinear	1D	0.0031	0.0044	0.0050	0.0174	0.0158	0.0392	0.0138	0.0098	0.0136
	3D	0.0153	0.0159	0.0202	0.0378	0.0349	0.0970	0.0383	0.0263	0.0357
	5D	0.0321	0.0298	0.0362	0.0561	0.0518	0.1392	0.0581	0.0433	0.0558
	7D	0.0520	0.0449	0.0526	0.0742	0.0677	0.1724	0.0754	0.0607	0.0750
	Avg.	0.0256	0.0238	0.0285	0.0464	0.0426	0.1120	0.0464	0.0350	0.0450
TCN	1D	0.0925	0.0196	0.0078	0.0383	0.0743	0.0452	0.0900	0.1310	0.0623
	3D	0.1029	0.1063	0.0244	0.0440	0.0804	0.1231	0.0514	0.1998	0.0916
	5D	0.1186	0.0304	0.0379	0.0681	0.1249	0.1691	0.0767	0.2183	0.1055
	7D	0.1905	0.1158	0.0575	0.0808	0.1354	0.1655	0.1613	0.1934	0.1375
	Avg.	0.1261	0.0680	0.0319	0.0578	0.1038	0.1257	0.0949	0.1856	0.0992
CNN	1D	0.0048	0.0050	0.0045	0.0208	0.0142	0.0451	0.0148	0.0100	0.0149
	3D	0.0219	0.0210	0.0187	0.0451	0.0345	0.1297	0.0455	0.0276	0.0430
	5D	0.0466	0.0406	0.0363	0.0757	0.0548	0.1949	0.0722	0.0451	0.0708
	7D	0.0710	0.0626	0.0527	0.1014	0.0705	0.2231	0.0981	0.0619	0.0927
	Avg.	0.0361	0.0323	0.0280	0.0607	0.0435	0.1482	0.0576	0.0362	0.0553
TimesNet	1D	0.0035	0.0052	0.0066	0.0215	0.0262	0.0586	0.0206	0.0163	0.0198
	3D	0.0159	0.0181	0.0214	0.0461	0.0445	0.1318	0.0553	0.0331	0.0458
	5D	0.0315	0.0351	0.0369	0.0654	0.0684	0.2087	0.0811	0.0509	0.0723
	7D	0.0514	0.0543	0.0537	0.0919	0.0871	0.2343	0.1012	0.0706	0.0931
	Avg.	0.0256	0.0282	0.0297	0.0562	0.0565	0.1584	0.0645	0.0427	0.0577
Transformer	1D	0.0035	0.0047	0.0050	0.0177	0.0157	0.0436	0.0140	0.0094	0.0142
	3D	0.0171	0.0171	0.0209	0.0403	0.0352	0.1103	0.0410	0.0255	0.0384
	5D	0.0343	0.0335	0.0356	0.0623	0.0527	0.1582	0.0674	0.0423	0.0608
	7D	0.0518	0.0522	0.0539	0.0807	0.0714	0.1897	0.0847	0.0590	0.0804
	Avg.	0.0267	0.0269	0.0288	0.0503	0.0437	0.1254	0.0518	<u>0.0341</u>	0.0485
PatchTST	1D	0.0032	0.0048	0.0044	0.0179	0.0154	0.0444	0.0144	0.0096	0.0143
	3D	0.0161	0.0205	0.0212	0.0420	0.0353	0.1100	0.0432	0.0266	0.0393
	5D	0.0385	0.0364	0.0374	0.0608	0.0531	0.1632	0.0668	0.0438	0.0625
	7D	0.0564	0.0578	0.0514	0.0832	0.0700	0.2020	0.0825	0.0598	0.0829
	Avg.	0.0285	0.0299	0.0286	0.0510	0.0435	0.1299	0.0517	0.0350	0.0498

Table 31: Benchmark(MSE) on part 3(Fort Myers area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.0183	0.0087	0.0109	0.0287	0.0217	0.0748	0.0274	0.0181	0.0261
	3D	0.0420	0.0271	0.0312	0.0588	0.0444	0.1455	0.0540	0.0381	0.0551
	5D	0.0624	0.0427	0.0506	0.0774	0.0630	0.1761	0.0717	0.0552	0.0749
	7D	0.0949	0.0621	0.0773	0.0985	0.0797	0.2013	0.0880	0.0714	0.0967
	Avg.	0.0544	0.0352	0.0425	0.0659	0.0522	0.1494	0.0603	0.0457	0.0632
RNN	1D	0.0408	0.0090	0.0108	0.0298	0.0224	0.0750	0.0266	0.0173	0.0290
	3D	0.0634	0.0251	0.0316	0.0568	0.0463	0.1353	0.0580	0.0370	0.0567
	5D	0.0682	0.0414	0.0506	0.0743	0.0642	0.1775	0.0754	0.0564	0.0760
	7D	0.0910	0.0569	0.0728	0.0969	0.0799	0.1947	0.0899	0.0703	0.0941
	Avg.	0.0659	0.0331	0.0414	0.0644	0.0532	0.1457	0.0625	0.0453	0.0639
DilatedRNN	1D	0.0172	0.0048	0.0056	0.0194	0.0162	0.0408	0.0175	0.0104	0.0165
	3D	0.0395	0.0193	0.0225	0.0442	0.0356	0.1128	0.0489	0.0300	0.0441
	5D	0.0583	0.0347	0.0401	0.0675	0.0560	0.1787	0.0786	0.0509	0.0706
	7D	0.0808	0.0583	0.0578	0.0958	0.0747	0.2167	0.0990	0.0714	0.0943
	Avg.	0.0489	0.0293	0.0315	0.0567	0.0457	0.1372	0.0610	0.0407	0.0564
GCN	1D	0.0108	0.0094	0.0060	0.3682	0.0982	0.0392	0.0227	0.0923	0.0809
	3D	0.0267	0.0246	0.0196	0.3854	0.1184	0.0918	0.0472	0.1217	0.1044
	5D	0.0434	0.0359	0.0339	0.4006	0.1337	0.1285	0.0674	0.1440	0.1234
	7D	0.0569	0.0499	0.0486	0.4154	0.1489	0.1565	0.0868	0.1794	0.1428
	Avg.	0.0344	0.0300	0.0270	0.3924	0.1248	0.1040	0.0560	0.1343	0.1129
GNN	1D	0.0037	0.0049	0.0063	0.0178	0.0153	0.0430	0.0145	0.0108	0.0145
	3D	0.0154	0.0166	0.0207	0.0379	0.0364	0.0938	0.0399	0.0268	0.0359
	5D	0.0298	0.0302	0.0359	0.0555	0.0496	0.1312	0.0609	0.0432	0.0545
	7D	0.0477	0.0434	0.0542	0.0769	0.0691	0.1594	0.0787	0.0591	0.0736
	Avg.	0.0241	0.0237	0.0293	0.0471	0.0426	0.1069	0.0485	0.0350	0.0446
StemGNN	1D	0.0390	0.0081	0.0070	0.0221	0.0281	0.0535	0.0164	0.0145	0.0236
	3D	0.0816	0.0405	0.0270	0.0609	0.0599	0.1179	0.0722	0.0861	0.0683
	5D	0.1100	0.0850	0.0433	0.1015	0.0809	0.1587	0.1540	0.1312	0.1081
	7D	0.1554	0.1173	0.1096	0.1232	0.1079	0.1982	0.1746	0.2072	0.1492
	Avg.	0.0965	0.0628	0.0467	0.0769	0.0692	0.1321	0.1043	0.1098	0.0873
GPT4TS	1D	0.0038	0.0050	0.0060	0.0202	0.0223	0.0542	0.0183	0.6054	0.0919
	3D	0.0151	0.0174	0.0205	0.0442	0.0474	0.1530	0.0496	0.0339	0.0476
	5D	0.0321	0.0319	0.0371	0.0648	0.0716	0.2071	0.0829	0.0526	0.0725
	7D	0.0494	0.0489	0.0528	0.0872	0.0841	0.2397	0.0996	0.0734	0.0919
	Avg.	0.0251	0.0258	0.0291	0.0541	0.0564	0.1635	0.0626	0.1913	0.0760
LLM	1D	0.0037	0.0052	0.0048	0.0183	0.0159	0.0447	0.0161	0.0106	0.0149
	3D	0.0159	0.0167	0.0189	0.0396	0.0342	0.1026	0.0409	0.0263	0.0369
	5D	0.0321	0.0314	0.0337	0.0564	0.0519	0.1485	0.0614	0.0429	0.0573
	7D	0.0522	0.0466	0.0508	0.0745	0.0661	0.1807	0.0781	0.0586	0.0760
	Avg.	0.0260	0.0250	0.0271	0.0472	0.0420	0.1191	0.0491	0.0346	0.0463

Table 32: Benchmark(SEDI10) on part 3(Fort Myers area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.	
MLP	1D	0.6714	0.9580	0.6989	0.7088	0.4910	0.6202	0.5661	0.8442	0.6948	
	3D	0.6107	0.8594	0.6242	0.7260	0.4937	0.4783	0.5238	0.8326	0.6436	
	5D	0.5833	0.7752	0.6230	0.6262	0.4892	0.5590	0.5791	0.7968	0.6290	
	7D	0.5649	0.7550	0.5619	0.5997	0.5046	0.4772	0.4864	0.6886	0.5798	
	Avg.	0.6076	0.8369	0.6270	0.6652	0.4946	0.5337	0.5389	0.7906	0.6368	
MLP	1D	0.6675	0.9189	0.6900	0.7345	0.5080	0.6620	0.6030	0.8160	0.7000	
	3D	0.5919	0.8285	0.6258	0.6505	0.4595	0.5203	0.4941	0.7342	0.6131	
	5D	0.5567	0.7866	0.5754	0.6006	0.4316	0.4412	0.4199	0.6803	0.5615	
	7D	0.5350	0.7498	0.5379	0.5651	0.4129	0.3812	0.3557	0.6375	0.5219	
	Avg.	0.5878	0.8209	0.6073	0.6377	0.4530	0.5012	0.4682	0.7170	0.5991	
NLinear	1D	0.6653	0.9466	0.7012	0.7387	0.5234	0.6204	0.6081	0.8296	0.7041	
	3D	0.5963	0.8470	0.6314	0.6528	0.4532	0.5048	0.4892	0.7448	0.6149	
	5D	0.5604	0.7925	0.5800	0.6023	0.4297	0.4342	0.4069	0.6914	0.5622	
	7D	0.5393	0.7539	0.5417	0.5660	0.4113	0.3787	0.3420	0.6494	0.5228	
	Avg.	0.5903	0.8350	0.6136	0.6399	0.4544	0.4845	0.4615	0.7288	0.6010	
TCN	1D	0.6607	0.9349	0.7126	0.6214	0.3251	0.6057	0.3817	0.4229	0.5831	
	3D	0.5849	0.7254	0.6607	0.6150	0.3200	0.3803	0.4778	0.3677	0.5165	
	5D	0.4883	0.8096	0.5672	0.5006	0.2824	0.3337	0.3708	0.4034	0.4695	
	7D	0.2490	0.6355	0.5764	0.6034	0.2939	0.3018	0.2916	0.3836	0.4169	
	Avg.	0.4957	0.7763	0.6292	0.5851	0.3054	0.4054	0.3805	0.3944	0.4965	
CNN	1D	0.6643	0.9457	0.6975	0.7506	0.5325	0.6309	0.5645	0.8346	0.7026	
	3D	0.5833	0.8013	0.6528	0.6401	0.4782	0.4488	0.4296	0.7353	0.5962	
	5D	0.5274	0.7259	0.5842	0.6140	0.4395	0.3732	0.3519	0.6687	0.5356	
	7D	0.4967	0.6819	0.5463	0.5468	0.4399	0.3177	0.2592	0.6290	0.4897	
	Avg.	0.5679	0.7887	0.6202	0.6379	0.4725	0.4426	0.4013	0.7169	0.5810	
TimesNet	1D	0.6628	0.8907	0.6824	0.7392	0.4722	0.5926	0.5199	0.7606	0.6651	
	3D	0.5922	0.8103	0.6122	0.6461	0.4452	0.4391	0.3654	0.6902	0.5751	
	5D	0.5666	0.7517	0.5491	0.6090	0.4134	0.3682	0.3255	0.6374	0.5276	
	7D	0.5199	0.7043	0.5105	0.5579	0.3969	0.2931	0.2576	0.5950	0.4794	
	Avg.	0.5854	0.7892	0.5885	0.6380	0.4319	0.4233	0.3671	0.6708	0.5618	
Transformer	iTransformer	1D	0.6753	0.9308	0.6948	0.7630	0.5149	0.6342	0.5745	0.8270	0.7018
	3D	0.5920	0.8284	0.6263	0.6781	0.4784	0.4697	0.4621	0.7383	0.6092	
	5D	0.5688	0.7619	0.5898	0.6217	0.4517	0.4084	0.3802	0.6824	0.5581	
	7D	0.5440	0.7203	0.5514	0.5852	0.4287	0.3532	0.3130	0.6412	0.5171	
	Avg.	0.5950	0.8103	0.6156	0.6620	0.4684	0.4664	0.4325	0.7222	0.5965	
PatchTST	1D	0.6779	0.9328	0.7007	0.7626	0.5236	0.6109	0.5828	0.8365	0.7035	
	3D	0.6072	0.8306	0.6327	0.6674	0.4839	0.4845	0.4406	0.7434	0.6113	
	5D	0.5537	0.7707	0.5892	0.6232	0.4570	0.4017	0.3710	0.6887	0.5569	
	7D	0.5374	0.7078	0.5492	0.5853	0.4358	0.3455	0.3136	0.6471	0.5152	
	Avg.	0.5941	0.8105	0.6180	0.6596	0.4751	0.4606	0.4270	0.7289	0.5967	

Table 33: Benchmark(SEDI10) on part 3(Fort Myers area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.6375	0.8919	0.6771	0.7127	0.5130	0.6141	0.4802	0.6979	0.6530
	3D	0.5603	0.7366	0.5969	0.7031	0.4666	0.4219	0.4553	0.6716	0.5765
	5D	0.5230	0.7399	0.5582	0.6459	0.4201	0.3505	0.4925	0.6755	0.5507
	7D	0.5015	0.7195	0.5452	0.5964	0.4224	0.3320	0.4663	0.6533	0.5296
	Avg.	0.5556	0.7720	0.5943	0.6645	0.4555	0.4296	0.4736	0.6746	0.5775
RNN	1D	0.6113	0.9191	0.5886	0.7494	0.5485	0.4949	0.4766	0.7039	0.6365
	3D	0.5456	0.7545	0.6393	0.6426	0.4689	0.3235	0.3871	0.7007	0.5578
	5D	0.5199	0.7361	0.5829	0.6406	0.3841	0.2802	0.2172	0.5790	0.4925
	7D	0.5267	0.7149	0.4846	0.5765	0.4471	0.2646	0.3546	0.5960	0.4956
	Avg.	0.5509	0.7811	0.5738	0.6523	0.4621	0.3408	0.3589	0.6449	0.5456
DilatedRNN	1D	0.6157	0.9270	0.7114	0.8285	0.4265	0.5347	0.4866	0.7628	0.6616
	3D	0.5809	0.7952	0.6612	0.5798	0.4546	0.3865	0.3726	0.7252	0.5695
	5D	0.5291	0.8040	0.5665	0.6098	0.4765	0.2496	0.2638	0.6925	0.5239
	7D	0.5392	0.6853	0.5523	0.5684	0.4365	0.2247	0.3043	0.5970	0.4885
	Avg.	0.5662	0.8029	0.6228	0.6466	0.4485	0.3489	0.3568	0.6944	0.5609
GCN	1D	0.6792	0.8309	0.7292	0.4914	0.4288	0.5120	0.4721	0.7129	0.6071
	3D	0.6144	0.8262	0.6910	0.4325	0.3749	0.4319	0.5107	0.6594	0.5676
	5D	0.5949	0.7921	0.6016	0.4126	0.3364	0.3577	0.3270	0.6403	0.5078
	7D	0.5728	0.7445	0.6076	0.3891	0.3311	0.2809	0.3782	0.6840	0.4985
	Avg.	0.6153	0.7984	0.6574	0.4314	0.3678	0.3956	0.4220	0.6741	0.5453
GNN	1D	0.6606	0.9310	0.7011	0.7420	0.5204	0.5268	0.6103	0.7723	0.6831
	3D	0.6044	0.8487	0.6062	0.6820	0.4898	0.5028	0.5337	0.7566	0.6281
	5D	0.5545	0.8082	0.6223	0.6375	0.4579	0.3712	0.5406	0.7085	0.5876
	7D	0.5737	0.7776	0.6044	0.6699	0.4793	0.4239	0.5228	0.6609	0.5891
	Avg.	0.5983	0.8414	<u>0.6335</u>	0.6828	0.4869	0.4562	0.5519	0.7246	0.6219
StemGNN	1D	0.6429	0.9027	0.6334	0.6619	0.4429	0.4493	0.5562	0.7738	0.6329
	3D	0.5120	0.8540	0.6633	0.6193	0.4460	0.3921	0.3833	0.6762	0.5683
	5D	0.4908	0.7262	0.6228	0.5595	0.3974	0.3235	0.3157	0.6449	0.5101
	7D	0.4009	0.6838	0.4941	0.5233	0.3831	0.2672	0.2345	0.5653	0.4440
	Avg.	0.5116	0.7917	0.6034	0.5910	0.4173	0.3580	0.3724	0.6650	0.5388
GPT4TS	1D	0.6635	0.9232	0.6942	0.7230	0.4875	0.6188	0.5335	0.7614	0.6757
	3D	0.6057	0.8277	0.6177	0.6298	0.4394	0.4481	0.4085	0.6951	0.5840
	5D	0.5684	0.7776	0.5751	0.5932	0.4164	0.3858	0.3212	0.6346	0.5340
	7D	0.5243	0.7301	0.5333	0.5432	0.3953	0.3318	0.2623	0.5767	0.4871
	Avg.	0.5905	0.8147	0.6051	0.6223	0.4347	0.4462	0.3814	0.6670	0.5702
AutoTimes	1D	0.6599	0.9250	0.6921	0.7510	0.5256	0.6094	0.5451	0.8114	0.6899
	3D	0.6089	0.8376	0.6380	0.6568	0.4840	0.4850	0.4504	0.7400	0.6126
	5D	0.5746	0.7828	0.5837	0.6040	0.4538	0.4243	0.3857	0.6793	0.5610
	7D	0.5422	0.7350	0.5560	0.5717	0.4415	0.3649	0.3277	0.6459	0.5231
	Avg.	0.5964	0.8201	0.6175	0.6459	0.4762	0.4709	0.4272	0.7191	0.5967

Table 34: Benchmark(SEDI5) on part 3(Fort Myers area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.4894	0.8078	0.7149	0.4916	0.3294	0.5618	0.3728	0.6667	0.5543
	3D	0.4773	0.7190	0.6372	0.5178	0.3379	0.4137	0.3164	0.6467	0.5083
	5D	0.4737	0.6523	0.5924	0.4466	0.3030	0.4697	0.3772	0.6306	0.4932
	7D	0.4645	0.6408	0.5297	0.4251	0.3127	0.3881	0.2769	0.5457	0.4479
	Avg.	0.4762	0.7050	0.6186	0.4703	0.3208	0.4583	0.3358	0.6224	0.5009
MLP	1D	0.4885	0.7318	0.7059	0.5263	0.3722	0.5500	0.3410	0.6631	0.5474
	3D	0.4713	0.6731	0.6384	0.4570	0.3260	0.3770	0.2068	0.5727	0.4653
	5D	0.4577	0.6479	0.5783	0.4103	0.3053	0.2971	0.1542	0.5068	0.4197
	7D	0.4474	0.6274	0.5311	0.3795	0.2879	0.2336	0.1456	0.4626	0.3894
	Avg.	0.4662	0.6700	0.6134	0.4433	0.3229	0.3644	0.2119	0.5513	0.4554
NLinear	1D	0.4908	0.8016	0.7131	0.5444	0.3801	0.5330	0.3531	0.6808	0.5621
	3D	0.4729	0.6957	0.6459	0.4658	0.3114	0.3720	0.2127	0.5928	0.4711
	5D	0.4581	0.6564	0.5865	0.4188	0.2919	0.2901	0.1535	0.5293	0.4231
	7D	0.4475	0.6307	0.5373	0.3836	0.2755	0.2271	0.1539	0.4864	0.3927
	Avg.	0.4673	0.6961	0.6207	0.4532	0.3147	0.3555	0.2183	0.5724	0.4623
TCN	1D	0.4885	0.5418	0.6754	0.4250	0.0903	0.5215	0.1972	0.1387	0.3848
	3D	0.2287	0.4575	0.4992	0.3864	0.0903	0.2572	0.2046	0.1727	0.2871
	5D	0.2222	0.6698	0.3806	0.3737	0.1045	0.2205	0.1504	0.1179	0.2799
	7D	0.2372	0.4134	0.4001	0.3923	0.1494	0.2020	0.1318	0.1100	0.2545
	Avg.	0.2941	0.5206	0.4888	0.3943	0.1086	0.3003	0.1710	0.1348	0.3016
CNN	1D	0.4916	0.8000	0.7168	0.5206	0.3806	0.5178	0.3296	0.6500	0.5509
	3D	0.4736	0.6654	0.6446	0.4359	0.3385	0.2951	0.1874	0.5540	0.4493
	5D	0.4546	0.6087	0.5675	0.3828	0.3163	0.2291	0.1094	0.4870	0.3944
	7D	0.4437	0.5747	0.5238	0.3442	0.2964	0.1796	0.0746	0.4491	0.3607
	Avg.	0.4659	0.6622	0.6132	0.4209	0.3329	0.3054	0.1752	0.5350	0.4388
TimesNet	1D	0.4887	0.7914	0.6931	0.5047	0.3416	0.4730	0.2862	0.5783	0.5196
	3D	0.4738	0.6845	0.6056	0.4265	0.3145	0.3008	0.1060	0.4963	0.4260
	5D	0.4572	0.6307	0.5365	0.3854	0.2857	0.2354	0.0782	0.4423	0.3814
	7D	0.4477	0.6095	0.4801	0.3425	0.2637	0.1745	0.0602	0.3952	0.3467
	Avg.	0.4669	0.6790	0.5788	0.4148	0.3013	0.2959	0.1327	0.4780	0.4184
Transformer	1D	0.4861	0.7835	0.7049	0.5526	0.3745	0.5381	0.3514	0.6485	0.5549
	3D	0.4722	0.6740	0.6211	0.4677	0.3346	0.3229	0.2136	0.5557	0.4577
	5D	0.4566	0.6520	0.5766	0.4145	0.3157	0.2654	0.1486	0.5018	0.4164
	7D	0.4463	0.6171	0.5237	0.3832	0.2963	0.2088	0.1013	0.4623	0.3799
	Avg.	0.4653	0.6817	0.6066	0.4545	0.3303	0.3338	0.2037	0.5421	0.4522
PatchTST	1D	0.4924	0.8199	0.7104	0.5501	0.3796	0.4983	0.3188	0.6643	0.5542
	3D	0.4747	0.6933	0.6454	0.4688	0.3416	0.3528	0.1760	0.5698	0.4653
	5D	0.4567	0.6503	0.5773	0.4337	0.3168	0.2647	0.1300	0.5138	0.4179
	7D	0.4544	0.6030	0.5288	0.3971	0.2896	0.2039	0.1030	0.4685	0.3810
	Avg.	0.4695	0.6916	0.6154	0.4624	0.3319	0.3299	0.1820	0.5541	0.4546

Table 35: Benchmark(SEDI5) on part 3(Fort Myers area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.4820	0.6970	0.6832	0.5265	0.3526	0.4093	0.2297	0.5224	0.4878
	3D	0.4696	0.6260	0.5874	0.4875	0.2963	0.1749	0.2572	0.4429	0.4177
	5D	0.4635	0.6474	0.3289	0.4312	0.2716	0.1569	0.0442	0.4783	0.3528
	7D	0.4340	0.6733	0.3075	0.3608	0.2654	0.1364	0.0542	0.4455	0.3347
	Avg.	0.4623	0.6609	0.4768	0.4515	0.2965	0.2194	0.1463	0.4723	0.3982
RNN	1D	0.4587	0.7348	0.5771	0.5112	0.3805	0.3081	0.1743	0.4682	0.4516
	3D	0.4104	0.6217	0.3898	0.4468	0.3524	0.1630	0.1703	0.4901	0.3806
	5D	0.3463	0.6626	0.3252	0.4434	0.2483	0.1350	0.0699	0.3937	0.3280
	7D	0.4636	0.6772	0.2954	0.4170	0.2511	0.1049	0.1605	0.4080	0.3472
	Avg.	0.4197	0.6741	0.3969	0.4546	0.3081	0.1778	0.1437	0.4400	0.3769
DilatedRNN	1D	0.4427	0.7950	0.7256	0.6408	0.2791	0.4482	0.2522	0.5895	0.5216
	3D	0.4861	0.6492	0.6471	0.3783	0.3198	0.2445	0.1764	0.5450	0.4308
	5D	0.3852	0.6652	0.5548	0.4024	0.3185	0.1632	0.1136	0.5060	0.3886
	7D	0.4685	0.5954	0.5216	0.3734	0.2855	0.1483	0.1444	0.4271	0.3705
	Avg.	0.4456	0.6762	0.6123	0.4487	0.3007	0.2510	0.1716	0.5169	0.4279
GCN	1D	0.4893	0.7551	0.7258	0.3366	0.3070	0.3908	0.2399	0.5583	0.4754
	3D	0.4817	0.7295	0.6715	0.2813	0.2968	0.2954	0.2091	0.5118	0.4346
	5D	0.4785	0.7154	0.5946	0.2677	0.2595	0.2635	0.0498	0.4793	0.3885
	7D	0.4697	0.6812	0.5678	0.2610	0.2722	0.2174	0.1484	0.5102	0.3910
	Avg.	0.4798	0.7203	0.6399	0.2867	0.2839	0.2918	0.1618	0.5149	0.4224
GNN	1D	0.4873	0.7843	0.7085	0.5126	0.3541	0.4430	0.4340	0.6063	0.5412
	3D	0.4800	0.7305	0.6125	0.4674	0.3419	0.3558	0.3325	0.5709	0.4864
	5D	0.4773	0.6881	0.5911	0.4507	0.3064	0.2782	0.3179	0.5243	0.4543
	7D	0.4705	0.6634	0.5482	0.4566	0.3343	0.3072	0.3234	0.4842	0.4485
	Avg.	0.4788	0.7166	0.6151	0.4718	0.3342	0.3460	0.3519	0.5464	0.4826
StemGNN	1D	0.4950	0.7993	0.6503	0.4581	0.3070	0.3161	0.3138	0.5819	0.4902
	3D	0.4878	0.6894	0.6336	0.4333	0.2762	0.2694	0.1468	0.4612	0.4247
	5D	0.4762	0.6203	0.6074	0.4065	0.2529	0.2048	0.1256	0.4329	0.3908
	7D	0.2821	0.5487	0.4545	0.3398	0.2205	0.1658	0.0749	0.3691	0.3069
	Avg.	0.4353	0.6644	0.5865	0.4094	0.2642	0.2390	0.1653	0.4613	0.4032
GPT4TS	1D	0.4909	0.7803	0.6876	0.4807	0.3624	0.4659	0.2754	0.6054	0.5186
	3D	0.4731	0.6916	0.6056	0.4087	0.3166	0.3103	0.1481	0.4977	0.4315
	5D	0.4595	0.6523	0.5701	0.3712	0.2935	0.2494	0.0887	0.4326	0.3896
	7D	0.4498	0.6217	0.5153	0.3324	0.2688	0.2017	0.0738	0.3889	0.3566
	Avg.	0.4683	0.6865	0.5947	0.3983	0.3103	0.3068	0.1465	0.4811	0.4241
LLM	1D	0.4910	0.7669	0.6932	0.5324	0.3777	0.5069	0.3207	0.6462	0.5419
	3D	0.4693	0.6760	0.6316	0.4674	0.3322	0.3641	0.2025	0.5692	0.4640
	5D	0.4547	0.6498	0.5702	0.4229	0.3073	0.2926	0.1335	0.5071	0.4173
	7D	0.4421	0.6195	0.5349	0.3920	0.2933	0.2368	0.1115	0.4639	0.3868
	Avg.	0.4643	0.6781	0.6075	0.4537	0.3277	0.3501	0.1920	0.5466	0.4525

Table 36: Benchmark(SEDI1) on part 3(Fort Myers area) stations(MLP,CNN,Transformer)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.4747	0.4387	0.5051	0.2137	0.1295	0.3272	0.0017	0.2836	0.2968
	3D	0.4444	0.4520	0.2868	0.2392	0.1628	0.2860	0.0006	0.3115	0.2729
	5D	0.4302	0.2950	0.1017	0.1766	0.1696	0.3612	0.0005	0.3690	0.2380
	7D	0.4116	0.2920	0.2236	0.1575	0.1613	0.2435	0.0004	0.2357	0.2157
	Avg.	0.4402	<u>0.3694</u>	0.2793	0.1968	0.1558	0.3045	0.0008	0.3000	0.2558
MLP	1D	0.4450	0.5180	0.4305	0.2132	0.1929	0.2698	0.0000	0.3516	0.3026
	3D	0.3848	0.3874	0.2667	0.1432	0.1651	0.1587	0.0000	0.2378	0.2180
	5D	0.3283	0.3044	0.2146	0.1247	0.1417	0.1347	0.0000	0.1959	0.1805
	7D	0.2849	0.2479	0.1902	0.1069	0.1197	0.1177	0.0000	0.1482	0.1519
	Avg.	0.3607	0.3644	0.2755	0.1470	0.1548	0.1702	0.0000	0.2334	0.2133
NLinear	1D	0.4657	0.5401	0.4711	0.2360	0.2096	0.3693	0.0017	0.3968	0.3363
	3D	0.4022	0.3957	0.2858	0.1465	0.1677	0.2077	0.0006	0.2691	0.2344
	5D	0.3454	0.3299	0.2238	0.1311	0.1556	0.1897	0.0003	0.2083	0.1980
	7D	0.3003	0.2835	0.1990	0.1112	0.1377	0.1638	0.0002	0.1777	0.1717
	Avg.	0.3784	0.3873	0.2949	0.1562	<u>0.1676</u>	0.2326	<u>0.0007</u>	0.2630	0.2351
TCN	1D	0.3817	0.3979	0.5681	0.1620	0.1078	0.2254	0.0001	0.2565	0.2624
	3D	0.3269	0.2261	0.2620	0.1106	0.1482	0.1507	0.0000	0.1577	0.1728
	5D	0.3281	0.3077	0.2576	0.1217	0.1144	0.1199	0.0000	0.1687	0.1773
	7D	0.2895	0.2433	0.2785	0.0881	0.0582	0.1289	0.0000	0.1664	0.1566
	Avg.	0.3315	0.2938	0.3415	0.1206	0.1072	0.1562	0.0000	0.1873	0.1923
CNN	1D	0.4700	0.3796	0.4606	0.1984	0.1834	0.2864	0.0003	0.3521	0.2913
	3D	0.4134	0.2972	0.3154	0.1244	0.1610	0.1089	0.0000	0.2540	0.2093
	5D	0.3802	0.2393	0.2666	0.0989	0.1319	0.0585	0.0000	0.1811	0.1696
	7D	0.3497	0.2258	0.2511	0.0857	0.1173	0.0431	0.0000	0.1418	0.1518
	Avg.	0.4033	0.2855	<u>0.3234</u>	0.1269	0.1484	0.1242	0.0001	0.2322	0.2055
TimesNet	1D	0.4591	0.3666	0.4525	0.1548	0.1686	0.1891	0.0000	0.2756	0.2583
	3D	0.4283	0.2930	0.2820	0.1182	0.1470	0.0856	0.0000	0.1680	0.1902
	5D	0.3777	0.2554	0.2337	0.1004	0.1092	0.0487	0.0000	0.1362	0.1577
	7D	0.3632	0.2254	0.1776	0.0801	0.0995	0.0378	0.0000	0.0923	0.1345
	Avg.	0.4071	0.2851	0.2865	0.1134	0.1311	0.0903	0.0000	0.1680	0.1852
Transformer	1D	0.4584	0.4340	0.4950	0.2389	0.1775	0.3983	0.0005	0.3726	0.3219
	3D	0.4305	0.2699	0.2939	0.1542	0.1563	0.1485	0.0001	0.2423	0.2120
	5D	0.3914	0.2541	0.2244	0.1336	0.1363	0.1145	0.0001	0.1856	0.1800
	7D	0.3506	0.2224	0.1918	0.1115	0.1161	0.0841	0.0000	0.1842	0.1576
	Avg.	0.4077	0.2951	0.3013	0.1595	0.1465	0.1863	0.0002	0.2462	0.2179
PatchTST	1D	0.4734	0.4614	0.4548	0.2418	0.1918	0.3446	0.0003	0.3847	0.3191
	3D	0.4278	0.3069	0.2981	0.1757	0.1645	0.1653	0.0001	0.2403	0.2223
	5D	0.3990	0.2060	0.2292	0.1506	0.1415	0.0919	0.0000	0.1697	0.1735
	7D	0.3603	0.1898	0.1968	0.1058	0.1200	0.0701	0.0000	0.1344	0.1472
	Avg.	<u>0.4151</u>	0.2910	0.2947	0.1685	0.1544	0.1680	0.0001	0.2323	0.2155

Table 37: Benchmark(SEDI1) on part 3(Fort Myers area) stations(RNN,GNN,LLM)

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
LSTM	1D	0.2500	0.3899	0.0246	0.1762	0.1810	0.1240	0.0000	0.2035	0.1686
	3D	0.2375	0.0599	0.0035	0.1100	0.1246	0.0055	0.0000	0.1008	0.0802
	5D	0.0000	0.0617	0.0000	0.0920	0.0927	0.0081	0.0000	0.0967	0.0439
	7D	0.2466	0.0486	0.0230	0.0938	0.0791	0.0256	0.0000	0.0555	0.0715
	Avg.	0.1835	0.1400	0.0128	0.1180	0.1194	0.0408	0.0000	0.1141	0.0911
RNN	1D	0.1603	0.0000	0.0998	0.1648	0.1890	0.1125	0.0000	0.1389	0.1082
	3D	0.2332	0.1744	0.0000	0.0798	0.0822	0.0015	0.0000	0.0887	0.0825
	5D	0.2488	0.0519	0.0000	0.1098	0.0928	0.0190	0.0000	0.0000	0.0653
	7D	0.0000	0.0091	0.0056	0.0657	0.0243	0.0088	0.0000	0.0961	0.0262
	Avg.	0.1606	0.0589	0.0263	0.1050	0.0971	0.0355	0.0000	0.0809	0.0705
DilatedRNN	1D	0.1049	0.3690	0.3560	0.3224	0.0969	0.3043	0.0003	0.2794	0.2291
	3D	0.2490	0.2811	0.0161	0.1159	0.1277	0.0744	0.0001	0.1527	0.1271
	5D	0.0949	0.2827	0.2367	0.1348	0.1246	0.0415	0.0000	0.2265	0.1427
	7D	0.2413	0.1618	0.2236	0.1230	0.1192	0.0497	0.0000	0.0805	0.1249
	Avg.	0.1725	0.2736	0.2081	0.1740	0.1171	0.1175	0.0001	0.1848	0.1560
GCN	1D	0.3817	0.3979	0.5681	0.1620	0.1078	0.2254	0.0001	0.2565	0.2624
	3D	0.3269	0.2261	0.2620	0.1106	0.1482	0.1507	0.0000	0.1577	0.1728
	5D	0.3281	0.3077	0.2576	0.1217	0.1144	0.1199	0.0000	0.1687	0.1773
	7D	0.2895	0.2433	0.2785	0.0881	0.0582	0.1289	0.0000	0.1664	0.1566
	Avg.	0.3315	0.2938	0.3415	0.1206	0.1072	0.1562	0.0000	0.1873	0.1923
GNN	1D	0.4363	0.3351	0.3967	0.2117	0.1928	0.2551	0.0018	0.2885	0.2647
	3D	0.4293	0.2745	0.2772	0.1733	0.2145	0.1698	0.0000	0.2810	0.2274
	5D	0.4124	0.3231	0.2661	0.1731	0.1441	0.2090	0.0000	0.2253	0.2191
	7D	0.3564	0.2544	0.2314	0.1561	0.2023	0.1808	0.0000	0.1790	0.1951
	Avg.	0.4086	0.2968	0.2928	0.1785	0.1884	0.2037	0.0005	0.2434	0.2266
StemGNN	1D	0.1616	0.4190	0.1862	0.1781	0.1094	0.1578	0.0002	0.2462	0.1823
	3D	0.2372	0.2514	0.1446	0.1443	0.0443	0.1068	0.0000	0.1309	0.1324
	5D	0.0956	0.1130	0.0178	0.1026	0.0790	0.0844	0.0000	0.1162	0.0761
	7D	0.0360	0.0911	0.0384	0.1245	0.0247	0.1175	0.0000	0.0726	0.0631
	Avg.	0.1326	0.2186	0.0968	0.1374	0.0644	0.1167	0.0001	0.1415	0.1135
GPT4TS	1D	0.4586	0.4502	0.4421	0.1424	0.1876	0.2122	0.0000	0.2792	0.2715
	3D	0.4088	0.3268	0.2713	0.1127	0.1530	0.0738	0.0000	0.1810	0.1909
	5D	0.3916	0.2650	0.2210	0.1011	0.1201	0.0412	0.0000	0.1263	0.1583
	7D	0.3606	0.2190	0.1939	0.0847	0.1091	0.0466	0.0000	0.1000	0.1392
	Avg.	0.4049	0.3153	0.2821	0.1102	0.1424	0.0935	0.0000	0.1716	0.1900
LLM	1D	0.4710	0.3713	0.4458	0.2398	0.1955	0.3082	0.0000	0.3205	0.2940
	3D	0.4297	0.2009	0.2890	0.1551	0.1653	0.1697	0.0000	0.2165	0.2033
	5D	0.3880	0.1895	0.2212	0.1363	0.1375	0.1271	0.0000	0.1561	0.1695
	7D	0.3494	0.1505	0.1984	0.1012	0.1175	0.0965	0.0000	0.1250	0.1423
	Avg.	0.4095	0.2280	0.2886	0.1581	0.1539	0.1753	0.0000	0.2045	0.2023

Table 38: Benchmark(MAE) on all stations

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.1324	0.1424	0.1441	0.1043	0.1098	0.1192	0.1423	0.1310	0.1282
	3D	0.2158	0.2352	0.2471	0.1838	0.1875	0.1903	0.2516	0.2082	0.2150
	5D	0.2810	0.2902	0.3178	0.2277	0.2437	0.2354	0.3228	0.2565	0.2719
	7D	0.3329	0.3374	0.3666	0.2661	0.2837	0.2719	0.3793	0.2915	0.3162
	Avg.	0.2405	0.2513	0.2689	0.1955	0.2062	0.2042	0.2740	0.2218	0.2328
TCN	1D	0.3079	0.2613	0.2634	0.2414	0.2079	0.1885	0.2957	0.2740	0.2550
	3D	0.4332	0.3215	0.3490	0.3061	0.2900	0.2554	0.3944	0.3465	0.3370
	5D	0.4324	0.3788	0.4040	0.3563	0.3457	0.3070	0.4547	0.3992	0.3848
	7D	0.4842	0.3951	0.4429	0.3915	0.3803	0.3397	0.4979	0.4242	0.4195
	Avg.	0.4144	0.3392	0.3648	0.3238	0.3060	0.2727	0.4107	0.3610	0.3491
LSTM	1D	0.1867	0.2613	0.2736	0.1595	0.2496	0.1697	0.3087	0.1836	0.2241
	3D	0.2862	0.3516	0.3751	0.2391	0.3291	0.2446	0.4267	0.2704	0.3153
	5D	0.3475	0.4012	0.4475	0.2892	0.3753	0.2944	0.4867	0.3172	0.3699
	7D	0.4174	0.4312	0.5125	0.3261	0.4193	0.3261	0.5173	0.3429	0.4116
	Avg.	0.3094	0.3613	0.4022	<u>0.2535</u>	0.3433	0.2587	0.4348	0.2785	0.3302
GCN	1D	0.1850	0.2317	0.2365	0.2061	0.1953	0.1681	0.1944	0.2152	0.2040
	3D	0.2668	0.3177	0.3426	0.2735	0.2782	0.2385	0.3119	0.2914	0.2901
	5D	0.3329	0.3673	0.4137	0.3184	0.3256	0.2796	0.3977	0.3336	0.3461
	7D	0.3863	0.3860	0.4624	0.3484	0.3646	0.3102	0.4265	0.3620	0.3808
	Avg.	<u>0.2928</u>	<u>0.3257</u>	<u>0.3638</u>	0.2866	<u>0.2909</u>	<u>0.2491</u>	<u>0.3326</u>	<u>0.3005</u>	<u>0.3053</u>

Table 39: Benchmark(MSE) on all stations

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.1144	0.1360	0.1284	0.0684	0.1124	0.0747	0.1674	0.1256	0.1159
	3D	0.2226	0.3080	0.2875	0.1478	0.3256	0.1424	0.5486	0.2405	0.2779
	5D	0.3117	0.4687	0.4318	0.2081	0.6883	0.1928	1.2099	0.3317	0.4804
	7D	0.3808	0.6967	0.6094	0.2648	1.0917	0.2369	1.8233	0.3899	0.6867
	Avg.	0.2574	0.4024	0.3643	0.1723	0.5545	0.1617	0.9373	0.2719	0.3902
TCN	1D	0.3964	0.5800	0.4714	0.2575	0.3783	0.2180	1.3093	0.5552	0.5208
	3D	0.6306	0.6849	0.6180	0.3337	0.5102	0.2900	1.5333	0.6517	0.6566
	5D	0.5709	0.7997	0.7216	0.3930	0.6323	0.3534	1.6885	0.7298	0.7362
	7D	0.6880	0.8404	0.7966	0.4639	0.7027	0.3941	1.8442	0.7434	0.8092
	Avg.	0.5715	0.7263	<u>0.6519</u>	0.3620	<u>0.5559</u>	0.3139	1.5938	0.6700	0.6807
LSTM	1D	0.2550	1.4732	1.0367	0.1620	5.2575	0.1497	6.8509	0.4084	1.9492
	3D	0.3871	1.3765	1.0735	0.2642	5.7927	0.2476	8.5420	0.5052	2.2736
	5D	0.4911	1.5590	1.4324	0.3373	5.8335	0.3032	8.3886	0.5652	2.3638
	7D	0.6198	1.6812	1.8663	0.3982	6.0814	0.3494	8.5648	0.6176	2.5224
	Avg.	0.4383	1.5225	1.3522	<u>0.2904</u>	5.7413	0.2625	8.0866	<u>0.5241</u>	2.2772
GCN	1D	0.1994	0.2886	0.7728	0.3590	0.2534	0.1165	0.2171	6.2444	1.0564
	3D	0.3102	0.4645	1.0667	0.4061	0.5848	0.1883	0.9433	3.7576	0.9652
	5D	0.4273	0.5908	1.3546	0.4490	0.8592	0.2408	2.4139	2.7351	1.1338
	7D	0.5166	0.5968	1.5264	0.4728	1.3609	0.2752	1.8411	2.3212	1.1139
	Avg.	0.3634	<u>0.4852</u>	1.1801	0.4217	0.7646	<u>0.2052</u>	<u>1.3539</u>	3.7646	1.0673

Table 40: Benchmark(SEDI10) on all stations

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.7865	0.8159	0.8152	0.7883	0.7313	0.7192	0.7116	0.7581	0.7658
	3D	0.7108	0.7527	0.7714	0.7639	0.6709	0.5924	0.6336	0.6414	0.6921
	5D	0.6823	0.6972	0.7482	0.7092	0.6185	0.5809	0.5830	0.6343	0.6567
	7D	0.6455	0.6726	0.7134	0.6753	0.6142	0.5620	0.5496	0.5815	0.6268
	Avg.	0.7063	0.7346	0.7621	0.7342	0.6587	0.6136	0.6195	0.6538	0.6853
TCN	1D	0.6211	0.7314	0.7264	0.6545	0.5807	0.6232	0.5594	0.4628	0.6199
	3D	0.4990	0.6284	0.6505	0.5697	0.4678	0.5020	0.4852	0.3738	0.5221
	5D	0.4997	0.6025	0.6164	0.5319	0.4059	0.4549	0.4289	0.3187	0.4824
	7D	0.4366	0.5680	0.5829	0.4747	0.3637	0.4181	0.3906	0.3059	0.4426
	Avg.	0.5141	0.6326	0.6440	0.5577	0.4545	0.4996	0.4660	0.3653	0.5167
LSTM	1D	0.7478	0.7266	0.7410	0.7615	0.6781	0.6422	0.6255	0.6716	0.6993
	3D	0.6290	0.6424	0.6606	0.6671	0.5615	0.5070	0.5091	0.5369	0.5892
	5D	0.5632	0.5683	0.6139	0.6053	0.5035	0.4434	0.4307	0.4694	0.5247
	7D	0.5249	0.5386	0.5845	0.5758	0.4605	0.3974	0.3886	0.4256	0.4870
	Avg.	0.6162	0.6190	0.6500	0.6524	0.5509	0.4975	0.4885	0.5259	0.5751
GCN	1D	0.7461	0.7112	0.7543	0.7031	0.6442	0.6466	0.6565	0.6686	0.6913
	3D	0.6725	0.6669	0.6912	0.6308	0.5846	0.5725	0.5800	0.6044	0.6254
	5D	0.6331	0.6218	0.6668	0.5908	0.5515	0.5247	0.5314	0.5519	0.5840
	7D	0.5989	0.5811	0.6531	0.5765	0.5183	0.4903	0.5025	0.5109	0.5539
	Avg.	0.6626	0.6452	0.6913	0.6253	0.5746	0.5585	0.5676	0.5840	0.6137

Table 41: Benchmark(SEDI5) on all stations

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.7136	0.7660	0.7338	0.7105	0.6322	0.6125	0.6060	0.6216	0.6745
	3D	0.6163	0.7050	0.6930	0.6926	0.5791	0.4788	0.5176	0.5137	0.5995
	5D	0.5987	0.6437	0.6679	0.6416	0.5329	0.4810	0.4710	0.5052	0.5678
	7D	0.5702	0.6306	0.6431	0.6119	0.5267	0.4753	0.4457	0.4662	0.5462
	Avg.	0.6247	0.6863	0.6845	0.6642	0.5678	0.5119	0.5101	0.5267	0.5970
TCN	1D	0.4196	0.6603	0.6184	0.5291	0.4095	0.4960	0.4138	0.3313	0.4848
	3D	0.3450	0.5523	0.5307	0.4549	0.2944	0.3870	0.3365	0.2404	0.3926
	5D	0.3236	0.5302	0.5064	0.4307	0.2343	0.3585	0.2869	0.2074	0.3597
	7D	0.2981	0.4912	0.4486	0.3805	0.1940	0.3273	0.2666	0.1886	0.3243
	Avg.	0.3466	0.5585	0.5260	0.4488	0.2831	0.3922	0.3259	0.2419	0.3904
LSTM	1D	0.6546	0.6720	0.6402	0.6737	0.5568	0.5009	0.4883	0.5126	0.5874
	3D	0.5112	0.5662	0.5342	0.5557	0.4398	0.3529	0.3555	0.3751	0.4613
	5D	0.4395	0.4974	0.4877	0.4993	0.3820	0.3008	0.2920	0.2929	0.3990
	7D	0.3879	0.4701	0.4458	0.4642	0.3109	0.2694	0.2556	0.2664	0.3588
	Avg.	0.4983	0.5514	0.5270	0.5482	0.4224	0.3560	0.3479	0.3618	0.4516
GCN	1D	0.6660	0.6406	0.6734	0.6192	0.5498	0.5445	0.5430	0.5278	0.5955
	3D	0.5885	0.6013	0.6062	0.5427	0.4961	0.4673	0.4629	0.4690	0.5292
	5D	0.5593	0.5528	0.5793	0.5059	0.4684	0.4301	0.4192	0.4291	0.4930
	7D	0.5471	0.5177	0.5610	0.4934	0.4320	0.4081	0.3855	0.3945	0.4674
	Avg.	0.5902	0.5781	0.6050	0.5403	0.4866	0.4625	0.4526	0.4551	0.5213

Table 42: Benchmark(SEDI1) on all stations

Method	T	S0	S1	S2	S3	S4	S5	S6	S7	Avg.
MLP	1D	0.5452	0.6313	0.5586	0.5222	0.3956	0.3222	0.3557	0.3511	0.4602
	3D	0.4265	0.5790	0.5332	0.5326	0.3359	0.2318	0.2916	0.2890	0.4025
	5D	0.4535	0.5114	0.5228	0.4968	0.3349	0.2440	0.2813	0.3036	0.3935
	7D	0.4738	0.5377	0.5065	0.4550	0.3195	0.2545	0.2705	0.2740	0.3864
	Avg.	0.4748	0.5649	0.5303	0.5017	0.3465	0.2631	0.2998	0.3044	0.4107
TCN	1D	0.1333	0.4546	0.4028	0.3001	0.0736	0.2119	0.1582	0.1412	0.2345
	3D	0.1118	0.3684	0.3182	0.2284	0.0440	0.1646	0.1282	0.0833	0.1809
	5D	0.1033	0.3531	0.3050	0.2328	0.0305	0.1499	0.1147	0.0757	0.1706
	7D	0.0802	0.3176	0.2611	0.2063	0.0249	0.1433	0.0978	0.0746	0.1507
	Avg.	0.1071	0.3734	0.3218	0.2419	0.0432	0.1674	0.1247	0.0937	0.1842
LSTM	1D	0.4140	0.4712	0.4004	0.4026	0.2813	0.1812	0.2226	0.2254	0.3249
	3D	0.2247	0.2719	0.2113	0.2167	0.1419	0.0936	0.1123	0.1012	0.1717
	5D	0.1839	0.1920	0.1746	0.1489	0.1184	0.0703	0.0868	0.0744	0.1312
	7D	0.1173	0.1752	0.1429	0.1642	0.0394	0.0573	0.0715	0.0608	0.1036
	Avg.	0.2350	0.2776	0.2323	0.2331	0.1453	0.1006	0.1233	0.1155	0.1828
GCN	1D	0.4770	0.4553	0.4873	0.4167	0.3108	0.2708	0.3098	0.2742	0.3752
	3D	0.4046	0.4419	0.3886	0.3418	0.2585	0.2237	0.2572	0.2331	0.3187
	5D	0.3840	0.3823	0.3712	0.3168	0.2323	0.1991	0.2144	0.2147	0.2894
	7D	0.3641	0.3583	0.3773	0.2930	0.2013	0.1843	0.1942	0.1889	0.2702
	Avg.	0.4074	0.4095	0.4061	0.3421	0.2507	0.2195	0.2439	0.2277	0.3134

Table 43: MAE results of input factors ablation study on S6. All, G, R, and C represent all factors, groundwater, rainfall, and human control(pump and gate).

Method	T	w/ All	w/o G	w/o R	w/o C	w/o GR	w/o RC	w/o WC	w/o WRC
iTTransformer	1D	0.0738	0.0735	0.0729	0.0733	0.0739	0.0737	0.0739	0.0742
	3D	0.1275	0.1279	0.1278	0.1275	0.1290	0.1280	0.1265	0.1276
	5D	0.1660	0.1662	0.1664	0.1662	0.1660	0.1660	0.1658	0.1671
	7D	0.1953	0.1953	0.1953	0.1948	0.1954	0.1961	0.1947	0.1955
	Avg.	0.1406	0.1407	0.1406	0.1405	0.1411	0.1410	0.1402	0.1411
PatchTST	1D	0.0726	0.0731	0.0713	0.0719	0.0732	0.0722	0.0727	0.0772
	3D	0.1252	0.1262	0.1252	0.1255	0.1273	0.1273	0.1254	0.1295
	5D	0.1615	0.1639	0.1624	0.1618	0.1649	0.1649	0.1634	0.1667
	7D	0.1910	0.1933	0.1923	0.1916	0.1938	0.1941	0.1927	0.1951
	Avg.	0.1376	0.1391	<u>0.1378</u>	0.1377	0.1398	0.1396	0.1385	0.1421
TSMixer	1D	0.1004	0.0781	0.1599	0.1443	0.0778	0.1789	0.0782	0.0782
	3D	0.1471	0.1296	0.2611	0.2216	0.1297	0.2506	0.1299	0.1299
	5D	0.1802	0.1658	0.3104	0.2350	0.1657	0.3018	0.1664	0.1663
	7D	0.2107	0.1941	0.2776	0.2435	0.1939	0.3914	0.1947	0.1944
	Avg.	0.1596	0.1419	0.2523	0.2111	0.1418	0.2807	0.1423	0.1422
NLinear	1D	0.0937	0.0977	0.0854	0.0933	0.0850	0.0804	0.0983	0.0780
	3D	0.1429	0.1461	0.1363	0.1422	0.1366	0.1328	0.1462	0.1314
	5D	0.1769	0.1796	0.1713	0.1764	0.1717	0.1689	0.1797	0.1678
	7D	0.2050	0.2074	0.2003	0.2043	0.2008	0.1978	0.2072	0.1970
	Avg.	0.1546	0.1577	0.1483	0.1540	0.1485	<u>0.1450</u>	0.1579	0.1435
TimesNet	1D	0.0986	0.0936	0.0986	0.0986	0.0928	0.0970	0.0924	0.0930
	3D	0.1513	0.1446	0.1511	0.1525	0.1465	0.1529	0.1448	0.1468
	5D	0.1883	0.1838	0.1889	0.1890	0.1843	0.1908	0.1830	0.1814
	7D	0.2187	0.2177	0.2262	0.2200	0.2131	0.2219	0.2111	0.2109
	Avg.	0.1642	0.1599	0.1662	0.1650	0.1592	0.1656	0.1578	0.1580

Table 44: MSE results of input factors ablation study on S6. All, G, R, and C represent all factors, groundwater, rainfall, and human control(pump and gate).

Method	T	w/ All	w/o G	w/o R	w/o C	w/o GR	w/o RC	w/o WC	w/o WRC
iTTransformer	1D	0.0400	0.0407	0.0398	0.0402	0.0405	0.0400	0.0405	0.0409
	3D	0.0817	0.0828	0.0821	0.0818	0.0827	0.0825	0.0818	0.0832
	5D	0.1175	0.1172	0.1170	0.1167	0.1166	0.1169	0.1164	0.1184
	7D	0.1419	0.1430	0.1407	0.1430	0.1429	0.1431	0.1407	0.1424
	Avg.	0.0953	0.0960	0.0949	0.0954	0.0957	0.0956	0.0949	0.0962
PatchTST	1D	0.0395	0.0393	0.0392	0.0393	0.0405	0.0399	0.0389	0.0432
	3D	0.0798	0.0807	0.0806	0.0797	0.0831	0.0819	0.0797	0.0850
	5D	0.1107	0.1128	0.1122	0.1105	0.1154	0.1140	0.1123	0.1173
	7D	0.1368	0.1400	0.1388	0.1367	0.1412	0.1393	0.1385	0.1428
	Avg.	0.0917	0.0932	0.0927	0.0916	0.0950	0.0938	0.0923	0.0971
TSMixer	1D	0.0562	0.0415	0.1891	0.2590	0.0414	0.4396	0.0419	0.0416
	3D	0.0957	0.0827	0.4979	0.2580	0.0830	0.4833	0.0825	0.0826
	5D	0.1243	0.1145	0.5722	0.3232	0.1145	0.4855	0.1139	0.1141
	7D	0.1557	0.1396	0.3653	0.2389	0.1394	1.2701	0.1389	0.1382
	Avg.	0.1080	0.0946	0.4061	0.2698	0.0946	0.6697	0.0943	0.0941
NLinear	1D	0.0477	0.0497	0.0450	0.0471	0.0459	0.0434	0.0494	0.0426
	3D	0.0878	0.0898	0.0848	0.0871	0.0856	0.0837	0.0895	0.0829
	5D	0.1178	0.1196	0.1154	0.1171	0.1160	0.1145	0.1191	0.1139
	7D	0.1434	0.1451	0.1411	0.1425	0.1417	0.1400	0.1444	0.1395
	Avg.	0.0992	0.1011	0.0966	0.0984	0.0973	<u>0.0954</u>	0.1006	<u>0.0947</u>
TimesNet	1D	0.0564	0.0518	0.0580	0.0573	0.0521	0.0550	0.0515	0.0517
	3D	0.1049	0.0970	0.1043	0.1049	0.0985	0.1086	0.0966	0.1015
	5D	0.1409	0.1357	0.1419	0.1416	0.1397	0.1485	0.1338	0.1338
	7D	0.1729	0.1770	0.1863	0.1770	0.1677	0.1826	0.1624	0.1624
	Avg.	0.1188	0.1154	0.1226	0.1202	0.1145	0.1237	0.1111	<u>0.1123</u>

Table 45: SEDI(10%) results of input factors ablation study on S6. All, G, R, and C represent all factors, groundwater, rainfall, and human control(pump and gate).

Method	T	w/ All	w/o G	w/o R	w/o C	w/o GR	w/o RC	w/o WC	w/o WRC
iTTransformer	1D	0.6508	0.6479	0.6510	0.6551	0.6525	0.6548	0.6512	0.6520
	3D	0.5526	0.5429	0.5485	0.5510	0.5393	0.5485	0.5513	0.5533
	5D	0.4856	0.4793	0.4831	0.4877	0.4786	0.4864	0.4824	0.4779
	7D	0.4384	0.4323	0.4408	0.4399	0.4301	0.4380	0.4320	0.4326
	Avg.	0.5318	0.5256	0.5308	0.5334	0.5251	<u>0.5319</u>	0.5292	0.5289
PatchTST	1D	0.6632	0.6629	0.6665	0.6644	0.6686	0.6618	0.6621	0.6561
	3D	0.5696	0.5614	0.5704	0.5717	0.5616	0.5586	0.5658	0.5455
	5D	0.5028	0.5013	0.4997	0.5043	0.4926	0.4945	0.4993	0.4857
	7D	0.4562	0.4518	0.4498	0.4544	0.4442	0.4465	0.4476	0.4364
	Avg.	0.5480	0.5443	0.5466	0.5487	0.5418	0.5404	0.5437	0.5309
TSMixer	1D	0.6052	0.6538	0.5202	0.5722	0.6475	0.5493	0.6499	0.6525
	3D	0.5267	0.5571	0.4251	0.4290	0.5580	0.4402	0.5581	0.5596
	5D	0.4732	0.4959	0.3523	0.4067	0.4963	0.3709	0.4932	0.4973
	7D	0.4245	0.4487	0.3516	0.3703	0.4497	0.3206	0.4454	0.4474
	Avg.	0.5074	0.5389	0.4123	0.4446	0.5379	0.4202	0.5367	0.5392
NLinear	1D	0.6477	0.6485	0.6485	0.6481	0.6506	0.6529	0.6602	0.6561
	3D	0.5476	0.5556	0.5521	0.5472	0.5521	0.5566	0.5517	0.5582
	5D	0.4868	0.4928	0.4905	0.4865	0.4904	0.4936	0.4960	0.4950
	7D	0.4356	0.4355	0.4383	0.4345	0.4381	0.4440	0.4345	0.4450
	Avg.	0.5294	0.5331	0.5324	0.5291	0.5328	<u>0.5368</u>	0.5356	0.5386
TimesNet	1D	0.5936	0.6093	0.5855	0.5884	0.6067	0.5939	0.6086	0.6089
	3D	0.4772	0.5059	0.4986	0.4925	0.5117	0.4854	0.5093	0.5046
	5D	0.4304	0.4489	0.4264	0.4305	0.4460	0.4202	0.4456	0.4396
	7D	0.3878	0.3959	0.3696	0.3837	0.4095	0.3839	0.4091	0.4082
	Avg.	0.4723	0.4900	0.4700	0.4738	0.4935	0.4709	<u>0.4931</u>	0.4903

Table 46: MAE results of temporal information ablation study on S6.

Method	T	6H	12H	1D	2D	3D	4D	5D	6D
iTTransformer	1D	0.1136	0.1123	0.1097	0.1123	0.1133	0.1198	0.1214	0.1209
	3D	0.1999	0.1978	0.1980	0.2025	0.2071	0.2121	0.2168	0.2137
	5D	0.2571	0.2537	0.2547	0.2586	0.2725	0.2776	0.2815	0.2779
	7D	0.3036	0.3007	0.3015	0.3059	0.3179	0.3242	0.3221	0.3261
	Avg.	0.2185	<u>0.2161</u>	0.2160	0.2198	0.2277	0.2334	0.2354	0.2346
PatchTST	1D	0.1266	0.1164	0.1135	0.1127	0.1094	0.1099	0.1106	0.1159
	3D	0.2123	0.2067	0.1988	0.1979	0.1963	0.1940	0.1979	0.2011
	5D	0.2693	0.2650	0.2566	0.2530	0.2554	0.2518	0.2522	0.2559
	7D	0.3143	0.3116	0.3039	0.2988	0.2984	0.2983	0.3011	0.3069
	Avg.	0.2306	0.2249	0.2182	0.2156	<u>0.2149</u>	0.2135	0.2155	0.2199
TSMixer	1D	0.1197	0.1216	0.1237	0.1269	0.1258	0.1269	0.1289	0.1266
	3D	0.2021	0.2033	0.2058	0.2087	0.2051	0.2073	0.2074	0.2052
	5D	0.2586	0.2592	0.2618	0.2639	0.2576	0.2604	0.2585	0.2582
	7D	0.3046	0.3049	0.3070	0.3092	0.3007	0.3029	0.2997	0.3009
	Avg.	0.2213	<u>0.2222</u>	0.2246	0.2272	0.2223	0.2244	0.2236	0.2227
NLinear	1D	0.1159	0.1180	0.1211	0.1462	0.1505	0.1508	0.1587	0.1586
	3D	0.1998	0.2011	0.2047	0.2220	0.2289	0.2305	0.2388	0.2382
	5D	0.2568	0.2577	0.2605	0.2743	0.2796	0.2870	0.2871	0.2925
	7D	0.3031	0.3036	0.3064	0.3186	0.3234	0.3293	0.3323	0.3355
	Avg.	0.2189	<u>0.2201</u>	0.2232	0.2403	0.2456	0.2494	0.2542	0.2562
TimesNet	1D	0.1175	0.1234	0.1357	0.1565	0.1750	0.1903	0.2066	0.2184
	3D	0.2017	0.2084	0.2235	0.2423	0.2618	0.2775	0.2880	0.3012
	5D	0.2602	0.2635	0.2757	0.2887	0.2989	0.3235	0.3386	0.3657
	7D	0.3086	0.3097	0.3254	0.3272	0.3570	0.3636	0.3804	0.4016
	Avg.	0.2220	<u>0.2262</u>	0.2401	0.2537	0.2732	0.2887	0.3034	0.3217

Table 47: MSE results of temporal information ablation study on S6.

Method	T	6H	12H	1D	2D	3D	4D	5D	6D
iTTransformer	1D	0.0829	0.0815	0.0793	0.0825	0.0844	0.0878	0.0899	0.0899
	3D	0.2093	0.2068	0.2082	0.2152	0.2218	0.2281	0.2379	0.2306
	5D	0.3186	0.3162	0.3174	0.3219	0.3472	0.3551	0.3556	0.3594
	7D	0.4170	0.4149	0.4139	0.4220	0.4499	0.4614	0.4614	0.4757
	Avg.	0.2569	<u>0.2549</u>	0.2547	0.2604	0.2758	0.2831	0.2862	0.2889
PatchTST	1D	0.0888	0.0860	0.0851	0.0827	0.0795	0.0787	0.0788	0.0835
	3D	0.2151	0.2104	0.2107	0.2078	0.2064	0.2050	0.2072	0.2089
	5D	0.3243	0.3214	0.3210	0.3156	0.3130	0.3256	0.3408	0.3187
	7D	0.4236	0.4187	0.4270	0.4193	0.4275	0.4548	0.4255	0.4348
	Avg.	0.2630	0.2592	0.2609	0.2563	0.2566	0.2660	<u>0.2631</u>	0.2615
TSMixer	1D	0.0899	0.0899	0.0888	0.0887	0.0860	0.0853	0.0855	0.0837
	3D	0.2152	0.2145	0.2141	0.2121	0.2047	0.2042	0.2034	0.1992
	5D	0.3241	0.3228	0.3226	0.3170	0.3060	0.3050	0.3035	0.3024
	7D	0.4235	0.4222	0.4214	0.4142	0.3999	0.3982	0.3968	0.3985
	Avg.	0.2632	0.2623	0.2617	0.2580	0.2491	0.2482	<u>0.2473</u>	0.2460
NLinear	1D	0.0852	0.0858	0.0873	0.1111	0.1135	0.1122	0.1196	0.1176
	3D	0.2111	0.2115	0.2148	0.2370	0.2444	0.2436	0.2535	0.2509
	5D	0.3201	0.3206	0.3236	0.3439	0.3490	0.3587	0.3563	0.3629
	7D	0.4199	0.4200	0.4235	0.4427	0.4480	0.4555	0.4586	0.4623
	Avg.	0.2591	<u>0.2595</u>	0.2623	0.2836	0.2887	0.2925	0.2970	0.2984
TimesNet	1D	0.0866	0.0937	0.1104	0.1357	0.1603	0.1750	0.2039	0.2297
	3D	0.2177	0.2333	0.2594	0.2953	0.3156	0.3607	0.3811	0.4114
	5D	0.3316	0.3372	0.3781	0.3997	0.4169	0.4597	0.5154	0.5691
	7D	0.4363	0.4407	0.4918	0.4891	0.5910	0.5837	0.5953	0.6566
	Avg.	0.2681	<u>0.2762</u>	0.3100	0.3299	0.3710	0.3948	0.4239	0.4667

Table 48: SEDI(10%) results of temporal information ablation study on S6.

Method	T	6H	12H	1D	2D	3D	4D	5D	6D
iTTransformer	1D	0.7825	0.7945	0.7941	0.7898	0.7703	0.7857	0.7728	0.7632
	3D	0.6792	0.6945	0.6987	0.6957	0.6847	0.6801	0.6691	0.6627
	5D	0.6220	0.6392	0.6420	0.6318	0.6201	0.6189	0.6178	0.6022
	7D	0.5797	0.5984	0.6006	0.5992	0.5867	0.5841	0.5738	0.5665
	Avg.	0.6659	0.6816	0.6838	0.6791	0.6654	0.6672	0.6584	0.6487
PatchTST	1D	0.7522	0.7946	0.8004	0.8033	0.8100	0.8118	0.8058	0.8099
	3D	0.6324	0.6425	0.6977	0.7010	0.7046	0.7030	0.7033	0.7032
	5D	0.5820	0.5796	0.6389	0.6427	0.6404	0.6467	0.6362	0.6326
	7D	0.5415	0.5314	0.5959	0.6024	0.5986	0.5957	0.5929	0.5982
	Avg.	0.6270	0.6370	0.6832	0.6874	<u>0.6884</u>	0.6893	0.6846	0.6860
TSMixer	1D	0.7812	0.7886	0.7954	0.7912	0.7931	0.7965	0.7905	0.7952
	3D	0.6861	0.6906	0.6971	0.7005	0.7016	0.7067	0.6946	0.6983
	5D	0.6307	0.6351	0.6397	0.6463	0.6445	0.6503	0.6400	0.6385
	7D	0.5917	0.5942	0.5984	0.6054	0.6003	0.6038	0.5959	0.5900
	Avg.	0.6724	0.6771	0.6827	<u>0.6859</u>	0.6849	0.6893	0.6803	0.6805
NLinear	1D	0.7915	0.8002	0.8029	0.7843	0.7891	0.7933	0.7929	0.7904
	3D	0.6956	0.6989	0.7019	0.6857	0.6958	0.6974	0.6925	0.6940
	5D	0.6359	0.6383	0.6447	0.6320	0.6329	0.6373	0.6422	0.6373
	7D	0.5944	0.5995	0.6016	0.5875	0.5889	0.5951	0.5973	0.5994
	Avg.	0.6794	<u>0.6842</u>	0.6878	0.6724	0.6767	0.6808	0.6812	0.6803
TimesNet	1D	0.7590	0.7610	0.7482	0.7130	0.6908	0.6854	0.6548	0.6392
	3D	0.6582	0.6606	0.6499	0.6148	0.5993	0.5889	0.5790	0.5731
	5D	0.5976	0.6096	0.5988	0.5779	0.5631	0.5461	0.5458	0.5246
	7D	0.5647	0.5692	0.5451	0.5447	0.5269	0.5263	0.5062	0.5060
	Avg.	0.6449	0.6501	0.6355	0.6126	0.5950	0.5867	0.5714	0.5607

Table 49: MAE results of spatial information ablation study on S6.

Method	T	MAE					MSE					SEDI(10%)				
		1.0R	1.2R	1.4R	1.6R	1.8R	1.0R	1.2R	1.4R	1.6R	1.8R	1.0R	1.2R	1.4R	1.6R	1.8R
iTTransformer	1D	0.1123	0.1135	0.1125	0.1099	0.1119	0.0825	0.0825	0.0818	0.0822	0.0807	0.7898	0.7810	0.7894	0.7863	0.7939
	3D	0.2025	0.2038	0.2020	0.2024	0.2021	0.2152	0.2144	0.2117	0.2136	0.2119	0.6957	0.7011	0.6952	0.7022	0.7036
	5D	0.2586	0.2650	0.2584	0.2538	0.2513	0.3219	0.3314	0.3241	0.3257	0.3186	0.6318	0.6294	0.6329	0.6353	0.6361
	7D	0.3059	0.3040	0.3017	0.3006	0.2995	0.4220	0.4239	0.4181	0.4176	0.4197	0.5992	0.5973	0.5989	0.5972	0.5965
	Avg.	0.2198	0.2216	0.2187	<u>0.2167</u>	0.2162	0.2604	0.2631	<u>0.2589</u>	0.2598	0.2577	0.6791	0.6772	0.6791	<u>0.6803</u>	0.6825
PatchTST	1D	0.1127	0.1131	0.1139	0.1113	0.1118	0.0827	0.0833	0.0829	0.0825	0.0821	0.8033	0.8030	0.8066	0.8068	0.8095
	3D	0.1979	0.1967	0.1990	0.1976	0.1976	0.2078	0.2060	0.2087	0.2082	0.2083	0.7010	0.7036	0.7048	0.7040	0.7073
	5D	0.2530	0.2531	0.2548	0.2556	0.2550	0.3156	0.3133	0.3151	0.3182	0.3160	0.6427	0.6440	0.6433	0.6437	0.6483
	7D	0.2988	0.2971	0.2994	0.3000	0.2991	0.4193	0.4102	0.4102	0.4115	0.4097	0.6024	0.6035	0.6049	0.6056	0.6065
	Avg.	0.2156	0.2150	0.2168	0.2161	0.2159	0.2563	0.2532	0.2542	0.2551	<u>0.2540</u>	0.6874	0.6885	0.6899	<u>0.6900</u>	0.6929
TSMixer	1D	0.1269	0.1259	0.1255	0.1271	0.1256	0.0887	0.0882	0.0876	0.0888	0.0879	0.7912	0.7896	0.7918	0.7895	0.7891
	3D	0.2087	0.2082	0.2072	0.2075	0.2064	0.2121	0.2125	0.2096	0.2093	0.2096	0.7005	0.6996	0.7026	0.7011	0.7001
	5D	0.2639	0.2638	0.2621	0.2632	0.2625	0.3170	0.3182	0.3136	0.3147	0.3146	0.6463	0.6451	0.6503	0.6484	0.6492
	7D	0.3092	0.3086	0.3074	0.3082	0.3062	0.4142	0.4151	0.4117	0.4114	0.4097	0.6054	0.6042	0.6078	0.6066	0.6055
	Avg.	0.2272	0.2266	<u>0.2255</u>	0.2265	0.2252	0.2580	0.2585	<u>0.2556</u>	0.2560	0.2554	0.6859	0.6846	0.6881	<u>0.6864</u>	0.6860
NLinear	1D	0.1462	0.1482	0.1473	0.1464	0.1449	0.1111	0.1133	0.1123	0.1113	0.1097	0.7843	0.7816	0.7807	0.7811	0.7821
	3D	0.2220	0.2238	0.2227	0.2213	0.2205	0.2370	0.2397	0.2381	0.2358	0.2347	0.6857	0.6836	0.6850	0.6865	0.6876
	5D	0.2743	0.2756	0.2746	0.2744	0.2735	0.3439	0.3461	0.3443	0.3437	0.3423	0.6320	0.6300	0.6315	0.6317	0.6310
	7D	0.3186	0.3199	0.3191	0.3181	0.3173	0.4427	0.4450	0.4435	0.4415	0.4402	0.5875	0.5888	0.5904	0.5930	0.5932
	Avg.	0.2403	0.2419	0.2409	<u>0.2400</u>	0.2390	0.2836	0.2860	0.2845	0.2831	0.2817	0.6724	0.6710	0.6719	<u>0.6731</u>	0.6735
TimesNet	1D	0.1565	0.1582	0.1597	0.1655	0.1661	0.1357	0.1423	0.1392	0.1458	0.1492	0.7130	0.7099	0.7155	0.7093	0.7007
	3D	0.2423	0.2451	0.2408	0.2424	0.2454	0.2953	0.3075	0.2847	0.2893	0.3104	0.6148	0.6168	0.6174	0.6202	0.6131
	5D	0.2887	0.2935	0.3038	0.2924	0.2985	0.3997	0.4087	0.4531	0.4028	0.4239	0.5779	0.5812	0.5651	0.5824	0.5674
	7D	0.3272	0.3399	0.3380	0.3422	0.3413	0.4891	0.5223	0.5136	0.5219	0.5451	0.5447	0.5377	0.5359	0.5426	0.5410
	Avg.	0.2537	<u>0.2592</u>	0.2606	0.2606	0.2628	0.3299	0.3452	0.3476	<u>0.3400</u>	0.3571	0.6126	0.6114	0.6085	0.6136	0.6056