

# Economics 103 – Statistics for Economists

Francis J. DiTraglia

University of Pennsylvania

Lecture 15

# Sampling Distributions and Estimation – Part II

# Unbiased means “Right on Average”

## Bias of an Estimator

Let  $\hat{\theta}_n$  be a sample estimator of a population parameter  $\theta_0$ . The *bias* of  $\hat{\theta}_n$  is  $E[\hat{\theta}_n] - \theta_0$ .

## Unbiased Estimator

A sample estimator  $\hat{\theta}_n$  of a population parameter  $\theta_0$  is called *unbiased* if  $E[\hat{\theta}_n] = \theta_0$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] =$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] =$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{1}{n} E \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right] =$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{1}{n} E \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{(n-1)\sigma^2}{n}$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 =$$



## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{1}{n} E \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{(n-1)\sigma^2}{n}$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 =$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{1}{n} E \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{(n-1)\sigma^2}{n}$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \frac{n\sigma^2}{n} =$$

## A Different Estimator of the Population Variance

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E[\hat{\sigma}^2] = E \left[ \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{1}{n} E \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right] = \frac{(n-1)\sigma^2}{n}$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \frac{n\sigma^2}{n} = -\sigma^2/n$$

# Procedure versus Result of the Procedure

# Procedure versus Result of the Procedure

Procedure = Random Variable

# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.

# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents procedure of taking sample mean

# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents procedure of taking sample mean

Sampling Dist. = Probabilistic Behavior of Procedure



# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents procedure of taking sample mean

Sampling Dist. = Probabilistic Behavior of Procedure

If I repeat the procedure of taking the mean of a random sample over and over for many samples, what relative frequencies do I get for the sample means?

# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents procedure of taking sample mean

Sampling Dist. = Probabilistic Behavior of Procedure

If I repeat the procedure of taking the mean of a random sample over and over for many samples, what relative frequencies do I get for the sample means?

Result of Procedure = Constant

# Procedure versus Result of the Procedure

Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents procedure of taking a random sample.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents procedure of taking sample mean

Sampling Dist. = Probabilistic Behavior of Procedure

If I repeat the procedure of taking the mean of a random sample over and over for many samples, what relative frequencies do I get for the sample means?

Result of Procedure = Constant

- ▶  $x_1, \dots, x_n$  is the result of sampling, the observed data.

# Procedure versus Result of the Procedure

## Procedure = Random Variable

- ▶  $X_1, \dots, X_n$  represents **procedure of taking a random sample**.
- ▶  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  represents **procedure of taking sample mean**

## Sampling Dist. = Probabilistic Behavior of Procedure

If I repeat the procedure of taking the mean of a random sample over and over for many samples, what relative frequencies do I get **for the sample means?**

## Result of Procedure = Constant

- ▶  $x_1, \dots, x_n$  is the **result of sampling**, the observed data.
- ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$  is the **result of taking sample mean**

## Procedure? Long-Run Relative Frequencies?

Why would I advise you not to play the lottery?

## Procedure? Long-Run Relative Frequencies?

Why would I advise you not to play the lottery?

- ▶ You may sometimes win, but if you play the lottery many times, on average you will lose money.

## Procedure? Long-Run Relative Frequencies?

Why would I advise you not to play the lottery?

- ▶ You may sometimes win, but if you play the lottery many times, on average you will lose money.
- ▶ Let  $X$  be a random variable representing lottery winnings. I am arguing that  $E[X] - \text{Cost of Ticket} < 0$

## Procedure? Long-Run Relative Frequencies?

Why would I advise you not to play the lottery?

- ▶ You may sometimes win, but if you play the lottery many times, on average you will lose money.
- ▶ Let  $X$  be a random variable representing lottery winnings. I am arguing that  $E[X] - \text{Cost of Ticket} < 0$

Procedure = Random Variable

Making a habit of playing the lottery. Expectation is negative.



## Procedure? Long-Run Relative Frequencies?

Why would I advise you not to play the lottery?

- ▶ You may sometimes win, but if you play the lottery many times, on average you will lose money.
- ▶ Let  $X$  be a random variable representing lottery winnings. I am arguing that  $E[X] - \text{Cost of Ticket} < 0$

### Procedure = Random Variable

Making a habit of playing the lottery. Expectation is negative.

### Result of that Procedure = Constant

How much you win in a *particular* lottery. Could be greater than or less than cost of ticket in any *individual* instance.

## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$

Choose  $n$  Students from Class List with Replacement

## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$

Choose  $n$  Students from Class List with Replacement



```
graph TD; A[Choose n Students from Class List with Replacement] --> B[Sample 1]
```

Sample 1


## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$

Choose  $n$  Students from Class List with Replacement


Sample 1

$\bar{x}_1$


## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$




## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$



## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$




## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$






## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$




## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$



Repeat  $M$  times  $\rightarrow$  get  $M$  different sample means

## Sampling Distribution of $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$



Repeat  $M$  times  $\rightarrow$  get  $M$  different sample means

Sampling Dist: long run relative frequencies of the  $\bar{x}_i$

# Height of Econ 103 Students





Figure : Left: Population, Right: Sampling distribution of  $\bar{X}_5$

## How Large is the Average Family?



How many brothers and sisters are in your family, including yourself?

The average number of children per family was about 2.0 twenty years ago.

# What's Going On Here?

# What's Going On Here?

Biased Sample!



# What's Going On Here?

Biased Sample!

- ▶ Zero children  $\Rightarrow$  didn't send any to college

# What's Going On Here?

Biased Sample!


- ▶ Zero children  $\Rightarrow$  didn't send any to college
- ▶ Sampling by *children* so large families **oversampled**

## Candy Weighing: 44 Estimates, Each With $n = 5$

$$\hat{\theta} = 20 \times (X_1 + \dots + X_5)$$

| Summary of Sampling Dist. |            |
|---------------------------|------------|
| Overestimates             | 42         |
| Exactly Correct           | 0          |
| Underestimates            | 2          |
| $E[\hat{\theta}]$         | 1269 grams |
| $SD(\hat{\theta})$        | 189 grams  |

Actual Mass:  $\theta_0 = 958$  grams



# What was in the bag?

100 Candies Total:

## What was in the bag?

100 Candies Total:

- ▶ 20 Fun Size Snickers Bars (large)

# What was in the bag?

100 Candies Total:

- ▶ 20 Fun Size Snickers Bars (large)
- ▶ 30 Reese's Miniatures (medium)

# What was in the bag?

100 Candies Total:

- ▶ 20 Fun Size Snickers Bars (large)
- ▶ 30 Reese's Miniatures (medium)
- ▶ 50 Tootsie Roll "Midgees" (small)

# What was in the bag?

100 Candies Total:

- ▶ 20 Fun Size Snickers Bars (large)
- ▶ 30 Reese's Miniatures (medium)
- ▶ 50 Tootsie Roll "Midgees" (small)

So What Happened?



# What was in the bag?

100 Candies Total:

- ▶ 20 Fun Size Snickers Bars (large)
- ▶ 30 Reese's Miniatures (medium)
- ▶ 50 Tootsie Roll "Midgees" (small)

## So What Happened?

Not a random sample! The Snickers bars were *oversampled*.

Could we have avoided this? How?



Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$  and define  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ . True or False:

*$\bar{X}_n$  is an unbiased estimator of  $\mu$*

- (a) True
- (b) False



Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$  and define  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ . True or False:

*$\bar{X}_n$  is an unbiased estimator of  $\mu$*

- (a) True
- (b) False

TRUE!



Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ . True or False:

*$X_1$  is an unbiased estimator of  $\mu$*

- (a) True
- (b) False



Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ . True or False:

*$X_1$  is an unbiased estimator of  $\mu$*

(a) True

(b) False

TRUE!

## How to choose between two unbiased estimators?

Suppose  $X_1, X_2, \dots, X_n \sim iid$  with mean  $\mu$  and variance  $\sigma^2$

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu$$

## How to choose between two unbiased estimators?

Suppose  $X_1, X_2, \dots, X_n \sim iid$  with mean  $\mu$  and variance  $\sigma^2$

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu$$

$$E[X_1] = \mu$$

## How to choose between two unbiased estimators?

Suppose  $X_1, X_2, \dots, X_n \sim iid$  with mean  $\mu$  and variance  $\sigma^2$

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu$$

$$E[X_1] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n Var(X_i) = \sigma^2/n$$



## How to choose between two unbiased estimators?

Suppose  $X_1, X_2, \dots, X_n \sim iid$  with mean  $\mu$  and variance  $\sigma^2$

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu$$

$$E[X_1] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n Var(X_i) = \sigma^2/n$$

$$Var(X_1) = \sigma^2$$

## Efficiency - Compare Unbiased Estimators by Variance

Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be unbiased estimators of  $\theta_0$ . We say that  $\hat{\theta}_1$  is *more efficient* than  $\hat{\theta}_2$  if  $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$ .

## Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

# Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

- ▶ Low bias estimators often have a high variance

# Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

- ▶ Low bias estimators often have a high variance
- ▶ Low variance estimators often have high bias

# Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

- ▶ Low bias estimators often have a high variance
- ▶ Low variance estimators often have high bias

Mean Squared Error (MSE):

# Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

- ▶ Low bias estimators often have a high variance
- ▶ Low variance estimators often have high bias

**Mean Squared Error (MSE):** Squared Bias plus Variance

$$MSE(\hat{\theta}) = \text{Bias}(\hat{\theta})^2 + \text{Var}(\hat{\theta})$$

# Mean-squared Error

Except in very simple situations, unbiased estimators are hard to come by. In fact, in many interesting applications there is a *tradeoff* between **bias** and **variance**:

- ▶ Low bias estimators often have a high variance
- ▶ Low variance estimators often have high bias


**Mean Squared Error (MSE):** Squared Bias plus Variance

$$\begin{aligned}MSE(\hat{\theta}) &= \text{Bias}(\hat{\theta})^2 + \text{Var}(\hat{\theta}) \\&= \left(E[\hat{\theta}] - \theta_0\right)^2 + \text{Var}(\hat{\theta})\end{aligned}$$



# Calculate MSE for Candy Experiment


|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |



Bias =

# Calculate MSE for Candy Experiment


|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |



$$\begin{aligned}\text{Bias} &= 1269 \text{ grams} - 958 \text{ grams} \\ &= \end{aligned}$$

# Calculate MSE for Candy Experiment

|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |




$$\text{Bias} = 1269 \text{ grams} - 958 \text{ grams}$$

$$= 311 \text{ grams}$$

$$\text{MSE} =$$

# Calculate MSE for Candy Experiment

|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |



$$\text{Bias} = 1269 \text{ grams} - 958 \text{ grams}$$


$$= 311 \text{ grams}$$

$$\text{MSE} = \text{Bias}^2 + \text{Variance}$$

=

# Calculate MSE for Candy Experiment

|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |




$$\begin{aligned}\text{Bias} &= 1269 \text{ grams} - 958 \text{ grams} \\ &= 311 \text{ grams}\end{aligned}$$

$$\begin{aligned}\text{MSE} &= \text{Bias}^2 + \text{Variance} \\ &= (311^2 + 189^2) \text{ grams}^2 \\ &= \end{aligned}$$

# Calculate MSE for Candy Experiment

|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |




$$\begin{aligned}\text{Bias} &= 1269 \text{ grams} - 958 \text{ grams} \\ &= 311 \text{ grams}\end{aligned}$$

$$\begin{aligned}\text{MSE} &= \text{Bias}^2 + \text{Variance} \\ &= (311^2 + 189^2) \text{ grams}^2 \\ &= 1.3244 \times 10^5 \text{ grams}^2\end{aligned}$$

# Calculate MSE for Candy Experiment

|                    |            |
|--------------------|------------|
| $E[\hat{\theta}]$  | 1269 grams |
| $\theta_0$         | 958 grams  |
| $SD(\hat{\theta})$ | 189 grams  |



$$\begin{aligned}\text{Bias} &= 1269 \text{ grams} - 958 \text{ grams} \\ &= 311 \text{ grams}\end{aligned}$$

$$\begin{aligned}\text{MSE} &= \text{Bias}^2 + \text{Variance} \\ &= (311^2 + 189^2) \text{ grams}^2 \\ &= 1.3244 \times 10^5 \text{ grams}^2\end{aligned}$$

$$\text{RMSE} = \sqrt{\text{MSE}} = 364 \text{ grams}$$

# Finite Sample versus Asymptotic Properties of Estimators

## Finite Sample Properties

For *fixed sample size*  $n$  what are the properties of the sampling distribution of  $\hat{\theta}_n$ ? (E.g. bias and variance.)



# Finite Sample versus Asymptotic Properties of Estimators

## Finite Sample Properties

For *fixed sample size  $n$*  what are the properties of the sampling distribution of  $\hat{\theta}_n$ ? (E.g. bias and variance.)

## Asymptotic Properties

What happens to the sampling distribution of  $\hat{\theta}_n$  *as the sample size  $n$  gets larger and larger?* (That is,  $n \rightarrow \infty$ ).

# Why Asymptotics?

## Law of Large Numbers

Make precise what we mean by “bigger samples are better.”

# Why Asymptotics?

## Law of Large Numbers

Make precise what we mean by “bigger samples are better.”

## Central Limit Theorem

As  $n \rightarrow \infty$  *nearly all* sampling distributions behave like a normal random variable!

# Consistency

## Consistency

If an estimator  $\hat{\theta}_n$  (which is a RV) *converges* to  $\theta_0$  (a constant) as  $n \rightarrow \infty$ , we say that  $\hat{\theta}_n$  *is consistent for  $\theta_0$* .

What does it mean for a *RV* to converge to a *constant*?

For this course we'll use *MSE Consistency*:

$$\lim_{n \rightarrow \infty} \text{MSE}(\hat{\theta}_n) = 0$$

This makes sense since  $\text{MSE}(\hat{\theta}_n)$  is a *constant*, so this is just an ordinary limit from calculus.

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ . Then the sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

is consistent for the population mean  $\mu$ .

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] =$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) =$$



## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \sigma^2/n$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \sigma^2/n$$

$$MSE(\bar{X}_n) = Bias(\bar{X}_n)^2 + Var(\bar{X}_n)$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \sigma^2/n$$

$$\begin{aligned} \text{MSE}(\bar{X}_n) &= \text{Bias}(\bar{X}_n)^2 + \text{Var}(\bar{X}_n) \\ &= (E[\bar{X}_n] - \mu)^2 + \text{Var}(\bar{X}_n) \end{aligned}$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \sigma^2/n$$

$$\begin{aligned} \text{MSE}(\bar{X}_n) &= \text{Bias}(\bar{X}_n)^2 + \text{Var}(\bar{X}_n) \\ &= (E[\bar{X}_n] - \mu)^2 + \text{Var}(\bar{X}_n) \\ &= 0 + \sigma^2/n \end{aligned}$$

## Law of Large Numbers (aka Law of Averages)

Let  $X_1, X_2, \dots, X_n \sim iid$  mean  $\mu$ , variance  $\sigma^2$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \mu$$

$$Var(\bar{X}_n) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \sigma^2/n$$

$$\begin{aligned} \text{MSE}(\bar{X}_n) &= \text{Bias}(\bar{X}_n)^2 + \text{Var}(\bar{X}_n) \\ &= (E[\bar{X}_n] - \mu)^2 + \text{Var}(\bar{X}_n) \\ &= 0 + \sigma^2/n \\ &\rightarrow 0 \end{aligned}$$

Hence  $\bar{X}_n$  is consistent for  $\mu$

# Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

# Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

## Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 =$$



## Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 =$$

## Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 = -\sigma^2/n$$

## Important!

An estimator *can* be biased but still consistent, as long as the bias disappears as  $n \rightarrow \infty$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

Bias of  $\hat{\sigma}^2$

$$E[\hat{\sigma}^2] - \sigma^2 = \frac{(n-1)\sigma^2}{n} - \sigma^2 = -\sigma^2/n \rightarrow 0$$



Suppose  $X_1, X_2, \dots, X_n \sim \text{iid } N(\mu, \sigma^2)$ . What is the sampling distribution of  $\bar{X}_n$ ?

- (a)  $\chi^2(n)$
- (b)  $t(n)$
- (c)  $F(n, n)$
- (d)  $N(\mu, \sigma^2/n)$
- (e) Not enough information to determine.

But still, how can something random  
converge to something constant?

## Sampling Distribution of $\bar{X}_n$ Collapses to $\mu$

Look at an example where we can directly calculate not only the mean and variance of the sampling distribution of  $\bar{X}_n$ , but the *sampling distribution itself*:

$$X_1, X_2, \dots, X_n \sim \text{iid } N(\mu, \sigma^2) \Rightarrow \bar{X}_n \sim N(\mu, \sigma^2/n)$$

## Sampling Distribution of $\bar{X}_n$ Collapses to $\mu$

$$X_1, X_2, \dots, X_n \sim \text{iid } N(\mu, \sigma^2) \Rightarrow \bar{X}_n \sim N(\mu, \sigma^2/n).$$





Figure : Sampling Distributions for  $\bar{X}_n$  where  $X_i \sim \text{iid } N(0, 1)$

## Another Visualization: Keep Adding Observations



| $n$   | $\bar{X}_n$ |
|-------|-------------|
| 1     | -2.69       |
| 2     | -3.18       |
| 3     | -5.94       |
| 4     | -4.27       |
| 5     | -2.62       |
| 10    | -2.89       |
| 20    | -5.33       |
| 50    | -2.94       |
| 100   | -1.58       |
| 500   | -0.45       |
| 1000  | -0.13       |
| 5000  | -0.05       |
| 10000 | 0.00        |

Figure : Running sample means:  $X_i \sim \text{iid } N(0, 100)$



# Important!

Although I showed two examples involving normal RVs, the LLN holds IN GENERAL!