



UNIVERSITAT POLITÈCNICA DE CATALUNYA
BARCELONATECH

Escola Superior d'Enginyeries Industrial,
Aeroespacial i Audiovisual de Terrassa

Interplanetary trajectories

Example: Earth to Mars case

Report

Degree: Master's degree in Aerospace Engineering

Course: 220301 - Aerodynamics, Flight and Orbital Mechanics

Delivery date: 15-01-2018

Students: Fontanes Molina, Pol; Martínez Viol, Víctor; Urbano González, Eva María

Contents

List of Tables	ii
List of Figures	iii
1 Figure example formats	1
2 Aim	2
3 Theoretical background	3
3.1 Planetary orbits and approximations analysis	3
3.1.1 Patched Conic Approximation (PCA)	3
3.1.1.1 1st. Geocentric stage	5
3.1.1.2 2nd. Heliocentric stage	5
3.1.1.3 3rd. Planetocentric stage	5
4 Calculations and results	6
4.1 Verification calculations	6
4.2 Main interplanetary orbit calculations	6
5 Conclusions	7
6 Bibliography	8

List of Tables

1.0.1	Thickness after the materials correction factor.	1
3.1.1	Radius of influence of the planets	4

List of Figures

1.0.1 Landing distance vs MTOW for the Boeing 777. 1

1 | Figure example formats

FIGURE




Figure 1.0.1: Landing distance vs MTOW for the Boeing 777.

TABLE

T_1	13 cm
T_2	21 cm
T_3	62 cm
T_t	95 cm

Table 1.0.1: Thickness after the materials correction factor.

2 | Aim

This project aims to compute an interplanetary trajectory which, for a given ecliptic rectangular positions of two planets in two known time instances, is able to carry a spaceship with a unique impulse, from the first planet to the second.

3 | Theoretical background

3.1 Planetary orbits and approximations analysis

In order to calculate the interplanetary trajectory between two planets, several approximations can be used. The simplest approximation, which can be called *aprox. 0* accepts the following hypothesis:

- Circular and coplanar orbits
- No analysis about the exit of the planet of start is done.
- No analysis about entering the planet of arrival is done.

Aprox. 0 is very basic and can be easily improved adding some parameters.

Another approximation widely used is **Patched Conic Approximation (PCA)**. This method improves significantly the results obtained with *aprox. 0* and represents a good starting point for a more precise numerical analysis of the mission. For this reason, in this project the PCA method will be used.

3.1.1 Patched Conic Approximation (PCA)

The Patched Conic Approximation (PCA) consist on the evaluation of an interplanetary trajectory dividing it into three stages. Considering the Earth as the planet of start, this stages are:

- Geocentric phase: Hyperbolic exit of the Earth. This phase takes place while the probe is going through the influence sphere of the Earth.
- Heliocentric phase: Elliptic trajectory with the Sun as main influencer.

Planetary orbits and approximations analysis

- Planet-centred phase: Hyperbolic arrival to the planet of destination. Similarly to the geocentric phase, this phase starts when the probe enters the sphere of influence of the planet.

The influence spheres mentioned are the space close to the planets where it can be considered that the influence of the Sun is negligible in comparison with that of the planet in question. The Laplace criteria will be considered to calculate this sphere. In Table 3.1.1 the radius of the sphere of influence of the solar system's planets are shown.

Planet	$R_I \times 10^6 \text{ km}$	$R_I \times 10^{-3} \text{ UA}$	R_I Radius of the planet
Mercury	0.111	0.740	45
Venus	0.616	4.11	100
Earth	0.924	6.16	145
Mars	0.577	3.85	170
Jupiter	48.157	321.0	677
Saturn	54.796	365.3	901
Uranus	91.954	346.4	2025
Neptune	80.196	534.6	3866

Table 3.1.1: Radius of influence of the planets

In order to set out the problem and begin with the resolution of it using the PCA method, the times and positions of the planets at the beginning and end of the trajectory are needed and some hypothesis are taken under consideration. The hypothesis are:

- The spheres of influences of the planets are not considered during the heliocentric phase. This hypothesis is admissible because the radius of the sphere are very small in comparison with the distance between planets.
- The spheres of influence are considered infinite from the point of view of the planet. This is assumed due to the fact that the radius of influence of the planets are much larger than the radius of the planet itself, as can be seen in Tabl 3.1.1.
- The duration of the trajectory can be taken as the duration of the heliocentric phase.

With this data the trajectory can be found through the orbital elements of the trajectories and the thrust required.

3.1.1.1 1st. Geocentric stage

3.1.1.2 2nd. Heliocentric stage

3.1.1.3 3rd. Planetocentric stage

4 | Calculations and results

4.1 Verification calculations

4.2 Main interplanetary orbit calculations

5 | Conclusions

6 | Bibliography

- [1] J. Calaf, "Trajectòries interplanetàries: Patched Conic Approximation," 2017.
- [2] —, "Trajectòries interplanetàries," 2017.
- [3] —, "Treballs de Mecànica Orbital," 2017.