

LSST Cadence Optimization for Disc-Intrinsic AGN Variability Science

Jackeline Moreno¹

jmoreno3663@gmail.com

and

Vishal Kasliwal^{2,3}

vishal.kasliwal@gmail.com

ABSTRACT

We propose a metric for the Metric Analysis Framework (MAF) that will help determine the utility of various OpSims for studying AGN accretion disc variability.

1. Introduction

Active Galactic Nuclei exhibit rapid stochastic (i.e., bereft of periodicity) flux variations at the $\sim 10 - 20$ % level on timescales ranging from hours to years across the electromagnetic spectrum (Ulrich et al. 1997; Peterson 1997). Variability may be observed in both the broad-line region (BLR) as well as in the continuum (Peterson et al. 1999). Continuum variability is thought to arise in the accretion disk of the AGN; however the exact origin of accretion disk variability is unknown (Maurizio et al. 2015). Probing the origins of accretion disk variability by matching phenomenological statistical models for the variability with physical models of accretion physics may prove to be of critical value in developing a deeper understanding of the nature of accretion.

The most popular statistical process used to describe AGN variability is the Continuous-time AutoRegressive Moving Average (C-ARMA) process (Kelly et al. 2014; Brockwell 2014) of which the damped random walk (DRW) process (Kelly et al. 2009) is the simplest kind. Kasliwal et al. (2016) argue in favor of a physical interpretation of C-ARMA processes that may reveal how

¹Drexel University
Department of Physics
3141 Chestnut Street
Philadelphia, PA 19104, USA

²University of Pennsylvania
Department of Physics & Astronomy
209 S. 33rd St.
Philadelphia, PA 19103, USA

³Princeton University
Department of Astrophysical Sciences
4 Ivy Ln
Princeton, NJ 08544, USA

flux perturbations in the accretion disk of an AGN evolve. Kasliwal et al. (2016) show that flux perturbations in the *Kepler* light curve of the Seyfert 1 AGN Zw 229-15 rapidly intensify over a period of ~ 5.5 d before exponentially decaying to the mean level with a decay timescale of ~ 70 d. The shape of the light curve of individual fluctuations is consistent with that expected from a 2nd-order differential equation suggesting that some combination of turbulent dissipation, thermal diffusion and viscous diffusion may be at work in the AGN accretion disk. The application of such analysis methods to the large samples of AGN light curves expected from LSST may tremendously enhance our knowledge of AGN by providing us with population statistics for the quantities estimated in Kasliwal et al. (2016).

Application of C-ARMA processes to AGN light curves requires *well-sampled* light curves. Kasliwal et al. (2016) demonstrate how two C-ARMA processes—a simple DRW and the 2nd-order damped harmonic oscillator (DHO) can have similar looking power spectral density (PSD) on the long timescales typically probed by ground-based surveys such as the SDSS Stripe 82 (Annis et al. 2014). On short timescales, the DRW contains excess power as compared to the DHO; a well-sampled light curve drawn from a DHO will look *smoother* than a light curve drawn from a DRW on short timescales even if they look indistinguishable on longer timescales. While this argument makes it clear that it is necessary to sample both short- and long-timescales with LSST, it does not provide us with a sampling strategy. *How should we allocate (in timesteps) the precious exposures available in each LSST filter band in order to unambiguously capture the nature of AGN variability?* It seems that uniform sampling is not desirable given a fixed exposure budget because it limits the shortest cadence interval severely and gives us no high frequency information. Non-uniform sampling promises to deliver information about variability on a multitude of time scales however it is not clear what the optimal pattern is. We outline a method for evaluating the optimality of each LSST cadence pattern produced by an OpSim for studying AGN accretion disc variability science.

2. AGN Lightcurve Simulation and MAF Metric

The project described here will evaluate how accurately proposed LSST OpSims will perform for AGN optical variability science goals. We propose to implement the cadence analysis discussed below into the Metric Analysis Framework (MAF) in order to automate the merit of each OpSim cadence pattern for AGN accretion disc science.

Light curves will be simulated for AGN across numerous fields in the projected LSST sky footprint including deep drilling fields (DDFs) using the Python language KĀLĪ software library of Kasliwal et al. (2016). Densely-sampled mock light curves, drawn from a range of C-ARMA models with different model orders and characteristic timescales, will be generated to simulate the underlying true population of AGN. Our prior for generating the dense mocks will be drawn from the results of the *K2* studies of AGN variability currently in progress. The frequency of visits for all bands in randomly selected fields will be used to generate realistic light curves for each LSST band by down-sampling from the dense mocks at various RA and Dec within the LSST footprint. The complete test ensemble of AGN targets will be simulated at different S/N levels to probe the effectiveness of binning rapidly-obtained observations of faint objects in order to boost the S/N.

The K $\bar{\text{A}}\text{L}\bar{\text{I}}$ software will then be used to infer what C-ARMA model fits each simulated light curve best.

To concretely evaluate the optimality of a given OpSim for accretion disc variability science, we propose the following metrics:

1. The fraction $f_{Comp.}$ of light curves for which we recover the original ‘complexity’ of the light curve variability. The complexity of a light curve is quantified by the number of features in the PSD of the light curve. Insufficient sampling can shift power from higher (poorly-sampled) frequencies to lower frequencies via aliasing. Aliasing can introduce false-periodicities into the PSD (Vaughan et al. 2016). In the case of C-ARMA processes, complexity is statistically parametrized by the order (p,q) of the C-ARMA process that best fits the mock light curve. Regardless of the utility of C-ARMA processes for modeling AGN variability, it is desirable that we accurately capture the behavior of the PSD of the light curve. We will look for any consistent bias that may overestimate or underestimate the complexity for the entire ensemble of mock light curves.
2. The fraction $f_{Amp.}$ of light curves for which we successfully recover the amplitude of variability to within 10 % of the simulated value and the median fractional discrepancy $f_{\delta Amp./Amp.}$ of the variability amplitude for the entire population of simulated AGN. The variability amplitude is correlated with the mass and anti-correlated with the intrinsic luminosity and Eddington ratio of the accreting supermassive black hole (MacLeod et al. 2010), making it very desirable to estimate the variability amplitude accurately.
3. The fraction f_T of light curves for which we recover the dominant variability time scale to within 10 % of the simulated value and the median fractional discrepancy $f_{\delta T/T}$ of the dominant timescale for the entire population of simulated AGN. MacLeod et al. (2010) have found a correlation between the dominant timescale and black hole mass. Kelly et al. (2009) have shown that inferences may be made about the nature of the physical processes operating in AGN accretion disks based on timescales

REFERENCES

- M.-H. Ulrich, L. Maraschi, and C. M. Urry. Variability of Active Galactic Nuclei. *ARA&A*, 35: 445–502, 1997. doi: 10.1146/annurev.astro.35.1.445.
- M. Peterson, Bradley. *An Introduction to Active Galactic Nuclei*. Cambridge University Press, 1997. ISBN 0521479118.
- B. M. Peterson, A. J. Barth, P. Berlind, R. Bertram, K. Bischoff, N. G. Bochkarev, A. N. Burenkov, F.-Z. Cheng, M. Dietrich, A. V. Filippenko, E. Giannuzzo, L. C. Ho, J. P. Huchra, J. Hunley, S. Kaspi, W. Kollatschny, D. C. Leonard, Y. F. Malkov, T. Matheson, M. Mignoli, B. Nelson, P. Papaderos, J. Peters, R. W. Pogge, V. I. Pronik, S. G. Sergeev, E. A. Sergeeva, A. I. Shapovalova, G. M. Stirpe, S. Tokarz, R. M. Wagner, I. Wanders, J.-Y. Wei, B. J. Wilkes,

- H. Wu, S.-J. Xue, and Z.-L. Zou. Steps toward Determination of the Size and Structure of the Broad-Line Region in Active Galactic Nuclei. XV. Long-Term Optical Monitoring of NGC 5548. *ApJ*, 510:659–668, January 1999. doi: 10.1086/306604.
- Falanga Maurizio, Tomaso Belloni, Piergiorgio Casella, M. Gilfanov, Peter Jonker, and Andrew King, editors. *The Physics of Accretion onto Black Holes*, volume 49 of *Space Sciences Series of ISSI*. Springer, 2015. ISBN 1493922262.
- B. C. Kelly, A. C. Becker, M. Sobolewska, A. Siemiginowska, and P. Uttley. Flexible and Scalable Methods for Quantifying Stochastic Variability in the Era of Massive Time-domain Astronomical Data Sets. *ApJ*, 788:33, June 2014. doi: 10.1088/0004-637X/788/1/33.
- P.J. Brockwell. Recent results in the theory and applications of carma processes. *Ann. Inst. Stat. Math.*, 66(4):647–685, 2014. ISSN 0020-3157. doi: 10.1007/s10463-014-0468-7. URL <http://dx.doi.org/10.1007/s10463-014-0468-7>.
- B. C. Kelly, J. Bechtold, and A. Siemiginowska. Are the Variations in Quasar Optical Flux Driven by Thermal Fluctuations? *ApJ*, 698:895, June 2009. doi: 10.1088/0004-637X/698/1/895.
- V. P. Kasliwal, M. S. Vogeley, and G. T. Richards. Extracting Information from AGN Variability. Submitted to MNRAS, 2016.
- J. Annis, M. Soares-Santos, M. A. Strauss, A. C. Becker, S. Dodelson, X. Fan, J. E. Gunn, J. Hao, Ž. Ivezić, S. Jester, L. Jiang, D. E. Johnston, J. M. Kubo, H. Lampeitl, H. Lin, R. H. Lupton, G. Miknaitis, H.-J. Seo, M. Simet, and B. Yanny. The Sloan Digital Sky Survey Coadd: 275 deg² of Deep Sloan Digital Sky Survey Imaging on Stripe 82. *ApJ*, 794:120, October 2014. doi: 10.1088/0004-637X/794/2/120.
- S. Vaughan, P. Uttley, A. G. Markowitz, D. Huppenkothen, M. J. Middleton, W. N. Alston, J. D. Scargle, and W. M. Farr. False periodicities in quasar time-domain surveys. *ArXiv e-prints*, June 2016.
- C. L. MacLeod, Ž. Ivezić, C. S. Kochanek, S. Kozłowski, B. Kelly, E. Bullock, A. Kimball, B. Sesar, D. Westman, K. Brooks, R. Gibson, A. C. Becker, and W. H. de Vries. Modeling the Time Variability of SDSS Stripe 82 Quasars as a Damped Random Walk. *ApJ*, 721:1014–1033, oct 2010. doi: 10.1088/0004-637X/721/2/1014.