THE MATHEMATICS OF DEEP LEARNING*

VISHAL P. KASLIWAL¹

Wave Computing
W. Campbell Ave, # 301
Campbell, CA 95008, USA

(Received December 28, 2017; Revised December 28, 2017; Accepted December 28, 2017)

ABSTRACT

Deep Learning is a branch of Machine Learning in which 'deep' neural networks are used for various purposes.

Keywords: neural networks, training

Corresponding author: Vishal Kasliwal vishal@wavecomp.com, vishal.kasliwal@gmail.com

^{*} This is a draft prepared for the Agent Library group at Wave Computing.

1. PRELIMINARIES

2. NOTATION

We shall use a to denote vectors and A to denote tensors.

Tensors are associated with layers. The $p \times q$ tensor \mathbf{A} when associated with layer l is denoted $\mathbf{A}_{p \times q}^{[l]}$. The transpose of this tensor is denoted by $(\mathbf{A}_{p \times q}^{[l]})^{\top}$. If the tensor is an input tensor to a layer, it may additionally have a minibatch number, m, and instance within the minibatch, i, associated with it. Thus $(\mathbf{A}_{p \times q}^{[l]\{m\}\{i\}})^{\top}$ is the transpose of the i-th training example from the m-th minibatch in the l-th layer from the input tensor \mathbf{A} which is p rows high by q columns long.

The $n^{[l]}$ operator returns the number of nodes, i.e. the width, of the l-th layer. Traditionally, the depth of the neural network is denoted by L, and hence we must have $0 \leq l \leq L$. The 0-th layer is a pure input layer i.e. it has no transformation associated with it.

3. ACTIVATION FUNCTIONS

3.1. Sigmoid Activation Function

The sigmoid activation function takes the form

$$g(z) = \sigma(z) = \frac{1}{1 + e^{-z}}.$$
 (1)

The derivative of the sigmoid activation function is

$$\frac{\mathrm{d}\sigma}{\mathrm{d}z} = \sigma(1 - \sigma). \tag{2}$$

3.1.1. Derivation

By definition

$$\sigma(z) = \frac{1}{1 + e^{-z}} = (1 + e^{-z})^{-1},$$

and so

$$\frac{d\sigma(z)}{dz} = -(1 + e^{-z})^{-2} \frac{d(1 + e^{-z})}{dz}.$$

But this is just

$$-(1 + e^{-z})^{-2}(-e^{-z}) = \sigma^2 e^{-z}.$$

Now $e^{-z} = \frac{1}{\sigma} - 1 = \frac{1-\sigma}{\sigma}$ and so

$$\frac{d\sigma(z)}{dz} = \sigma^2 e^{-z} = \sigma^2 \frac{1-\sigma}{\sigma} = \sigma(1-\sigma),$$

Q.E.D.

- 4. OUTPUT LAYERS
- 5. FEED-FORWARD LAYERS
- 6. CONVOLUTIONAL LAYERS
 - 7. POOLING LAYERS

APPENDIX

REFERENCES