

Introduction

Hello everyone.

Today I'd like to talk about Axion electrodynamics.

You might be surprised that there is a picture of a laundry detergent here.

Don't worry, that will become clear later.

Overview

More precisely, I will explain what an Axion is and where it comes from.

To understand this, we need to talk about the strong CP problem, as well as the Peccei-Quinn mechanism which, in turn, postulates the Axion field.

Then I will introduce the topic of Axion electrodynamics in general, by looking at the work of Pierre Sikivie in this regard.

For the last part of the presentation, we will look at experiments that he proposed and which are currently under development or already returning data.

What is an Axion

According to Wikipedia:

The axion is a hypothetical elementary particle postulated by the Peccei–Quinn theory to resolve the strong CP problem in quantum chromodynamics (QCD).

I will describe the strong CP problem and the Peccei-Quinn theory in future slides.

Why are Axions interesting?

If axions exist and have low mass within a specific range, they are of interest as a possible component of cold dark matter.

CP symmetry

CP-symmetry states that the laws of physics should be the same if a particle were interchanged with its antiparticle (C symmetry, as charges of antiparticles are the negative of the corresponding particle), and then left and right were swapped (P symmetry).

Strong CP problem

This is the Lagrangian density of quantum chromodynamics.

Quantum chromodynamics is the quantum field theory of the strong interaction concerned with quarks, gluons and their color charge.

As shown by Gerard 't Hooft, pictured on the right, the Lagrangian density includes a potentially CP symmetry violating term due to its non-trivial vacuum structure.

Combined with effects generated by weak interactions, this effective periodic term, Θ (theta), appears as a Standard Model input, which means that its value is not predicted by the theory and must thus be measured.

Strong CP problem

Neutron electric dipole moment

However, large CP-violating interactions originating from QCD would induce a large electric dipole moment (EDM) for the neutron.

Experimental constraints on the currently unobserved EDM implies CP violation from QCD must be extremely tiny and thus Θ must itself be extremely small.

The current limit on the nEDM constrains this angle to be less than 10^{-10} radians.

Since Θ could have any value between 0 and 2π , this presents a "naturalness" problem for the standard model.

Why should this parameter find itself so close to zero? Or why should QCD find itself CP-preserving?

This question constitutes what is known as the strong CP problem.

Strong CP problem

Massless quarks

One simple solution exists: If at least one of the quarks of the standard model is massless, CP symmetry violation becomes unobservable.

However, empirical evidence strongly suggests that none of the quarks are massless.

Consequently, particle theorists sought other resolutions to the problem of inexplicably conserved CP.

Peccei–Quinn mechanism

Postulated by Helen Quinn and Roberto Peccei the therefore called Peccei–Quinn theory predicts that the small value of the θ parameter is explained by a dynamic field, rather than a constant value.

The potential which this field carries causes it to have a value that naturally cancels, making the θ parameter uneventfully zero.

On the left you can see them showing their theory with a sombrero modeling the graph of Goldstone's "Mexican Hat" potential, which is an example of a spontaneously broken symmetry.

Axion field

Peccei–Quinn symmetry presents θ as a functional component. This symmetry is spontaneously broken by the vacuum expectation value.

According to the Goldstone theorem a discrete, spontaneously broken symmetry corresponds to a new particle.

The Boson for the Peccei–Quinn symmetry was independently proposed by Frank Wilczek and Steven Weinberg, which are pictured on the right here.

Frank Wilczek named it after the laundry detergent called Axion, because it cleaned up a profound physical problem.

Possible axion models

The original Wilczek/Weinberg axion has since been ruled out but more recent axion models still hold up.

The properties of the axion depend mainly on the magnitude v of the vacuum expectation value that spontaneously breaks the Peccei–Quinn symmetry.

The axion mass and its couplings to ordinary particles are all inversely proportional to v .

As far as the solution to the strong CP problem is concerned, the value of v is arbitrary.

Values for v with less than ten to the power of eight Giga electron volt have since been ruled out by experiments and theoretical considerations. These axions are so weakly coupled that they have been called "invisible".

Axion

If the "invisible axion" exists it has the following properties:

No electric charge and no spin.

A mass between one hundred-thousandth and one thousandth electron volt.

It interacts only gravitationally and electromagnetically but the electromagnetic interaction is very weak.

Due to its mass and weak electromagnetic interaction it is a good candidate for dark matter.

Axion electrodynamics

Pierre Sikivie published a modification of Maxwell's equations that arise from a light, stable axion in 1983, making the invisible axions visible.

In his paper he showed that the axions should be observable by multiple experimental setups.

All of which rely on the fact that the axion could transform into a photon (and viceversa) in the presence of electromagnetic fields. This property of the axion is crucial for most of the experimental strategies of axion detection.

Axion electrodynamics

Modified Maxwell's equation

As you can see the Maxwell's equations look very similar to the unmodified ones.

The expressions for the E- and B- fields are replaced by an expression including the axion field Θ coupled with a coupling constant κ .

Axion electrodynamics

Rotating E and B into one another

Incorporating the axion has the effect of rotating the electric and magnetic fields into each other, where the mixing angle χ depends on the coupling constant κ and the axion field strength Θ .

In addition we also get a new differential equation - the axion law - which is simply the Klein–Gordon equation with an E and B source term.

The Klein-Gordon equation is the quantum field theory equation for massive spin-zero particles, as the axion is massless and spin zero as well.

Axion electrodynamics

With the modified equations Sikivie showed that the photons can be converted into axions and vice-versa in the presence of a magnetic field.

This is the basis for all the proposed experiments in his paper.

Here is how that process looks like in a Feynman diagram.

A is the axion and γ the converted photon. The doubled lines at the bottom together with the second wavy line indicate the magnetic field pushing the conversion.

Cosmological considerations

Axions may have played a critical cosmological role with regard to the problem of galaxy formation.

First, the primordial density perturbations from which galaxies evolved may have been produced by the presence of axionic domain walls for a limited time period in the early universe. Domain walls are topologically protected sheet-like surfaces that form when the potential of a field has a discrete symmetry that is spontaneously broken as is the case in the Peccei-Quinn mechanism.

Secondly, axions may be the stuff the dark halos of galaxies are made of. Because of their very large primordial phase-space density, axions cluster easily and if they have been produced abundant enough, they could even provide all of the halo matter.

They might be produced by the Primakoff effect. The effect, named after the physicists Henry Primakoff, describes the production of two pseudoscalar mesons, which the Axion classifies as, by the interaction of high energy photons with an atomic nucleus.

Lastly, due to the aforementioned effect they should also be continually produced inside the sun.

Experiments

Sikivie proposed essentially two types of possible experiments:

If the axion exist and it is the main component of dark matter, the very relic axions that would be bombarding us continuously could be detected using microwave cavities, immersed in powerful magnetic fields, that are resonant to the axion mass.

Another promising detection technique, this one independent of the axion being the dark matter, is that of the axion helioscope, aiming to detect axions produced at the solar interior. These could be detected, once again, using a powerful magnet, but this time equipped with low background x-ray detectors.

Experiments

The axion haloscope

The axion haloscope would Sikivie proposed consists of a

Using previously calculated estimates for the axion flux on earth based on the cosmological considerations described previously, Sikivie

Current experiments

The axion haloscope

Summary

The axion is an hypothetical particle that appears in extensions of the Standard Model of Particle Physics that include the so-called Peccei-Quinn mechanism. This mechanism was postulated already 35 years ago to explain an standing problem of the Standard Model: the strong-CP problem.

The Peccei-Quinn mechanism was proposed to solve this problem in a natural way, without required parameter fine-tuning. As a collateral effect, however, a new particle appears, the axion, which may have important observable consequences.

In the first place, the axion is a neutral and very light (but not massless) particle, and it does not interact (or does it very weakly) with conventional matter. In some way one can see the axion as a "strange photon".

In fact, theory predicts that the axion, if it exists, could transform into a photon (and viceversa) in the presence of electromagnetic fields. This property of the axion is crucial for most of the experimental strategies of axion detection.

One of the most suggestive properties of axions is that, in a natural way, they could be produced in huge numbers soon after the Big Bang. This population of axions would still be present today and could compose the Dark Matter of the Universe. The existence of Dark Matter is widely accepted in the scientific community, but its nature is still a mystery. Together with WIMPs, the axions are among the most searched candidates in the context of the nature of Dark Matter.