

Lagrangian Duality and Convex Optimization

David Rosenberg

New York University

February 6, 2017


Introduction

Why Convex Optimization?

- Historically:
 - **Linear programs** (linear objectives & constraints) were the focus
 - **Nonlinear programs**: some easy, some hard
- More Recently:
 - Main distinction is between **convex** and **non-convex** problems
 - Convex problems are the ones we know how to solve efficiently
- Many techniques that are well understood for convex problems are applied to non-convex problems
 - e.g. SGD is routinely applied to neural networks, which are not convex

Your Reference for Convex Optimization

- Boyd and Vandenberghe (2004)
 - Very clearly written, but has a ton of detail for a first pass.
 - See my “Extreme Abridgement of Boyd and Vandenberghe”.



Notation from Boyd and Vandenberghe

- $f : \mathbf{R}^p \rightarrow \mathbf{R}^q$ to mean that f maps from some *subset* of \mathbf{R}^p
 - namely **dom** $f \subset \mathbf{R}^p$, where **dom** f is the domain of f


Convex Sets and Functions

Convex Sets

Definition

A set C is **convex** if for any $x_1, x_2 \in C$ and any θ with $0 \leq \theta \leq 1$ we have

$$\theta x_1 + (1 - \theta)x_2 \in C.$$




KPM Fig. 7.4

Convex and Concave Functions

Definition

A function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is **convex** if $\mathbf{dom} f$ is a convex set and if for all $x, y \in \mathbf{dom} f$, and $0 \leq \theta \leq 1$, we have

$$f(\theta x + (1-\theta)y) \leq \theta f(x) + (1-\theta)f(y).$$



KPM Fig. 7.5

Examples of Convex Functions on \mathbf{R}

Examples

- $x \mapsto ax + b$ is both convex and concave on \mathbf{R} for all $a, b \in \mathbf{R}$.
- $x \mapsto |x|^p$ for $p \geq 1$ is convex on \mathbf{R}
- $x \mapsto e^{ax}$ is convex on \mathbf{R} for all $a \in \mathbf{R}$

Convex Functions and Optimization

Definition

A function f is **strictly convex** if the line segment connecting any two points on the graph of f lies **strictly** above the graph (excluding the endpoints).

Consequences for optimization:

- **convex**: if there is a local minimum, then it is a **global** minimum
- **strictly convex**: if there is a local minimum, then it is the **unique global** minimum

The General Optimization Problem

General Optimization Problem: Standard Form

General Optimization Problem: Standard Form

$$\begin{array}{ll}
 \text{minimize} & f_0(x) \\
 \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\
 & h_i(x) = 0, \quad i = 1, \dots, p,
 \end{array}$$

where $x \in \mathbf{R}^n$ are the **optimization variables** and f_0 is the **objective function**.

Assume **domain** $\mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \cap \bigcap_{i=1}^p \text{dom } h_i$ is nonempty.

General Optimization Problem: More Terminology

- The set of points satisfying the constraints is called the **feasible set**.
- A point x in the feasible set is called a **feasible point**.
- If x is feasible and $f_i(x) = 0$,
 - then we say the inequality constraint $f_i(x) \leq 0$ is **active** at x .

- The **optimal value** p^* of the problem is defined as

$$p^* = \inf \{f_0(x) \mid x \text{ satisfies all constraints}\}.$$

- x^* is an **optimal point** (or a solution to the problem) if x^* is feasible and $f(x^*) = p^*$.

Do We Need Equality Constraints?

- Note that

$$h(x) = 0 \iff (h(x) \geq 0 \text{ AND } h(x) \leq 0)$$

- So any equality-constrained problem

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & h(x) = 0 \end{array}$$

can be rewritten as

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & h(x) \leq 0 \\ & -h(x) \leq 0. \end{array}$$

- For simplicity, we'll drop equality constraints from this presentation.

Lagrangian Duality: Convexity not required

The Lagrangian

The general [inequality-constrained] optimization problem is:

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \end{array}$$

Definition

The **Lagrangian** for this optimization problem is

$$L(x, \lambda) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x).$$

- λ_i 's are called **Lagrange multipliers** (also called the **dual variables**).

The Lagrangian Encodes the Objective and Constraints

- Supremum over Lagrangian gives back objective and constraints:

$$\begin{aligned} \sup_{\lambda \succeq 0} L(x, \lambda) &= \sup_{\lambda \succeq 0} \left(f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) \right) \\ &= \begin{cases} f_0(x) & \text{when } f_i(x) \leq 0 \text{ all } i \\ \infty & \text{otherwise.} \end{cases} \end{aligned}$$

- Equivalent **primal form** of optimization problem:

$$p^* = \inf_x \sup_{\lambda \succeq 0} L(x, \lambda)$$

The Primal and the Dual

- Original optimization problem in **primal form**:

$$p^* = \inf_x \sup_{\lambda \succeq 0} L(x, \lambda)$$

- The **Lagrangian dual problem**:

$$d^* = \sup_{\lambda \succeq 0} \inf_x L(x, \lambda)$$

- We will show **weak duality**: $p^* \geq d^*$ for any optimization problem

Weak Max-Min Inequality

Theorem

For **any** $f : W \times Z \rightarrow \mathbf{R}$, we have

$$\sup_{z \in Z} \inf_{w \in W} f(w, z) \leq \inf_{w \in W} \sup_{z \in Z} f(w, z).$$

Proof.

For any $w_0 \in W$ and $z_0 \in Z$, we clearly have

$$\inf_{w \in W} f(w, z_0) \leq f(w_0, z_0) \leq \sup_{z \in Z} f(w_0, z).$$

Since this is true for all w_0 and z_0 , we must also have

$$\sup_{z_0 \in Z} \inf_{w \in W} f(w, z_0) \leq \inf_{w_0 \in W} \sup_{z \in Z} f(w_0, z).$$



Weak Duality

- For any optimization problem (**not just convex**), weak max-min inequality implies **weak duality**:

$$\begin{aligned}
 p^* &= \inf_x \sup_{\lambda \succeq 0} \left[f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) \right] \\
 &\geq \sup_{\lambda \succeq 0, v} \inf_x \left[f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) \right] = d^*
 \end{aligned}$$

- The difference $p^* - d^*$ is called the **duality gap**.
- For *convex* problems, we often have **strong duality**: $p^* = d^*$.

The Lagrange Dual Function

- The **Lagrangian dual problem**:

$$d^* = \sup_{\lambda \succeq 0} \underbrace{\inf_x L(x, \lambda)}_{\text{Lagrange dual function}}$$

Definition

The **Lagrange dual function** (or just **dual function**) is

$$g(\lambda) = \inf_x L(x, \lambda) = \inf_x \left(f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) \right).$$

- The dual function may take on the value $-\infty$ (e.g. $f_0(x) = x$).

The Lagrange Dual Problem

- In terms of Lagrange dual function, we can write weak duality as

$$p^* \geq \sup_{\lambda \geq 0} g(\lambda) = d^*$$

- So for any λ with $\lambda \geq 0$, **Lagrange dual function gives a lower bound on optimal solution:**

$$g(\lambda) \leq p^*$$

The Lagrange Dual Problem

- The **Lagrange dual problem** is a search for best lower bound:

$$\begin{array}{ll}\text{maximize} & g(\lambda) \\ \text{subject to} & \lambda \succeq 0.\end{array}$$

- λ **dual feasible** if $\lambda \succeq 0$ and $g(\lambda) > -\infty$.
- λ^* **dual optimal** or **optimal Lagrange multipliers** if they are optimal for the Lagrange dual problem.
- Lagrange dual problem often easier to solve (simpler constraints).
- d^* can be used as stopping criterion for primal optimization.
- Dual can reveal hidden structure in the solution.

Convex Optimization

Convex Optimization Problem: Standard Form

Convex Optimization Problem: Standard Form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m\end{array}$$

where f_0, \dots, f_m are convex functions.

Strong Duality for Convex Problems

- For a convex optimization problems, we **usually** have strong duality, but not always.
 - For example:

$$\begin{array}{ll}\text{minimize} & e^{-x} \\ \text{subject to} & x^2/y \leq 0 \\ & y > 0\end{array}$$

- The additional conditions needed are called **constraint qualifications**.

Slater's Constraint Qualifications for Strong Duality

- Sufficient conditions for strong duality in a **convex** problem.
- Roughly: the problem must be **strictly** feasible.
- Qualifications when problem domain $\mathcal{D} \subset \mathbf{R}^n$ is an open set:
 - (\mathcal{D} is the set where all functions are defined, NOT the feasible set.)
 - **Strict feasibility is sufficient.** ($\exists x \ f_i(x) < 0$ for $i = 1, \dots, m$)
 - For any affine inequality constraints, $f_i(x) \leq 0$ is sufficient.
- Otherwise, see notes or BV Section 5.2.3, p. 226.

Complementary Slackness

Complementary Slackness

- Consider a general optimization problem (i.e. not necessarily convex).
- If we have **strong duality**, we get an interesting relationship between
 - the optimal Lagrange multiplier λ_i and
 - the i th constraint at the optimum: $f_i(x^*)$
- Relationship is called “**complementary slackness**”:

$$\lambda_i^* f_i(x^*) = 0$$

- Lagrange multiplier is zero unless constraint is active at optimum.

Complementary Slackness Proof

- Assume strong duality: $p^* = d^*$ in a general optimization problem
- Let x^* be primal optimal and λ^* be dual optimal. Then:

$$\begin{aligned}
 f_0(x^*) &= g(\lambda^*) \\
 &= \inf_x \left(f_0(x) + \sum_{i=1}^m \lambda_i^* f_i(x) \right) \\
 &\leq f_0(x^*) + \sum_{i=1}^m \underbrace{\lambda_i^* f_i(x^*)}_{\leq 0} \\
 &\leq f_0(x^*).
 \end{aligned}$$

Each term in sum $\sum_{i=1}^m \lambda_i^* f_i(x^*)$ must actually be 0. That is

$$\boxed{\lambda_i^* f_i(x^*) = 0, \quad i = 1, \dots, m.}$$

This condition is known as **complementary slackness**.