


Instructions for use

Emgality 120 mg solution for injection in pre-filled syringe

Galcanezumab

For subcutaneous use



Before using your pre-filled syringe:


Important Information

- Your doctor or nurse should show you how to prepare and inject Emgality using the syringe. Do not inject yourself or someone else until you have been shown how to inject Emgality.
- Keep these instructions and refer to them as needed.
- Each syringe is for **ONE-TIME USE ONLY**. Do not share or reuse your syringe. Otherwise, you may give or get an infection.
- Your doctor, pharmacist or nurse can help you decide where on your body to inject your dose. You can also read the “**Choose your injection site**” section of these instructions to help you choose which area can work best for you.
- If you have vision problems, **do not** use Emgality syringe without help from a caregiver.

INSTRUCTIONS FOR USE

Before you use the EMGALITY syringe, read and carefully follow all the step-by-step instructions.

Parts of the Emgality Syringe



Before You Get Started

Take the syringe from the refrigerator

Put the original package with any unused syringes back in the refrigerator.

Leave the needle cap on until you are ready to inject.

Do not shake.

For a more comfortable injection, leave the syringe at room temperature for 30 minutes before injecting.

Do not microwave the syringe, run hot water over it, or leave it in direct sunlight.


Inspect the syringe and the medicine

Make sure you have the right medicine. The medicine inside should be clear. It may be colourless to slightly yellow.

Do not use the syringe, and dispose of as directed by your doctor, pharmacist or nurse if:

- it looks damaged
- the medicine is cloudy, is discolored, or has small particles
- the expiration date printed on the label has passed
- the medicine is frozen


Expiration Date



Prepare for injection

Wash your hands with soap and water before you inject your Emgality. Make sure a sharps disposal container is close by.

Choose your injection site




Your doctor, pharmacist or nurse can help you choose the injection site that is best for you.

- **You** may inject the medicine into your stomach area (abdomen) or thigh. Do not inject within 5 centimeters of the belly button (navel).
- **Another person** may give you the injection in the back of your upper arm or buttock.
- **Do not** inject in the same spot as before. For example, if your first injection was in your abdomen, your next injection could be in another area of your abdomen.
- **Clean and dry the injection site before you inject.**


1 Uncap

- Leave the needle cap on until you are ready to inject.
- When you are ready to inject, pull the needle cap off and throw it away in the bin.
- **Do not** put the needle cap back on – you could damage the needle or injure yourself by accident.
- **Do not** touch the needle.




2 Insert

- Gently pinch and hold a fold of skin where you will inject.
- Insert the needle at a 45 degree angle.




3 Inject

- Slowly push on the thumb pad to push the plunger all the way in until all the medicine is injected.
- The grey syringe plunger should be pushed all the way to end of the syringe.



- You should see the teal plunger rod show through the syringe body when the injection is complete as shown.
- Remove the needle from your skin and gently let go of your skin.
- **Do not** put the needle cap back on the syringe.



Teal plunger rod

Grey syringe plunger

After You Inject Your Medicine

Dispose of the syringe

DO NOT put the needle cap back on.
Dispose of the syringe in a sharps disposal container or as directed by your doctor, pharmacist or nurse.



When you dispose of the syringe and the sharps disposal container:

- Do not recycle the filled sharps container.
- Ask your doctor, pharmacist or nurse about how to dispose of medicines you no longer use.

Commonly Asked Questions

Q. What if I see air bubbles in my Emgality syringe?

A. It is normal to have air bubbles in the syringe. Emgality is injected under your skin (subcutaneous injection).

Q. What if there is a drop of liquid on the tip of the needle when I remove the needle cap?

A. It is okay to see a drop of liquid on the tip of the needle.

Q. What if I cannot push in the plunger?

A. If the plunger is stuck or damaged:

- **Do not** continue to use the syringe
- Remove the needle from your skin
- Dispose of the syringe and get a new one

Q. What if there is a drop of liquid or blood on my skin after my injection?

A. This is normal. Press a cotton ball or gauze over the injection site. Do not rub the injection site.

Q. How can I tell if my injection is complete?

A. When your injection is complete:

- The teal plunger rod should show through the body of the syringe.
- The grey syringe plunger should be pushed all the way to end of the syringe.

Read the full Package Leaflet for Emgality inside this box to learn more about your medicine.