

## **Package leaflet: Information for the patient**

### **Emgality 120 mg solution for injection in pre-filled syringe**

galcanezumab

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Emgality is and what it is used for
2. What you need to know before you use Emgality
3. How to use Emgality
4. Possible side effects
5. How to store Emgality
6. Contents of the pack and other information

#### **1. What Emgality is and what it is used for**

Emgality contains galcanezumab, a medicine that stops the activity of a naturally occurring substance in the body called calcitonin gene-related peptide (CGRP). People with migraine may have increased levels of CGRP.

Emgality is used to prevent migraine in adult patients who have at least 4 migraines days per month.

Emgality can reduce the frequency of migraine headache and improve your quality of life. It starts working in about a week.

#### **2. What you need to know before you use Emgality**

##### **Do not use Emgality:**

- if you are allergic to galcanezumab or any of the other ingredients of this medicine (listed in section 6).

##### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before or during treatment with Emgality if:

- you have a serious cardiovascular disease. Emgality has not been studied in patients with serious cardiovascular diseases.

##### **Look out for allergic reactions**

Emgality can potentially cause serious allergic reactions. Serious allergic reactions happen mainly within 1 day after having taken Emgality, but some reactions can be delayed (happen more than 1 day to 4 weeks after having taken Emgality). Some allergic reactions can be prolonged in duration. You must look out for signs of these reactions while you are using Emgality. Stop using Emgality and tell

your doctor or seek medical help immediately if you notice any signs of a serious allergic reaction. Such signs are listed under “Serious side effects” in section 4.

### **Children and adolescents**

This medicine should not be given to children and adolescents under 18 years of age because it has not been studied in this age group.

### **Other medicines and Emgality**

Tell your doctor, pharmacist or nurse if you are using, have recently used or might use any other medicines.

### **Pregnancy and breast-feeding**

If you are a woman able to have children, you are advised to avoid becoming pregnant while using Emgality.

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. It is preferable to avoid the use of Emgality in pregnancy as the effects of this medicine in pregnant women are not known.

If you are breast-feeding or are planning to breast-feed, talk to your doctor before using this medicine. You and your doctor should decide if you should breast feed and use Emgality .

### **Driving and using machines**

Galcanezumab could have a minor effect on your ability to drive and use machines. Some patients have had vertigo whilst using Emgality.

### **Emgality contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per 120 mg dose, that is to say essentially “sodium-free”.

## **3. How to use Emgality**

Always use this medicine exactly as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Emgality pre-filled syringe is for single use only and contains one dose of Emgality (120 mg).

- The first time you receive Emgality your doctor or nurse will inject two syringes (total 240 mg).
- After the first dose, you will use one syringe (120 mg) every month.

Your doctor will decide for how long you should use Emgality.

Emgality is given by injection under your skin (subcutaneous injection). You and your doctor or nurse should decide if you can inject Emgality yourself.

It is important not to try to inject yourself until you have been trained by your doctor or nurse. A caregiver may also give you your Emgality injection after proper training.

The syringe must not be shaken.

Read the “Instructions for Use” for the syringe carefully before using Emgality.

### **If you use more Emgality than you should**

If you have injected more Emgality than you should, e.g. if after the first dose of 240 mg, you have injected it twice in a single month, or if anyone else has accidentally used Emgality, contact your doctor immediately.

**If you forget to use Emgality**

Do not take a double dose to make up for forgotten injection

If you have forgotten to inject a dose of Emgality, inject the missed dose as soon as possible and then inject the next dose after a month from that date.

**If you stop using Emgality**

You should not stop using Emgality without speaking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Serious side effects**

Allergic reactions with Emgality are usually mild to moderate (such as rash or itching). Serious allergic reaction may occur rarely (may affect up to 1 in 1 000 people) and the signs may include:

- difficulty breathing or swallowing,
- low blood pressure, which can cause dizziness or light-headedness,
- swelling of the neck, face, mouth, lips, tongue or throat which may develop rapidly,
- severe itching of the skin, with a red rash or raised bumps.

Tell your doctor or get emergency medical help straight away if you notice any of those signs.

**Other side effects that have been reported.**

**Very common side effects** (may affect more than 1 in 10 people):

- Injection site pain
- Injection site reactions (e.g. red skin, itching, bruising, swelling)

**Common side effects** (may affect up to 1 in 10 people):

- Vertigo (a feeling of dizziness or “spinning”)
- Constipation
- Itching
- Rash

**Uncommon side effects** (may affect up to 1 in 100 people):

- Hives (raised itchy areas of skin)

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Emgality**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and on the carton after “EXP.” The expiry date refers to the last day of that month.

Store in a refrigerator (2 °C to 8 °C). Do not freeze.

Store in the original package in order to protect from light.

Emgality may be stored unrefrigerated for a single period up to 7 days when stored at temperatures up to 30 °C. If the syringe is stored at a higher temperature or for a longer period it must be discarded.

Do not use this medicine if you notice that the syringe is damaged, or the medicine is cloudy or has particles in it.

Do not throw away any medicines via wastewater or household waste. Ask your doctor, nurse or pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Emgality contains**

The active substance is galcanezumab. . Each pre-filled syringe contains 120 mg of galcanezumab in 1 mL solution.

The other ingredients are: L-histidine, L-histidine hydrochloride monohydrate, polysorbate 80, sodium chloride and water for injections.

### **What Emgality looks like and contents of the pack**

Emgality is a solution for injection in a clear glass single-dose syringe. Its colour may vary from colourless to slightly yellow. Pack sizes of 1, 2 or 3 pre-filled syringes.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Eli Lilly Nederland B. V., Papendorpseweg 83, 3528 BJ Utrecht, The Netherlands.

### **Manufacturer:**

Eli Lilly Italia S.p.A., Via Gramsci 731/733, 50019, Sesto Fiorentino (FI), Italy.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **Belgique/België/Belgien**

Eli Lilly Benelux S.A./N.V.  
Tél/Tel: + 32-(0)2 548 84 84

#### **България**

ТП "Ели Лили Недерланд" Б.В. - България  
тел. + 359 2 491 41 40

#### **Česká republika**

ELI LILLY ČR, s.r.o.  
Tel: + 420 234 664 111

#### **Danmark**

Eli Lilly Danmark A/S  
Tlf: +45 45 26 60 00

#### **Lietuva**

Eli Lilly Lietuva  
Tel. +370 (5) 2649600

#### **Luxembourg/Luxemburg**

Eli Lilly Benelux S.A./N.V.  
Tél/Tel: + 32-(0)2 548 84 84

#### **Magyarország**

Lilly Hungária Kft.  
Tel: + 36 1 328 5100

#### **Malta**

Charles de Giorgio Ltd.  
Tel: + 356 25600 500

**Deutschland**

Lilly Deutschland GmbH  
Tel: +49-(0) 6172 273 2222

**Eesti**

Eli Lilly Nederland B. V.  
Tel: +372 6 817 280

**Ελλάδα**

ΦΑΡΜΑΣΕΡΒ-ΛΙΑΛΛΥ Α.Ε.Β.Ε.  
Τηλ: +30 210 629 4600

**España**

Organon Salud, S.L.  
Tel: +34 91 591 12 79

**France**

Organon France  
Tél: +33-(0) 1 57 77 32 00

**Hrvatska**

Eli Lilly Hrvatska d.o.o.  
Tel: +385 1 2350 999

**Ireland**

Eli Lilly and Company (Ireland) Limited  
Tel: + 353-(0) 1 661 4377

**Ísland**

Icepharma hf.  
Sími + 354 540 8000

**Italia**

Eli Lilly Italia S.p.A.  
Tel: + 39- 055 42571

**Κύπρος**

Phadisco Ltd  
Τηλ: +357 22 715000

**Latvija**

Eli Lilly (Suisse) S.A Pārstāvniecība Latvijā  
Tel: +371 67364000

**Nederland**

Eli Lilly Nederland B.V.  
Tel: + 31-(0) 30 60 25 800

**Norge**

Eli Lilly Norge A.S.  
Tlf: + 47 22 88 18 00

**Österreich**

Eli Lilly Ges.m.b.H.  
Tel: + 43-(0) 1 711 780

**Polska**

Eli Lilly Polska Sp. z o.o.  
Tel: +48 22 440 33 00

**Portugal**

Lilly Portugal Produtos Farmacêuticos, Lda  
Tel: + 351-21-4126600

**România**

Eli Lilly România S.R.L.  
Tel: + 40 21 4023000

**Slovenija**

Eli Lilly farmacevtska družba, d.o.o.  
Tel: +386 (0)1 580 00 10

**Slovenská republika**

Eli Lilly Slovakia s.r.o.  
Tel: + 421 220 663 111

**Suomi/Finland**

Oy Eli Lilly Finland Ab  
Puh/Tel: + 358-(0) 9 85 45 250

**Sverige**

Eli Lilly Sweden AB  
Tel: + 46-(0) 8 7378800

**United Kingdom (Northern Ireland)**

Eli Lilly and Company (Ireland) Limited  
Tel: + 353-(0) 1 661 4377

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>