Daniel Kim

Theory of the Evolution without Oxygen

The study of **evolution without oxygen**, also known as anaerobic evolution, focuses on the fascinating adaptations and evolutionary processes that have shaped life in environments where oxygen is absent or in limited supply. Anaerobic organisms have developed unique metabolic pathways and strategies to thrive in such conditions, providing insights into the early stages of life on Earth before the oxygen-rich atmosphere we know today. Exploring the genetic innovations and survival mechanisms of anaerobes sheds light on the diversity and resilience of life forms in a variety of anoxic settings, from deep-sea hydrothermal vents to the digestive tracts of animals. From this the question arises, “How do organisms that evolved in environments without oxygen adapt to survive and thrive in anaerobic conditions? What are the key genetic and metabolic adaptations involved in this process?”

The existence of oxygen is essential to life as we know it and is a basic component of aerobic respiration. However, the development and persistence of life in anaerobic conditions have captured the attention of scientists, resulting in important new understandings and discoveries. (Figure 1)



Figure 1: Evolutionary timeline illustrating the emergence and diversification of anaerobic life forms alongside the oxygenation of Earth's atmosphere.

Simple bacteria to more sophisticated eukaryotes are examples of anaerobic creatures. They have developed special metabolic and biochemical routes to get around the lack of oxygen. From simple bacteria to more complex eukaryotes, a wide range of anaerobic organisms have successfully adapted to the difficulties presented by oxygen-deficient habitats. These varied organisms have developed complex metabolic and biochemical processes, demonstrating an amazing capacity to avoid oxygen deprivation and continue to exist in other ways. Simple anaerobic bacteria have evolved streamlined metabolic pathways that rely on anaerobic respiration or fermentation to obtain energy from available substrates. These bacteria are frequently the first to inhabit anaerobic habitats. The adaptations get more complicated and sophisticated as complexity rises to include eukaryotic species, such as some protists and fungi.

These anaerobic eukaryotes have honed their cellular machinery and developed novel ways to produce energy, highlighting the adaptability of life and its ability to use a variety of metabolic pathways under oxygen-limited circumstances. (Figure 2)



Figure 2: Metabolic pathways employed by anaerobic organisms, showcasing the diversity of strategies for energy production in the absence of oxygen.

The study of oxygen-free evolution has provided important new understandings of the many tactics and adaptations used by organisms that survive under anaerobic conditions. Anaerobic microbes from antiquity have revealed a pre-oxygen Earth populated by microorganisms, offering important insights into the origins of life. Methanogenic archaea are vital to the global carbon cycle because they produce methane as a metabolic byproduct. illustrating how anaerobic life affects Earth's ecosystems. (Figure 3)



Figure 3: Interactions between anaerobic methane oxidizers performing N-dAOM with anaerobic ammonia oxidation (anammox) incorporated into partial-nitritation systems (PN/A)

With unexpected findings of anaerobic eukaryotes upend preconceived ideas by demonstrating the capacity of some protists and fungi to thrive in oxygen-free conditions. The metabolic flexibility of living organisms has also been brought to light by the study of anaerobic evolution, which has demonstrated a variety of energy-generating processes such fermentation and chemolithotrophy.

In conclusion, The study of oxygen-free evolution has revealed an amazing picture of life that goes against conventional wisdom. From the earliest anaerobic microorganisms to the sophisticated eukaryotic adaptations, organisms have shown an amazing capacity for adaptability and survival in oxygen-deficient conditions. These findings show the diversity and adaptability inherent in the evolutionary process, challenging our beliefs regarding aerobic respiration's universality as a fundamental aspect of life.

Bibliography

1) Awai K (2014) Oxygenic photosynthesis without galactolipids | PNAS. *Oxygenic photosynthesis without galactolipids*. Available at: https://www.pnas.org/doi/full/10.1073/pnas.1403708111 [Accessed October 27, 2023].
2) Blake RE (2001) *Oxygen isotope ratios of PO4: An inorganic indicator of . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.051515898 [Accessed November 30, 2023].
3) Bristow LA (2016) *Ammonium and nitrite oxidation at nanomolar oxygen . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1600359113 [Accessed November 30, 2023].
4) Canfield DE (2021) *Petrographic carbon in ancient sediments constrains proterozoic . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.2101544118 [Accessed November 30, 2023].
5) Castello PR (2008) *Oxygen-regulated isoforms of cytochrome - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.0709461105 [Accessed November 30, 2023].
6) Fenchel T (1994) The evolution of life without oxygen - JSTOR. *The Evolution of Life without Oxygen*. Available at: https://www.jstor.org/stable/29775097 [Accessed October 27, 2023].
7) He H (2023) *A mineral-based origin of earth???s initial hydrogen peroxide . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.2221984120 [Accessed November 30, 2023].
8) Huang P (2023) *Dynamic orbital hybridization triggered spin-disorder . - PNASPeilin Huang*. Available at: https://www.pnas.org/doi/abs/10.1073/pnas.2219661120 [Accessed November 30, 2023].
9) Lenton TM (2016) *Earliest land plants created modern levels of atmospheric oxygen | PNAS*. Available at: https://www.pnas.org/doi/full/10.1073/pnas.1604787113 [Accessed November 30, 2023].
10) Maisch T (2007) *The role of singlet oxygen and oxygen concentration in . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.0611328104 [Accessed November 30, 2023].
11) Markhard AL (2022) *A genetically encoded system for oxygen generation in living cells | PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.2207955119 [Accessed November 30, 2023].
12) Meng Y (2019) *Highly active oxygen evolution integrated with efficient co - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1915319116 [Accessed November 30, 2023].
13) Ratcliff WC (2012) *Experimental evolution of multicellularity | PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1115323109 [Accessed November 30, 2023].
14) Seekell RP (2016) *Oxygen delivery using engineered microparticles | PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1608438113 [Accessed November 30, 2023].
15) Strasser B (2010) *Arabidopsis thaliana life without phytochromes | PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.0910446107 [Accessed November 30, 2023].
16) Trifunovic A (2005) *Somatic mtdna mutations cause aging phenotypes without affecting . - PNAS*. Available at: https://www.pnas.org/doi/full/10.1073/pnas.0508886102 [Accessed November 30, 2023].
17) Wulff P (2014) *How oxygen reacts with oxygen-tolerant respiratory [NIFE . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1322393111 [Accessed November 30, 2023].
18) Zhang S (2016) *Sufficient oxygen for animal respiration 1,400 million years ago - PNAS*. Available at: https://www.pnas.org/doi/abs/10.1073/pnas.1523449113 [Accessed October 27, 2023].

19) Zhang X (2019) Highly active oxygen evolution integrated with efficient CO2 to Co ... *Highly active oxygen evolution integrated with efficient CO2 to CO electroreduction*. Available at: https://www.pnas.org/doi/10.1073/pnas.1915319116 [Accessed October 27, 2023].
20) Zhang X-S (2018) *Reactive oxygen species extend insect life span using components . - PNAS*. Available at: https://www.pnas.org/doi/10.1073/pnas.1711042114 [Accessed November 30, 2023].